

planetmath.org

Math for the people, by the people.

terminating reduction

Canonical name TerminatingReduction
Date of creation 2013-03-22 17:57:41
Last modified on 2013-03-22 17:57:41

Owner CWoo (3771)

Owner CWoo (3771) Last modified by CWoo (3771)

Numerical id 11

Author CWoo (3771) Entry type Definition Classification msc 68Q42

Related topic Normalizing Reduction

Related topic Confluence
Related topic DiamondLemma
Defines terminating

Defines descending chain condition

Defines DCC

Defines convergent reduction

Defines acyclic

Let X be a set and \rightarrow a reduction (binary relation) on X. A chain with respect to \rightarrow is a sequence of elements x_1, x_2, x_3, \ldots in X such that $x_1 \rightarrow x_2, x_2 \rightarrow x_3$, etc... A chain with respect to \rightarrow is usually written

$$x_1 \to x_2 \to x_3 \to \cdots \to x_n \to \cdots$$
.

The length of a chain is the cardinality of its underlying sequence. A chain is finite if its length is finite. Otherwise, it is infinite.

Definition. A reduction \rightarrow on a set X is said to be *terminating* if it has no infinite chains. In other words, every chain *terminates*.

Examples.

- If \rightarrow is reflexive, or non-trivial symmetric, then it is never terminating.
- Let X be the set of all positive integers greater than 1. Define \to on X so that $a \to b$ means that a = bc for some $c \in X$. Then \to is a terminating reduction. By the way, \to is also a normalizing reduction.
- In fact, it is easy to see that a terminating reduction is normalizing: if a has no normal form, then we may form an infinite chain starting from a.
- On the other hand, not all normalizing reduction is terminating. A canonical example is the set of all non-negative integers with the reduction \rightarrow defined by $a \rightarrow b$ if and only if
 - either $a, b \neq 0$, $a \neq b$, and a < b,
 - or $a \neq 0$ and b = 0.

The infinite chain is given by $1 \to 2 \to 3 \to \cdots$, so that \to is not terminating. However, $n \to 0$ for every positive integer n. Thus every integer has 0 as its normal form, so that \to is normalizing.

Remarks.

- A reduction is said to be *convergent* if it is both terminating and confluent.
- A relation is terminating iff the transitive closure of its inverse is well-founded.

To see this, first let R be terminating on the set X. And let S be the transitive closure of R^{-1} . Suppose A is a non-empty subset of X that contains no S-minimal elements. Pick $x_0 \in A$. Then we can find $x_1 \in A$ with $x_1 \neq x_0$, such that x_1Sx_0 . By the assumption on A, this process can be iterated indefinitely. So we have a sequence x_0, x_1, x_2, \ldots such that $x_{i+1}Sx_i$ with $x_i \neq x_{i+1}$. Since each pair (x_i, x_{i+1}) can be expanded into a finite chain with respect to R, we have produced an infinite chain containing elements x_0, x_1, x_2, \ldots , contradicting the assumption that R is terminating.

On the other hand, suppose the transitive closure S of R^{-1} is well-founded. If the chain $x_0Rx_1Rx_2R\cdots$ is infinite, then the set $\{x_0, x_1, x_2, \ldots\}$ has no S-minimal elements, as x_iSx_j whenever i > j, and j arbitrary.

• The reflexive transitive closure of a terminating relation is a partial order.

A closely related concept is the descending chain condition (DCC). A reduction \to on X is said to satisfy the descending chain condition (DCC) if the only infinite chains on X are those that are eventually constant. A chain $x_1 \to x_2 \to x_3 \to \cdots$ is eventually constant if there is a positive integer N such that for all $n \geq N$, $x_n = x_N$. Every terminating relation satisfies DCC. The converse is obviously not true, as a reflexive reduction illustrates.

Another related concept is acyclicity. Let \to be a reduction on X. A chain $x_0 \to x_1 \to \cdots x_n$ is said to be cyclic if $x_i = x_j$ for some $0 \le i < j \le n$. This means that there is a "closed loop" in the chain. The reduction \to is said to be *acyclic* if there are no cyclic chains with respect to \to . Every terminating relation is acyclic, but not conversely. The usual strict inequality relation on the set of positive integers is an example of an acyclic but non-terminating relation.