

Characterization of the sequential nature of neuronal dynamics: Experimental recordings, computational models and novel stimulation neurotechnologies

Alicia Garrido Peña

Advisor: Pablo Varona Martínez

**Escuela Politécnica Superior.
Universidad Autónoma de Madrid**

Alicia Garrido Peña. PhD thesis Seminar
Tuesday 23rd July, 2024



Universidad Autónoma
de Madrid




Escuela
Politécnica Superior

Contents


- 1 Introduction
- 2 Motivation and Objectives
- 3 Sequential constraints in CPG circuits: Dynamical invariants
- 4 CW-NIR laser stimulation as an effective neuromodulation technique
- 5 Conclusions

Introduction


Neuroscience




Neurobiology




Clinical Neuroscience



Neurotechnology



Cognitive Neuroscience



Computational Neuroscience

All images here are licensed under the Creative Commons Attribution 1, 2.5 or 4.0 Generic licenses

Approach


- Neurocomputational Perspective

Approach

- Neurocomputational Perspective
- Bottom-up approach

Approach


- Neurocomputational Perspective
- Bottom-up approach




From ionic channels

Approach

- Neurocomputational Perspective
- Bottom-up approach



From ionic channels



To minimal circuits

Approach

- Neurocomputational Perspective
- Bottom-up approach
- Combining electrophysiology

Approach

- Neurocomputational Perspective
- Bottom-up approach
- Combining electrophysiology, computational work

Approach

- Neurocomputational Perspective
- Bottom-up approach
- Combining electrophysiology, computational work and novel neurotechnology.

Neuronal and Networks Dynamics

Neuronal electrical activity is often described in terms of the evolution of membrane voltage caused by the flow of ionic channels between the inside and outside of the cell.


Neuronal and Networks Dynamics

Depending on the channels that constitute the neuron and the circuit it is immersed in, the activity varies.

Neuronal and Networks Dynamics

Depending on the channels that constitute the neuron and the circuit it is immersed in, the activity varies.


In terms of the spike waveform:




Neuronal and Networks Dynamics

Depending on the channels that constitute the neuron and the circuit it is immersed in, the activity varies.

In terms of the spike waveform:



And the type of spiking activity: tonic firing, bursting, etc.




The sequential nature of neural dynamics

- There are sequential processes at different time-scales.


The sequential nature of neural dynamics

- There are sequential processes at different time-scales.




The sequential nature of neural dynamics

- There are sequential processes at different time-scales.
- Many behaviors and actions are governed by sequential processes




The sequential nature of neural dynamics

- There are sequential processes at different time-scales.
- Many behaviors and actions are governed by sequential processes
- In humans: Motor control, speech, decision making, etc.



The sequential nature of neural dynamics

- There are sequential processes at different time-scales.
- Many behaviors and actions are governed by sequential processes
- In humans: Motor control, speech, decision making, etc.
- Study of sequential activity might have a crucial role in neural coordination to autonomously establish a balance between the robustness and flexibility required for effective motor function.



Studying neural dynamics in computational models

- Computational models are powerful tools to study neural dynamics

Studying neural dynamics in computational models

- Computational models are powerful tools to study neural dynamics
- Advantages

Studying neural dynamics in computational models

- Computational models are powerful tools to study neural dynamics
- Advantages
 - Full accessibility to the system

Studying neural dynamics in computational models

- Computational models are powerful tools to study neural dynamics
- Advantages
 - Full accessibility to the system
 - Detailed reproduction of the living activity

Studying neural dynamics in computational models

- Computational models are powerful tools to study neural dynamics
- Advantages
 - Full accessibility to the system
 - Detailed reproduction of the living activity
- Limitations

Studying neural dynamics in computational models

- Computational models are powerful tools to study neural dynamics
- Advantages
 - Full accessibility to the system
 - Detailed reproduction of the living activity
- Limitations
 - Restricted variability

Studying neural dynamics in computational models

- Computational models are powerful tools to study neural dynamics
- Advantages
 - Full accessibility to the system
 - Detailed reproduction of the living activity
- Limitations
 - Restricted variability
 - The description may depend on the information about the living system

Studying neural dynamics in computational models

- Computational models are powerful tools to study neural dynamics
- Advantages
 - Full accessibility to the system
 - Detailed reproduction of the living activity
- Limitations
 - Restricted variability
 - The description may depend on the information about the living system

Studying neural dynamics in computational models

Studying neural dynamics in computational models

Conductance-based models

Studying neural dynamics in computational models

Conductance-based models

Voltage equation	$C \frac{dV}{dt} = I - g_K n^4(V - E_K) - g_{Na} m^3 h(V - E_{Na}) - g_L(V - E_L)$		
	Activation variables	Inactivation variable	
gating variables	$\frac{dm(t)}{dt} = \frac{m_\infty(V(t)) - m(t)}{\tau_m(V(t))}$	$\frac{dn(t)}{dt} = \frac{n_\infty(V(t)) - n(t)}{\tau_n(V(t))}$	$\frac{dh(t)}{dt} = \frac{h_\infty(V(t)) - h(t)}{\tau_h(V(t))}$

Studying neural dynamics in computational models

Conductance-based models


By different combinations of ionic channels we can achieve different activities and waveform shapes

Studying neural dynamics in computational models


Conductance-based models

By different combinations of ionic channels we can achieve different activities and waveform shapes

a)



b)



Studying neural dynamics in computational models


Conductance-based models

By **modeling synapses** we can model whole **circuits** by connecting single modeled neurons.

Studying neural dynamics in computational models

Conductance-based models

By **modeling synapses** we can model whole **circuits** by connecting single modeled neurons.



Vertebrate and invertebrate animal studies

- In addition to the well known vertebrate models, there have been invaluable findings using invertebrates.

Vertebrate and invertebrate animal studies

- In addition to the well known vertebrate models, there have been invaluable findings using invertebrates.
- There are many universal characteristics of nervous systems and behaviors that can be extrapolated to humans.

Vertebrate and invertebrate animal studies

- In addition to the well known vertebrate models, there have been invaluable findings using invertebrates.
- There are many universal characteristics of nervous systems and behaviors that can be extrapolated to humans.
- Using invertebrates as animal models have different advantages:

Vertebrate and invertebrate animal studies

- In addition to the well known vertebrate models, there have been invaluable findings using invertebrates.
- There are many universal characteristics of nervous systems and behaviors that can be extrapolated to humans.
- Using invertebrates as animal models have different advantages:
 - Ease of accessibility to the nervous system.

Vertebrate and invertebrate animal studies

- In addition to the well known vertebrate models, there have been invaluable findings using invertebrates.
- There are many universal characteristics of nervous systems and behaviors that can be extrapolated to humans.
- Using invertebrates as animal models have different advantages:
 - Ease of accessibility to the nervous system.
 - Longer survival times of the preparations.

Vertebrate and invertebrate animal studies

- In addition to the well known vertebrate models, there have been invaluable findings using invertebrates.
- There are many universal characteristics of nervous systems and behaviors that can be extrapolated to humans.
- Using invertebrates as animal models have different advantages:
 - Ease of accessibility to the nervous system.
 - Longer survival times of the preparations.
 - Ease of breeding and reproduction.

Vertebrate and invertebrate animal studies

- In addition to the well known vertebrate models, there have been invaluable findings using invertebrates.
- There are many universal characteristics of nervous systems and behaviors that can be extrapolated to humans.
- Using invertebrates as animal models have different advantages:
 - Ease of accessibility to the nervous system.
 - Longer survival times of the preparations.
 - Ease of breeding and reproduction.
 - Full description of their systems.

Vertebrate and invertebrate animal studies



- In this thesis we work with the neural system of *Lymanea stagnalis*.

Vertebrate and invertebrate animal studies



- In this thesis we work with the neural system of *Lymnea stagnalis*.
- A pond snail whose system is well studied and described.

Neural stimulation

When studying neural systems, we need neuromodulatory technics.

- To study different conditions.

Neural stimulation

When studying neural systems, we need neuromodulatory techniques.

- To study different conditions.
- Test their robustness.

Neural stimulation

When studying neural systems, we need neuromodulatory technics.

- To study different conditions.
- Test their robustness.
- Simulate external inputs.

Neural stimulation

We can classify them based on:

Neural stimulation

We can classify them based on:

- The change in the system: Invasive / non-invasive

Neural stimulation

We can classify them based on:

- The change in the system: Invasive / non-invasive
- The source of the stimulation:

Neural stimulation

We can classify them based on:

- The change in the system: Invasive / non-invasive
- The source of the stimulation:
 - Chemical

Neural stimulation

We can classify them based on:

- The change in the system: Invasive / non-invasive
- The source of the stimulation:
 - Chemical
 - Electrical

Neural stimulation

We can classify them based on:

- The change in the system: Invasive / non-invasive
- The source of the stimulation:
 - Chemical
 - Electrical
 - Magnetic

Neural stimulation

We can classify them based on:

- The change in the system: Invasive / non-invasive
- The source of the stimulation:
 - Chemical
 - Electrical
 - Magnetic
 - Mechanical

Neural stimulation

We can classify them based on:

- The change in the system: Invasive / non-invasive
- The source of the stimulation:
 - Chemical
 - Electrical
 - Magnetic
 - Mechanical
 - Optical

Neural stimulation

We can classify them based on:

- The change in the system: Invasive / non-invasive
- The source of the stimulation:
 - Chemical
 - Electrical
 - Magnetic
 - Mechanical
 - Optical

Neural stimulation

We can classify them based on:

- The change in the system: Invasive / **non-invasive**
- The source of the stimulation:
 - Chemical
 - Electrical
 - Magnetic
 - Mechanical
 - Optical

Motivation and Objectives

Motivation and Objectives

1. To explore the **sequential nature of neuronal dynamics** at distinct description levels.

Motivation and Objectives

1. To explore the **sequential nature of neuronal dynamics** at distinct description levels.
2. To study **sequence interval variability** constraints and relationships in neural models and living circuits.

Motivation and Objectives

1. To explore the **sequential nature of neuronal dynamics** at distinct description levels.
2. To study **sequence interval variability** constraints and relationships in neural models and living circuits.
3. To analyze the feeding CPG of *Lymnaea stagnalis* to provide evidence of the **universality of sequential dynamical invariants** found in the pyloric CPG by:

Motivation and Objectives

1. To explore the **sequential nature of neuronal dynamics** at distinct description levels.
2. To study **sequence interval variability** constraints and relationships in neural models and living circuits.
3. To analyze the feeding CPG of *Lymnaea stagnalis* to provide evidence of the **universality of sequential dynamical invariants** found in the pyloric CPG by:
 - 3.1 Characterizing the sequential invariants cycle-by-cycle in a bursting CPG model.

Motivation and Objectives

1. To explore the **sequential nature of neuronal dynamics** at distinct description levels.
2. To study **sequence interval variability** constraints and relationships in neural models and living circuits.
3. To analyze the feeding CPG of *Lymnaea stagnalis* to provide evidence of the **universality of sequential dynamical invariants** found in the pyloric CPG by:
 - 3.1 Characterizing the sequential invariants cycle-by-cycle in a bursting CPG model.
 - 3.2 Analyzing intracellular recordings with spontaneous activity and with different rhythm initiation stimulation protocols.

Motivation and Objectives

1. To explore the **sequential nature of neuronal dynamics** at distinct description levels.
2. To study **sequence interval variability** constraints and relationships in neural models and living circuits.
3. To analyze the feeding CPG of *Lymnaea stagnalis* to provide evidence of the **universality of sequential dynamical invariants** found in the pyloric CPG by:
 - 3.1 Characterizing the sequential invariants cycle-by-cycle in a bursting CPG model.
 - 3.2 Analyzing intracellular recordings with spontaneous activity and with different rhythm initiation stimulation protocols.
4. To illustrate the possible **functionality** of sequential dynamical invariants in biohybrid robotics.

Motivation and Objectives

5. To test the capability of **CW-NIR laser stimulation to modulate** neuronal dynamics by:

Motivation and Objectives

5. To test the capability of **CW-NIR laser stimulation to modulate** neuronal dynamics by:
 - 5.1 Characterizing the CW-NIR effect in the spike waveform dynamics.

Motivation and Objectives

5. To test the capability of **CW-NIR laser stimulation to modulate** neuronal dynamics by:
 - 5.1 Characterizing the CW-NIR effect in the spike waveform dynamics.
 - 5.2 Analyzing the ability of this neurotechnology to change the spiking rate and the circuit dynamics.

Motivation and Objectives

5. To test the capability of **CW-NIR laser stimulation to modulate** neuronal dynamics by:
 - 5.1 Characterizing the CW-NIR effect in the spike waveform dynamics.
 - 5.2 Analyzing the ability of this neurotechnology to change the spiking rate and the circuit dynamics.
6. To study the possible **biophysical candidates** underlying the CW-NIR effect in model simulations.



Motivation and Objectives

5. To test the capability of **CW-NIR laser stimulation to modulate** neuronal dynamics by:
 - 5.1 Characterizing the CW-NIR effect in the spike waveform dynamics.
 - 5.2 Analyzing the ability of this neurotechnology to change the spiking rate and the circuit dynamics.
6. To study the possible **biophysical candidates** underlying the CW-NIR effect in model simulations.
7. To design and implement a **new technique** for CW-NIR stimulation in **closed-loop**.

Sequential constraints in CPG circuits: Dynamical invariants


Central Pattern Generators

- Neural circuits generating robust sequences of neural activity




Central Pattern Generators

- Neural circuits generating robust sequences of neural activity
- Control motor rhythms in an autonomous manner




Central Pattern Generators

- Neural circuits generating robust sequences of neural activity
- Control motor rhythms in an autonomous manner
- Present in vertebrates and invertebrates





Central Pattern Generators

- Neural circuits generating robust sequences of neural activity
- Control motor rhythms in an autonomous manner
- Present in vertebrates and invertebrates
- Flexible enough to adapt the rhythm to the variability keeping a robust sequential activity




Central Pattern Generators

- Non-open topologies
- Based on mutual inhibition
- Temporal sequences maintained cycle-by-cycle



Temporal constraints: Sequential Dynamical invariants


- Recently found in pyloric CPG (experimental study). (Elices et al.)
- Specific intervals that build the sequence are highly correlated cycle-by-cycle.
- Consistent under high variability induced situations (Ethanol).



Feeding CPG of *Lymnaea stagnalis*

Triphasic rhythm:

- N1 phase → Protraction
- N2 phase → Rasp
- N3 phase → Swallow



Computational and Experimental approach



Buccal Ganglia of *Lymnaea stagnalis* Microscope Image





Lymnaea stagnalis

$$I = C_M \frac{dV}{dt} + \bar{g}_K n^4 (V - V_K) + \bar{g}_{Na} m^3 h (V - V_{Na}) + \bar{g}_I (V - V_t)$$




$$\begin{aligned} dn/dt &= \alpha_n(1-n) - \beta_n n \\ dm/dt &= \alpha_m(1-m) - \beta_m m \\ dh/dt &= \alpha_h(1-h) - \beta_h h \end{aligned}$$



Characterization of the time-intervals cycle-by-cycle



Characterization of the time-intervals cycle-by-cycle

To analyze the dynamics of the sequential activity cycle-by-cycle we need to define time references to characterize the time-intervals in the sequence.




Characterization of the time-intervals cycle-by-cycle

To analyze the dynamics of the sequential activity cycle-by-cycle we need to define time references to characterize the time-intervals in the sequence.




Characterization of the time-intervals cycle-by-cycle

We can also define intervals combining the time references of two bursts




Characterization of the time-intervals cycle-by-cycle

We can also define intervals combining the time references of two bursts




Characterization of the time-intervals cycle-by-cycle

We can also define intervals combining the time references of two bursts





Characterization of the time-intervals cycle-by-cycle

We can also define intervals combining the time references of two bursts




Computational Approach: Model description


- (Vavoulis et al.)
- Conductance-based model.
- Specific waveforms and gradual synapse.
- Two compartments.
- Feeding CPG Model.




Computational Approach: Ramp stimulation protocol




Cycle-by-cycle restrictions



N1M stimulation



SO stimulation




Models with chaotic activity

Variability in models is usually limited. The variability can be induced by a current ramp as in this case, but with a chaotic activity the variability could be more realistic.

Models with chaotic activity

Variability in models is usually limited. The variability can be induced by a current ramp as in this case, but with a chaotic activity the variability could be more realistic.


Neuronal recording variability




Models with chaotic activity

Variability in models is usually limited. The variability can be induced by a current ramp as in this case, but with a chaotic activity the variability could be more realistic.

Neuronal recording variability



Model simulation with chaotic activity



Experimental approach


- Are sequential invariants present also in the experimental recordings?

Experimental approach

- Are sequential invariants present also in the experimental recordings?
- Do the invariants change under different stimulation conditions?


Experimental approach

Previous studies point to the linear relation between some phases and the period




Elliott, C. J. H., & Andrew, T. (1991). Temporal Analysis of Snail Feeding Rhythms: A Three-Phase Relaxation Oscillator. *Journal of Experimental Biology*, 157(1), 391–408. Figure 5.

Distributed neural system of *L. stagnalis*




Benjamin, P. R. (2012). Distributed network organization underlying feeding behavior in the mollusk *Lymnaea*. *Neural Systems & Circuits*, 2(1), 1–16. <https://doi.org/10.1186/2042-1001-2-4>

Distributed neural system of *L. stagnalis*




Benjamin, P. R. (2012). Distributed network organization underlying feeding behavior in the mollusk *Lymnaea*. *Neural Systems & Circuits*, 2(1), 1–16. <https://doi.org/10.1186/2042-1001-2-4>

Distributed neural system of *L. stagnalis*




Benjamin, P. R. (2012). Distributed network organization underlying feeding behavior in the mollusk *Lymnaea*. *Neural Systems & Circuits*, 2(1), 1–16. <https://doi.org/10.1186/2042-1001-2-4>

Distributed neural system of *L. stagnalis*



Benjamin, P. R. (2012). Distributed network organization underlying feeding behavior in the mollusk *Lymnaea*. *Neural Systems & Circuits*, 2(1), 1–16. <https://doi.org/10.1186/2042-1001-2-4>


Distributed neural system of *L. stagnalis*




Benjamin, P. R. (2012). Distributed network organization underlying feeding behavior in the mollusk *Lymnaea*. *Neural Systems & Circuits*, 2(1), 1–16. <https://doi.org/10.1186/2042-1001-2-4>

Spontaneous activity

Case 1




Case 2




Spontaneous activity


Case 1




Case 2



SO driven activity




SO spontaneous driven




SO driven activity


SO driven by stimulation





SO spontaneous driven



MLN driven activity




MLN driven activity




CV1 driven activity

CV1a driven by electrical stimulation. CV1a driven by electrical stimulation.

Case 1




Case 2




CV1 driven activity


CV1a driven by electrical stimulation. Case 1





CV1a driven by electrical stimulation. Case 2



Study of functional the role of dynamical invariants in a Hybot



Study of functional the role of dynamical invariants in a Hybrot



Summary of results


Study of the Sequential nature in neuronal dynamics

Case of study

Feeding CPG *Lymnaea stagnalis*

Experimental approach

1.




Presence of dynamical invariants under different cases of stimulation

Universality of sequential dynamical invariants

The variability distribution is dependent on the CPG activity context


Computational approach

2.




3.

Transformation of sequential intervals into effective robot movement



4.

Importance of reproducing the functional variability in computational models



CW-NIR laser stimulation as an effective neuromodulation technique

Current findings

- Laser stimulation accelerates neural activity.
- Most studies use pulsed laser and general dynamics
- Open questions:
 - Source of the effect: Photothermal, Photomechanical ...
 - Biophysical candidates: Capacitance, TRPV-4

Clinical applications



Figure 2
from Saucedo, C. L. et al (2021). Brain Stimulation, 14(2), 440–449.
<https://doi.org/10.1016/j.brs.2021.02.011>






Figure 6A from Liang, S. et al. (2009). Cell Biochemistry and Biophysics, 53(1), 33–42.
<https://doi.org/10.1007/s12013-008-9035-2>

Experimental setup




Sustained stimulation protocol

- The neuron was illuminated during the intracellular recording.
- The illumination lasted 1-3 min.
- We recorded a control before and after the illumination.




Spike Waveform Characterization



To characterize the change in the spike dynamics we defined waveform metrics.




Modulation of the spike waveform




Effect on the firing rate



Effect on a minimal circuit




The stimulation of electrically coupled neurons, showed a first approximation to the capacity of the laser illumination to modulate neural dynamics.




Model analysis to explore biophysical candidates

$$C_m \frac{dV}{dt} = I_{inj} - I_{NaT} - I_{NaP} - I_A - I_D - I_{LVA} - I_{HVA} \quad (1)$$




Model analysis to explore biophysical candidates

Each candidate generated a change in the waveform, but each one fitted the experimental change only partly.



Model analysis to explore biophysical candidates


Each candidate generated a change in the waveform, but each one fitted the experimental change only partly.



The computational model study shows that no modulation of an individual candidate but a combination of them can explain the experimental quantification


Model with temperature description

Including temperature dependency in the model by a Q_{10} factor showed the closest representation of the observed experimental modulation.




Model with temperature description

Including temperature dependency in the model by a Q_{10} factor showed the closest representation of the observed experimental modulation.




Temperature change plays a key role to explain the laser stimulation effect.


Activity-dependent stimulation protocol




Activity-dependent stimulation protocol




Activity-dependent stimulation protocol



Activity-dependent stimulation protocol




Activity-dependent stimulation effect



Summary of results

Study of the Sequential nature in neuronal dynamics



Conclusions

Conclusions

- The study of neural sequences at multiple spatial and temporal scales can provide **novel insights to relate neural dynamics and behavior.**

Conclusions

- The study of neural sequences at multiple spatial and temporal scales can provide **novel insights to relate neural dynamics and behavior.**
- We have **hinted the universality of sequential dynamical invariants** in experimental and modeling data in the feeding CPG.

Conclusions

- The study of neural sequences at multiple spatial and temporal scales can provide **novel insights to relate neural dynamics and behavior**.
- We have **hinted the universality of sequential dynamical invariants** in experimental and modeling data in the feeding CPG.
- Dynamical invariants can be **indicators of the functional distribution** of variability in the CPG.

Conclusions

- The study of neural sequences at multiple spatial and temporal scales can provide **novel insights to relate neural dynamics and behavior**.
- We have **hinted the universality of sequential dynamical invariants** in experimental and modeling data in the feeding CPG.
- Dynamical invariants can be **indicators of the functional distribution** of variability in the CPG.
- **Novel neurotechnologies** can contribute to identifying and exploiting the sequential nature of neural information processing.

Conclusions

- Sustained CW-NIR laser illumination **asymmetrically accelerates action potential dynamics and the spiking rate** on single neurons.

Conclusions

- Sustained CW-NIR laser illumination **asymmetrically accelerates action potential dynamics and the spiking rate** on single neurons.
- A model study of the effect of CW-NIR showed that **no biophysical candidate alone could fully reproduce the observed modulation** and the global modulation through **temperature change** in the simulation was **the closest approximation** to it.

Conclusions

- Sustained CW-NIR laser illumination **asymmetrically accelerates action potential dynamics and the spiking rate** on single neurons.
- A model study of the effect of CW-NIR showed that **no biophysical candidate alone could fully reproduce the observed modulation** and the global modulation through **temperature change** in the simulation was **the closest approximation** to it.
- The **closed-loop protocol** unveiled the CW-laser **effect at different phases of the neuron dynamics**.

Conclusions

- Sustained CW-NIR laser illumination **asymmetrically accelerates action potential dynamics and the spiking rate** on single neurons.
- A model study of the effect of CW-NIR showed that **no biophysical candidate alone could fully reproduce the observed modulation** and the global modulation through **temperature change** in the simulation was **the closest approximation** to it.
- The **closed-loop protocol** unveiled the CW-laser **effect at different phases of the neuron dynamics**.
- The results discussed in this thesis in the context of neural sequences can also have **strong implications** in the field of neurorhabilitation, robotics, and artificial intelligence.



Universidad Autónoma
de Madrid



Escuela
Politécnica Superior