

Sviluppo di un'applicazione Web per la visualizzazione e l'analisi di dati dell'esperimento AEgIS

Author:

Andrea G.B. DAMIOLI

Supervisor:

Dr. Germano BONOMI

***Co-supervisor:
Bianchini***



Master Thesis

Abstract

Master degree in Computer Engineering

Sviluppo di un'applicazione Web per la visualizzazione e l'analisi di dati dell'esperimento AEgIS

by Andrea G.B. DAMIOLI

The AEgIS Experiment at the CERN aims to verify the weak interaction principle for antimatter. This ~~document talks~~ ^{thesis presents} about "gAnWeb", a web application designed to simplify the analysis of physical data under the AEgIS experiment. ~~This analysis can be performed using Root Data Analysis Framework by the Linux Terminal,~~ ^{called "gAnWeb"} but a graphical interface can ensure a better user experience, ease the user training and improve the productivity. A web application is a smart way to implement the interface because ~~allows~~ ^{it} users to avoid installations, and centralizes all the eventual modifications. This document explains the choices made during the development of this application ~~related to the goal of the user friendly data analysis,~~ and shows the design process that led to the final product.

run via command lines in a unix terminal,

Contents

Abstract	iii
1 Introduction	1
1.1 The AEgIS experiment	1
2 gAn, gAn Web, and their goal	7
2.1 User friendly Data analysis: gAn Web	7
2.2 What is data analysis?	11
2.3 Root - Physical Data Analysis	14
3 General organization of the project	19
3.1 The roles of the designer and of the users	19
3.2 General organization of the design process	20
3.3 The stages of the design project	21
3.4 The role of aesthetics	22
3.5 Analysis with inspection methods and Tests with users	22
3.6 Prototyping and mock-up	23
3.7 Ambiguities and doubts	24
3.8 How the various stages are presented	24
4 First version	25
4.1 Functional requirements	25
4.2 Non-functional requirements	27
4.3 Scenario based functional analysis	28
4.4 Prototyping	28
4.5 How the early version can be improved	31
5 Intermediate version	33
5.1 Functional requirements	33
5.1.1 Ambiguities (and related solutions)	36
5.2 Scenario based functional analysis	36
5.3 Prototyping	38
5.3.1 Modified pages	39
5.3.2 Added pages	47
6 Final version	49

6.1	Functional requirements	49
6.2	Non-functional requirements	51
6.3	Scenario based functional analysis	52
6.4	Prototyping - Implementation	54
7	Users and Tests with them	71
7.1	User's Profile: why?	71
7.2	User's Profile: classification	71
7.2.1	Generic Characteristics	71
7.2.2	Education Background	72
7.2.3	Models in the user's mind	73
7.2.4	User's psychological characteristics	74
7.2.5	User's level of expertise	74
7.2.6	Environment	75
7.3	Relation developer-user and progressive sematization .	75
7.4	How the tests take place	75
7.4.1	Preliminary observations	77
7.4.2	What the test is evaluating?	77
7.4.3	Execution and Results	78
7.4.4	Is the testing finished here?	81
8	Patterns	83
8.1	Used patterns	83
8.1.1	Clear entry point	83
8.1.2	Wizard	84
8.1.3	Spatial memory	84
8.1.4	Grid of equals	85
8.1.5	Responsive enabling	85
8.1.6	Progress indicator	86
8.1.7	Go Back to a Safe Place	86
8.1.8	Liquid layout	86
8.1.9	Hover tools	87
8.1.10	Harmless default	87
8.1.11	Same-page error messages	88
8.1.12	Alternative views	88
9	Web Design Development	89
9.1	Requirements	89
9.2	Why a web solution?	92
9.3	MVC	93
9.4	Used Technologies	95
9.5	General Schema of the project	97
10	Conclusions	99

10.1 Conclusions	99
----------------------------	----

List of Figures

1.1	AEgIS's Logo	1
1.2	Aegis Collaboration institutes	2
1.3	AEgIS's Scheme, taken from "AEgIS experiment at CERN: measuring antihydrogen free-fall in Earth's gravitational field to test WEP with antimatter" TODO INSERT-bibliographical-reference	3
1.4	Moiré Deflectometer's Scheme, taken from "A http://www.nature.com/articles/ TODO INSERT-bibliographical-reference	4
1.5	AEgIS apparatus set up, taken from "AEgIS experiment at CERN: measuring antihydrogen free-fall in Earth's gravitational field to test W.E.P. with antimatter" TODO INSERT-bibliographical-reference	5
2.1	gAn - gAn Web simple scheme	10
2.2	Root format input files generation, DAQ System	11
2.3	Here is visible a basic schema of data processes and analysis	12
2.4	ROOT structure	17
3.1	An overview of the timeline of the project	22
4.1	First version: the first homepage of gAn Web	29
4.2	First version: the page related to the textual output of gAn Web	30
4.3	First version: the page able to show the output images in the early prototype	30
4.4	First version: the configuration page	31
5.1	Intermediate version: the homepage of gAn Web without input	39
5.2	Intermediate version: the homepage of gAn Web ready to start	39
5.3	Intermediate version: the "Start" button is red if the values in the input field are invalid	40
5.4	Intermediate version: the progress bar aims to make more comfortable the user's waiting	40

5.5	Intermediate version: the modal opened by clicking the button "Add runs by range"	41
5.6	Intermediate version: the edit-configurator page of gAn Web	42
5.7	Intermediate version: the page who shows the textual output of gAn	43
5.8	Intermediate version: by this "navbar" the user can choose the run results to show	43
5.9	Intermediate version: this page shows all the images that gAn produces in output	44
5.10	Intermediate version: moving the cursor the system shows the value of this histogram in the selected point	45
5.11	Intermediate version: the user can modify numerous settings in the generated image	45
5.12	Intermediate version: the user can show the histogram not only in traditional format, but also in a numerical format where the numbers are the value of the function in their position	46
5.13	Intermediate version: another solution is to generate a 3d image in lego-style of the histogram	46
5.14	Intermediate version: simple login page	47
5.15	Intermediate version: page where the user can choose the Root version to use	48
5.16	Intermediate version: page where the user can choose the Branch of gAn to use	48
6.1	Final version: the Login Page of gAn Web	54
6.2	Final version: the first thing that a logged user sees	55
6.3	Final version: how the user can change his choice	55
6.4	Final version: the Homepage if the user chooses Single run analysis	56
6.5	Final version: a label with some information about the current status is always visible in the homepage	57
6.6	Final version: how error messages work	57
6.7	Final version: grouped dropdown menu	58
6.8	Final version: the Homepage if the user chooses multiple runs analysis (first implementation)	59
6.9	Final version: modal using which the users insert different ranges of runs (first implementation)	60
6.10	Final version: here is if the user selects "By Range" (second implementation)	60
6.11	Final version: here is if the user selects "By InputArea" (second implementation)	60

6.12	Final version: how the homepage appears (second implementation)	61
6.13	Final version: the page related to edit configurations . .	61
6.14	Final version: an example of exponential scale (the first) followed by a non-exponential scale (the second)	62
6.15	Final version: a group of dropdown menus	63
6.16	Final version: the Navbar	64
6.17	Final version: the output in textual version	64
6.18	Final version: formatted vs unformatted output	65
6.19	Final version: the output images	65
6.20	Final version: how the user can choose which images to show	66
6.21	Final version: Images that are meant to be compared . .	67
6.22	Final version: images with info	67
6.23	Final version: other image with special info in a clear position	68
6.24	Final version: some options	69
6.25	Final version: other option	69
6.26	Final version: label to inform the user about a feature .	70
6.27	Final version: not frequently used controls	70
9.1	The general structure of the MVC pattern	94

Chapter 1

Introduction

First of all it is important to understand at least generically what is the AEgIS experiment at the CERN and ~~what~~ ^{which} are its goals. The acronym AEgIS stands for "Antimatter Experiment: gravity, Interferometry, Spectroscopy", this research aims to ~~measure~~ ^{verify} the weak equivalence principle for antimatter. In the first part of this chapter some particulars are explained about this experiment, in the second part is introduced gAn Web, the main topic of this document, the application that allows the physicists to do data analysis in the AEgIS experiment environment easily, through a web interface.

1.1 The AEgIS experiment

A E \bar{g} I S

FIGURE 1.1: AEgIS's Logo

The weak equivalence principle, also known as universality of free fall, states that in the same field all bodies fall with the same acceleration, regardless of the mass and the composition. This principle has been thoroughly tested for the matter, but not for the antimatter: the most important goal of AEgIS experiment is to measure the weak equivalence principle for the ~~anti-matter~~ ^{antimatter}. to test the universality of free fall, AEgIS ~~measures~~ ^{will attempt to measure the} gravitational interaction between matter (the ~~earth~~ ^{Earth}) and anti-matter (anti-hydrogen). Anti-hydrogen is the antimatter counterpart of hydrogen, it is the simplest atom

earth -> Earth

built with antimatter (see below for a description of its components). The first anti-hydrogen in an electromagnetic trap has been produced by the ATHENA experiment in 2012 (<http://www.nature.com/nature/journal/v419/n6906/full/419439a.html>) with the contribution of a group of the University of Brescia.

The AEGIS experiment is the result of a wide and international scientific collaboration, as visible in the figure 1.2.



FIGURE 1.2: Aegis Collaboration institutes

In the context of neutral antimatter, the gravitational interaction is of high interest, because it can potentially reveal new forces that violate the weak equivalence principle. Thomas Phillips, from Duke University, says: "If antimatter fell down faster, it would mean the discovery of at least one new force, probably two. If it fell up, it would mean our understanding of general relativity is incorrect". In a practical point of view AEGIS tries to measure the time of flight and the vertical displacement of anti-hydrogen, ~~by~~ ^{with} a moiré deflectometer: this process is quite complex, and it is easier to explain it ~~by~~ ^{using} the following two figures (1.3 and 1.4).




FIGURE 1.3: AEgIS's Scheme, taken from "AEgIS experiment at CERN: measuring antihydrogen free-fall in Earth's gravitational field to test WEP with antimatter" **TODO INSERT-bibliographical-reference**

1.3
In this first figure we can see the process that allows to create some anti-hydrogen. To correctly explain this process it is better to start with some definitions:

1. Positron: it is the correspondent of the electron in the anti-matter. It is an anti-electron, **that is** so an electron with positive electrical charge. It is indicated by " e^+ ".
2. Positronium: it is an unstable system consisting of an electron and a positron, bound together into an exotic atom. It is indicated with Ps .
3. **Antiproton**: it is the antiparticle of the proton. Antiprotons are stable, but they are typically short-lived since any collision with a proton will cause both particles to be annihilated **in a burst of energy**. It is indicated with \bar{p} (pronounced pbar).
4. **Antihydrogen**: it is the antimatter counterpart of hydrogen. Whereas the common hydrogen atom is composed of an electron and a proton, the antihydrogen atom is made up of a positron and an antiproton. It is indicated with \bar{H} (pronounced hbar).

5. Antiproton trap: a device that uses an axial magnetic field to **transversely** confine charged particles, in this case antiprotons.

Antiprotons are provided in this way: protons collide with nuclei inside a metal cylinder called "target". About four proton-antiproton

Figura del sistema di acceleratori del CERN con in evidenza AD, e bisogna dire che AEgIS è dentro AD e che riceve antiprotoni da AD

(*) In few words, to create antihydrogen, two ingredients are needed: antiprotons and positrons. The antiprotons are provided by CERN, while positrons are produced by the experiment with a radioactive source.

**sempre anti-hydrogen
oppure sempre
antihydrogen**

radially

and an electric field to confine them axially

by the CERN accelerator system

pairs are produced in every million collisions, and it is possible to separate antiprotons using magnetic fields. The following step is to guide antiprotons toward the AD (Antiproton Decelerator) where they are slowed down. To execute AEGIS experiment, antiproton must be trapped and held inside an antiproton trap, where magnetic fields force the charged antiparticles to spiral around the magnetic field lines, and electric fields confine them along the magnetic axis.

For what concerns positrons, as represented in Fig. 1.3

The process shown in the figure 1.3 is the following: a beam of positrons (that comes from a ^{22}Na radioactive source) is accelerated and driven to collide against a "positron-positronium converter" (that is a mesoporous silica film). This process creates positronium, that needs to be excited by lasers, to reach an excited state called Rydberg State. The positronium in Rydberg state is indicated by Ps^* , it has a longer life than the unexcited positronium, and can be driven to fly into an antiproton trap.

When Ps^* is excited by lasers it can combine itself with \bar{p} to generate excited antihydrogen (\bar{H}^*) and electrons. The antihydrogen beam is accelerated using an electric field towards a moiré deflectometer. Then, during the travel it decays to ground state.




FIGURE 1.4: Moiré Deflectometer's Scheme, taken from "A <http://www.nature.com/articles/ncomms5538>"
TODO INSERT-bibliographical-reference

In the second figure (1.4) a schematic view of a moiré deflectometer is presented. An antihydrogen beam is thrown toward two subsequent gratings that restrict the transmitted particles to well-defined trajectories. The trajectories are deflected by a force (in this case the force related to $m * g$) and follow a parabolic path. At the final part of the deflectometer there is a detector that shows where the antimatter annihilates, so it is possible to compare the expected trajectories without forces with the obtained trajectories, and measure the force. The

proof of principle that such system can be used with antimatter has been realized and operated with \bar{p} s. It showed that a displacement of few μm can be detected with this technique **This result has been published in a Nature [reference]**




FIGURE 1.5: AEgIS apparatus set up, taken from "AEgIS experiment at CERN: measuring antihydrogen free-fall in Earth's gravitational field to test W.E.P. with antimatter" **TODO INSERT-bibliographical-reference**