

“A p-value of < 0.05 was considered statistically significant”: a content analysis of published statistical methods sections

Nicole M White *

Australian Centre for Health Services Innovation and Centre for Healthcare Transformation
School of Public Health and Social Work, Queensland University of Technology

and

Thiru Balasubramaniam, Richi Nayak

Centre for Data Science and School of Computer Science

Queensland University of Technology

and

Adrian G Barnett

Australian Centre for Health Services Innovation and Centre for Healthcare Transformation
School of Public Health and Social Work, Queensland University of Technology

June 28, 2021

Abstract

The text of your abstract. 200 or fewer words.

Keywords: 3 to 6 keywords, that do not appear in the title

*The authors gratefully acknowledge computational resources and services used in this work provided by the eResearch Office, Queensland University of Technology, Brisbane, Australia.

1 Introduction

An ideal statistical analysis will use appropriate methods to draw insights from the data and inform the research questions. Unfortunately many current statistical analyses are far from ideal, with researchers often using the wrong methods, misinterpreting the results, or failing to adequately check their assumptions (Leek et al. 2017). Some researchers take a “mechanistic” approach to statistics, copying the few methods they know regardless of their appropriateness, and then going through the motions of the analysis (Stark & Saltelli 2018). Acceptance of methodological illiteracy in favour of research quantity and perceived novelty is at odds with principles of scientific inquiry, yet continues to pervade published scientific research (Van Calster et al. 2021). This paradox has been exemplified during the COVID-19 pandemic, which has led to unprecedented levels of published research of largely poor quality due to biases in conduct and reporting (Glasziou et al. 2020, Wynants et al. 2020).

Many researchers lack adequate training in research methods, and statistics is something they do with trepidation and even ignorance (Altman 1994, King et al. 2019). However, using the wrong statistical methods can cause real harm (Altman 1994, Brown et al. 2018) and bad statistical practices are being used to weaken science (Stark & Saltelli 2018). Statistical mistakes are a key source of research waste and are contributing to the current reproducibility crisis in science (Allison et al. 2016). Even when the correct methods are used, many researchers fail to describe them adequately, making it difficult to reproduce the results (Ernst & Albers 2017, Zhou & Skidmore 2018). Poor statistical methods might not be caught by reviewers, as they may not be qualified to judge the statistics. A recent survey of editors found that only 23% of health and medical journals used expert statistical review for all articles (Hardwicke & Goodman 2020), which was little different from a survey from 22 years ago (Goodman et al. 1998).

There is guidance for researchers on how to write up their statistical methods and results. The International Committee of Medical Journal Editors recommend that researchers should: “Describe statistical methods with enough detail to enable a knowledgeable reader with access to the original data to judge its appropriateness for the study and to verify the reported results” (ICJME 2019). More detailed guidance is given by the SAMPL and

EQUATOR guidelines (Lang & Altman 2013, Altman & Simera 2016) with the latter covering all aspects of the paper tailored to different study designs. Both of these guidelines were led by Doug Altman, who spoke often and for many years about the need for better statistical reporting. The awareness and use of these guidelines could be improved. There were 256 Google Scholar citations to the SAMPL paper (as at 15 March 2021) which is a good citation statistic for most papers, but is low considering the millions of papers that use statistical analysis.

A potential contributor to poor statistical reporting is the temptation for researchers to re-use descriptions of statistical methods, in an effort to make their papers resemble those of their peers and increase perceived chances of publication (Diong et al. 2018). As descriptions become more common, rebuke by reviewers and journal editors can be difficult, as their past use may be argued by researchers as offering precedent for the conduct of analysis within their discipline (Altman 2002). Two statisticians on this paper (AB and NW) have heard researchers admit that they have copied-and-pasted their statistical methods sections from other papers, regardless of whether they are appropriate. To investigate the extent of this practice, we applied a text-based clustering method to analyse the content of published statistical methods sections. Clustering results are then evaluated to estimate the extent that researchers are using cut-and-paste or ‘boilerplate’ statistical methods sections. Boilerplate text is that “which can be reused in new contexts or applications without significant changes to the original” (Wikipedia 2021). Use of these methods sections indicates that little thought has gone into the statistical analysis.

2 Methods

2.1 Data sources

We used two openly available data sources to find statistical methods sections: study protocols registered on the Australian and New Zealand Clinical Trials Registry (ANZCTR) and research articles published in *PLOS ONE*. Data sources were chosen as examples of common research outputs that include descriptions of statistical methods that were planned or used for analysing study outcomes.

2.1.1 Australia and New Zealand Clinical Trials Registry (ANZCTR)

The ANZCTR was established in 2005 as part of a coordinated global effort to improve research quality and transparency in clinical trials reporting; observational studies can also be registered. All studies registered on ANZCTR are publicly available and can be searched via an online portal (<https://www.anzctr.org.au>).

Details required for registration follow a standardised template (ANZCTR 2019), which covers participant eligibility, the intervention(s) being evaluated, study design and outcomes. The information provided must be in English. Studies are not peer reviewed.

For the statistical methods section, researchers are asked to provide a brief description of all sample size calculations, statistical methods and planned analyses, although this section is not compulsory (ANZCTR 2019). Studies are reviewed by ANZCTR staff for completeness of key information, which does not include the completeness of the statistical methods sections.

All studies available on ANZCTR were downloaded on 1 February 2020 in XML format. For our analysis, we used all text available in the “Statistical methods” section. We also collated basic information about the study including the study type (interventional or observational), submission date, number of funders and target sample size. These variables were chosen as we believed they might influence the completeness of the statistical methods section. For example, we hypothesised that larger studies and those with funding to be more complete. We were also interested in changes over time.

Studies prior to 2013 were excluded as the statistical methods section appeared to be introduced in 2013. Some studies were first registered on the alternative trial database *clinicaltrials.gov* and then also posted to ANZCTR. We excluded these studies because they almost all had no completed statistical methods section as this section is not included in *clinicaltrials.gov*.

Statistical methods sections were missing for some studies downloaded from ANZCTR, including sections labelled as “Not applicable”, “Nil” or “None”. We examined if there were particular studies where the statistical methods section was more likely to be missing. Analysis considered a logistic regression model estimated in the Bayesian framework [Rue et al. (2009); www.r-inla.org], with missing statistical methods section (yes/no) as the

dependent variable. The independent variables were date, study type (observational or interventional), number of funders and target sample size. Results were reported as odds ratios with 95% credible intervals (CI).

2.1.2 Public Library of Science (PLOS ONE)

PLOS ONE is a open access mega-journal that publishes original research across a wide range of scientific fields. Article submissions are handled by an academic editor who selects peer reviewers based on their self-nominated area(s) of expertise. Currently there are 324 academic editors out of 9,648 (3%) with the keywords of "statistics (mathematics)" or "statistical methods" in their expertise list (web search on 25-May-2021, <https://journals.plos.org/plosone/static/editorial-board>). Submissions do not undergo formal statistical review. Instead, reviewers are required to assess submissions against several publication criteria, including whether: "Experiments, statistics, and other analyses are performed to a high technical standard and are described in sufficient detail" (PLOS 2021). All reviewers are asked the question: "Has the statistical analysis been performed appropriately and rigorously?", with the possible responses of "Yes", "No" and "I don't know".

Authors are encouraged to follow published reporting guidelines such as EQUATOR, to ensure that chosen statistical methods are appropriate for the study design, and adequate details are provided to enable independent replication of results.

All *PLOS ONE* articles are freely accessible via the PLOS Application Programming Interface (API). This enabled us to conduct searches of full-text articles and analyse data on articles' text content and general attributes such as publication date and field(s) of research. Since *PLOS ONE* does not prescribe a set article format, we applied a two-step approach to identify statistical methods sections:

Step 1: Targeted API searches were completed using the R package 'rplos' (Chamberlain et al. 2020). Search queries targeted analysis-related terms, combining the words "data" or "statistical" with one of: "analysis", "analyses", "method", "methodology" or "model(l)ing". Terms could appear anywhere within the main body of the article, to account for the placement of relevant text in different sections, for example, in the *Material*

and *Methods* section versus *Results*. Search results were indexed by a unique Digital Object Identifier (DOI). Attribute data collected per DOI included journal volume and subject classification(s).

Step 2: PLOS ONE does not use standardised headings to preface statistical methods sections. To address this, we performed partial matching on available headings against frequently used terms in initial search results: ‘Statistical analysis’, ‘Statistical analyses’, ‘Statistical method’, ‘Statistics’, ‘Data analysis’ and ‘Data analyses’.

Data were downloaded on 3 July 2020. For records that did not pass Step 2, we reviewed where initial search terms appeared in the full-text, to estimate the proportion of statistical methods sections that were missed under our chosen approach.

2.2 Full-text processing

Text cleaning aimed to standardise notation and statistical terminology, whilst minimising changes to article style and formatting. *R* code used for data extraction and cleaning is available from https://github.com/agbarnett/stats_section.

Mathematical notation was converted from Unicode characters to plain text. For example, the Unicode character corresponding to θ (<U+03B8>) was replaced with ‘theta’. Common symbols outside of Unicode blocks including ‘%’ (percent) and ‘<’ (‘less-than’) were similarly converted into plain text. General formatting was removed, including carriage returns, punctuation marks, in-text references (e.g. “[42]”) centred equations, and other non-ASCII characters. Bracketed text was retained with brackets removed to maximise content for analysis. Common stop words including pronouns, contractions and selected prepositions were removed. We retained selected stop words that, if excluded, may have changed the context of statistical methods being described, for example ‘between’ and ‘against’.

We compiled an extensive list of statistical terms to standardise reported descriptions of statistical methods. An initial list was compiled by calculating individual word frequencies and identifying relevant terms that appeared at least 100 times. Further terms were sourced from index searches of three statistics textbooks (Diggle et al. 2013, Bland 2015, Dobson & Barnett 2018). Plurals (e.g., ‘chi-squares’) unhyphenated terms (e.g., ‘chi square’) and

combined terms (e.g. ‘chisquare’) were transformed to singular, hyphenated form (e.g., ‘chi-square’). Common statistical tests were also hyphenated (e.g., ‘hosmer lemeshow’ to ‘hosmer-lemeshow’). The final list is provided in Supplementary File 1.

2.3 Clustering algorithm

Text from statistical methods sections was analysed using Non-Negative Matrix Factorization (NMF). NMF is an established approach for topic modelling, and provides an effective solution for text-based clustering when dealing with high-dimensional data (Kim et al. 2014, Luong & Nayak (2019)).

For N studies, let $P \in R^{M \times N}$ denote a content matrix of text from statistical methods sections, comprising of M unique terms. Text clustering algorithms for identifying common topics across studies requires P to be represented with a vector space model. In our case, unique terms in P are modelled using the tf-idf (term frequency \times inverse document frequency) weighting schema, to account for the relative importance of common and rare terms.

A common problem facing text clustering algorithms is the curse of dimensionality due to the high number of terms in the doc \times term matrix representation (Aggarwal & Zhai 2012, Sutanto & Nayak (2018)). Applying text-based methods based on distance, density or probability therefore face difficulties in high-dimensional settings (Park et al. 2018, Mohotti & Nayak (2018a), Mohotti et al. (2019)). Specifically, distances between near and far points becomes negligible (Aggarwal & Zhai 2012). This behaviour directly affects the performance of distance-based clustering methods such as k -means (Jain 2010) in accurately identifying subgroups (topics) present in the data. Furthermore, sparseness associated with high-dimensional matrix representations does not allow for differentiation between topics based on density differences (Mohotti & Nayak 2018a, Mohotti & Nayak (2018b)).

To address these limitations, NMF deals with high-dimensional data by mapping it to a lower-dimensional space. This mapping is achieved by approximating P with two factor matrices: $W \in R^{M \times g}$ and $H \in R^{N \times g}$ (Aggarwal & Zhai 2012), such that $P \approx WH^T$. The number of subgroups of common topics inferred from the data is given by g .

The matrix factorization process approximates the lower dimensional non-negative factor matrices W and H such that they can represent high dimensional P with the least error. Estimation of W and H is achieved by optimising an objective function; for NMF, the Frobenius norm is used, equivalent to minimising the sum of squares for all elements of P :

$$\min \frac{1}{2} \|P - WH\|^2 = \sum_{i=1}^M \sum_{j=1}^N \left(P_{i,j} - (WH)_{i,j} \right)^2 \quad (1)$$

Following estimation, H contains the information regarding topic membership for all studies. In our case, topic membership $(1, \dots, g)$ for a statistical methods section is inferred from the maximum coefficient value in the corresponding row of H , also known as the topic coherence score. For our two datasets, we applied NMF with $g = 10$ topics.

2.4 Content analysis

Results were visualised by word clouds to summarise frequently occurring terms associated with topic membership. Follow up analysis considered similarities in text between statistical methods sections assigned to the same topic.

The Jaccard similarity index was used to compare the statistical methods section with the highest topic coherence score with all other sections assigned to the same topic. Indices were calculated at the sentence level, to account for instances of boilerplate text within larger methods sections. We chose the Jaccard index as an easy to interpret measure, which summarises the similarity between two sentences by the number of words common to each sentence (intersection), divided by the number of words that appeared in either sentence (union). Instances of boilerplate text were defined by a Jaccard index of 0.9 or higher and a difference in word count of plus or minus three words.

3 Results

3.1 ANZCTR

We downloaded 28,008 studies. The numbers of excluded studies are shown in Figure 1. Of the 12,700 included studies, 9,523 (75%) had a statistical methods section. The median length of sections was 129 words with an inter-quartile range of 71 to 219 words.




Figure 1: Search results for ANZCTR (left) and PLOS ONE (right).

Odds ratios and 95% credible intervals for study characteristics associated with missing statistical methods sections are in Table 1. Observational studies were less likely to have a missing statistical methods section compared with interventional studies. Missing sections became less likely over time. Studies with more funders and a larger target sample size were less likely to have a missing statistical methods section.

Figure 2 shows word clouds for the ten topics identified by the NMF algorithm. Frequent words related to the largest topic (topic 2, $n = 1834$) reflected descriptions of sample size calculations and represented 19% of studies analysed. Other topics indicated pilot studies (topic 5, $n = 834$), safety/tolerability studies (topic 6, $n = 524$), descriptive statistics (topic 7, $n = 333$), intervention studies (topic 8, $n = 1020$) and repeated measures ANOVA (topic 10 = 852). A review of word counts by topic identified statistical methods sections that were only one word, including “ANOVA”, “t-test”, “SPSS” and even “SSPS”.

There were cases where the exact same method section had been re-used in a different



Table 1: Logistic regression results for study characteristics associated with missing statistical methods sections in ANZCTR

Variable	Odds ratio	95% CI
Study type = Observational	0.78	(0.69, 0.89)
Date (per year)	0.90	(0.88, 0.91)
Number of funders	0.80	(0.74, 0.86)
Target sample size (per doubling)	0.90	(0.88, 0.92)

study. For example, in topic 7 ($n = 333$), 232 sections stated ‘descriptive statistics’ or ‘descriptive statistics used’ with no additional details provided. Other instances with expanded descriptions of methods included topic 3 ($n = 1277$) outlining descriptive analyses only, and topic 6 ($n = 524$), for hypothesis testing methods, statistical significance and software. In other cases, text had been slightly modified to account for changes in primary and secondary outcomes. Examples of these text changes were found in topics 2 and 4; identified instances related to sample size calculations for patient recruitment to different studies.

Examples of boilerplate text are provided in Table 2.

Since studies in this dataset described planned analyses, we hypothesised that some studies did not specify statistical methods because they had yet to consult with a statistician. Targeted searches for “statistician” across all topics returned 1,277 matching studies, with examples including “A statistician employed by hospital was used” and “Pilot study at this point will use a statistician professionally to determine sample size calculations as required”.

Table 2: Example boilerplate text from ANZCTR dataset

Topic	Statistical methods section
3	<p>Comparisons between categorical variables will be made either using chi square or Fisher exact test. Continuous data will be compared using the Student's t-test or Mann-Whitney U test. Two sided p values of less than 0.05 will be considered statistically significant.</p> <p>The Mann-Whitney U, Student t, 1-way ANOVA, and Kruskal-Wallis tests will be used to compare continuous variables where relevant. The Fisher exact and Pearson's Chi-square test will be used to compare proportions as appropriate.</p>
5	<p>Pilot study</p> <p>No formal sample size calculation was performed</p>
7	<p>Descriptive statistics</p> <p>Descriptive statistics used</p>
9	<p>Linear mixed models will be used to analyse the data.</p>
10	<p>Repeated measures of ANOVA</p> <p>Pre-, during, post- and follow-up variables will be subjected to mixed methods and repeated measures analyses to determine significant changes over (group and) time.</p>

3.2 PLOS ONE

API searches returned 131,847 papers (DOIs) (Figure 1). After partial matching, 111,731 (85%) statistical methods sections were identified. In the final sample, 95,518 (85%) DOIs returned an exact match against common section headings: 64,133 for ‘statistical analysis’, 13,380 for ‘statistical analyses’ and 13,627 for ‘data analysis’.




Figure 3: A: Search results by PLOS ONE volume; B: Subject classifications assigned to full-text recorded included in the analysis

Search results varied by journal volume (Figure 3A). The total number of API search results peaked at volumes 8 ($n = 19,045$) and 9 ($n = 19,045$) in years 2013 and 2014. This trend aligned with the total number of papers published in *PLOS ONE* over the same period. The percentage of records that included a statistical methods section varied between 64% (volume 2) and 86% (volume 9). All DOIs included “Biology and life sciences”

(n = 107,584), “Earth sciences” (n = 7,605) and/or “Computer and information sciences” (n = 5,190) in their top 3 subject classifications (Figure 3B).

Statistical methods sections had a median length of 127 words and inter-quartile range of 61 to 254 words. 7,450 articles (7%) had a statistical methods section of 500 words or more. 19,461 articles (17%) had statistical methods sections with 50 words or less, equal to the length of this paragraph.

For studies excluded based on section headings (n = 20,116), analysis-related search terms for 2,136 DOIs appeared in the general text of the Materials and Methods section. For studies without a matching Materials and Methods section, search terms often appeared in the Introduction (n = 728) and Results and/or Discussion (n = 981) sections. 4,010 DOIs included at least 1 section heading that contained the terms “analysis” or “analyses” without direct reference to statistical method(s) (e.g. Microarray analysis).

The word clouds for ten topics are in Figure 4. Frequently occurring words reflected the use of statistical software (topics 3 and 5), descriptive statistics (topic 6), group based hypothesis testing (topics 1 and 4) and definitions of statistical significance (topics 1 and 9). Topics related to regression (topic 2) and meta-analysis (topic 7) were also identified. Examples of boilerplate text for selected topics are in Table 3.

Topics related to statistical software differentiated between Prism GraphPad (topic 3: n = 9974) and SPSS (topic 5: n = 9648). Within topic 3, the sentence “GraphPad Prism (Graphpad Software, San Diego, CA) was used for all analyses” was matched in 1,941 papers based on our definition of boilerplate text. Similarly, for topic 5, “SPSS software version 17.0 (SPSS, Chicago, IL, USA) was used for statistical analysis” returned matches for 441 papers.

Table 3: Example boilerplate text from PLOS ONE papers (sentence level). The number of matching to each sentence was based on a Jaccard score of 0.9 or higher and a difference in word count of ± 3 words

Topic	Statistical methods text	Matching papers
Topic 1	Students t-test was used for statistical analysis.	29
	A p value of < 0.05 was considered statistically significant.	1883
Topic 3	GraphPad Prism (Graphpad Software, San Diego, CA) was used for all analyses.	1941
Topic 4	Significant differences were determined using analysis of variance (ANOVA) followed by tukey post-hoc tests for multiple comparisons.	2216
Topic 5	SPSS software version 17.0 (SPSS, Chicago, IL, USA) was used for statistical analysis.	441
Topic 6	All results are expressed as means \pm standard deviation (\pm SD).	102
Topic 9	Statistical significance was determined by Students t-tests.	49
	*p < 0.05 , **p < 0.01 , ***p < 0.001 .	6

Definitions of statistical significance featured strongly in topic 1 ($n = 3784$) and topic 9 ($n = 6195$). Topic 1 reflected applications of two-tailed and unpaired Student’s t-tests, however, instances of boilerplate text emphasised the 5% level of statistical significance (Table 3). In contrast, Topic 9 focused on multiple thresholds for declaring statistical significance by asterisk: “ $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$ ”, a practice that has been criticised (Wasserstein et al. 2019).

Group-based hypothesis testing was a recurring theme across topics, with text descriptions varying based on method(s) used. One-way analysis of variance also featured strongly in topic 4 ($n = 10212$), combined with common methods for performing post-hoc multiple comparisons. Based on the Jaccard index, 1 in 5 studies were matched to the sentence “Significant differences were determined using analysis of variance (ANOVA) followed by Tukey post-hoc tests for multiple comparisons”. Frequently occurring words in topic 6 ($n = 4764$) reflected mentions of descriptive statistics for summarising continuous variables, namely means with standard deviations or standard errors.

4 Discussion

The first line in many statistical analysis sections in *PLOS ONE* was the software used and some entire sections in ANZCTR only stated the software, implying that the software is the most important detail. As Doug Altman said, “Many people think that all you need to do statistics is a computer and appropriate software” (Altman 1994). This is far from the truth, and whilst it is important for researchers to mention the software and version used for reproducibility purposes, it is a minor detail compared with detailing what methods were used and why.

A frequent theme in the boilerplate statistical methods is the definition of statistical significance, nearly always using a p-value at the 5% level. This widespread use of statistical significance is troubling giving the bright-line thinking it engenders (McShane et al. 2019) and the common misinterpretations of p-values (Goodman 2008).

Despite the extensive array of statistical tests available, many authors are reporting the same few methods.

One reason these inadequate sections get published is that most journals do not use

statistical reviewers, despite empirical evidence showing they improve manuscript quality (Hardwicke & Goodman 2020).

A related paper has criticised vague statistical methods sections because they deprive readers and reviewers for the opportunity to confirm that the appropriate methods were used (Weissgerber Tracey et al. 2018). These authors checked hundreds of papers using ANOVA and found that 95% did not contain the information needed to determine what type of ANOVA was performed. This lack of information could well be because the authors used a boilerplate statistical methods section that was missing key details.

If authors shared their code then this would provide an alternative route for checking what statistical methods were used. This is not a perfect solution, as we still want authors to accurately report their methods, but it does increase transparency. However, a recent paper found that code sharing was very low in biomedical papers, with just 2% of a sample of over 6,000 papers sharing code (Serghiou et al. 2021).

Many researchers are using lazy practice by copying a standard “boilerplate” statistical methods section, likely cut-and-pasting from other researchers or projects. This is a strong sign of the ritualistic practice of statistics where researchers go through the motions rather than using conscientious practice (Stark & Saltelli 2018). This is concerning because using the wrong statistical methods can reduce the value of study, or worse, invalidate the entire study. These mistakes are avoidable and are wasting of thousands of hours of researchers’ time and the time of patients and volunteers. Poor statistical practice is a key driver of the ongoing reproducibility crisis in science (Ioannidis et al. 2014).

4.1 Limitations

We did not check whether papers used the correct methods, and for some simple studies a ‘boilerplate’ statistical methods might be adequate.

We examined papers where there was a statistics section, and we missed papers that used statistical analysis but did not include a statistical analysis section. Reiterate outcomes of random sample checking here.

We only examined one large journal and one trial registry and hence our results may not be generalisable to all journals or registries, especially those that consistently use a

statistical reviewer.

We searched the full text of *PLOS ONE* papers but not the supporting information which may contain statistical methods sections for some papers. The search terms we used to find statistical methods appeared in the supporting information titles for xxx papers (x%). We did not include the supporting information because it is less structured than the paper and could be in PDF or Word format.

References

- Aggarwal, C. C. & Zhai, C. (2012), *Mining text data*, Springer Science & Business Media.
- Allison, D. B., Brown, A. W., George, B. J. & Kaiser, K. A. (2016), ‘Reproducibility: A tragedy of errors’, *Nature* **530**(7588), 27–29.
URL: <https://doi.org/10.1038/530027a>
- Altman, D. G. (1994), ‘The scandal of poor medical research’, *BMJ* **308**(6924), 283–284.
URL: <https://doi.org/10.1136/bmj.308.6924.283>
- Altman, D. G. (2002), ‘Poor-quality medical research: what can journals do?’, *Jama* **287**(21), 2765–2767.
- Altman, D. G. & Simera, I. (2016), ‘A history of the evolution of guidelines for reporting medical research: the long road to the EQUATOR Network’, *Journal of the Royal Society of Medicine* **109**(2), 67–77.
URL: <https://doi.org/10.1177/0141076815625599>
- ANZCTR (2019), ANZCTR data field definitions v25, Technical report.
URL: <https://www.anzctr.org.au/docs/ANZCTR%20Data%20field%20explanation.pdf>
- Bland, M. (2015), *An Introduction to Medical Statistics*, Oxford medical publications, Oxford University Press.
- Brown, A. W., Kaiser, K. A. & Allison, D. B. (2018), ‘Issues with data and analyses: Errors, underlying themes, and potential solutions’, *Proceedings of the National Academy*

of Sciences **115**(11), 2563–2570.

URL: <https://www.pnas.org/content/115/11/2563>

Chamberlain, S., Boettiger, C. & Ram, K. (2020), *rplos: Interface to the Search API for 'PLOS' Journals*. R package version 0.9.0.

URL: <https://CRAN.R-project.org/package=rplos>

Diggle, P., Heagerty, P., Liang, K. & Zeger, S. (2013), *Analysis of Longitudinal Data*, Oxford Statistical Science Series, OUP Oxford.

Diong, J., Butler, A. A., Gandevia, S. C. & Héroux, M. E. (2018), 'Poor statistical reporting, inadequate data presentation and spin persist despite editorial advice', *PloS one* **13**(8), e0202121.

Dobson, A. & Barnett, A. (2018), *An Introduction to Generalized Linear Models*, Chapman & Hall/CRC Texts in Statistical Science, CRC Press.

Ernst, A. F. & Albers, C. J. (2017), 'Regression assumptions in clinical psychology research practice—a systematic review of common misconceptions', *PeerJ* **5**, e3323.

URL: <https://doi.org/10.7717/peerj.3323>

Glasziou, P. P., Sanders, S. & Hoffmann, T. (2020), 'Waste in covid-19 research', *The BMJ* **369**, m1847.

Goodman, S. (2008), 'A dirty dozen: Twelve p-value misconceptions', *Seminars in Hematology* **45**(3), 135–140.

URL: <https://doi.org/10.1053/j.seminhematol.2008.04.003>

Goodman, S. N., Altman, D. G. & George, S. L. (1998), 'Statistical reviewing policies of medical journals', *Journal of General Internal Medicine* **13**(11), 753–756.

URL: <https://doi.org/10.1046/j.1525-1497.1998.00227.x>

Hardwicke, T. E. & Goodman, S. (2020), 'How often do leading biomedical journals use statistical experts to evaluate statistical methods? The results of a survey'.

URL: osf.io/preprints/metaarxiv/z27u4

ICJME (2019), ‘Recommendations for the conduct, reporting, editing, and publication of scholarly work in medical journals’.

URL: <http://www.icmje.org/icmje-recommendations.pdf>

Ioannidis, J. P. A., Greenland, S., Hlatky, M. A., Khoury, M. J., Macleod, M. R., Moher, D., Schulz, K. F. & Tibshirani, R. (2014), ‘Increasing value and reducing waste in research design, conduct, and analysis’, *The Lancet* **383**(9912), 166–175.

URL: [https://doi.org/10.1016/s0140-6736\(13\)62227-8](https://doi.org/10.1016/s0140-6736(13)62227-8)

Jain, A. K. (2010), ‘Data clustering: 50 years beyond K-means’, *Pattern recognition letters* **31**(8), 651–666.

Kim, J., He, Y. & Park, H. (2014), ‘Algorithms for nonnegative matrix and tensor factorizations: a unified view based on block coordinate descent framework’, *Journal of Global Optimization* **58**(2), 285–319.

King, K. M., Pullmann, M. D., Lyon, A. R., Dorsey, S. & Lewis, C. C. (2019), ‘Using implementation science to close the gap between the optimal and typical practice of quantitative methods in clinical science’, *Journal of Abnormal Psychology* **128**(6), 547–562.

URL: <https://doi.org/10.1037/abn0000417>

Lang, T. & Altman, D. (2013), Basic statistical reporting for articles published in clinical medical journals: the SAMPL guidelines, *in* P. Smart, H. Maisonneuve & A. Polderman, eds, ‘Science Editors’ Handbook’, European Association of Science Editors.

Leek, J., McShane, B. B., Gelman, A., Colquhoun, D., Nuijten, M. B. & Goodman, S. N. (2017), ‘Five ways to fix statistics’, *Nature* **551**(7682), 557–559.

URL: <https://doi.org/10.1038/d41586-017-07522-z>

Luong, K. & Nayak, R. (2019), Clustering multi-view data using non-negative matrix factorization and manifold learning for effective understanding: A survey paper, *in* ‘Linking and Mining Heterogeneous and Multi-view Data’, Springer, pp. 201–227.

- McShane, B. B., Gal, D., Gelman, A., Robert, C. & Tackett, J. L. (2019), ‘Abandon statistical significance’, *The American Statistician* **73**(sup1), 235–245.
URL: <https://doi.org/10.1080/00031305.2018.1527253>
- Mohotti, W. A., Lukas, D. C. & Nayak, R. (2019), Concept mining in online forums using self-corpus-based augmented text clustering, in ‘Pacific Rim International Conference on Artificial Intelligence’, Springer, pp. 397–402.
- Mohotti, W. A. & Nayak, R. (2018*a*), Corpus-based augmented media posts with density-based clustering for community detection, in ‘2018 IEEE 30th International Conference on Tools with Artificial Intelligence (ICTAI)’, IEEE, pp. 379–386.
- Mohotti, W. A. & Nayak, R. (2018*b*), An efficient ranking-centered density-based document clustering method, in ‘Pacific-Asia Conference on Knowledge Discovery and Data Mining’, Springer, pp. 439–451.
- Park, A., Conway, M. & Chen, A. T. (2018), ‘Examining thematic similarity, difference, and membership in three online mental health communities from reddit: a text mining and visualization approach’, *Computers in human behavior* **78**, 98–112.
- PLOS (2021), Plos one: accelerating the publication of peer-reviewed science, Technical report.
URL: <https://journals.plos.org/plosone/s/criteria-for-publication>
- Rue, H., Martino, S. & Chopin, N. (2009), ‘Approximate Bayesian inference for latent Gaussian models using integrated nested Laplace approximations (with discussion).’, *Journal of the Royal Statistical Society B* **71**, 319–392.
- Serghiou, S., Contopoulos-Ioannidis, D. G., Boyack, K. W., Riedel, N., Wallach, J. D. & Ioannidis, J. P. A. (2021), ‘Assessment of transparency indicators across the biomedical literature: How open is open?’, *PLOS Biology* **19**(3), 1–26.
URL: <https://doi.org/10.1371/journal.pbio.3001107>
- Stark, P. B. & Saltelli, A. (2018), ‘Cargo-cult statistics and scientific crisis’, *Significance*

15(4), 40–43.

URL: <https://rss.onlinelibrary.wiley.com/doi/abs/10.1111/j.1740-9713.2018.01174.x>

Sutanto, T. & Nayak, R. (2018), ‘Fine-grained document clustering via ranking and its application to social media analytics’, *Social Network Analysis and Mining* **8**(1), 29.

Van Calster, B., Wynants, L., Riley, R. D., van Smeden, M. & Collins, G. S. (2021), ‘Methodology over metrics: Current scientific standards are a disservice to patients and society’, *Journal of Clinical Epidemiology* .

Wasserstein, R. L., Schirm, A. L. & Lazar, N. A. (2019), ‘Moving to a world beyond p < 0.05’, *The American Statistician* **73**(sup1), 1–19.

URL: <https://doi.org/10.1080/00031305.2019.1583913>

Weissgerber Tracey, L., Garcia-Valencia, O., Garovic, V. D., Milic, N. M. & Winham, S. J. (2018), ‘Why we need to report more than ’Data were analyzed by t-tests or ANOVA’’, *eLife* **7**.

URL: <https://gateway.library.qut.edu.au/login?url=https://search.proquest.com/docview/2174217344?>

Wikipedia (2021), Boilerplate text, Technical report.

URL: https://en.wikipedia.org/wiki/Boilerplate_text

Wynants, L., Van Calster, B., Collins, G. S., Riley, R. D., Heinze, G., Schuit, E., Bonten, M. M., Dahly, D. L., Damen, J. A., Debray, T. P. et al. (2020), ‘Prediction models for diagnosis and prognosis of covid-19: systematic review and critical appraisal’, *BMJ* **369**, m1328.

Zhou, Y. & Skidmore, S. (2018), ‘A reassessment of ANOVA reporting practices: A review of three APA journals’, *Journal of Methods and Measurement in the Social Sciences* **8**(1), 3–19.

URL: <https://journals.uair.arizona.edu/index.php/jmmss/article/view/22019>