

Module 3: Cross-recurrence quantification analysis of dyadic interaction

With the five MATLAB scripts in this module you can perform a simple cross-recurrence quantification analysis (CRQA) on categorical data. Start MATLAB and select the folder where the provided material (M-functions and data file) is located. That is, make MATLAB's current folder equal to the folder to which you downloaded the GitHub folder of this module (e.g. 'C:\...\My Documents'). After this you can call these functions on the command line.

1. First run the following piece of code:

```
load('example_data.mat')
```

Hint: Either copy-paste to an empty MATLAB script file and then run it, or copy-paste the lines in each individual assignment to the command line followed by 'enter'.

You have just loaded two time series, *PP1* and *PP2*, into the Workspace, but the data could be any two nominal data streams of equal length. These time series consist of nominal data having integer values from 1 to 5. The time series come from a dyadic interaction study, collaborative behaviors of two children was coded from video at 1 Hz by using the same five specific behavioral categories for each.

2. Plot these time series to see what they look like. Run:

```
plotTS(PP1, PP2)
```

3. Next step is to create the cross-recurrence matrix, *rec*, by running the following:

```
rec = CatCRMMatrix(PP1,PP2);
```

Have a look at the M-function *CatCRMMatrix* and try to understand what it does. Also, have a look at the matrix *rec* consisting of 1630×1630 points. The analysis distinguishes between two types of behavioral matches, which represent different combinations of behaviors of the interaction partners. Each combination corresponds to a different value (+1 or -1) in the recurrence matrix *rec*. All other combinations are considered to be non-matching and obtain the value 0.

Note: For nominal data you can directly build-up the cross-recurrence matrix by comparing the values in the two time series. In the cross-recurrence plot (CRP) a dot (i.e. recurrent point) is plotted whenever these values match in a pre-specified way. This first creates the main diagonal (Line-of-Synchrony) of the CRP, containing the equal-time behavioral matches. But by shifting the time series with respect to each other in both directions, lines parallel to the Line-of-Synchrony on both sides are created. Each of these lines reflects the behavioral

matches with a specific (increasing) delay between the occurrence of those behaviors in the two time series.

4. Now let's plot the CRP to study its structure, by running this:

PlotCRP(rec)

Note: This is a typical checkerboard pattern you would expect for nominal CRQA. There are three colors in this CRP representing the three different types of states of the dyadic system, based on the numerical values in *rec*: red for the value +1, blue for the value -1, and white for the value 0. All together the CRP nicely displays the rich coordinative structure of the dyadic interaction across all possible timescales. The color-coded analysis performed here is called **Chromatic CRQA**.

5. Finally, calculate the non-diagonal CRQA measures to quantify this structure. For now we will ignore the different types of behavioral matches (i.e. colors in the CRP) and treat them all as equal. Run the following function:

output = CRQA_out(rec)

This function performs **Anisotropic CRQA** by quantifying both the vertical and horizontal patterns. It produces the following measures:

<i>LAM</i>	=	Proportion of matches in vertical/horizontal patterns
<i>TT</i>	=	Average length of vertical/horizontal patterns
<i>Max_L</i>	=	Length of the longest vertical/horizontal pattern
<i>Ent_L</i>	=	Shannon entropy of vertical/horizontal length distribution

You can have a look at the results in the command window or by opening the file *output* from the Workspace. There are two rows in *output*. The upper row gives the values for the vertical patterns; the lower row those for the horizontal patterns.

Note: By calculating these measures in both the vertical and horizontal orientation and comparing them, Anisotropic CRQA enables you to study asymmetries in the dynamics. Differences between these measures reflect differences in relative contribution and dominance between the interaction partners. For more information about this see:

Cox, R.F.A., Van der Steen, S., De Jonge-Hoekstra, L., Guevara, M., & Van Dijk, M. (2016). Chromatic and anisotropic cross-recurrence quantification analysis of interpersonal behaviour. In C. Webber, C. Ioana, & N. Marwan (Eds). *Recurrence Plots and Their Quantifications: Expanding Horizons* (pp. 209-225). Springer Proceedings in Physics. ([link](#))