



GHRP-2

PEPTIDE RESEARCH COMPOUND

Dilution Guidelines

- Minimally 1–1.5 mL BAC water per 5 mg vial
- Tilt the vial and allow BAC water to flow slowly along the inner wall—do not inject directly onto the powder
- Avoid under-dilution, which may lead to clumping or reduced solubility
- Allow solution to reach room temperature before mixing
- Swirl gently—do not shake to maintain peptide integrity

Appearance After Mixing

- Typically forms a clear to slightly hazy solution
- Mild foaming may occur—allow to settle before use
- Full dissolution generally occurs within 1–2 minutes at room temperature

Storage

- Store at **2–8°C (refrigerated)** after reconstitution
- Do not freeze and protect from light and air exposure
- Use within 10–21 days for best stability

DISCLAIMER

This compound is provided for laboratory research purposes only. It is not intended for human or veterinary use, and no therapeutic or diagnostic claims are made.

Note

While both GHRP-2 and GHRP-6 are part of the growth hormone-releasing peptide family, research often distinguishes them based on specific focus areas. **GHRP-2** is commonly studied for its **stronger GH response** and potential **anti-inflammatory properties**, while **GHRP-6** is more frequently explored in relation to **appetite stimulation** and **ghrelin activity**. Selection may vary based on the intended area of research.