

# Existence of parabolic minimizers to the total variation flow on metric measure spaces

Cintia Pacchiano Camacho

joint work with

Vito Buffa,


Michael Collins (University of Erlangen)

AGENT Forum 2021

October 27th, 2021



# Setting



## Definition of Variational Solutions

We assume  $\Omega$  to be open and bounded,  $\Omega^*$  open and bounded with  $\Omega \Subset \Omega^*$  and

$$u_0 \in L^2(\Omega^*) \cap \text{BV}(\Omega^*). \quad (0.1)$$

### Definition 1

Assume that the Cauchy-Dirichlet datum  $u_0$  fulfills (0.1). A map  $u : \Omega_T^* \rightarrow \mathbb{R}$ ,  $T \in (0, \infty)$  in the class

$$L_w^1(0, T; \text{BV}_{u_0}(\Omega)) \cap C^0([0, T]; L^2(\Omega^*))$$

will be referred to as a *variational solution* on  $\Omega_T$  to the Cauchy-Dirichlet problem for the total variation flow if and only if the variational inequality

$$\begin{aligned} \int_0^T \|Du(t)\|(\Omega^*) \, dt &\leq \int_0^T \left[ \int_{\Omega^*} \partial_t v(v - u) \, d\mu + \|Dv(t)\|(\Omega^*) \right] dt \\ &\quad - \frac{1}{2} \|(v - u)(T)\|_{L^2(\Omega^*)}^2 + \frac{1}{2} \|v(0) - u_0\|_{L^2(\Omega^*)}^2 \end{aligned} \quad (0.2)$$

holds true for any  $v \in L_w^1(0, T; \text{BV}_{u_0}(\Omega))$  with  $\partial_t v \in L^2(\Omega_T^*)$  and  $v(0) \in L^2(\Omega^*)$ .

## Main results

### Theorem 2

*Suppose that the Cauchy-Dirichlet datum  $u_0$  fulfills the requirements of (0.1). Then, there exists a unique global variation solution in the sense of Definition 1.*

### Theorem 3

*Suppose that the Cauchy-Dirichlet datum  $u_0$  fulfills the requirements of (0.1). Then, any variational solution in the sense of Definition 1 on  $\Omega_T$  with  $T \in (0, \infty]$  satisfies*

$$\partial_t u \in L^2(\Omega^*) \text{ and } u \in C^{0, \frac{1}{2}}([0, \tau]; L^2(\Omega^*)) \text{ for all } \tau \in \mathbb{R} \cap (0, T].$$

*Furthermore, for the time derivative  $\partial_t u$  there holds the quantitative bound*

$$\int_0^T \int_{\Omega^*} |\partial_t u|^2 d\mu dt \leq \|Du_0\|(\Omega^*).$$

*Finally, for any  $t_1, t_2 \in \mathbb{R}$  with  $0 \leq t_1 < t_2 \leq T$  one has the energy estimate*

$$\frac{1}{t_2 - t_1} \int_{t_1}^{t_2} \|Du(t)\|(\Omega^*) dt \leq \|Du_0\|(\Omega^*). \quad (0.3)$$

# What's next?

Variational  
solutions



Existence  
Theory



Parabolic  
minimizers



Regularity  
Theory

Thanks!



Aalto University