



HELSINGIN YLIOPISTO  
HELSINGFORS UNIVERSITET  
UNIVERSITY OF HELSINKI

# **Singular stochastic integral operators and their applications to SPDEs**

Emiel Lorst

Joint work with M.C. Veraar



October 28, 2021

**University of Helsinki**  
**Department of Mathematics and Statistics**







# Polycrystalline materials







## Polycrystalline materials




The Allen-Cahn equation is a prototype for the growth of grains:




$t = 0$



$t = 1$




$t = 2$




$t = 50$




## The Allen-Cahn equation




$t = 0$



$t = 1$



$t = 2$




$t = 50$

Given an initial state  $u_0: \mathbb{T}^2 \rightarrow \mathbb{R}$ , look for a  $u: \mathbb{R}_+ \times \mathbb{T}^2 \rightarrow \mathbb{R}$  satisfying


$$\begin{cases} \frac{du}{dt} - \Delta u = -\Psi'(u) & \text{in } \mathbb{R}_+ \times \mathbb{T}^2, \\ u(0, \cdot) = u_0, \end{cases}$$

■  $\Psi$  is a double well potential:







# The Allen-Cahn equation




$t = 0$



$t = 1$



$t = 2$




$t = 50$

Given an initial state  $u_0: \mathbb{T}^2 \rightarrow \mathbb{R}$ , look for a  $u: \Omega \times \mathbb{R}_+ \times \mathbb{T}^2 \rightarrow \mathbb{R}$  satisfying

$$\begin{cases} du - \Delta u dt = -\Psi'(u) dt + B(u) dW & \text{in } \mathbb{R}_+ \times \mathbb{T}^2, \\ u(0, \cdot) = u_0, \end{cases}$$

- $\Psi$  is a double well potential:
- $W$  is a Brownian motion.
- For example  $B(u) = \varepsilon$ .





## Stochastic evolution equations

“Hide” the space-variable in a Banach space  $X$  to obtain an SDE:

Look for a function  $u: \Omega \times \mathbb{R}_+ \rightarrow X$  satisfying

$$\begin{cases} du + Au \, dt = F(u) \, dt + G(u) \, dW & \text{in } \mathbb{R}_+, \\ u(0) = u_0. \end{cases}$$



## Stochastic evolution equations

“Hide” the space-variable in a Banach space  $X$  to obtain an SDE:

Look for a function  $u: \Omega \times \mathbb{R}_+ \rightarrow X$  satisfying

$$\begin{cases} du + Au \, dt = F(u) \, dt + G(u) \, dW & \text{in } \mathbb{R}_+, \\ u(0) = u_0. \end{cases}$$

For Allen-Cahn on  $\mathbb{T}^2$ :

- $X = L^q(\mathbb{T}^2)$
- $A = -\Delta$
- $F(u) = -\Psi'(u)$   
 $G(u) = B(u)$
- $u_0 \in W^{2,q}(\mathbb{T}^2)$



## Stochastic evolution equations

“Hide” the space-variable in a Banach space  $X$  to obtain an SDE:

Look for a function  $u: \Omega \times \mathbb{R}_+ \rightarrow X$  satisfying

$$\begin{cases} du + Au \, dt = F(u) \, dt + G(u) \, dW & \text{in } \mathbb{R}_+, \\ u(0) = u_0. \end{cases}$$

For Allen-Cahn on  $\mathbb{T}^2$ :

- $X = L^q(\mathbb{T}^2)$
- $A = -\Delta$
- $F(u) = -\Psi'(u)$   
 $G(u) = B(u)$
- $u_0 \in W^{2,q}(\mathbb{T}^2)$

### Question

Under what conditions does a unique solution exist?





## Abstract Cauchy problem

We first study the problem with  $F$  and  $G$  independent of  $u$  and  $u_0 = 0$ .

Take  $f, g: \Omega \times \mathbb{R}_+ \rightarrow X$ . We look for a function  $u: \Omega \times \mathbb{R}_+ \rightarrow X$  satisfying

$$\begin{cases} du + Au \, dt = f \, dt + g \, dW, \\ u(0) = 0, \end{cases}$$



## Abstract Cauchy problem

We first study the problem with  $F$  and  $G$  independent of  $u$  and  $u_0 = 0$ .

Take  $f, g: \Omega \times \mathbb{R}_+ \rightarrow X$ . We look for functions  $u, v: \Omega \times \mathbb{R}_+ \rightarrow X$

$$\begin{cases} \frac{du}{dt} + Au = f, \\ u(0) = 0. \end{cases} \qquad \begin{cases} dv + Av \, dt = g \, dW, \\ v(0) = 0. \end{cases}$$



## Abstract Cauchy problem

We first study the problem with  $F$  and  $G$  independent of  $u$  and  $u_0 = 0$ .

Take  $f, g: \Omega \times \mathbb{R}_+ \rightarrow X$ . We look for functions  $u, v: \Omega \times \mathbb{R}_+ \rightarrow X$

$$\begin{cases} \frac{du}{dt} + Au = f, \\ u(0) = 0. \end{cases} \quad \begin{cases} dv + Av \, dt = g \, dW, \\ v(0) = 0. \end{cases}$$

The solutions  $u$  and  $v$  are given by the variation of constants formulas:

$$u(t) = \int_0^t e^{-(t-s)A} f(s) \, ds, \quad v(t) = \int_0^t e^{-(t-s)A} g(s) \, dW(s),$$

where  $(e^{-tA})_{t \geq 0}$  is a semigroup of bounded operators on  $X$ .



## Abstract Cauchy problem

We first study the problem with  $F$  and  $G$  independent of  $u$  and  $u_0 = 0$ .

Take  $f, g: \Omega \times \mathbb{R}_+ \rightarrow X$ . We look for functions  $u, v: \Omega \times \mathbb{R}_+ \rightarrow X$

$$\begin{cases} \frac{du}{dt} + Au = f, \\ u(0) = 0. \end{cases} \quad \begin{cases} dv + Av \, dt = g \, dW, \\ v(0) = 0. \end{cases}$$

The solutions  $u$  and  $v$  are given by the variation of constants formulas:

$$u(t) = \int_0^t e^{-(t-s)A} f(s) \, ds, \quad v(t) = \int_0^t e^{-(t-s)A} g(s) \, dW(s),$$

where  $(e^{-tA})_{t \geq 0}$  is a semigroup of bounded operators on  $X$ .

### Deterministic question

If

$$f \in L^p(\mathbb{R}_+; X),$$

do we have

$$Au \in L^p(\mathbb{R}_+; X)?$$

$$f \in L^p(\mathbb{R}_+; X)$$

$\Leftrightarrow$

$$\|f\|_{L^p(\mathbb{R}_+; X)} = \left( \int_0^\infty \|f(t)\|_X^p \, dt \right)^{\frac{1}{p}} < \infty$$



## Abstract Cauchy problem

We first study the problem with  $F$  and  $G$  independent of  $u$  and  $u_0 = 0$ .

Take  $f, g: \Omega \times \mathbb{R}_+ \rightarrow X$ . We look for functions  $u, v: \Omega \times \mathbb{R}_+ \rightarrow X$

$$\begin{cases} \frac{du}{dt} + Au = f, \\ u(0) = 0. \end{cases} \quad \begin{cases} dv + Av dt = g dW, \\ v(0) = 0. \end{cases}$$

The solutions  $u$  and  $v$  are given by the variation of constants formulas:

$$u(t) = \int_0^t e^{-(t-s)A} f(s) ds, \quad v(t) = \int_0^t e^{-(t-s)A} g(s) dW(s),$$

where  $(e^{-tA})_{t \geq 0}$  is a semigroup of bounded operators on  $X$ .

### Deterministic question

If

$$f \in L^p(\mathbb{R}_+; X),$$

do we have

$$Au \in L^p(\mathbb{R}_+; X)?$$

### Stochastic question

If

$$g \in L^p(\Omega \times \mathbb{R}_+; X),$$

do we have

$$A^{\frac{1}{2}} v \in L^p(\mathbb{R}_+; X)?$$



## From SPDEs to harmonic analysis

Note that

$$Au(t) = \int_0^t Ae^{-(t-s)A}f(s) \, ds, \quad A^{\frac{1}{2}}v(t) = \int_0^t A^{\frac{1}{2}}e^{-(t-s)A}g(s) \, dW(s).$$



## From SPDEs to harmonic analysis

Note that

$$Au(t) = \int_0^t Ae^{-(t-s)A} f(s) \, ds, \quad A^{\frac{1}{2}} v(t) = \int_0^t A^{\frac{1}{2}} e^{-(t-s)A} g(s) \, dW(s).$$

So we can reformulate our questions as:

### Deterministic question

Does

$$T_K f(t) = \int_0^\infty K(t-s) f(s) \, ds$$

for  $K(t) = Ae^{-tA} \mathbf{1}_{t>0}$

define a bounded operator on

$$L^p(\mathbb{R}_+; X)?$$

### Stochastic question

Does

$$S_K g(t) = \int_0^\infty K(t-s) g(s) \, dW(s)$$

for  $K(t) = A^{\frac{1}{2}} e^{-tA} \mathbf{1}_{t>0}$

define a bounded operator on

$$L^p(\Omega \times \mathbb{R}_+; X)?$$



## From SPDEs to harmonic analysis

Note that

$$Au(t) = \int_0^t Ae^{-(t-s)A} f(s) \, ds, \quad A^{\frac{1}{2}} v(t) = \int_0^t A^{\frac{1}{2}} e^{-(t-s)A} g(s) \, dW(s).$$

So we can reformulate our questions as:

### Deterministic question

Does

$$T_K f(t) = \int_0^\infty K(t-s) f(s) \, ds$$

for  $K(t) = Ae^{-tA} \mathbf{1}_{t>0}$

define a bounded operator on

$$L^p(\mathbb{R}_+; X)?$$

### Stochastic question

Does

$$S_K g(t) = \int_0^\infty K(t-s) g(s) \, dW(s)$$

for  $K(t) = A^{\frac{1}{2}} e^{-tA} \mathbf{1}_{t>0}$

define a bounded operator on

$$L^p(\Omega \times \mathbb{R}_+; X)?$$

These kernels  $K$  are singular:

$$\|K(t)\| \leq \frac{C}{t}$$

$$\|K(t)\| \leq \frac{C}{t^{\frac{1}{2}}}$$





## (Stochastic) singular integral operators

The operator

$$T_K f(t) = \int_{\mathbb{R}^d} K(t-s) f(s) \, ds$$

on  $L^p$  for a singular kernel  $K$  has been studied thoroughly:

- Hilbert transform  $K(t) = \frac{1}{\pi} \frac{1}{t}$  ('28)
- Regularity theory for elliptic PDE ('40s)
- Calderón–Zygmund and Fourier multiplier theory ('50s)
- Regularity theory for parabolic PDE ('00s)



## (Stochastic) singular integral operators

The operator

$$T_K f(t) = \int_{\mathbb{R}^d} K(t-s) f(s) \, ds$$

on  $L^p$  for a singular kernel  $K$  has been studied thoroughly:

- Hilbert transform  $K(t) = \frac{1}{\pi} \frac{1}{t}$  ('28)
  - Regularity theory for elliptic PDE ('40s)
  - Calderón–Zygmund and Fourier multiplier theory ('50s)
  - Regularity theory for parabolic PDE ('00s)
- 

The operator

$$S_K g(t) = \int_0^\infty K(t-s) g(s) \, dW(s)$$

on  $L^p$  for a singular kernel  $K$  has only been studied in a few special cases.



## Stochastic Calderón–Zygmund theory

Theorem (L., Veraar, Analysis & PDE '21)

- Let  $X$  be a Banach space with certain geometric properties.
- Let  $K: \mathbb{R}_+ \rightarrow \mathcal{L}(X)$  be a smooth singular kernel.

Suppose that  $S_K$  is bounded on  $L^{p_0}(\mathbb{R}_+ \times \Omega; X)$  for **some**  $p_0 \in [2, \infty)$ .

Then  $S_K$  is bounded on  $L^p(\mathbb{R}_+ \times \Omega; X)$  for **all**  $p \in (2, \infty)$ .



## Stochastic Calderón–Zygmund theory

### Theorem (L., Veraar, Analysis & PDE '21)

- Let  $X$  be a Banach space with certain geometric properties.
- Let  $K: \mathbb{R}_+ \rightarrow \mathcal{L}(X)$  be a smooth singular kernel.

Suppose that  $S_K$  is bounded on  $L^{p_0}(\mathbb{R}_+ \times \Omega; X)$  for **some**  $p_0 \in [2, \infty)$ .

Then  $S_K$  is bounded on  $L^p(\mathbb{R}_+ \times \Omega; X)$  for **all**  $p \in (2, \infty)$ .

- $L^q$  is allowed for  $q \in [2, \infty)$ .
- It suffices to have

$$\|K(t)\| \leq \frac{C}{t^{\frac{1}{2}}}, \quad \|K'(t)\| \leq \frac{C}{t^{\frac{3}{2}}}.$$



## Stochastic Calderón–Zygmund theory

### Theorem (L., Veraar, Analysis & PDE '21)

- Let  $X$  be a Banach space with certain geometric properties.
- Let  $K: \mathbb{R}_+ \rightarrow \mathcal{L}(X)$  be a smooth singular kernel.

Suppose that  $S_K$  is bounded on  $L^{p_0}(\mathbb{R}_+ \times \Omega; X)$  for **some**  $p_0 \in [2, \infty)$ .  
Then  $S_K$  is bounded on  $L^p(\mathbb{R}_+ \times \Omega; X)$  for **all**  $p \in (2, \infty)$ .

- $L^q$  is allowed for  $q \in [2, \infty)$ .
- It suffices to have

$$\|K(t)\| \leq \frac{C}{t^{\frac{1}{2}}}, \quad \|K'(t)\| \leq \frac{C}{t^{\frac{3}{2}}}.$$

### Corollary

If our stochastic question holds for **some**  $p_0 \in [2, \infty)$ , then it holds for **all**  $p \in (2, \infty)$ .

For our deterministic question this was shown in:

(Dore, Adv. Differential Equations, '00).



## Back to Allen-Cahn

Given an initial state  $u_0 \in L^q(\mathbb{T}^2)$ , look for a  $u: \Omega \times \mathbb{R}_+ \rightarrow L^q(\mathbb{T})$  satisfying

$$\begin{cases} du - \Delta u \, dt = -\Psi'(u) \, dt + B(u) \, dW & \text{in } \mathbb{R}_+, \\ u(0, \cdot) = u_0. \end{cases}$$



## Back to Allen-Cahn

Given an initial state  $u_0 \in L^q(\mathbb{T}^2)$ , look for a  $u: \Omega \times \mathbb{R}_+ \rightarrow L^q(\mathbb{T})$  satisfying

$$\begin{cases} du - \Delta u dt = -\Psi'(u) dt + B(u) dW & \text{in } \mathbb{R}_+, \\ u(0, \cdot) = u_0. \end{cases}$$

Positive answers to our  
det. and stoch. question

$\Rightarrow$

Banach fixed  
point theorem

$\Rightarrow$

Unique solution  $u$  in  
 $L^p(\Omega \times \mathbb{R}_+; L^q(\mathbb{T}))$



## Back to Allen-Cahn

Given an initial state  $u_0 \in L^q(\mathbb{T}^2)$ , look for a  $u: \Omega \times \mathbb{R}_+ \rightarrow L^q(\mathbb{T})$  satisfying

$$\begin{cases} du - \Delta u dt = -\Psi'(u) dt + B(u) dW & \text{in } \mathbb{R}_+, \\ u(0, \cdot) = u_0. \end{cases}$$

Positive answers to our  
det. and stoch. question

$\Rightarrow$

Banach fixed  
point theorem

$\Rightarrow$

Unique solution  $u$  in  
 $L^p(\Omega \times \mathbb{R}_+; L^q(\mathbb{T}))$

Our theorem allows one to study the stochastic question in

$$L^q(\Omega \times \mathbb{R}_+; L^q(\mathbb{T}^2)) = L^q(\Omega \times \mathbb{R}_+ \times \mathbb{T}^2),$$

which is significantly simpler than  $p \neq q$ .





## Back to Allen-Cahn

Given an initial state  $u_0 \in L^q(\mathbb{T}^2)$ , look for a  $u: \Omega \times \mathbb{R}_+ \rightarrow L^q(\mathbb{T})$  satisfying

$$\begin{cases} du - \Delta u dt = -\Psi'(u) dt + B(u) dW & \text{in } \mathbb{R}_+, \\ u(0, \cdot) = u_0. \end{cases}$$

Positive answers to our  
det. and stoch. question

$\Rightarrow$

Banach fixed  
point theorem

$\Rightarrow$

Unique solution  $u$  in  
 $L^p(\Omega \times \mathbb{R}_+; L^q(\mathbb{T}))$

Our theorem allows one to study the stochastic question in

$$L^q(\Omega \times \mathbb{R}_+; L^q(\mathbb{T}^2)) = L^q(\Omega \times \mathbb{R}_+ \times \mathbb{T}^2),$$

which is significantly simpler than  $p \neq q$ .

Why not just analyze  $p = q = 2$ ?

- More classical smoothness through Sobolev embeddings with large  $p, q$
- Scaling of the nonlinearities can dictate  $p \neq q$ .



**Thank you for your attention!**