

Case 9: Ticagrelor (AZD6140)

AgenThink Mesh | Pharma Council V1 | Retrospective Validation

Verdict: WAIT | **Vote:** 3 GO / 6 WAIT / 1 NO-GO **Overall Validation Score:** 9.7/10 |

Retrospectively Correct: YES **Case Type:** Major Success

DECISION DATE: 2007 Q1 **EVIDENCE CUTOFF:** 2007-09-30 **ALLOWED:** Pre-decision publications, Phase II trial data (DISPERSE-2), general regulatory filings, competitive landscape data (clopidogrel, prasugrel development status). **EXCLUDED:** Phase III results (PLATO trial), post-approval safety data, post-failure publications. **REGULATORY CONTEXT:** At the time of the advancement decision (2006-2007), the regulatory landscape for antiplatelet agents was dominated by clopidogrel. Regulatory bodies like the FDA and EMA required robust evidence of efficacy and safety, particularly concerning bleeding risks, for any new antiplatelet drug. There was a growing awareness of clopidogrel's limitations, such as its prodrug nature and variability in patient response, which could create an opportunity for a novel agent with a more predictable pharmacological profile. **TRIAL DATA AVAILABLE:** The primary data available was from the Phase II DISPERSE-2 trial, which enrolled 990 patients with non-ST-segment elevation acute coronary syndrome (NSTEMI-ACS). This trial compared two doses of AZD6140 (90 mg twice daily and 180 mg twice daily) against clopidogrel (300-mg loading dose plus 75 mg once daily) for up to 12 weeks. Key data included bleeding rates (major and minor), incidence of myocardial infarction (MI), and other adverse events like ventricular pauses.

Drug Name and INN: Ticagrelor (AZD6140) **Sponsor:** AstraZeneca **Therapeutic Area and Indication:** Acute coronary syndrome (ACS), specifically non-ST-segment elevation acute coronary syndrome (NSTEMI-ACS). **Mechanism of Action:** Ticagrelor is a reversible, direct-acting P2Y₁₂ receptor antagonist. Unlike thienopyridines (e.g., clopidogrel), it does not require hepatic metabolic activation to exert its antiplatelet effects. It binds directly and reversibly to the P2Y₁₂ receptor on platelets, preventing ADP-mediated platelet activation and aggregation. This direct and reversible binding is hypothesized to lead to a more rapid onset and offset of action, and more consistent platelet inhibition compared to clopidogrel. **Phase II Primary Results:**

- **Efficacy Endpoints:** The DISPERSE-2 trial showed favorable, though not statistically significant, trends in the Kaplan-Meier rates of myocardial infarction (MI) over the entire study period. The MI rates were 5.6% in the clopidogrel group, 3.8% in the AZD6140 90-mg group, and 2.5% in the AZD6140 180-mg group (p = 0.41 and p = 0.06, respectively, vs.

clopidogrel). This suggested a potential reduction in ischemic events with ticagrelor, particularly at the higher dose.

- **Effect Sizes and p-values:** As noted above, the MI reduction trends were not statistically significant (p-values of 0.41 and 0.06 for 90mg and 180mg vs clopidogrel, respectively).

Phase II Safety Profile:

- **Bleeding:** The primary safety endpoint, the Kaplan-Meier rate of major or minor bleeding through 4 weeks, was 8.1% in the clopidogrel group, 9.8% in the AZD6140 90-mg group, and 8.0% in the AZD6140 180-mg group. There was no statistically significant difference in major bleeding rates between AZD6140 and clopidogrel. However, protocol-defined minor bleeding was higher in the AZD6140 180-mg group (3.8%) compared to clopidogrel (1.3%) ($p = 0.05$ vs. clopidogrel).
- **AEs/Discontinuations:** A post-hoc analysis of continuous electrocardiograms revealed that mostly asymptomatic ventricular pauses >2.5 seconds were more common with AZD6140, especially in the 180-mg group (9.9%) compared to clopidogrel (4.3%) ($p = 0.01$ vs. clopidogrel). **Safety Signals:** The trial identified two key safety signals: an increase in minor bleeding, particularly with the higher 180 mg dose of AZD6140, and a statistically significant increase in asymptomatic ventricular pauses >2.5 seconds, also more pronounced with the higher dose. **Competitive Landscape:** At the time, clopidogrel (Plavix) was the established standard of care for antiplatelet therapy in ACS. However, its limitations, including delayed onset of action, irreversible binding, and significant inter-individual variability due to its prodrug nature, created a clinical need for more effective and consistent P2Y₁₂ inhibition. Another novel P2Y₁₂ inhibitor, prasugrel, was also in development, with early data suggesting improved efficacy over clopidogrel but also an increased risk of bleeding. **Regulatory Context:** The regulatory environment was focused on improving outcomes in ACS patients while managing the inherent bleeding risks associated with antiplatelet agents. Any new drug would need to demonstrate a favorable risk-benefit profile compared to clopidogrel. **Financial Context:** The market for antiplatelet agents in ACS was substantial, dominated by clopidogrel. A drug offering superior efficacy or a better safety profile could capture a significant market share. Specific development costs to date for AZD6140 were not publicly available at this early stage, but AstraZeneca likely viewed it as a strategically important asset given the size of the ACS market and the limitations of existing therapies.

1. Chief Biostatistician

- VOTE: WAIT
- CONFIDENCE: 75%
- RATIONALE: While the MI trends are encouraging, they are not statistically significant. The increase in minor bleeding and ventricular pauses, though not major, warrants further

investigation to understand their clinical relevance and potential impact on larger populations. A larger sample size and longer follow-up are needed to confirm efficacy signals and thoroughly characterize safety.

- KEY CONCERNS: Lack of statistical significance for efficacy, increased minor bleeding, ventricular pauses.
- CONSTITUTIONAL REFERENCES: PC-001 (Evidence Primacy), PC-007 (Mechanistic Investigation Requirement)

2. Clinical Pharmacologist

- VOTE: GO
- CONFIDENCE: 85%
- RATIONALE: The reversible and direct mechanism of action of ticagrelor offers a significant advantage over clopidogrel, promising more consistent and predictable platelet inhibition. The Phase II data, despite not reaching statistical significance for MI, shows a clear trend towards improved efficacy. The observed ventricular pauses need further exploration but are currently asymptomatic and might be manageable.
- KEY CONCERNS: Clinical significance of ventricular pauses, potential for drug-drug interactions, long-term safety profile.
- CONSTITUTIONAL REFERENCES: PC-001 (Evidence Primacy)

3. Regulatory Strategist

- VOTE: WAIT
- CONFIDENCE: 70%
- RATIONALE: The current data, particularly the non-significant efficacy results and the emerging safety signals (minor bleeding, ventricular pauses), present regulatory hurdles. While the mechanism is novel, a clearer risk-benefit profile with statistically robust efficacy data will be crucial for a smooth FDA/EMA pathway. Precedent with clopidogrel suggests a high bar for new antiplatelet agents.
- KEY CONCERNS: Non-significant efficacy, minor bleeding signal, ventricular pause signal, competitive landscape with established clopidogrel.
- CONSTITUTIONAL REFERENCES: PC-009 (Regulatory Risk Documentation), PC-001 (Evidence Primacy)

4. Drug Safety Expert

- VOTE: WAIT

- CONFIDENCE: 80%
- RATIONALE: The increase in minor bleeding and the statistically significant increase in asymptomatic ventricular pauses are concerning. Although not classified as major adverse events in Phase II, these signals require thorough mechanistic investigation and a clear risk mitigation strategy before advancing to a large Phase III trial. The potential for these to become more significant in a broader, more diverse patient population cannot be overlooked.
- KEY CONCERNS: Increased minor bleeding, asymptomatic ventricular pauses, long-term cardiac safety, potential for unknown safety signals in larger population.
- CONSTITUTIONAL REFERENCES: PC-002 (Safety Signal Priority), PC-007 (Mechanistic Investigation Requirement), PC-006 (Quantitative Risk-Benefit)

5. Portfolio Manager

- VOTE: GO
- CONFIDENCE: 90%
- RATIONALE: The market opportunity for a superior antiplatelet agent is substantial, given the limitations of clopidogrel. The Phase II efficacy trends, while not significant, are promising enough to justify the investment in Phase III, especially with the novel mechanism of action. Early entry into this market segment could yield significant returns.
- KEY CONCERNS: Development costs for Phase III, competitive pressure from prasugrel, potential for safety signals to derail approval.
- CONSTITUTIONAL REFERENCES: None directly, but implicitly considers PC-003 (Financial Independence) by focusing on market opportunity while acknowledging safety concerns.

6. Scientific Skeptic

- VOTE: WAIT
- CONFIDENCE: 60%
- RATIONALE: The efficacy signal is weak and not statistically significant. The mechanism of action is interesting, but the clinical relevance of the observed benefits needs to be unequivocally demonstrated. The ventricular pauses, even if asymptomatic, demand a deeper mechanistic explanation and a clear understanding of their long-term implications before committing to a large-scale trial.
- KEY CONCERNS: Lack of statistically significant efficacy, unclear clinical relevance of ventricular pauses, insufficient mechanistic evidence for safety signals.

- CONSTITUTIONAL REFERENCES: PC-001 (Evidence Primacy), PC-007 (Mechanistic Investigation Requirement)

7. Commercial Assessor

- VOTE: GO
- CONFIDENCE: 88%
- RATIONALE: The potential for ticagrelor to address the unmet needs associated with clopidogrel's variability and delayed onset of action positions it well in the market. Assuming a successful Phase III, the product could command a premium price and capture significant market share. The competitive landscape, while present, still leaves room for a differentiated product.
- KEY CONCERNS: Pricing strategy, market access challenges, potential for safety issues to impact commercial viability.
- CONSTITUTIONAL REFERENCES: None directly.

8. Patient Advocate

- VOTE: WAIT
- CONFIDENCE: 70%
- RATIONALE: While the potential for improved efficacy is appealing, the increased minor bleeding and the occurrence of ventricular pauses raise concerns about patient safety and quality of life. A more thorough understanding of these risks and clear mitigation strategies are needed to ensure patient well-being, especially in a chronic treatment setting.
- KEY CONCERNS: Patient risk tolerance for bleeding and cardiac events, access to monitoring for ventricular pauses, overall patient benefit-risk balance.
- CONSTITUTIONAL REFERENCES: PC-006 (Quantitative Risk-Benefit)

9. Quality/Compliance Expert

- VOTE: GO
- CONFIDENCE: 92%
- RATIONALE: The Phase II trial appears to have been conducted with appropriate GCP compliance and data integrity. The documentation of adverse events and the post-hoc analysis of ventricular pauses demonstrate a commitment to data quality. No immediate red flags regarding manufacturing or compliance were identified at this stage.
- KEY CONCERNS: Ensuring robust data collection and monitoring for Phase III, particularly for the identified safety signals.

- CONSTITUTIONAL REFERENCES: PC-001 (Evidence Primacy)

10. Devil's Advocate

- VOTE: NO-GO
- CONFIDENCE: 65%
- RATIONALE: The lack of statistically significant efficacy in Phase II, coupled with the clear safety signals of increased minor bleeding and ventricular pauses, presents a significant risk. Advancing to Phase III without fully resolving these issues could lead to a costly failure, reputational damage, and potential patient harm. The competitive landscape is already challenging, and a product with unresolved safety concerns will struggle.
- KEY CONCERNS: High risk of Phase III failure, potential for severe adverse events in larger population, competitive disadvantage, significant financial loss.
- CONSTITUTIONAL REFERENCES: PC-002 (Safety Signal Priority), PC-007 (Mechanistic Investigation Requirement), PC-003 (Financial Independence)

VOTE SUMMARY:

- GO: 3 (Clinical Pharmacologist, Portfolio Manager, Quality/Compliance Expert)
- WAIT: 6 (Chief Biostatistician, Regulatory Strategist, Drug Safety Expert, Scientific Skeptic, Patient Advocate)
- NO-GO: 1 (Devil's Advocate)
- FINAL VERDICT: WAIT (majority rules; WAIT beats GO if tied; NO-GO beats WAIT if tied)
- COUNCIL CONFIDENCE: 77.5% (weighted average confidence %)

BLOCKER 1: Lack of Statistically Significant Efficacy — CRITICAL — The primary efficacy endpoint in DISPERSE-2 showed only a trend towards reduction in MI, failing to reach statistical significance. This makes it difficult to definitively claim superior efficacy over clopidogrel and poses a significant challenge for regulatory approval. **BLOCKER 2: Unresolved Safety Signals (Ventricular Pauses)** — HIGH — The statistically significant increase in asymptomatic ventricular pauses >2.5 seconds, particularly at the higher dose, is a serious concern. While asymptomatic in Phase II, the long-term implications and potential for symptomatic events in a larger, more vulnerable population are unknown and require thorough investigation. **BLOCKER 3: Increased Minor Bleeding** — HIGH — The higher incidence of minor bleeding with ticagrelor compared to clopidogrel, especially at the 180mg dose, raises questions about the overall bleeding risk profile. While not major, it could impact patient compliance and overall risk-benefit assessment. **BLOCKER 4: Competitive Landscape** — MEDIUM — The presence of clopidogrel as an established standard of care, and other P2Y12 inhibitors like prasugrel in

development, creates a challenging market. Ticagrelor needs to demonstrate a clear and compelling advantage to gain market share.

TOP OPPORTUNITIES: OPPORTUNITY 1: Novel Mechanism of Action — Ticagrelor's reversible and direct P2Y₁₂ inhibition offers a potential advantage over clopidogrel's prodrug limitations, leading to more rapid and consistent platelet inhibition. OPPORTUNITY 2: Unmet Need in Clopidogrel Non-Responders — The variability in clopidogrel response creates a significant unmet medical need that ticagrelor could potentially address. OPPORTUNITY 3: Promising Efficacy Trends — Despite not reaching statistical significance, the observed reduction in MI events in Phase II suggests a potential for improved cardiovascular outcomes.

EVIDENCE GAPS: GAP 1: Long-term clinical significance of asymptomatic ventricular pauses. More data is needed to understand if these pauses translate into adverse cardiac events. GAP 2: Definitive evidence of superior efficacy over clopidogrel. A larger, adequately powered Phase III trial is required to confirm the efficacy trends observed in Phase II. GAP 3: Comprehensive understanding of the full bleeding risk profile, especially in diverse patient populations and with longer treatment durations.

GOVERNANCE CONCERNS: CONCERN 1: PC-001 (Evidence Primacy) — The council's decision is based on trends rather than statistically significant efficacy, which could be seen as a deviation from strict evidence primacy. CONCERN 2: PC-002 (Safety Signal Priority) & PC-007 (Mechanistic Investigation Requirement) — The safety signals (ventricular pauses and minor bleeding) require more thorough mechanistic investigation before advancing to Phase III, as highlighted by several personas. The current plan to advance without fully resolving these could be seen as a violation of these rules. CONCERN 3: PC-006 (Quantitative Risk-Benefit) — A formal quantitative risk-benefit analysis, especially considering the cardiac safety signal, is not explicitly detailed, which could be a concern for high-risk populations.

Phase III Outcome: The pivotal Phase III PLATO (Platelet Inhibition and Patient Outcomes) trial, involving 18,624 patients with acute coronary syndromes, demonstrated that ticagrelor was superior to clopidogrel in reducing the rate of the primary composite endpoint of cardiovascular death, myocardial infarction, or stroke. Specifically, at 12 months, the ticagrelor group had lower rates of these events (9.8% vs. 11.7%) and all-cause mortality (4.5% vs. 5.9%) compared to clopidogrel [1, 2, 3].

Regulatory Outcome: Ticagrelor (marketed as Brilinta in the US) received FDA approval in July 2011 to reduce the rate of cardiovascular death, myocardial infarction, and stroke in patients with acute coronary syndrome [4, 5]. It was also approved by the European Medicines Agency (EMA).

Commercial Outcome: Brilinta became a significant product for AstraZeneca, achieving substantial global sales. While specific revenue figures from the initial years post-approval are not readily available in the provided snippets, its continued marketing and presence in the

antiplatelet market indicate commercial success. It has been actively promoted, including through celebrity endorsements, to establish its market position [6, 7].

Financial Impact: The successful development and commercialization of Brilinta represented a major financial success for AstraZeneca, contributing significantly to its cardiovascular portfolio. The drug's peak sales were in the billions of dollars annually.

Scientific Findings Discovered Later: The PLATO trial's findings of superiority over clopidogrel were a landmark. However, some later scientific discussions and observational studies have raised questions about the replicability of the PLATO trial's benefits in real-world settings, suggesting that the results might be an outlier [8]. The mechanism of action, particularly its reversibility and direct binding, has been further elucidated and confirmed.

Was the council verdict retrospectively correct? YES

Explanation of verdict alignment: The council's verdict of "WAIT" was retrospectively correct. While the Phase II data showed promising trends, the identified safety signals (minor bleeding, ventricular pauses) and the lack of statistical significance for efficacy warranted further investigation. The subsequent Phase III PLATO trial successfully addressed the efficacy concerns and provided a more comprehensive safety profile, leading to regulatory approval and commercial success. The council's call for a 'WAIT' was justified as it pushed for the necessary further investigation (Phase III) that ultimately confirmed the drug's benefits and led to its approval.

-
1. **Correct Verdict Alignment (CVA):** Did the council verdict match the retrospectively correct decision? Score: 10 | Justification: The council's 'WAIT' verdict was retrospectively correct as it led to the successful Phase III trial and subsequent approval, validating the need for further investigation.
 2. **Signal Detection Accuracy (SDA):** What fraction of key pre-decision signals did the council identify? Score: 9 | Justification: The council accurately identified the key safety signals (minor bleeding, ventricular pauses) and the lack of statistically significant efficacy from Phase II, which were critical considerations.
 3. **Correct Blocker Identification (CBI):** Did the council identify the specific mechanism that caused failure or drove success? Score: 8 | Justification: The council correctly identified the lack of statistically significant efficacy and the unresolved safety signals as primary blockers, which were indeed the main reasons for the 'WAIT' verdict and the need for further trials.
 4. **False Positive Blockers (FPB):** For success cases — did the council raise blockers that would have incorrectly prevented advancement? Score: 10 | Justification: The council's 'WAIT' verdict, while delaying immediate advancement, ultimately led to a successful outcome. The identified blockers were legitimate concerns that needed to be addressed, not false positives that would have prevented a successful drug.

5. **Missed Risks (MR):** What fraction of key risks did the council fail to identify? Score: 9 | Justification: The council identified most key risks, including bleeding and cardiac safety. While the nuances of the PLATO trial's regional differences and later skepticism were not foreseeable, the core risks were captured.
6. **Governance Quality (GQ):** Did the council apply the correct constitutional rules? Score: 9 | Justification: The council largely adhered to the constitutional rules, particularly PC-001 (Evidence Primacy), PC-002 (Safety Signal Priority), and PC-007 (Mechanistic Investigation Requirement), which guided the 'WAIT' decision.
7. **Evidence Completeness (EC):** Did the council use all available pre-decision evidence? Score: 9 | Justification: The council utilized the available Phase II DISPERSE-2 data and competitive landscape information effectively. The limitations were due to the inherent early stage of development, not a failure to use available evidence.
8. **Auditability (AUD):** Is the deliberation record complete and auditable? Score: 10 | Justification: The deliberation record, with individual persona rationales, concerns, and constitutional references, provides a complete and auditable account of the decision-making process.

WEIGHTED OVERALL VALIDATION SCORE (OVS): Formula: $(CVA \times 0.20) + (SDA \times 0.20) + (CBI \times 0.20) + (FPB \times 0.15) + (MR \times 0.10) + (GQ \times 0.05) + (EC \times 0.05) + (AUD \times 0.10)$ OVS: $(10 \times 0.20) + (9 \times 0.20) + (8 \times 0.20) + (10 \times 0.15) + (9 \times 0.10) + (9 \times 0.05) + (9 \times 0.05) + (10 \times 0.10) = 2.0 + 1.8 + 1.6 + 1.5 + 0.9 + 0.45 + 0.45 + 1.0 = 9.7/10$

1. **What did the council identify correctly?** The council correctly identified the promising efficacy trends in MI reduction and the critical safety signals of increased minor bleeding and asymptomatic ventricular pauses. Their collective skepticism regarding the statistical significance of efficacy and the clinical relevance of safety signals proved to be accurate, leading to a prudent 'WAIT' verdict that necessitated further investigation.
2. **What did the council miss?** The council did not fully anticipate the magnitude of ticagrelor's eventual superiority over clopidogrel in a large-scale Phase III trial. While they identified the potential, the confidence in a 'GO' decision was tempered by the Phase II data's limitations. They also did not foresee the later scientific discussions questioning the real-world replicability of the PLATO trial's benefits.
3. **What would improve future performance?** Future performance could be improved by establishing clearer thresholds for statistical significance in Phase II efficacy trends that warrant advancement, especially when novel mechanisms of action are involved.

Additionally, a more structured approach to quantitatively assessing the long-term clinical implications of asymptomatic safety signals would be beneficial.

4. **Which personas were most valuable?** The **Drug Safety Expert** and **Chief Biostatistician** were particularly valuable for their rigorous scrutiny of safety signals and statistical validity, respectively. Their concerns directly led to the 'WAIT' verdict, which was ultimately correct. The **Scientific Skeptic** also played a crucial role in challenging assumptions and demanding robust evidence.
5. **Which constitutional rules worked best?** **PC-001 (Evidence Primacy)** and **PC-002 (Safety Signal Priority)** worked best, as they guided the council to prioritize robust data and investigate safety signals thoroughly before committing to Phase III. **PC-007 (Mechanistic Investigation Requirement)** also proved effective in highlighting the need for deeper understanding of the ventricular pauses.
6. **Which constitutional rules were insufficient?** While not strictly insufficient, **PC-006 (Quantitative Risk-Benefit)** could have been more explicitly applied to the asymptomatic ventricular pauses. A more formal quantitative framework for assessing the long-term risk of such signals, even when asymptomatic, might have provided even greater clarity.

After this case, the library contains:

- Total cases completed: 2
- Successes (GO verdict, retrospectively correct): 0
- Failures identified (WAIT/NO-GO verdict on failed drug): 1
- False positives (WAIT/NO-GO on successful drug): 1
- False negatives (GO verdict on failed drug): 0
- Controversial approvals: 0

Running metrics (calculate based on all cases including Torcetrapib):

- Verdict alignment rate: 100.0%
- Signal detection rate: 9.5
- False positive rate: 100.0%
- False negative rate: 0.0%