

A Modular Systems-Biology Framework for nsPEF-Enabled Extracellular Vesicle Engineering

Journal Title
XX(X):1–24
©The Author(s) 2016
Reprints and permission:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/ToBeAssigned
www.sagepub.com/

SAGE

Arash Ghorbannia^{1,2}, Mihir Mantena³, Ruben M. L. Colunga-Biancatelli^{4,5}

Abstract

Nanosecond pulsed electric fields (nsPEF) can stimulate extracellular vesicle (EV) release through coupled membrane electrodynamics, organelle perturbation, Ca^{2+} redistribution, membrane remodeling, repair, vesicle biogenesis, and stress-dependent cargo sorting. However, these processes span nanosecond electrical events, millisecond-to-second Ca^{2+} and organelle signaling, and minute-to-hour EV release and cargo phenotypes, making whole-cell simulation impractical for early model development. Here we define a modular, cell-type-adaptable systems-biology framework for nsPEF-enabled EV engineering. The framework organizes the mechanism into linked layers: pulse delivery and dosimetry; plasma membrane and organelle electrodynamics; ion transport, Ca^{2+} mobilization, ROS, and bioenergetics; Ca^{2+} -dependent remodeling and repair; EV biogenesis and subtype release; cargo sorting, EV composition, and potency; injury, debris, and quality gating; and manufacturing, isolation, and quality-control interfaces. Cell-state and disease-state modifiers act as cross-cutting parameter layers that adapt the shared model to cancer cells, endothelial cells, immune cells, stromal cells, and other EV-producing systems. This structure provides a transparent framework for connecting pulse parameters to EV yield, subtype distribution, cargo state, functional potency, purity, and cell viability.

Keywords

nanosecond pulsed electric fields; electro-exocytosis; extracellular vesicles; EV engineering; systems biology; compartmental modeling; calcium signaling; electroporation; cargo modulation; biomanufacturing

Abbreviations and Nomenclature

Table 1 summarizes the core biological abbreviations used throughout the manuscript. Variable symbols and parameter notation are listed separately in Appendix Table A17.

Introduction

Nanosecond pulsed electric fields (nsPEF) are ultrashort, high-intensity electric pulses capable of perturbing plasma membranes and intracellular membrane-bound organelles. Unlike conventional electroporation, which typically uses microsecond-to-millisecond pulses that primarily charge the plasma membrane, nsPEF can access intracellular membranes because the pulse duration is comparable to, or shorter than, relevant membrane-charging times^{9,99,129}. This physical distinction allows nsPEF to trigger rapid Ca^{2+} mobilization, organelle stress responses, phospholipid redistribution, cytoskeletal remodeling, and regulated or stress-associated vesicle release^{8,68,83}. nsPEF have emerged as a bioelectric modality in cancer and other biomedical applications, and recent experimental evidence suggests that they can also stimulate robust extracellular vesicle (EV) release, a process referred to here as *electro-exocytosis*⁶⁰.

EVs are heterogeneous membrane-bound particles involved in intercellular communication, immune modulation, vascular remodeling, tissue repair, and disease progression. In accordance with MISEV2023 nomenclature principles, this manuscript uses the umbrella term EV when vesicle origin has not been experimentally resolved¹³⁸.

Mechanistic terms such as exosome, microvesicle, and apoptotic body are used when referring to biogenesis-specific model pathways. This distinction is important for modeling because measured EV populations often reflect mixtures of small EVs, medium/large EVs, apoptotic bodies, damaged organellar fragments, protein aggregates, and other co-isolated materials. A useful nsPEF-EV model must therefore predict not only total particle release, but also subtype weighting, cargo state, functional potency, purity, and the boundary between controlled EV release and injury-associated debris.

At the cellular level, nsPEF stimulation initiates a cascade that bridges membrane electrodynamics with vesicle biogenesis. Rapid transmembrane-potential changes and nanopore

¹Health, Education, AI and Learning, Office of Enterprise Research and Innovation, Old Dominion University, Norfolk, VA 23508, USA

²National Center for Collaboration in Medical Modeling and Simulation, Macon and Joan Brock Virginia Health Sciences at Old Dominion University, Norfolk, VA 23507, USA

³Eastern Virginia Medical Center, Old Dominion University, Norfolk, VA 23508, USA

⁴Frank Reidy Research Center for Bioelectrics, Old Dominion University, Norfolk, VA 23508, USA

⁵Division of Pulmonary and Critical Care Medicine, Department of Medicine, School of Medicine, Macon & Joan Brock Virginia Health Sciences at Old Dominion University, Norfolk, VA 23507, USA

Corresponding author:

Arash Ghorbannia, Old Dominion University, Norfolk, VA, USA.

Email: [Corresponding author email to be added]

Table 1. Core abbreviations used throughout the manuscript.

Abbr.	Definition
nsPEF	nanosecond pulsed electric field
EV	extracellular vesicle
sEV	small extracellular vesicle
m/IEV	medium/large extracellular vesicle
AB	apoptotic body
MVB	multivesicular body
ILV	intraluminal vesicle
ER	endoplasmic reticulum
ROS	reactive oxygen species
ATP	adenosine triphosphate
PS	phosphatidylserine
ESCRT	endosomal sorting complex required for transport
QC	quality control

formation enable ionic fluxes, most notably Ca^{2+} influx and redistribution, with patch-clamp and imaging studies showing minute-scale permeability recovery and Ca^{2+} thresholds that depend on pulse width and cell type^{74,80,81,99}. Intracellular membranes such as the ER, mitochondria, endosomes, and MVBs may also be perturbed, creating routes for store release, mitochondrial stress, ROS generation, trafficking changes, and cargo-sorting modulation^{8,71,100}. Local Ca^{2+} signals then activate effectors including annexins, calpain, scramblases, phospholipases, synaptotagmin-like sensors, and SNARE-associated fusion machinery. These signals drive phosphatidylserine externalization, cytoskeletal reorganization, membrane curvature, MVB docking and fusion, microvesicle budding, and membrane repair^{12,68,69,140}.

From a computational systems-biology perspective, nsPEF-induced EV engineering is best formulated as a modular pathway-level perturbation rather than as a complete whole-cell simulation. In this framework, the electrical pulse is represented as a controllable external input that modifies plasma membrane and organelle permeability, producing time-dependent Ca^{2+} , ROS, ATP, repair, remodeling, trafficking, and injury states. These upstream variables are then mapped to EV release rates, subtype weighting, cargo composition, functional potency, and quality-control outputs. Such a structure preserves mechanistic interpretability while allowing the same core model to be instantiated in cancer cells, immune cells, endothelial cells, mesenchymal stromal cells, or isolated EVs.

The present manuscript therefore focuses on the modeling problem: how to connect an externally defined nsPEF exposure to membrane and organelle perturbation, Ca^{2+} -dependent remodeling, repair, EV biogenesis, cargo modulation, injury-quality gating, and manufacturing-facing EV outputs through a transparent and literature-curatable computational framework.

Methods

We formulated nsPEF-enabled EV engineering as a modular systems-biology framework that links pulse exposure to membrane and organelle perturbation, Ca^{2+} signaling, membrane remodeling, repair, vesicle release, cargo state, injury-quality gating, and manufacturing-facing outputs. Each biological process is represented as a model module with defined inputs, state variables, outputs, and

candidate mathematical forms. Paired appendix tables [A1–A16](#) summarize biological evidence and computational implementation options.

- (i) Define the minimal set of biological and engineering layers required to connect nsPEF exposure to EV yield, subtype distribution, cargo state, potency, purity, and viability.
- (ii) Assign each layer a computational representation appropriate to its dominant timescale, state variables, measurable outputs, and available modeling literature.
- (iii) Separate biological evidence curation from computational model curation for each module.
- (iv) Distinguish cell-based electro-exocytosis from direct post-isolation EV engineering.
- (v) Represent cell type, disease state, activation state, and population heterogeneity as cross-cutting modifiers rather than as a downstream pathway.

Results

The resulting framework is a cell-type-adaptable module map rather than a single parameterized model. [Table 2](#) provides the concise navigation map. [Table 3](#) defines the mathematical modeling strategy and module-interface logic. The appendix tables then provide the biological and computational evidence-curation structure needed for future model implementation and parameterization.

Preliminary results

To demonstrate how the current implementation behaves before formal calibration, we generated a set of targeted example simulations aligned to the first five layers of the framework. These examples are not intended as validation studies. Rather, they show that the selected mathematical forms already produce interpretable regime structure: dosimetry responds to conductivity and pulse-train timing, electrodynamics responds to pulse width and cell size, compartmental ion-transport equations generate coupled Ca^{2+} /ROS/ATP trajectories, remodeling rules separate buffering from repair-amplification effects, and the EV-release pool model distinguishes productive secretory windows from injury-dominant vesiculation.

The dosimetry example therefore illustrates the value of a selectable Layer 1 representation rather than a single absorbed-energy scalar. The legacy, adiabatic, and lumped-retention variants produce similar exposure ordering but different thermal interpretations, which is exactly the modeling role intended in [Table A2](#). The electrodynamics example then shows the consequence of the Layer 2 reduced charging model: pulse width is not merely a protocol label, but an explicit determinant of how close the membrane gets to its Schwan-limit voltage during each pulse. This matters downstream because the same voltage state drives the permeability and pore-density surrogates that feed the Ca^{2+} and organelle modules.

The Layer 3 example is especially useful because it shows how the compartmental flux-balance formulation behaves under qualitatively different assumptions. Increasing pulse severity amplifies cytosolic Ca^{2+} , ROS, and downstream stress, whereas calcium-limited conditions blunt the Ca^{2+}




Figure 1. Integrated mechanisms linking nanosecond pulsed electric field stimulation to extracellular vesicle (EV) engineering. Nanosecond pulsed electric fields induce membrane and organelle perturbation, Ca^{2+} redistribution, phosphatidylserine externalization, cytoskeletal remodeling, and repair activation. These processes can promote EV release through endosomal/MVB fusion, plasma membrane budding, and injury-associated vesiculation. In an EV-engineering framework, model outputs include EV yield, size/subtype distribution, cargo composition, functional potency, purity, and cell viability.

Abbreviations: EV, extracellular vesicle; ILV, intraluminal vesicle; MVB, multivesicular body; ESCRT, endosomal sorting complex required for transport; VGCC, voltage-gated calcium channel; PLD2, phospholipase D2; ARF6, ADP-ribosylation factor 6; SNARE, soluble N-ethylmaleimide-sensitive factor attachment protein receptor; LAMP1, lysosome-associated membrane protein 1; HIF-1 α , hypoxia-inducible factor 1- α ; IP $_3$ R, inositol 1,4,5-trisphosphate receptor; PS, phosphatidylserine.




Figure 2. Preliminary Layer 1–2 example behavior. The dosimetry comparison shows how the lumped thermal framework separates absorbed electrical dose from retained heating: low-conductivity conditions remain weakly heated, while high-conductivity and high-repetition-rate pulse trains accumulate larger temperature rises even under similar nominal amplitudes.




Figure 3. Preliminary Layer 1–2 example behavior. The reduced electrodynamics model shows first-order membrane charging toward the Schwan limit, so longer pulses and larger cells reach higher induced membrane voltages and cross a permeability-relevant threshold more readily than ultrashort or smaller-cell cases.

response even when the same electrodynamics trigger is applied. Slowing ion recovery extends osmotic imbalance, which in turn sustains damage pressure despite a similar initial poration event. These behaviors are consistent with the purpose of Table A6: the module is meant to translate a brief electroporation event into longer-lived signaling, organelle, and metabolic trajectories.

The Layer 4 example then shows why the remodeling layer is not redundant with Layer 3. In the present Hill-activation and repair-state framework, buffered microdomains reduce submembrane Ca^{2+} amplification and therefore suppress phosphatidylserine exposure and downstream remodeling. Calpain inhibition preferentially reduces




Figure 4. Preliminary Layer 3–4 example behavior. The compartmental ion-transport/bioenergetics example compares baseline, higher-dose, calcium-limited, and slow-recovery settings; the chosen ODE framework converts the electroporation state into different Ca^{2+} transients, osmotic stress histories, mitochondrial stress, and ROS accumulation.




Figure 5. Preliminary Layer 3–4 example behavior. The remodeling/repair example shows how the reduced Ca^{2+} -sensor module separates phosphatidylserine exposure, actin disruption, and resealing competence across buffered, calpain-inhibited, and annexin/lysosome-supported regimes.

actin disruption without removing the upstream Ca^{2+} transient itself, while stronger annexin/lysosomal support preserves a larger resealing state over time. This is the intended function of Table A8: to resolve whether a given Ca^{2+} /stress history is interpreted as recoverable remodeling or escalating injury.

The Layer 5 example is particularly important because it demonstrates the value of the explicit MVB/ILV/docking/fusion/budding state structure. Rab/ESCRT limitation reduces small-EV release without requiring a generic global suppression of all vesiculation, whereas a ceramide-favored secretory bias increases the release-competent MVB route by reducing lysosomal routing. In contrast, the injury-shift scenario sharply increases apoptotic-body production and collapses viability. This is exactly the interpretive gain intended by the Table A10 framework: subtype release becomes a competition between secretory trafficking, membrane budding, and apoptotic disassembly rather than a single EV-rate output.

The integrated storyboard provides a compact narrative view of the framework. In the mild reversible window, the model predicts threshold-level charging, modest temperature




Figure 6. Preliminary Layer 5 example behavior. The EV-biogenesis comparison shows that the pool-balance release model distinguishes Rab/ESCRT-limited, ceramide-favored secretory, and injury-shifted regimes. In this framework, MVB pool size and secretory bias respond differently from final apoptotic-body output, allowing the model to separate productive small-EV routing from high-stress vesiculation.

Abbreviations: EV, extracellular vesicle; MVB, multivesicular body; ESCRT, endosomal sorting complex required for transport.

rise, limited Ca^{2+} amplification, and predominantly repair-compatible remodeling with low apoptotic output. In the productive secretory window, larger Ca^{2+} loading and stronger repair support coexist with a more favorable secretory routing bias, increasing small-EV output without a catastrophic loss of viability. In the injury-dominant window, the same layered structure yields a different interpretation: stronger charging and thermal load amplify Ca^{2+} and ROS, repair remains incomplete, and apoptotic vesiculation becomes a dominant endpoint. This row-wise transition is encouraging because it shows that the current framework already produces coherent scenario-level behavior while remaining modular enough for later biological calibration.

The Layer 6–8 comparison illustrates why the implementation now carries cargo, quality, and manufacturing states beyond total particle release. Productive and direct-loading scenarios can yield similar process objectives through different mechanisms—endogenous cargo sorting versus permeability-assisted loading—whereas the injury-dominant scenario shows that a high particle output is not automatically a useful EV product. This behavior follows the report’s computational tables: subtype-weighted cargo and stress sorting feed potency, damage and contaminant mixtures determine the quality gate, and recovery, purity, batch consistency, and potency jointly determine the manufacturing objective.

Discussion

The proposed structure shifts the conceptual endpoint of electro-exocytosis modeling from total EV release alone to nsPEF-enabled EV engineering. In this formulation, EV output is not a single scalar quantity but a multi-dimensional product state that includes particle yield, subtype weighting, cargo composition, potency, purity, and producer-cell viability. This distinction is essential because nsPEF can plausibly increase bona fide EV release, alter cargo sorting, stimulate repair-associated shedding, or generate injury-associated particles depending on pulse dose,





Figure 7. Integrated preliminary storyboard across Layers 1–5. Each row represents one illustrative scenario and each column summarizes one model layer from dosimetry through EV output. The top row shows a mild reversible window, the middle row a more productive secretory window, and the bottom row an injury-dominant window. The figure makes the multiscale logic of the framework visible: stronger Layer 1 dose and Layer 2 charging drive larger Layer 3 Ca^{2+} /ROS perturbations, which then reshape Layer 4 repair states and finally alter Layer 5 subtype weighting and viability.

Abbreviations: Ca^{2+} , calcium ion; ROS, reactive oxygen species; ATP, adenosine triphosphate; EV, extracellular vesicle; sEV, small extracellular vesicle; AB, apoptotic body.



Abbreviations: EV, extracellular vesicle.

Figure 8. Preliminary Layer 6–8 downstream engineering behavior. The downstream comparison extends the preliminary examples from vesicle release into product interpretation. A productive cargo window increases protein, RNA, lipid, and antigen cargo scores through secretory and ESCRT-linked terms; a direct-loading mode adds a permeability-driven cargo term while retaining acceptable viability; and an injury-dominant condition may still produce particles but fails the quality gate because apoptosis, necrosis, and contaminant fractions reduce bona fide EV purity and process objective.

Abbreviations: EV, extracellular vesicle; RNA, ribonucleic acid; ESCRT, endosomal sorting complex required for transport.

cell state, and recovery kinetics. A systems model that does not separate these outputs risks conflating useful EV production with nonspecific stress debris.

A second key feature of the framework is the explicit treatment of organelle electrodynamics. nsPEF effects on the ER, mitochondria, endosomes, and MVBs provide a mechanistic route by which pulse parameters may influence not only Ca^{2+} dynamics but also exosome-biogenesis machinery, cargo sorting, lysosomal routing, and mitochondrial stress. These processes are difficult to resolve experimentally in a single study, but modular modeling allows them to be represented as testable submodels with defined inputs and outputs.

Finally, the framework distinguishes two related but separate applications. In cell-based electro-exocytosis, nsPEF is applied to producer cells to stimulate vesicle release and modify cargo through endogenous signaling and sorting pathways. In direct EV engineering, nsPEF is applied to isolated EVs to transiently permeabilize vesicle membranes for cargo loading or surface modification. These modes share membrane-electroporation principles but require different state variables, size scales, and quality-control outputs. Separating them prevents overextension of a single model while preserving a shared bioelectric modeling backbone.

Appendix

Appendix organization and model-curation strategy

The appendix is organized as paired biological and computational tables for each modeling layer. The biological tables document the experimental and mechanistic rationale for including each submodule. The computational tables document candidate mathematical representations, variables, inputs, outputs, and parameterization strategies. Evidence confidence is determined by literature review using the following working categories: established, supported, hypothesized, or model assumption. Table pairs are referenced locally in each module subsection so that biological claims and candidate model forms can be traced back to their evidence and implementation tables.

Pulse Delivery, Exposure Geometry, and Dosimetry

Pulse delivery, exposure geometry, and dosimetry define the engineered exposure that initiates electro-exocytosis. In this layer, the pulse waveform denotes the externally imposed electric-field history, including pulse amplitude, duration, number, polarity, and repetition rate. Early nsPEF studies show that these variables set the transition from plasma-membrane poration to intracellular Ca^{2+} mobilization, PS exposure, and cell-fate responses^{9,81,99,129}. Exposure geometry denotes the physical arrangement of electrodes, chamber, cuvette, culture format, or flow path that converts a programmed voltage waveform into the electric field experienced by the sample. Because cells may receive different effective exposures depending on electrode gap, edge effects, medium conductivity, cell density, sample volume, and position within the device, geometry is central to reproducibility^{5,42,46}. A nominal pulse

protocol is therefore not sufficient unless the exposure configuration, field uniformity, and measurement method are also defined^{52,91}. The biological/process evidence for these exposure variables is summarized in Table A1, and the corresponding computational abstractions are listed in Table A2.

For the present compartmental modeling framework, the spatial field calculation should be treated as an upstream dosimetry and calibration step rather than as a spatially resolved biological state. Device settings and exposure geometry can be collapsed into lumped descriptors such as an effective field amplitude, a cumulative pulse-dose index, a heating constraint or temperature history, an exposure-uniformity factor, and the recovery or harvest interval. These descriptors define controllable engineering inputs and batch-to-batch exposure conditions for the downstream ODE modules, while avoiding the need to track chamber position as a dynamic state variable. This separation is also biologically important: energy deposition and Joule heating must be monitored so that field-driven signaling can be distinguished from thermal artifacts, and the recovery interval must be specified because EV yield, cargo composition, and viability are measured over minutes to hours after pulsing rather than at the instant of exposure. Recent nsPEF spheroid work explicitly links pulse width, repetition rate, buffer conductivity, absorbed energy density, and measured temperature rise, showing that high-conductivity media and compressed pulse trains can produce substantially different thermal histories even when nominal pulse amplitudes are similar⁷⁶. Related cancer-ablation studies emphasize the same practical distinction between absorbed electrical dose and heat retained during the treatment interval³². Figure A1 summarizes this interface between device-level pulse delivery and reduced model inputs.

We therefore use a tiered dosimetry strategy in which increasingly detailed pulse-energy models can be selected according to the available calibration data. The simplest baseline preserves a legacy absorbed-energy descriptor,

$$u_{\text{legacy}} = \sigma E_{\text{peak}}^2 t_p N, \quad (\text{A1})$$

where σ is the medium conductivity, E_{peak} is the nominal peak electric field, t_p is pulse width, and N is pulse number. This expression is useful as a transparent dose index and maintains continuity with early exploratory simulations, but it does not distinguish square, bipolar, or exponentially decaying waveforms and treats geometry correction only as a coarse scalar.

A waveform-aware Joule model retains the same physical skeleton but weights the pulse by the normalized squared waveform integral,

$$\chi_w = \frac{1}{t_p} \int_0^{t_p} \left(\frac{E(t)}{E_{\text{peak}}} \right)^2 dt, \quad (\text{A2})$$

$$u_J = \sigma E_{\text{peak}}^2 t_p N \chi_w \langle g_E^2 \rangle, \quad (\text{A3})$$

$$\Delta T_{\text{ad}} = \frac{u_J}{C_v}. \quad (\text{A4})$$

Here χ_w captures waveform shape, $\langle g_E^2 \rangle$ represents the geometry-averaged squared-field factor, and C_v is the volumetric heat capacity. This adiabatic estimate is

Table 2. Concise module map linking nsPEF-enabled EV engineering biology to appendix model-specification tables.

Layer	Module	Model interface/output	Biology table	Computational table
1	Pulse delivery, exposure geometry, and dosimetry	$E(t)$, pulse width, pulse number, repetition rate, energy, temperature	Table A1	Table A2
2	Plasma membrane and organelle electrodynamics	ΔV_m , ΔV_{ER} , ΔV_{mito} , ΔV_{MVB} , $P_i(t)$	Table A3	Table A4
3	Ion transport, Ca^{2+} mobilization, ROS, and bioenergetics	J_{Ca} , C_i , C_{ER} , C_{mito} , ROS, ATP	Table A5	Table A6
4	Ca^{2+} -dependent remodeling and repair	PS exposure, calpain, annexins, actin state, resealing	Table A7	Table A8
5	EV biogenesis and subtype release	R_{sEV} , $R_{m/lEV}$, R_{AB} , subtype weighting	Table A9	Table A10
6	Cargo sorting, EV composition, and potency	protein/RNA/lipid cargo, antigen enrichment, functional potency	Table A11	Table A12
7	Injury, debris, and quality gate	viability, apoptosis, necrosis, debris fraction, contamination markers	Table A13	Table A14
8	Manufacturing, isolation, and QC interface	yield, recovery, purity, batch consistency, scalability	Table A15	Table A16
Cross-cutting	Cell-state and disease modifiers	parameter scalars, phenotype priors, heterogeneity	Tables A1–A15	Tables A2–A16

Abbreviations: nsPEF, nanosecond pulsed electric field; EV, extracellular vesicle; sEV, small extracellular vesicle; m/lEV, medium/large extracellular vesicle; AB, apoptotic body; PS, phosphatidylserine; ROS, reactive oxygen species; QC, quality control.

Table 3. Modeling framework and module-interface logic for the integrated nsPEF-EV engineering model.

Module	Mathematical representation	Inputs	Outputs	Rationale
Pulse delivery and dosimetry	External forcing, exposure geometry, energy/thermal constraints	pulse protocol, chamber geometry, conductivity, cell density	$E(t)$, dose metrics, temperature	Defines controllable engineering input and batch-to-batch exposure.
Membrane and organelle electrodynamics	Circuit model plus pore-density/pore-radius kinetics	$E(t)$, geometry, membrane properties, organelle size	ΔV_m , ΔV_o , G_m , P_i	Converts nsPEF exposure into plasma-membrane and organelle permeability.
Ion transport, Ca^{2+} , ROS, ATP	Compartmental flux-balance ODEs with optional GHK/Nernst–Planck fluxes	permeability, ion gradients, pumps, channels, stores	C_i , C_{sub} , C_{ER} , C_{mito} , ROS, ATP	Links electroporation to signaling, organelle stress, and recovery capacity.
Remodeling and repair	Reaction–diffusion, buffer kinetics, Hill activation, repair ODEs	Ca^{2+} transient, ROS, membrane damage	effector activation, PS exposure, actin state, resealing	Defines reversible remodeling, budding competence, and membrane recovery.
EV release	Pool-balance and Ca^{2+} -dependent rate models	MVB pool, budding sites, Ca^{2+} , PS, cytoskeletal state	R_{sEV} , $R_{m/lEV}$, R_{AB}	Maps molecular states to size/subtype-resolved vesicle release.
Cargo and potency	Cargo-sorting weights, reaction networks, or phenotype-weighted output models	ESCRT, Rab, tetraspanins, stress, state, EV subtype	protein/RNA/lipid cargo, potency score	Captures EV engineering beyond yield alone.
Injury and quality gate	Threshold rules, hybrid state transitions, contamination models	residual conductance, Ca^{2+} , ATP, ROS, caspases	viable/stressed/apoptotic fractions, debris burden	Separates useful EV release from injury-associated particles.
Manufacturing and QC	Process-yield balances, recovery models, mixed-effects batch models	harvest time, isolation method, QC markers, scale	yield, recovery, purity, reproducibility	Connects cell-level biology to translational EV production.
Cell-state modifiers	Parameter scaling, priors, mixed-effects or ensemble models	cell type, disease state, activation, stiffness, lipid state	module-specific sensitivity factors	Adapts the shared model to different producer cells and disease contexts.

Note: The table defines modeling form at the module level. Appendix tables provide submodule-specific biological evidence and computational implementation options.

Abbreviations: nsPEF, nanosecond pulsed electric field; EV, extracellular vesicle; MVB, multivesicular body; ESCRT, endosomal sorting complex required for transport; PS, phosphatidylserine; ROS, reactive oxygen species; ATP, adenosine triphosphate; QC, quality control; GHK, Goldman-Hodgkin-Katz.

intentionally conservative: it represents the temperature rise that would occur if deposited energy remained locally

retained. It is appropriate for ranking protocols by electrical energy deposition, but it should not be interpreted as

a measured sample temperature when heat diffusion, convection, electrode heat exchange, or long inter-pulse intervals are relevant.

When temperature history is needed, the adiabatic estimate can be coupled to a reduced thermal-retention model,

$$\Delta T_{\text{end}} = \eta \Delta T_{\text{ad}} \frac{\tau_{\text{loss}}}{T_{\text{train}}} \left(1 - e^{-T_{\text{train}}/\tau_{\text{loss}}}\right), \quad (\text{A5})$$

where $T_{\text{train}} \approx N/f_{\text{rep}}$ is the pulse-train duration, τ_{loss} is an empirical thermal relaxation time, and η is a setup-dependent retention factor. This reduced form is not a substitute for finite-element thermal modeling, but it captures the main experimental observation that identical absorbed energy can lead to different end-of-train temperature rises when pulse repetition rate, train duration, buffer conductivity, and exposure geometry differ^{32,76}. Because temperature can itself modulate membrane fluidity, permeabilization thresholds, and cell fate, thermal terms should be reported as independent dosimetry outputs rather than folded silently into an electrical dose scalar^{43,83,112,113,144}.

Plasma Membrane and Organelle Electrodynamics

A central goal of the framework is to connect nsPEF exposure to downstream Ca^{2+} -dependent EV release and cargo modulation without requiring a complete whole-cell reconstruction. The membrane and organelle-electrodynamics layer therefore serves as the upstream bioelectric perturbation module. The applied pulse induces transient plasma-membrane polarization, electropore formation, increased conductance, and ion-specific permeability, all of which have been observed or modeled in single-cell electroporation studies^{45,47,80}. Short pulses can also couple to intracellular membranes, supporting reduced organelle-voltage terms for ER, mitochondrial, endosomal, and MVB-related perturbation^{5,8,99,129}. These outputs provide the primary interface between nsPEF exposure and downstream Ca^{2+} , repair, exocytosis, EV-biogenesis, and cargo-sorting modules.

Computational electroporation literature clarifies how much of this upstream module can remain reduced while still being mechanistically anchored. Molecular dynamics studies show field-driven water defects progressing into hydrophilic pores on nanosecond time scales, with pore formation, water-channel growth, voltage removal, and resealing all occurring as distinct processes^{118,120}. Vesicle electromechanics adds the continuum constraint that breakdown voltage depends on membrane tension, area expansivity, and lipid composition⁷³. Together, these papers support the present choice to keep pore nucleation, pore-radius/tension effects, permeability, and resealing as separable calibrated states rather than one binary permeabilization switch. Table A3 summarizes the biological electrodynamics evidence, and Table A4 lists the reduced computational forms used for this layer.

The current implementation uses a reduced electrodynamics ladder rather than a full spatial pore-kinetic simulation. Plasma-membrane polarization is represented by a Schwan-style limiting voltage scaled by the effective field, cell radius, and a cell-state membrane modifier, with a first-order

charging correction for finite pulse width:

$$\begin{aligned} \Delta V_m^\infty &= f_s E_{\text{eff}} a M_m, \\ \Delta V_m(t_p) &= \Delta V_m^\infty (1 - \exp[-t_p/\tau_m]). \end{aligned}$$

where f_s is the Schwan geometry factor, E_{eff} is the geometry-corrected electric field, a is the configurable cell radius, M_m is the membrane-state modifier, t_p is pulse width, and τ_m is an effective membrane charging time^{42,45,46}. This formulation resolves the previous fixed-radius placeholder by making ΔV_m explicitly cell-size dependent; larger cells reach higher induced membrane voltages at the same field, while very short pulses may terminate before the membrane reaches its Schwan limit.

Electroporation is then represented by reduced permeability and pore-density closures:

$$\begin{aligned} P_m &= [1 + \exp[-s(\Delta V_m - V_{\text{th}})]]^{-1}, \\ N_p &= N_{\text{max}} \frac{\Delta V_m^n}{V_{1/2}^n + \Delta V_m^n}. \end{aligned}$$

These terms are phenomenological surrogates for voltage-dependent pore creation, pore growth, conductance, and resealing models^{47,127}. Patch-clamp studies provide direct calibration targets because nsPEF-induced conductance can persist for minutes and recover in a cell-type-dependent manner^{74,80,81}. The reduced pore-density variable should therefore be interpreted as an effective permeability state, not an absolute physical pore count, until it is calibrated against conductance, dye uptake, ion flux, or recovery data. Organelle perturbation is currently represented by configurable fractions of the plasma-membrane voltage, $\Delta V_o = \alpha_o \Delta V_m$ for ER, mitochondria, and MVB/endosomal membranes. This preserves an explicit route from nsPEF exposure to intracellular membrane stress while marking the organelle layer as calibration-dependent until multi-shell, geometry-resolved, or organelle-specific simulations are incorporated^{5,8,99}. The same pore state also feeds the lipid-remodeling layer because experiments and MD simulations connect nanosecond-pulse nanopores with voltage-driven phosphatidylserine translocation^{130,131}.

Ion Transport, Ca^{2+} Mobilization, ROS, and Bioenergetics

The ion-transport and organelle-state module translates nsPEF-induced membrane permeability into intracellular ionic, metabolic, and stress responses. The minimal model treats electropore-mediated Ca^{2+} influx and ER release as source terms in a cytosolic Ca^{2+} balance. More detailed formulations include voltage-gated or leak-channel entry, ER release through IP_3 receptors or ryanodine receptors, SERCA-mediated refilling, PMCA and NCX extrusion, mitochondrial uptake and release, ROS generation, ATP dependence, and osmotic recovery. Table A5 organizes the biological evidence for these submodules, while Table A6 lists the corresponding reduced flux-balance and organelle-state model forms.

For the current computational framework, this layer is represented as a reduced flux-balance ODE system rather than a full electrodiffusion model. The pore term is treated as a transient extracellular Ca^{2+} source gated by the




Figure A1. Conceptual interface for pulse delivery, exposure geometry, and dosimetry. The programmed nsPEF waveform and physical exposure configuration determine the actual sample exposure. In a reduced compartmental framework, spatial field information is used for upstream calibration and then summarized as lumped descriptors such as effective field amplitude, cumulative pulse dose, heating constraint, exposure-uniformity factor, and recovery interval. These descriptors provide reproducible inputs to downstream biological ODE modules without requiring a spatially resolved core model.

Abbreviations: nsPEF, nanosecond pulsed electric field; ODE, ordinary differential equation.

Table A1. Biological evidence table for pulse delivery, exposure geometry, and dosimetry.

Submodule	Biological/process phenomenon	Evidence to curate	Relevance to EV engineering	Confidence
Pulse waveform	nsPEF duration, amplitude, number, repetition rate	Parameter-response studies across pulse width, field strength, repetition, and pulse number	Defines controllable input space for EV yield and cargo modulation	Established ^{9,76,81,99,129}
Exposure geometry	Electrode/chamber/cuvette/geometry	Field uniformity, edge effects, cell density, medium conductivity	Controls reproducibility and batch exposure	Supported; implementation remains lumped ^{5,42,46,52,76}
Energy and heating	Joule heating and thermal artifact risk	Absorbed energy density, waveform energy factor, temperature traces, duty-cycle and repetition-rate effects	Separates bioelectric effects from heat-induced stress	Supported; setup-specific calibration needed ^{32,43,76,112,113}
Recovery interval	Post-pulse harvest window	EV release and repair kinetics over seconds to hours	Determines when yield, cargo, and viability are measured	Supported ^{12,68,104,115,140}
Scale-up mode	Static versus flow-through treatment	Flow exposure, residence time, process monitoring, and sterility constraints	Connects mechanism to manufacturing implementation	Hypothesized ^{6,13,30,53,56}

Abbreviations: nsPEF, nanosecond pulsed electric field; EV, extracellular vesicle.

Table A2. Computational framework table for pulse delivery, exposure geometry, and dosimetry.

Submodule	Model form	Representative relation	Inputs	Outputs
Pulse input	External forcing function ^{9,81,99,129}	$E(t) = E_0 u(t; t_p, N, f)$	amplitude, width, number, repetition	$E(t)$
Dose metric	Selectable absorbed-energy model ^{32,76}	$u = \sigma E_{\text{peak}}^2 t_p N \chi_w \langle g_E^2 \rangle$	pulse train, conductivity, waveform, geometry	energy density, dose index
Field distribution	Laplace/quasi-static field model ^{5,42,46,52}	$\nabla \cdot (\sigma \nabla \phi) = 0, E = -\nabla \phi$	chamber geometry, conductivity	local $E(\mathbf{x}, t)$
Thermal constraint	Adiabatic or lumped heat-loss model ^{32,43,76,112,113}	$\Delta T_{\text{end}} = F_{\text{ret}} \Delta T_{\text{ad}}$	field, medium, cooling, pulse train	$T(t), \Delta T_{\text{end}}$
Flow-through exposure	Residence-time model ^{13,30,53,56}	$D_E(\mathbf{x}) = \int_{\tau} E(\mathbf{x}(t))^2 dt$	flow rate, chamber field	exposure distribution

Abbreviations: EV, extracellular vesicle.

electrodynamic permeability state. This choice is supported by studies in which nsPEF rapidly elevates cytosolic Ca^{2+}

and the response depends on extracellular Ca^{2+} , pulse width, and membrane pathway ^{4,129,146}. In CHO cells, a 60-ns pulse

Table A3. Biological evidence table for plasma membrane and organelle electrodynamics.

Submodule	Biological phenomenon	Evidence to curate	Relevance to EV engineering	Confidence
Membrane polarization	Rapid transmembrane-potential rise	Measurements or simulations of ΔV_m under nsPEF exposure	Determines whether local membrane regions reach electroporation range	Established; Schwan/charging reduced implementation 5,42,45,46
Plasma-membrane nanopores	Transient nanoscale membrane defects	Conductance, dye uptake, ion permeability, recovery kinetics	Creates pathways for Ca^{2+} and other ion entry	Established; proxy remains phenomenological 46,47,80,81,127
Organelle electroporation	ER, mitochondrial, nuclear, endosomal, and MVB membrane perturbation	Ca^{2+} release, mitochondrial depolarization, intracellular membrane access	Provides route for store release, ROS, and altered MVB/cargo dynamics	Supported; organelle fractions need calibration 5,8,99,129
Permeability recovery	Conductance decay and resealing	Patch clamp, uptake assays, dye exclusion	Sets duration and magnitude of post-pulse ion flux	Established; kinetics are cell-type dependent 74,80,81,83
Isolated EV permeabilization	Direct nsPEF effect on EV membranes	Vesicle integrity, cargo loading, aggregation, marker retention	Defines second mode for post-isolation EV engineering	Hypothesized; not implemented in the current cell-based layer 27,29,37,44,106

Abbreviations: nsPEF, nanosecond pulsed electric field; ER, endoplasmic reticulum; EV, extracellular vesicle; MVB, multivesicular body.

Table A4. Computational framework table for plasma membrane and organelle electrodynamics.

Submodule	Model form	Representative relation	Inputs	Outputs
Membrane charging	Reduced Schwan/circuit model 42,45,46	$\Delta V_m = f_s E_{\text{eff}} a (1 - e^{-t_p/\tau_m})$	$E(t)$, geometry, cell radius, membrane state	ΔV_m
Organelle charging	Reduced multi-shell surrogate 5,8,99	$\Delta V_o = \alpha_o \Delta V_m$	organelle factors, plasma-membrane voltage	ΔV_o
Pore nucleation	Voltage-dependent permeability proxy 118,120	$P_m = [1 + \exp[-s(\Delta V_m - V_{\text{th}})]]^{-1}$	ΔV_m , threshold parameters	P_m
Pore-radius evolution	Pore-energy or reduced kinetic model 73,118	$\frac{dr_p}{dt} = F(r_p, \Delta V, \gamma, \sigma)$	pore energy, tension	$r_p(t)$
Permeability closure	Saturating pore-density closure 47,108	$N_p = N_{\text{max}} \Delta V_m^n / (V_{1/2}^n + \Delta V_m^n)$	pore-density scale, voltage scale, Hill exponent	N_p, P_i
Isolated EV charging	Vesicle-scale circuit model 37,44,106	$C_{EV} \frac{d\Delta V_{EV}}{dt} + G_{EV} \Delta V_{EV} = f(E, r_{EV})$	EV radius, medium conductivity	$P_{EV}(t)$, integrity state

Abbreviations: ER, endoplasmic reticulum; EV, extracellular vesicle; MVB, multivesicular body.

produced separable external- Ca^{2+} and ER-release thresholds and a critical Ca^{2+} range for amplification, making these data especially useful for tuning reduced Ca^{2+} source terms [99](#).

ER mobilization is modeled as an electrically triggered store-release pathway with SERCA refilling, while $\text{IP}_3\text{R/RyR}$ -mediated amplification is retained as a cell-type-specific extension for excitable or store-sensitive cells [3,100,123,124,135](#). The computational calcium literature gives useful bounds for that extension: Li and Rinzel reduced a detailed IP_3 receptor model to a Hodgkin–Huxley-like form with IP_3 regulation, Ca^{2+} activation, and Ca^{2+} inactivation, while Shannon and coworkers used junctional, subsarcolemmal, and bulk compartments with realistic buffering, reversible SR uptake, $\text{Na}^+ - \text{Ca}^{2+}$ exchange, and load-dependent release [54,101](#). We therefore keep the default ER term compact, but expose channel opening, store content, pump capacity, and exchanger recovery as parameters when cell-type-specific data are available.

Na^+ , K^+ , and Cl^- fluxes are included as lumped electropore/channel-mediated perturbations that relax toward baseline and drive an osmotic-stress state. This abstraction is justified by evidence that nsPEF increases leak conductance, depolarizes membrane potential, alters voltage-gated channels, and changes cell volume [17,81,109,145](#).

Na^+/K^+ -ATPase sensitivity is retained as a recovery modifier because ATP1A1 silencing attenuates membrane disruption after nanosecond pulses [105](#).

Mitochondrial Ca^{2+} uptake/release and depolarization connect cytosolic Ca^{2+} to ROS and ATP availability. nsPEF studies support mitochondrial membrane perturbation and ROS generation, while broader calcium-electroporation data show that Ca^{2+} overload can deplete ATP and shift the cell toward injury [8,31,71,75,84,89](#). On the computational side, mitochondrial modules can be scaled from a minimal Ca^{2+} -handling block with respiration, ATP synthase, proton leak, adenine nucleotide exchange, uniporter, and $\text{Na}^+ - \text{Ca}^{2+}$ exchange [59](#), to a simplified ATP-production model driven by glycolytic and Ca^{2+} inputs [11](#), or to a thermodynamically balanced oxidative-phosphorylation model that conserves mass, charge, and free energy [7](#). ROS is represented with Ca^{2+} -, mitochondrial-, osmotic-, and pulse-dependent source terms plus first-order clearance; this reduced form is consistent with models in which ROS production, scavenging, and ROS-sensitive inner-membrane conductance shape mitochondrial potential, NADH, and redox dynamics [23,70,148](#). ATP is kept as a normalized production-consumption state coupled to mitochondrial potential, pump activity, repair demand, and oxidative stress.

$$\frac{d[\text{Ca}^{2+}]_i}{dt} = J_{\text{Ca,pore}} + J_{\text{VGCC}} + J_{\text{ER,rel}} - J_{\text{SERCA}} - J_{\text{PMCA}} - J_{\text{NCX}} - J_{\text{mito}} - J_{\text{buffer}}. \quad (\text{A6})$$

Ca²⁺-Dependent Remodeling and Repair

This module connects global and submembrane Ca²⁺ signals to local membrane remodeling and repair. Although cytosolic Ca²⁺ may be represented as a well-mixed compartment in reduced models, electro-exocytosis is likely governed by local Ca²⁺ domains near pores, damaged membrane regions, budding sites, repair complexes, and fusion machinery. Reaction–diffusion theory provides a practical reason to distinguish these domains from the bulk pool: mobile and immobile buffers can confine Ca²⁺ elevations over nanometer-to-submicrometer distances, with local gradients forming on submillisecond time scales near an open channel⁷². Ca²⁺-dependent resealing occurs on experimentally measurable time scales, with nano- and microsecond electroporation studies reporting fast and slow repair components that are sensitive to extracellular Ca²⁺¹². Local Ca²⁺ therefore serves not only as an influx variable but also as the organizing signal for lipid asymmetry, cytoskeletal mechanics, curvature, scission, fusion competence, and resealing. The paired evidence and modeling assumptions for these repair and remodeling variables are tabulated in Tables A7 and A8.

For the current computational framework, this layer is implemented as a reduced Ca²⁺-sensor and repair-state module. The upstream ion-transport state is converted into a submembrane Ca²⁺ microdomain proxy amplified by pore activation and osmotic stress, with buffer strength and shell depth treated as tunable parameters rather than fixed geometry. This choice reflects evidence that ESCRT and repair machinery assemble at damaged membrane domains rather than responding only to bulk cytosolic Ca²⁺^{96,102}. Annexin recruitment is represented as a Ca²⁺-dependent membrane-binding and curvature-stabilization state because annexin A4/A6/A7 can support membrane constriction, ESCRT-associated repair, and repair-linked vesicle secretion^{15,69,111,140}. Molecular and biophysical annexin studies then provide the curvature and lipid-composition rationale for keeping annexin effects separate from generic repair activity^{34,67}.

The lipid-remodeling branch maps local Ca²⁺ to scramblase activation, flippase suppression, and PS externalization. This branch is treated as established because nsPEF activates TMEM16F-dependent PS exposure and blebbing, while newer TIRF studies show delayed, spatially punctate PS externalization after 400-ns pulses^{68,104}. The Vernier studies add a second, faster computationally tractable route: nanopore-facilitated, voltage-driven PS movement can occur along electropore walls and appears experimentally as localized, anode-facing PS externalization after nanosecond pulses^{130,131}. The model therefore keeps an immediate pore/voltage PS term separate from slower Ca²⁺–scramblase dynamics. Chemically controlled TMEM16F vesiculation, lipid-asymmetry loss, and Ca²⁺-regulated flippases provide

complementary support for linking Ca²⁺ and PS state to EV-like budding competence^{35,98,134}.

The cytoskeletal branch maps local Ca²⁺ to calpain activity, actin disruption, actomyosin tension, and repair-associated shedding. Recent nsPEF actin and swelling studies are especially useful here because Ca²⁺-dependent actin shrinkage and swelling can be tested directly against the model's actin-disruption and osmotic-stress terms^{36,82,90}. Calpain and ezrin-cleavage studies support the protease component, and membrane-repair studies show that calpain can orchestrate annexin-containing microvesicle release^{18,92,141}.

Finally, lysosomal repair is represented as a Ca²⁺-triggered vesicle-fusion and membrane-resealing support state. Lysosomal exocytosis, extracellular protease release, sphingolipid-dependent EV biology, lysosomal-channel regulation, and ESCRT-mediated lysosome repair support this reduced repair branch^{2,21,88,97,128}. The resulting variables—PS exposure, calpain activity, annexin activity, lysosomal repair, actin state, resealing state, and repair-associated shedding—serve as interpretable coupling terms for medium/large EV budding, membrane recovery, and injury-quality gating.

EV Biogenesis and Subtype Release

The EV-release module converts nsPEF-induced intracellular and membrane responses into measurable vesicle outputs. It integrates endosomal/MVB-derived small-EV release, plasma-membrane budding of medium/large EVs, and apoptotic body or injury-associated vesiculation. The model should retain mechanistic terms such as exosome and microvesicle internally, but report outputs using EV nomenclature compatible with the level of experimental characterization. The biological release mechanisms are organized in Table A9, with candidate pool-balance and rate-law implementations listed in Table A10.

Small-EV production begins with endosome maturation and ILV biogenesis. Recent Rab-conversion studies indicate that Rab5-to-Rab7 transition is not an isolated timing event, but is coordinated with ESCRT-mediated cargo corraling through the Mon1/Ccz1 axis, so maturation and secretory competence should be treated as coupled states rather than as independent switches^{14,77,110}. Within maturing endosomes, canonical ESCRT machinery supplies one major ILV route: ESCRT-0 captures ubiquitinated cargo through Hrs/STAM, ESCRT-I/II concentrate cargo and deform the bud neck, and ESCRT-III/Vps4 completes constriction and scission^{87,116,142}. However, exosome-biased ILV formation is not exclusively canonical. ALIX-ESCRT-III, ceramide, and tetraspanin-rich membrane domains provide selective alternative routes that can redirect cargo and alter whether MVBs are routed toward secretion or lysosomal degradation^{22,25,50,93,132}.

Release of those ILVs as extracellular small EVs then depends on trafficking, docking, and fusion at the cell cortex. Rab27, Rab11, and Rab35 families regulate MVB positioning and membrane access, while Munc13-4 provides a Ca²⁺-responsive trafficking and priming branch that helps generate release-competent MVBs^{38,40,61,65,117}. The terminal fusion step is mediated by cell-context-dependent SNARE assemblies, with syntaxin-4/SNAP-23/VAMP-7 and VAMP5-based complexes now supported by direct exosome-secretion

Table A5. Biological evidence table for ion transport, Ca^{2+} mobilization, ROS, and bioenergetics.

Submodule	Biological phenomenon	Evidence to curate	Relevance to EV engineering	Confidence
Electropore influx	Ca^{2+} Rapid extracellular entry	Ca^{2+} imaging, chelation, Ca^{2+} -free media	Primary trigger for downstream release and repair mechanisms	Established ^{4,12,19,66,146}
ER Ca^{2+} release	Intracellular store mobilization	$\text{IP}_3\text{R/RyR}$ involvement, ER store depletion	Amplifies cytosolic Ca^{2+} transients and fusion signals	Established ^{3,100,123,124,135}
Na^+ , K^+ , Cl^- fluxes	Ionic/osmotic perturbation	Membrane-potential recovery, swelling, ionic imbalance	Modulates volume, tension, and repair state	Supported ^{17,68,105,109,145}
Mitochondrial Ca^{2+}	Uptake/release and depolarization	Mitochondrial Ca^{2+} , $\Delta\Psi_m$, mPTP evidence	Links Ca^{2+} to metabolic stress and fate decisions	Supported ^{8,70,71}
ROS generation	Oxidative stress after nsPEF	ROS probes, antioxidant rescue, mitochondrial source evidence	Modifies EV secretion, cargo, and injury transition	Supported ^{51,75,84,139,148}
ATP/bioenergetics	Energy availability for pumps and trafficking	ATP depletion, pump recovery, metabolic assays	Controls repair, cargo sorting, and EV-release capacity	Hypothesized ^{28,31,89}

Abbreviations: EV, extracellular vesicle; ER, endoplasmic reticulum; IP_3R , inositol 1,4,5-trisphosphate receptor; RyR, ryanodine receptor; ROS, reactive oxygen species; ATP, adenosine triphosphate; mPTP, mitochondrial permeability transition pore.

Table A6. Computational framework table for ion transport, Ca^{2+} mobilization, ROS, and bioenergetics.

Submodule	Model form	Representative relation	Inputs	Outputs
Pore-mediated Ca^{2+} flux	GHK/Ohmic flux ^{118,120}	$J_{\text{Ca,pore}} = P_{\text{Ca}}(t)\Phi_{\text{Ca}}(C_e, C_i, V_m)$	P_{Ca} , gradients, voltage	Ca^{2+} source term
ER release	Store-release ODE ^{54,101}	$J_{\text{ER,rel}} = k_{\text{rel}}O_{\text{IP3R/RyR}}(C_i, C_{\text{ER}})$	channel state, store content	ER-to-cytosol flux
Pump/exchanger recovery	Saturating fluxes ¹⁰¹	$J_{\text{PMCA}} = V_{\text{max}} \frac{C_i^n}{K^n + C_i^n}$	Ca^{2+} , Na^+ , ATP, voltage	Ca^{2+} clearance
Mitochondrial buffering	Coupled uptake/release ODE ^{11,59}	$\frac{dC_m}{dt} = J_{\text{MCU}} - J_{\text{NCLX}} - J_{\text{mPTP}}$	C_i , $\Delta\Psi_m$, ROS	mitochondrial Ca^{2+} state
ROS dynamics	Source-clearance ODE ²³	$\frac{dR}{dt} = k_{\text{ROS}}f(C_m, \Delta\Psi_m) - k_{\text{clear}}R$	mitochondrial state, antioxidants	ROS state variable
ATP state	Production-consumption ODE ^{7,11}	$\frac{dA}{dt} = k_{\text{prod}}f(\Delta\Psi_m) - k_{\text{use}}(J_{\text{pump}} + R_{\text{repair}})$	mitochondria, pumps, repair	ATP availability

Abbreviations: ER, endoplasmic reticulum; GHK, Goldman-Hodgkin-Katz; PMCA, plasma membrane Ca^{2+} ATPase; MCU, mitochondrial calcium uniporter; NCLX, $\text{Na}^+/\text{Ca}^{2+}/\text{Li}^+$ exchanger; ROS, reactive oxygen species; ATP, adenosine triphosphate; mPTP, mitochondrial permeability transition pore.

Table A7. Biological evidence table for Ca^{2+} -dependent remodeling and repair.

Submodule	Biological phenomenon	Evidence to curate	Relevance to EV engineering	Confidence
Ca^{2+} microdomains	Local Ca^{2+} elevation near pores and membranes	High-speed/local Ca^{2+} imaging, buffer sensitivity	Localizes activation to budding, repair, and fusion sites	Supported ^{96,102}
Annexin recruitment	Ca^{2+} -dependent membrane binding	Annexin translocation and repair studies	Links Ca^{2+} to membrane repair and curvature stabilization	Supported ^{15,34,67,69,111}
Scramblase/PS exposure	Loss of lipid asymmetry	Annexin V staining, scramblase/flippase studies	Promotes MV-like budding and EV surface phenotype	Established ^{35,68,98,104,134}
Calpain activation	Ca^{2+} -dependent protease activity	Calpain activation/inhibition, cytoskeletal remodeling	Supports blebbing, scission, and membrane remodeling	Supported ^{18,92,141}
Actomyosin remodeling	Tension, contraction, and scission	RhoA/ROCK, myosin, actin perturbation	Determines budding probability and vesicle detachment	Supported ^{36,82,90}
Lysosomal repair	Ca^{2+} -triggered vesicle fusion	LAMP1 exposure, lysosome fusion	Adds membrane and may release vesicle-like material	Supported ^{2,21,88,97,128}

Abbreviations: EV, extracellular vesicle; PS, phosphatidylserine; ESCRT, endosomal sorting complex required for transport.

studies, and with synaptotagmin-7-like Ca^{2+} sensing providing a plausible fusion-pore stabilization mechanism under strong submembrane Ca^{2+} signals ^{24,57,62,143}. In parallel,

outward budding of medium/large EVs is supported by ARF-, RhoA/ROCK-, phosphatidylserine-, and actomyosin-dependent membrane remodeling, whereas apoptotic bodies

Table A8. Computational framework table for Ca^{2+} -dependent remodeling and repair.

Submodule	Model form	Representative relation	Inputs	Outputs
Ca^{2+} microdomain	Reaction–diffusion shell ODE ^{72,101}	$\frac{\partial C}{\partial t} = D_{Ca} \nabla^2 C + J_{Ca} - R_{buffer} - R_{effector}$	local flux, diffusion, buffers	$C_{sub}(t)$
Generic Ca^{2+} sensor	Hill activation ^{72,96,101}	$A_x = \frac{C_{sub}^n}{K_x^n + C_{sub}^n}$	local Ca^{2+} , sensitivity	activated fraction
PS exposure	Activation plus lipid-state ODE ^{68,130,131}	$\frac{dPS_{out}}{dt} = k_{scr} A_{scr} (1 - PS_{out}) - k_{flip} PS_{out}$	Ca^{2+} , scramblase, flip-pase	PS externalization
Calpain/cytoskeleton	Protease activation and substrate loss ^{18,92,141}	$\frac{dF_{actin}}{dt} = -k_{calp} A_{calp} F_{actin} + k_{poly}$	Ca^{2+} , calpain state	cortical remodeling
Membrane resealing	Conductance-recovery ODE ^{73,118}	$\frac{dG_m}{dt} = -k_{seal} A_{repair} G_m$	residual conductance, repair activity	resealed membrane state
Damaged-domain shedding	Damage-removal rate ⁷³	$R_{shed} = k_{shed} D_m f_{Ca} f_{tension}$	damaged membrane, Ca^{2+} , tension	repair-associated shedding

Abbreviations: PS, phosphatidylserine; ODE, ordinary differential equation.

arise from caspase-linked ROCK1 activation and repetitive bleb fragmentation during higher-injury states ^{26,63,78,85,94,121}. These mechanistic branches justify treating subtype release as a competition between secretory MVB fate, plasma-membrane budding competence, and apoptotic disassembly rather than as one generic EV-rate variable.

For the current computational framework, this layer is implemented as a reduced pool-balance model with explicit MVB, ILV, docked-MVB, budding-precursor, and apoptotic-commitment states. Endosome maturation is represented as a Rab-conversion-supported source term for a secretory MVB pool, ILV loading is split into ESCRT-dependent and ceramide/tetraspanin-assisted branches, and MVB fate is biased between secretion and lysosomal loss by an acidification-sensitive routing term ^{22,77,132}. Small-EV release is then written as a Ca^{2+} - and SNARE-sensitive fusion flux from the docked MVB pool, while medium/large EV release is represented as a phosphatidylserine-, calpain-, and actomyosin-weighted budding/scission process. Apoptotic-body release is modeled separately as a damage- and blebbing-weighted branch so that injury-associated vesiculation is not conflated with productive secretory EV release ^{57,65,85,121}. This reduced formulation matches the current code implementation while preserving a direct path for later replacement with cell-type-specific Rab, SNARE, lipid, or apoptosis submodels as quantitative data become available.

State-Conditioned EV Size–Composition and Observation Bridge

The release states above specify biological routes, whereas particle-sizing assays report operational diameter classes. These descriptions should remain distinct because endosomal, plasma-membrane, and apoptotic pathways can generate overlapping size ranges, and diameter alone does not establish vesicle origin ¹³⁸. We therefore introduce an inter-layer bridge rather than redefining the existing sEV, m/IEV, and AB states. A biological mode q denotes ESCRT-enriched MVB/ILV release, ceramide/tetraspanin-enriched MVB/ILV release, plasma-membrane shedding or repair-associated release, or apoptotic fragmentation. The first two modes sum to the existing R_{sEV} output, the plasma-membrane mode contributes to $R_{m/IEV}$, and the apoptotic mode contributes to R_{AB} . This separation is consistent with

evidence that activation can differentially regulate distinct nanosized populations released by CD4^+ T cells ¹²⁶.

Let $\mathbf{x}(t)$ denote the upstream cell-state vector, d vesicle diameter, and $\mathbf{c}$ the composition vector. The joint product kernel for biological mode q is factorized as

$$K_q(d, \mathbf{c} | \mathbf{x}) = f_q(d | \theta_q[\mathbf{x}]) g_q(\mathbf{c} | d, \mathbf{x}),$$

where f_q is a nonnegative birth-size distribution and g_q represents conditional cargo allocation. Positive right-skewed distributions are appropriate starting points. For MVB-derived small EVs, a dynamic-scaling distribution derived from membrane-domain growth and ILV budding provides a directly implementable candidate; its tail exponent was sensitive to signaling perturbation ⁸⁶. More detailed ESCRT/cargo-crowding membrane models can constrain plausible ILV sizes and coupling signs but introduce unobserved mechanical parameters that are unlikely to be identifiable from sparse longitudinal particle counts ⁵⁵. Broad lognormal or gamma distributions are therefore retained as reduced alternatives for plasma-membrane and apoptotic modes. Their baseline median and geometric standard deviation should be cell- and assay-specific priors rather than universal subtype constants.

State dependence can be introduced parsimoniously through distribution parameters,

$$\log d_{50,q}(t) = \alpha_q + \beta_q^T [\mathbf{z}_q(t) - \mathbf{z}_{q,0}],$$

where $\mathbf{z}_q$ contains only pathway-matched states. Candidate inputs include ESCRT and ceramide/secretory routing for MVB modes; PS exposure, calpain, actomyosin, membrane tension, and repair shedding for plasma-membrane release; and damage, apoptosis, and blebbing for the apoptotic mode. The nested null case $\beta_q = \mathbf{0}$ changes pathway mixture weights while retaining fixed within-pathway size distributions.

For true-diameter bin $b = [d_b^-, d_b^+]$, the per-cell birth source and extracellular stock are

$$S_{qb}(t) = R_q(t) \int_{d_b^-}^{d_b^+} f_q(d | \theta_q[\mathbf{x}(t)]) dd,$$

$$\frac{dN_{qb}}{dt} = N_{\text{cells}} S_{qb}(t) - k_{\text{loss},q}(d_b) N_{qb} + \sum_{b'} T_{bb'} N_{qb'}.$$

This sectional population balance can use many numerical bins while inferring few shared parameters. It also preserves

Table A9. Biological evidence table for EV biogenesis and subtype release.

Submodule	Biological phenomenon	Evidence to curate	Relevance to EV engineering	Confidence
Endosome/MVB maturation	Early-to-late endosome and MVB formation	Rab5/Rab7, endosome maturation, ILV evidence	Defines small-EV-producing compartment pool	Established ^{14,77,110}
ESCRT-dependent ILV formation	Cargo sorting and ILV biogenesis	ESCRT, ALIX, TSG101 perturbation	Generates exosome-enriched precursors	Established ^{50,87,116,142}
ESCRT-independent ILV formation	Ceramide/tetraspanin/lipid-domain sorting	nSmase2, ceramide, CD63/CD81/CD9 evidence	Alternative small-EV biogenesis route	Supported ^{22,25,93}
MVB trafficking/docking	Rab-dependent transport to plasma membrane	Rab27A/B, Rab11, Rab35 evidence	Controls release-ready MVB pool	Established ^{38,40,61,65,117}
MVB fusion	SNARE-mediated fusion with plasma membrane	SNARE and Ca ²⁺ -sensitive fusion evidence	Converts ILVs into secreted small EVs	Supported ^{24,57,62,143}
Plasma membrane budding	MV-like outward budding	ARF6/RhoA/PS/cytoskeleton studies	Produces medium/large EV precursors	Supported ^{26,63,85}
Apoptotic body formation	Blebbing and fragmentation	Caspase/ROCK1/apoptosis evidence	Captures high-dose or injury-associated vesicles	Established ^{78,94,121}

Abbreviations: EV, extracellular vesicle; MVB, multivesicular body; ILV, intraluminal vesicle; ESCRT, endosomal sorting complex required for transport.

Table A10. Computational framework table for EV biogenesis and subtype release.

Submodule	Model form	Representative relation	Inputs	Outputs
MVB pool	Compartmental pool balance ^{14,39,77,110}	$\frac{dM}{dt} = k_{mat}E_{endo} - k_{dock}M - k_{deg}M$	endosome pool, maturation	MVB number
ILV loading	Production-rate law ^{22,25,50,87,116,142}	$\frac{dI}{dt} = k_{ILV}f_{ESCRT}M - k_{release}I$	ESCRT/ceramide state, MVB pool	ILV load
MVB docking	Trafficking/docking ODE ^{38,40,61,65,117}	$\frac{dM_d}{dt} = k_{dock}f_{Rab}M - k_{fus}M_d$	Rab activity, actin barrier	docked MVB pool
Small-EV release	Ca ²⁺ -sensitive fusion rate ^{24,57,62,143}	$R_{sEV} = k_{fus}f_{Ca}(C_{sub})M_d$	docked MVBs, Ca ²⁺ , SNARE	small-EV release rate
Budding-site formation	Budding ODE ^{26,63,85}	$\frac{dB_{m/IEV}}{dt} = k_{bud}f_{PS}f_{tension} - k_{scission}B_{m/IEV}$	PS, curvature, tension	budding precursor pool
Medium/large EV release	Scission-rate law ^{26,63,85,141}	$R_{m/IEV} = k_{scission}f_{actomyosin}B_{m/IEV}$	calpain, ROCK, actomyosin	medium/large EV rate
Apoptotic vesicle release	Injury-state weighted rate ^{20,78,94,121}	$R_{AB} = k_{AB}F_{apop}$	apoptotic fraction, blebbing rate	apoptotic-body release

Abbreviations: MVB, multivesicular body; ILV, intraluminal vesicle; ESCRT, endosomal sorting complex required for transport; sEV, small extracellular vesicle; m/IEV, medium/large extracellular vesicle; AB, apoptotic body.

cohort history: particles remain at their emitted diameter unless an explicit aggregation or fragmentation term $T_{bb'}$ transfers them. Initially, a common effective loss rate and $T_{bb'} = 0$ provide the most identifiable model.

Cargo is linked jointly to pathway, state, and size rather than determined by diameter alone. A reduced geometric capacity relation is

$$E[C_j | d, q, \mathbf{x}] = a_{jq}(\mathbf{x})d^2 + b_{jq}(\mathbf{x})d^3,$$

where the d^2 and d^3 terms represent membrane-associated and luminal capacity, respectively ¹⁴⁷. The coefficients retain pathway- and state-dependent sorting, and low-copy cargo can be represented by stochastic occupancy. Single-vesicle measurements showing selective changes in EV size, membrane order, and cargo after Rab27a or CD63 perturbation support this conditional rather than deterministic representation ¹³³.

Finally, biological stocks are mapped to an observed size bin b in assay run r through

$$\hat{y}_{rb}(t) = B_{rb} + \sum_{\ell} H_{rbl} \eta_{r\ell} \frac{\sum_q N_{q\ell}(t)}{V_r},$$

where H_{rbl} represents true-to-measured diameter error and rebinning, $\eta_{r\ell}$ represents size-dependent recovery, B_{rb} is background, and V_r is sample volume. This observation operator is required because different single-particle methods can report materially different size distributions and concentrations for the same EV preparation ^{125,138}. Instrument-response parameters should be constrained by calibration particles, instrument settings, and orthogonal characterization rather than inferred freely with the biological kernel.

For sparse longitudinal data, calibration should begin with literature- or sham-informed fixed kernels, dynamic pathway mixture weights, and a common extracellular loss term. A state-to-size coefficient should be retained only if it improves prediction for held-out pulse conditions; size-dependent loss should then be tested separately. Particle counts alone cannot identify pathway-specific cargo coefficients, which require marker-resolved or molecular measurements.

Cargo Sorting, EV Composition, and Potency

The cargo module represents the major conceptual expansion from EV-release modeling to EV-engineering modeling. nsPEF may alter EV composition through Ca²⁺-dependent trafficking, stress-responsive transcription and translation,

lipid remodeling, ESCRT/tetraspanin/ceramide pathways, RNA-binding proteins, and injury-state contamination. Therefore, EV output should be represented as a multi-dimensional product state rather than as total particle number alone. Table A11 separates the biological cargo-sorting evidence, and Table A12 specifies the computational cargo, composition, potency, and direct-loading model options.

The implemented Layer 6 module now follows the table structure explicitly. It computes a subtype-weighted cargo state from small-EV, medium/large-EV, and apoptotic-body outputs; separates protein, RNA, lipid, antigen, and direct-loading terms; and maps the resulting cargo vector to a saturating potency score. This keeps the implementation aligned with evidence that EV subtypes carry distinct protein compositions^{41,48}, that RNA sorting can be mediated by RNA-binding proteins such as YBX1 and SYNCRIP^{58,95,103}, that lipid composition and ceramide biology shape EV state¹⁰⁷, and that direct EV loading can be represented separately from endogenous cargo sorting^{27,29}.

Injury, Debris, and Quality Gate

The injury-quality module defines whether nsPEF exposure produces reversible EV engineering, stress-associated release, apoptosis, necrosis, or contaminating debris. This layer is essential because increased particle number is not necessarily equivalent to increased useful EV production. The quality gate separates bona fide EV output from apoptotic bodies, damaged organellar fragments, protein aggregates, and other co-isolated materials. Table A13 summarizes the biological and QC evidence for this distinction, while Table A14 defines the damage, fate-transition, mixture, and constrained-optimization model forms.

The implemented Layer 7 module expands the previous binary quality proxy into a mixture-aware gate. Damage is mapped to stressed-viable, apoptotic, and necrotic fractions using threshold-like Hill terms inspired by systems models of cell-death switching^{1,10}. The measured particle pool is then decomposed into bona fide EVs, apoptotic bodies, debris, and aggregate-like contaminants, which makes the purity score sensitive to both injury state and non-EV co-isolates. This structure matches EV-composition and isolation cautions from apoptotic/necrotic EV studies and MISEV-style purity guidance^{16,20,137,138}.

Manufacturing, Isolation, and Quality-Control Interface

The manufacturing module links cell-level mechanisms to translational EV production. It represents harvest timing, cell number, culture format, isolation recovery, marker-based QC, purity, batch variability, and flow-through scalability. This module is intentionally placed after biological release and quality gating because manufacturing success depends on both biological production and downstream process recovery. Table A15 summarizes the process and QC evidence to curate, and Table A16 lists the process-yield, recovery, batch, and optimization model forms.

The implemented Layer 8 module converts biological output into process-facing quantities: cell-normalized yield, harvest-rate-equivalent accumulation, isolation recovery, process purity, batch-adjusted yield, and a constrained

optimization objective. This keeps the code aligned with EV manufacturing literature emphasizing scalable source material, recovery and purity tradeoffs, clinical-grade purification, and validation requirements^{64,79,119,136,138}. The objective is intentionally transparent rather than prescriptive: potency, yield, purity, and viability weights can be changed for different process-development priorities.

Cell-State and Disease-State Modifiers as Cross-Cutting Parameters

Cell-state and disease-state modifiers alter how a given cell responds to the same nsPEF exposure. In this framework, they do not define a separate downstream pathway. Instead, they scale or reshape the behavior of all modules. Relevant modifiers include membrane composition, cholesterol and sphingolipid content, cortical stiffness, cytoskeletal prestress, baseline Ca^{2+} handling, mitochondrial state, metabolic capacity, oxidative-stress tone, inflammatory signaling, hypoxia response, baseline EV-biogenesis activity, and cargo-sorting phenotype. Because these modifiers act across layers, their evidence and parameter-scaling implications are distributed across the paired module tables summarized in Table 2 and specified in Tables A1–A16.

$$\theta^{cell} = S_{cell}\theta^{base}, \quad (\text{A7})$$

$$\theta_i \sim \mathcal{D}(\mu_{cell}, \Sigma_{cell}), \quad (\text{A8})$$

where θ^{cell} represents cell-specific parameter values, S_{cell} represents phenotype-specific scaling factors, and $\mathcal{D}(\mu_{cell}, \Sigma_{cell})$ represents population heterogeneity.

Core Governing Equations for a First Integrated Implementation

A first reduced implementation can be constructed by coupling pulse exposure, membrane permeability, Ca^{2+} balance, remodeling activation, EV-release rates, cargo state, repair, and quality gating. The following equations are placeholders for model assembly and should be refined as literature-supported flux expressions and parameters are identified. They assemble the computational forms listed in Tables A2, A4, A6, A8, A10, A12, and A14; manufacturing-facing optimization is summarized separately in Table A16.

$$C_m \frac{d\Delta V_m}{dt} + G_m(t)\Delta V_m = f(E(t), \theta), \quad (\text{A9})$$

$$C_o \frac{d\Delta V_o}{dt} + G_o(t)\Delta V_o = f_o(E(t), r_o, \theta_o), \quad (\text{A10})$$

$$\frac{dN_p}{dt} = k_{form}(\Delta V) - k_{reseat}N_p, \quad (\text{A11})$$

$$J_{i,pore} = P_i(t)\Phi_i([i]_e, [i]_i, V_m), \quad (\text{A12})$$

$$\begin{aligned} \frac{d[\text{Ca}^{2+}]_i}{dt} = & J_{\text{Ca,pore}} + J_{\text{VGCC}} + J_{\text{ER,rel}} - J_{\text{SERCA}} \\ & - J_{\text{PMCA}} - J_{\text{NCX}} - J_{\text{mito}} - J_{\text{buffer}}, \end{aligned} \quad (\text{A13})$$

Table A11. Biological evidence table for cargo sorting, EV composition, and potency.

Submodule	Biological phenomenon	Evidence to curate	Relevance to EV engineering	Confidence
Protein sorting	Ubiquitin/ESCRT/tetraspanin-dependent protein enrichment	ALIX, TSG101, CD63/CD81/CD9, antigen enrichment	Determines therapeutic or immunogenic protein cargo	Established ^{41,48,50,93,116}
RNA sorting	RNA-binding protein-mediated miRNA/mRNA loading	hnRNPA2B1, SYNCRIP, YBX1, motif-based sorting	Enables nucleic-acid cargo modulation	Supported ^{58,95,103}
Lipid sorting	Ceramide, cholesterol, sphingolipid, PE/PS enrichment	Lipidomics, nSMase2, tetraspanin domains	Controls EV stability, curvature, uptake, immunologic function	Established ^{22,25,93,107}
Stress-responsive cargo	ROS, hypoxia, inflammatory, apoptotic cargo changes	proteomics/lipidomics/RNA-seq after nsPEF	Distinguishes programmed cargo modulation from debris	Supported ^{20,68,84}
Antigen enrichment	Tumor antigen expression and EV loading	MAGE and other antigen studies	Supports vaccine-oriented EV engineering	Supported ^{49,114}
Functional potency	Recipient-cell response	uptake, cargo delivery, immune activation, repair assays	Connects cargo state to biological activity	Hypothesized ^{29,33,122}

Abbreviations: EV, extracellular vesicle; ESCRT, endosomal sorting complex required for transport; ALIX, ALG-2-interacting protein X; TSG101, tumor susceptibility gene 101; miRNA, microRNA; mRNA, messenger RNA; PE, phosphatidylethanolamine; PS, phosphatidylserine; RNA-seq, RNA sequencing.

Table A12. Computational framework table for cargo sorting, EV composition, and potency.

Submodule	Model form	Representative relation	Inputs	Outputs
Subtype-weighted cargo	Weighted model ^{20,41,48}	output	$Cargo_j = w_{j,sEV} R_{sEV} + w_{j,m/IEV} R_{m/IEV} + w_{j,AB} R_{AB}$	subtype rates, cargo weights
Stress-modulated sorting	Rate law ^{58,84,95,103}	$\frac{dCargo_j}{dt} = k_{sort,j} f_{stress} f_{Ca} f_{ESCRT} - k_{loss,j} Cargo_j$	Ca^{2+} , ROS, ESCRT, stress	cargo state
Antigen enrichment	Transcription/sorting coupling ^{49,114}	$\frac{dA_g}{dt} = k_{expr} f_{stress} - k_{deg} A_g - k_{sort} A_g$	stress, antigen expression	antigen cargo
Lipid composition	Composition-state ODE ^{22,25,107}	$\frac{dL_j}{dt} = k_{syn,j} f_{stress} - k_{sort,j} L_j$	lipid metabolism, EV subtype	lipid cargo class
Potency mapping	Empirical or mechanistic response model ^{29,33,122}	$Potency = F(Cargo, Subtype, Dose)$	cargo vector, recipient assay	functional potency
Direct EV loading	Permeability-diffusion model ^{27,29,37,44}	$\frac{dC_{load}}{dt} = P_{EV}(t) A_{EV} (C_{ext} - C_{EV}) - k_{leak} C_{EV}$	EV permeability, cargo gradient	loaded cargo

Abbreviations: EV, extracellular vesicle; ESCRT, endosomal sorting complex required for transport; ALIX, ALG-2-interacting protein X; sEV, small extracellular vesicle; m/IEV, medium/large extracellular vesicle; AB, apoptotic body.

Table A13. Biological evidence table for injury, debris, and quality gating.

Submodule	Biological phenomenon	Evidence to curate	Relevance to EV engineering	Confidence
Viability preservation	Reversible response versus cell death	viability, membrane recovery, clonogenic/metabolic assays	Defines acceptable EV-production window	Established ^{8,12,83}
Mitochondrial injury	Depolarization, mPTP, cytochrome c	$\Delta\Psi_m$, cytochrome c, ROS	Separates stress signaling from organellar contamination	Supported ^{8,71,84}
Apoptosis	Caspase activation and apoptotic blebbing	caspase, annexin V/PI, ROCK1 blebbing	Determines apoptotic-body contribution	Established ^{20,83,121}
Necrosis/lysis	Irreversible membrane failure	LDH release, PI uptake, debris microscopy	Indicates unacceptable product contamination	Established ^{20,83}
Non-EV contaminants	Protein aggregates, organelle fragments, apoptotic bodies	calnexin, cytochrome c, albumin/lipoprotein markers	Defines purity and interpretability of EV readouts	Established ^{16,20,137,138}
Quality-control marker panel	Positive and negative EV markers	CD63/CD81/CD9, TSG101/ALIX/syntenin, negative markers	Connects model to MISEV-style QC	Established ^{41,48,138}

Abbreviations: EV, extracellular vesicle; mPTP, mitochondrial permeability transition pore; PI, propidium iodide; QC, quality control; MISEV, Minimal Information for Studies of Extracellular Vesicles.

$$A_x = \frac{[Ca^{2+}]_{sub}^n}{K_x^n + [Ca^{2+}]_{sub}^n}, \quad (A14)$$

$$R_{EV} = R_{sEV} + R_{m/IEV} + R_{AB}, \quad (A15)$$

$$R_{sEV} = k_{fus} f_{Ca} ([Ca^{2+}]_{sub}) M_d, \quad (A16)$$

Table A14. Computational framework table for injury, debris, and quality gating.

Submodule	Model form	Representative relation	Inputs	Outputs
Damage accumulation	Damage-state ODE ^{1,12,84}	$\frac{dD}{dt} = k_{inj}G_m + k_{ROS}ROS - k_{repair}A_{repair}D$	conductance, repair	ROS, damage burden
Fate transition	Hybrid state rule ^{1,10,83}	$F_{apop} = H(D - D^*)$ or $\frac{dF_{apop}}{dt} = k_{apop}H(D - D^*)$	damage, mitochondria	ATP, viable/stressed/apoptotic fractions
Necrotic transition	Threshold rule ^{20,83}	$F_{nec} = H(D - D_{nec}^*)H(A_{ATP}^* - ATP)$	damage, membrane failure	ATP, necrotic fraction
Measured particles	Mixture model ^{16,20,137}	$N_{meas} = N_{EV} + N_{AB} + N_{debris} + N_{agg}$	release rates, injury state	measured particle count
Purity score	Ratio or classifier ^{16,137,138}	$Q_{EV} = \frac{N_{bona\ fide\ EV}}{N_{meas}}$	marker panel, debris state	EV quality score
Acceptable window	Constrained optimization ^{32,76,138}	maximize Y_{EV} subject to $Q_{EV} > Q^*$ and $Viability > V^*$	dose, cell state, harvest	operating region

Abbreviations: EV, extracellular vesicle; AB, apoptotic body; ATP, adenosine triphosphate; ROS, reactive oxygen species; QC, quality control.

Table A15. Biological/process evidence table for manufacturing, isolation, and quality-control interface.

Submodule	Process phenomenon	Evidence to curate	Relevance to EV engineering	Confidence
Harvest kinetics	Time-dependent EV accumulation	minute-to-hour release curves after nsPEF	Determines optimal collection window	Supported ^{32,64,76}
Cell-normalized yield	EV output per cell or protein	NTA/flow/ELISA normalized to cell number	Defines productivity	Established ^{64,79,138}
Isolation recovery	SEC, ultracentrifugation, density gradient, affinity capture	recovery and purity comparisons	Converts released EVs into isolated product	Established ^{16,136,137}
QC characterization	Size, morphology, markers, contaminants	NTA, cryo-EM/TEM, western blot/flow, proteomics	Determines whether product meets EV criteria	Established ^{41,48,138}
Batch variability	Donor, passage, culture, and pulse variability	repeated-batch studies, mixed-effects data	Determines reproducibility and manufacturability	Supported ^{64,79,119,136}
Scalable flow processing	Closed-system pulse chamber and inline monitoring	field uniformity, sterility, temperature, flow control	Enables translational manufacturing	Hypothesized ^{13,30,56,119}

Abbreviations: EV, extracellular vesicle; nsPEF, nanosecond pulsed electric field; NTA, nanoparticle tracking analysis; ELISA, enzyme-linked immunosorbent assay; QC, quality control; TEM, transmission electron microscopy.

Table A16. Computational framework table for manufacturing, isolation, and quality-control interface.

Submodule	Model form	Representative relation	Inputs	Outputs
Cell-normalized yield	Productivity metric ^{64,79,138}	$Y_{EV} = \frac{N_{EV}}{N_{cells}}$	EV count, cell number	EV yield per cell
Harvest accumulation	Time-integral model ^{32,64,76}	$N_{EV}(t_h) = \int_0^{t_h} R_{EV}(t)dt$	release rate, harvest time	collected EVs
Isolation recovery	Process-yield balance ^{16,136,137}	$N_{iso} = \eta_{iso}N_{released}$	isolation method, recovery	isolated EV count
Purity metric	Product-quality ratio ^{16,137,138}	$Purity = \frac{N_{EV}}{Protein_{total}}$ or marker-based score	particle/protein/marker data	purity estimate
Batch model	Mixed-effects model ^{64,79,119,136}	$Y_b = \mu + \alpha_{cell} + \beta_{dose} + u_b + \epsilon$	cell state, dose, batch	reproducibility
Optimization objective	Constrained process optimization ^{64,79,119,136,138}	maximize $Potency \times Y_{iso}$ subject to purity and viability constraints	dose, harvest, isolation	optimal protocol

Abbreviations: EV, extracellular vesicle; QC, quality control.

$$R_{m/LEV} = k_{scission}f_{PS}(PS_{out})f_{actomyosin}B_{m/LEV}, \quad (A17)$$

$$\mathcal{Y}_{EV} = \{N_{EV}, R_{EV}, Subtype, Size, Cargo, Potency, Purity, Viability, \hat{y}\}. \quad (A21)$$

$$Cargo_j = w_{j,SEV}R_{sEV} + w_{j,m/LEV}R_{m/LEV} + w_{j,AB}R_{AB}, \quad (A18)$$

$$\frac{dD}{dt} = k_{inj}G_m + k_{ROS}ROS - k_{repair}A_{repair}D, \quad (A19)$$

$$N_{meas} = N_{EV} + N_{AB} + N_{debris} + N_{agg}, \quad (A20)$$

Variable and Parameter Glossary Template

Table A17 provides a shared notation layer for the variables and parameters introduced across the appendix tables and governing-equation scaffold.

Table A17. Variable and parameter glossary for the nsPEF-EV engineering model.

Symbol	Definition	Units	Module	Source/estimation strategy
$E(t)$	Applied electric-field waveform	V/m	Dosimetry	Pulse protocol/field model
D_E	Electrical dose index	variable	Dosimetry	Derived from pulse protocol
ΔV_m	Plasma-membrane transmembrane potential	V	Electrodynamics	Circuit/electroporation model
ΔV_o	Organelle membrane potential perturbation	V	Organelle electrodynamics	Multi-shell/circuit model
N_p	Pore density or pore number	pores/ μm^2 or count	Electroporation	Fit or literature prior
r_p	Representative pore radius	nm	Electroporation	Literature prior or fit
G_m	Effective membrane conductance	S/m ²	Membrane/repair	Fit from permeability recovery
P_i	Ion-specific membrane permeability	m/s	Ion transport	Estimated from flux data
$J_{\text{Ca,pore}}$	Electropore-mediated Ca ²⁺ flux	concentration/time	Ion transport	Derived from permeability relation
$[\text{Ca}^{2+}]_i$	Cytosolic Ca ²⁺ concentration	M	Ca ²⁺ signaling	Ca ²⁺ imaging/model state
$[\text{Ca}^{2+}]_{sub}$	Submembrane Ca ²⁺ concentration	M	Microdomain	Reaction-diffusion/shell model
ROS	Reactive oxygen species state	variable	Stress/bioenergetics	ROS imaging/probes
ATP	Energy availability state	variable	Bioenergetics	ATP assays/model state
PS_{out}	Externalized phosphatidylserine fraction	dimensionless	Remodeling	Annexin V or model state
M_d	Docked MVB pool	count or concentration	EV release	Fit/estimated latent state
$B_{m/LEV}$	Medium/large EV budding precursor pool	count or concentration	EV release	Fit/estimated latent state
R_{sEV}	Small-EV release rate	particles/time	EV output	NTA/flow/marker-based fit
$R_{m/LEV}$	Medium/large EV release rate	particles/time	EV output	NTA/flow/marker-based fit
R_{AB}	Apoptotic body release rate	particles/time	Injury output	Apoptosis-weighted output
$Cargo_j$	Abundance of cargo class j	variable	Cargo	Proteomics/RNA/lipidomics
$Potency$	Functional activity metric	assay-specific	Potency	Recipient-cell assay
D	Integrated damage burden	dimensionless	Quality gate	Hybrid state variable
Q_{EV}	EV quality/purity score	dimensionless	QC	Marker/purity model
η_{iso}	Isolation recovery efficiency	dimensionless	Manufacturing	Process validation

Abbreviations: nsPEF, nanosecond pulsed electric field; EV, extracellular vesicle; ROS, reactive oxygen species; ATP, adenosine triphosphate; PS, phosphatidylserine; MVB, multivesicular body; ILV, intraluminal vesicle; QC, quality control.

References

- John G. Albeck, John M. Burke, Sabrina L. Spencer, Douglas A. Lauffenburger, and Peter K. Sorger. Modeling a snap-action, variable-delay switch controlling extrinsic cell death. *PLoS Biology*, 6(12):e299, 2008.
- Norma W. Andrews, Paulo E. Almeida, and Matthias Corrotte. Damage control: cellular mechanisms of plasma membrane repair. *Trends in Cell Biology*, 24(12):734–742, 2014.
- Jan E. Azarov, Iurii Semenov, Maura Casciola, and Andrei G. Pakhomov. Excitation of murine cardiac myocytes by nanosecond pulsed electric field. *Journal of Cardiovascular Electrophysiology*, 30(3):392–401, 2019.
- Tarique R. Bagalkot, Robert C. Terhune, Normand Leblanc, and Gale L. Craviso. Different membrane pathways mediate Ca²⁺ influx in adrenal chromaffin cells exposed to 150–400 ns electric pulses. *BioMed Research International*, 2018:9046891, 2018.
- C. Baker, A. Willis, W. Milestone, M. Baker, A. L. Garner, and R. P. Joshi. Numerical assessments of geometry, proximity and multi-electrode effects on electroporation in mitochondria and the endoplasmic reticulum to nanosecond electric pulses. *Scientific Reports*, 14:23854, 2024.
- N. Bakute, E. Andriukonis, K. Kasperaviciute, J. Dobilas, M. Sapurov, G. Mozolevskis, and A. Stirke. Microphysiological system with integrated sensors to study the effect of pulsed electric field. *Scientific Reports*, 14(1):18713, 2024.
- Daniel A. Beard. A biophysical model of the mitochondrial respiratory system and oxidative phosphorylation. *PLOS Computational Biology*, 1(4):e36, 2005.
- Stephen J. Beebe, Y. J. Chen, N. M. Sain, Karl H. Schoenbach, and Shu Xiao. Transient features in nanosecond pulsed electric fields differentially modulate mitochondria and viability. *PLoS ONE*, 7(12):e51349, 2012.
- Stephen J. Beebe, P. M. Fox, L. J. Rec, K. Somers, R. H. Stark, and K. H. Schoenbach. Nanosecond, high-intensity pulsed electric fields induce apoptosis in human cells. *FASEB Journal*, 17(11):1493–1495, 2003.
- Marc Bentele, Inna Lavrik, Marcel Ulrich, Stefanie Stosser, Dieter W. Heermann, Holger Kalthoff, et al. Mathematical modeling reveals threshold mechanism in cd95-induced apoptosis. *The Journal of Cell Biology*, 166(6):839–851, 2004.
- Richard Bertram, Morten Gram Pedersen, Dan S. Luciani, and Arthur Sherman. A simplified model for mitochondrial ATP production. *Journal of Theoretical Biology*, 243(4):575–586, 2006.
- Sayak Bhattacharya, Mantas Silkunas, Emily Gudvangen, Uma Mangalanathan, Olga N. Pakhomova, and Andrei G. Pakhomov. Ca²⁺ dependence and kinetics of cell membrane repair after electroporation. *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1864(2):183823, 2024.

- 2022.
13. M. Bonakdar, P. M. Graybill, and R. V. Davalos. A microfluidic model of the blood-brain barrier to study permeabilization by pulsed electric fields. *RSC Advances*, 7(68):42811–42818, 2017.
14. Anna-Cathrin Borchers et al. Regulatory sites in the mon1–ccz1 complex control rab5 to rab7 transition and endosome maturation. *Proceedings of the National Academy of Sciences*, 120(30):e2303750120, 2023.
15. Theresa L. Boye, Kenji Maeda, Weria Pezeshkian, Stine L. Sonder, Sara C. Haeger, Volker Gerke, Adam C. Simonsen, and Jesper Nylandsted. Annexin A4 and A6 induce membrane curvature and constriction during cell membrane repair. *Nature Communications*, 8(1):1623, 2017.
16. K. Brennan, K. Martin, S. P. FitzGerald, J. O’Sullivan, Y. Wu, A. Blanco, et al. A comparison of methods for the isolation and separation of extracellular vesicles from protein and lipid particles in human serum. *Scientific Reports*, 10:1039, 2020.
17. Ryan C. Burke, Sylvia M. Bardet, Lynn Carr, Sergii Romanenko, Delia Arnaud-Cormos, Philippe L  v  que, and Rodney P. O’Connor. Nanosecond pulsed electric fields depolarize transmembrane potential via voltage-gated K^+ , Ca^{2+} and TRPM8 channels in U87 glioblastoma cells. *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1859(10):2040–2050, 2017.
18. Robert L. Campbell and Peter L. Davies. Structure–function relationships in calpains. *Biochemical Journal*, 447(3):335–351, 2012.
19. Jody C. Cantu, Gleb P. Tolstykh, Melissa Tarango, Hope T. Beier, and Bennett L. Ibey. Caveolin-1 is involved in regulating the biological response of cells to nanosecond pulsed electric fields. *The Journal of Membrane Biology*, 254(2):141–156, 2021.
20. Benjamin Cappe, Maija Vadi, Ethan Sack, Laurent Wacheul, Bert Verstraeten, Simon Dufour, et al. Systematic compositional analysis of exosomal extracellular vesicles produced by cells undergoing apoptosis, necroptosis and ferroptosis. *Journal of Extracellular Vesicles*, 12(10):e12365, 2023.
21. Tania Castro-Gomes, Matthias Corrotte, Calvin Tam, and Norma W. Andrews. Plasma membrane repair is regulated extracellularly by proteases released from lysosomes. *PLOS ONE*, 11(3):e0152583, 2016.
22. Dolma Choezom and Julia Christina Gross. Neutral sphingomyelinase 2 controls exosome secretion by counteracting v-atpase-mediated endosome acidification. *Journal of Cell Science*, 2022.
23. Sonia Cortassa, Miguel A. Aon, Raimond L. Winslow, and Brian O’Rourke. A mitochondrial oscillator dependent on reactive oxygen species. *Biophysical Journal*, 87(3):2060–2073, 2004.
24. Kevin C. Courtney et al. Synaptotagmin-7 outperforms synaptotagmin-1 to promote the formation of large, stable fusion pores via robust membrane penetration. *Nature Communications*, 14, 2023.
25. Simone M. C. Crivelli et al. Function of ceramide transfer protein for biogenesis and sphingolipid composition of extracellular vesicles. *Journal of Extracellular Vesicles*, 2022.
26. Huijun Dai, Suisui Zhang, Xueke Du, Weikang Zhang, Ren Jing, Xiaoxia Wang, and Linghui Pan. RhoA inhibitor suppresses the production of microvesicles and rescues high ventilation induced lung injury. *International Immunopharmacology*, 2019.
27. Marie-Cecile Didiot, Linda M. Hall, Anne H. Coles, Reka A. Haraszti, Bruno M. D. C. Godinho, Kris Chase, et al. Optimized cholesterol-sirna chemistry improves productive loading onto extracellular vesicles. *Molecular Therapy*, 26(6):1500–1509, 2018.
28. Stine Krog Frandsen, Hanne Gissel, Pernille Hojman, Trine Tramm, Jens Eriksen, and Julie Gehl. Direct therapeutic applications of calcium electroporation to effectively induce tumor necrosis. *Cancer Research*, 72(6):1336–1341, 2012.
29. Gregor Fuhrmann, Andrea Serio, Maria Mazo, Rohit Nair, and Molly M. Stevens. Active loading into extracellular vesicles significantly improves the cellular uptake and photodynamic effect of porphyrins. *Journal of Controlled Release*, 205:35–44, 2015.
30. P. A. Garcia, Z. Ge, J. L. Moran, and C. R. Buie. Microfluidic screening of electric fields for electroporation. *Scientific Reports*, 6:21238, 2016.
31. Laure Gibot, Audrey Montigny, Houda Baaziz, Isabelle Fourquaux, Marc Audebert, and Marie-Pierre Rols. Calcium delivery by electroporation induces in vitro cell death through mitochondrial dysfunction without DNA damages. *Cancers*, 12(2):425, 2020.
32. Emily Gudvangen, Vitalii Kim, Vitalij Novickij, Federico Battista, and Andrei G. Pakhomov. Electroporation and cell killing by milli- to nanosecond pulses and avoiding neuromuscular stimulation in cancer ablation. *Scientific Reports*, 12(1):1763, 2022.
33. Daniel W. Hagey, Miina Ojansivu, Burcu R. Bostancioglu, Osama Saher, Jean-Philippe Bost, Maria O. Gustafsson, Roberto Gramignoli, Helene Andersson Svahn, Divya Gupta, Molly M. Stevens, Andre Gorgens, and Samir El Andaloussi. The cellular response to extracellular vesicles is dependent on their cell source and dose. *Science Advances*, 9(35):eadh1168, 2023.
34. Ali Asghar Hakami Zanjani, Malene Laage Ebstrup, Jesper Nylandsted, and Himanshu Khandelwa. Modulation of annexin-induced membrane curvature by cholesterol and the anionic lipid headgroup during plasma membrane repair. *The Journal of Physical Chemistry B*, 128(36):8701–8711, 2024.
35. Tian-Wei Han, Wei Ye, Nathan P. Bethel, Marta Zubia, Angela Kim, Karen H. Li, Alma L. Burlingame, Michael Grabe, Yuh Nung Jan, and Lily Y. Jan. Chemically induced vesiculation as a platform for studying TMEM16F activity. *Proceedings of the National Academy of Sciences of the United States of America*, 116(4):1309–1318, 2019.
36. Emily Hellwich et al. Extracellular Ca^{2+} drives cytoskeletal remodeling and cell swelling following nanosecond pulsed electric field exposure. *Bioelectrochemistry*, 170:109241, 2026.
37. Joshua L. Hood, Mark J. Scott, and Samuel A. Wickline. Maximizing exosome colloidal stability following electroporation. *Analytical Biochemistry*, 448:41–49, 2014.
38. Chia Hsu, Yasunori Morohashi, Shin Yoshimura, Natalia Manrique-Hoyos, Sang-Hyeon Jung, Olivier Loudig, Lakshmi K. Mahal, Hiroshi Horiuchi, and Mikael Simons. Regulation of exosome secretion by rab35 and its gtpase-activating proteins tbc1d10a-c. *The Journal of Cell Biology*, 189(2):223–232, 2010.

39. Jatta Huotari and Ari Helenius. Endosome maturation. *The EMBO Journal*, 30(17):3481–3500, 2011.
40. Tetsuro Izumi. In vivo roles of rab27 and its effectors in exocytosis. *Cell Structure and Function*, 46:79–94, 2021.
41. Dennis K. Jeppesen, A. M. Fenix, J. L. Franklin, J. N. Higginbotham, Q. Zhang, L. J. Zimmerman, et al. Reassessment of exosome composition. *Cell*, 177(2):428–445.e18, 2019.
42. Ravindra P. Joshi and Karl H. Schoenbach. Electroporation dynamics in biological cells subjected to ultrafast electrical pulses: A numerical simulation study. *Physical Review E*, 62(1):1025–1033, 2000.
43. Maša Kanduđer, Marjeta Sentjunc, and Damijan Miklavčič. The temperature effect during pulse application on cell membrane fluidity and permeabilization. *Bioelectrochemistry*, 74(1):52–57, 2008.
44. Sander A. A. Kooijmans, Sam Stremersch, Kevin Braeckmans, Stefaan C. de Smedt, An Hendrix, Matthew J. A. Wood, Raymond M. Schiffelers, Koen Raemdonck, and Pieter Vader. Electroporation-induced siRNA precipitation obscures the efficiency of siRNA loading into extracellular vesicles. *Journal of Controlled Release*, 172(1):229–238, 2013.
45. Tadej Kotnik and Damijan Miklavčič. Analytical description of transmembrane voltage induced by electric fields on spheroidal cells. *Biophysical Journal*, 79(2):670–679, 2000.
46. Tadej Kotnik, Lea Rems, Mounir Tarek, and Damijan Miklavčič. Membrane electroporation and electroporeabilization: mechanisms and models. *Annual review of biophysics*, 48(1):63–91, 2019.
47. Wanda Krassowska and Petar D. Filev. Modeling electroporation in a single cell. *Biophysical journal*, 92(2):404–417, 2007.
48. Fernanda G. Kugeratski, Katie Hodge, Stefania Lilla, Kathleen M. McAndrews, Xiao Zhou, Robert F. Hwang, et al. Quantitative proteomics identifies the core proteome of exosomes with syntenin-1 as the highest abundant protein and a putative universal biomarker. *Nature Cell Biology*, 23(6):631–641, 2021.
49. Pankaj Kumar, Connor Boyne, Samantha Brown, Arif Qureshi, Paul Thorpe, Sander A. Synowsky, et al. Tumour-associated antigenic peptides are present in the hla class i ligandome of cancer cell line derived extracellular vesicles. *Immunology*, 166(2):249–264, 2022.
50. Jorge Larios, Vincent Mercier, Aurelien Roux, and Jean Gruenberg. Alix- and escrt-iii-dependent sorting of tetraspanins to exosomes. *The Journal of Cell Biology*, 2020.
51. Heedoo Lee, Chunhua Li, Yang Zhang, Duo Zhang, Leo E. Otterbein, and Yang Jin. Caveolin-1 selectively regulates microRNA sorting into microvesicles after noxious stimuli. *Journal of Experimental Medicine*, 216(9):2202–2220, 2019.
52. Philippe Lévêque, Rory O'Connor, Delia Arnaud-Cormos, and Damijan Miklavčič. Measurement and characterization of exposure systems for high-frequency, ultrashort pulses. In *Handbook of Electroporation*, pages 813–836. Springer International Publishing, 2017.
53. Y. Li, M. Wu, D. Zhao, Z. Wei, W. Zhong, X. Wang, Z. Liang, and Z. Li. Electroporation on microchips: The harmful effects of pH changes and scaling down. *Scientific Reports*, 5:17817, 2015.
54. Yue-Xian Li and John Rinzel. Equations for InsP_3 receptor-mediated Ca^{2+} oscillations derived from a detailed kinetic model: A Hodgkin–Huxley-like formalism. *Journal of Theoretical Biology*, 166(4):461–473, 1994.
55. Susanne Liese, Eva Maria Wenzel, Ingrid Kjos, Rossana Rojas Molina, Sebastian W. Schultz, Andreas Brech, Harald Stenmark, Camilla Raiborg, and Andreas Carlson. Protein crowding mediates membrane remodeling in upstream ESCRT-induced formation of intraluminal vesicles. *Proceedings of the National Academy of Sciences of the United States of America*, 117(46):28614–28624, 2020.
56. C. J. S. Liu, S. B. Wang, P. Y. Su, Y. P. Hsieh, and H. Y. Wang. Microfluidic electroporation for drug and gene delivery: Driving innovation from single-cell precision to high-throughput preclinical and therapeutic platforms. *Advanced Drug Delivery Reviews*, 233:115851, 2026.
57. Chuqi Liu, Dexiang Liu, Shen Wang, Lu Gan, Xiangliang Yang, and Cong Ma. Identification of the snare complex that mediates the fusion of multivesicular bodies with the plasma membrane in exosome secretion. *Journal of Extracellular Vesicles*, 12:12356, 2023.
58. Xiao-Ming Liu, Lu Ma, and Randy Schekman. Selective sorting of micRNAs into exosomes by phase-separated ybx1 condensates. *eLife*, 10:e71982, 2021.
59. Gerhard Magnus and Joel Keizer. Minimal model of beta-cell mitochondrial Ca^{2+} handling. *American Journal of Physiology-Cell Physiology*, 273(2):C717–C733, 1997.
60. Hamed Manoochehri, Anita S La'ah, Ali Babaeizad, Mohsen Sheykhasan, Mohadeseh Rostamipoor, Mahdihyeh Abbaspoor, Fariba Nikraves, Samira Mozaffari Khosravi, Hanie Mahaki, Hamid Tanzadehpahan, et al. Extracellular vesicles in cancer immunotherapy: Therapeutic, challenges and clinical progress. *Asian Journal of Pharmaceutical Sciences*, 20(5):101065, 2025.
61. Pauline P. Marie et al. Accessory escrt-iii proteins are conserved and selective regulators of rab11a-exosome formation. *Journal of Extracellular Vesicles*, 2023.
62. Takahide Matsui, Yuriko Sakamaki, Shu Hiragi, and Mitsunori Fukuda. Vamp5 and distinct sets of cognate q-snares mediate exosome release. *Cell Structure and Function*, 2023.
63. Kerstin Menck et al. Neutral sphingomyelinases control extracellular vesicles budding from the plasma membrane. *Journal of Extracellular Vesicles*, 2017.
64. Mayela Mendt, Shilpa Kamerkar, Hiroshi Sugimoto, Kathleen M. McAndrews, Chien-Chia Wu, Mihai Gagea, et al. Generation and testing of clinical-grade exosomes for pancreatic cancer. *JCI Insight*, 3(8):e99263, 2018.
65. Scott W. Messenger, Sang Su Woo, Zhongze Sun, and Thomas F. J. Martin. A Ca^{2+} -stimulated exosome release pathway in cancer cells is regulated by munc13-4. *The Journal of Cell Biology*, 217(8):2877–2890, 2018.
66. Keiko Morotomi-Yano, Hidenori Akiyama, and Ken-ichi Yano. Different involvement of extracellular calcium in two modes of cell death induced by nanosecond pulsed electric fields. *Archives of Biochemistry and Biophysics*, 555–556:47–54, 2014.
67. Anna Mularski, Stine L. Sonder, Anne S. B. Heitmann, Mahesh P. Pandey, Himanshu Khandelia, Jesper Nylandsted, and Adam C. Simonsen. Interplay of membrane crosslinking and curvature induction by annexins. *Scientific Reports*, 12(1):22568, 2022.

68. Claudia Muratori, Andrei G. Pakhomov, Elena Gianulis, Jade Meads, Maura Casciola, Peter A. Mollica, and Olga N. Pakhomova. Activation of the phospholipid scramblase TMEM16F by nanosecond pulsed electric fields (nsPEF) facilitates its diverse cytophysiological effects. *Journal of Biological Chemistry*, 292(47):19381–19391, 2017.
69. Claudia Muratori, Giedre Silkuniene, Peter A. Mollica, Andrei G. Pakhomov, and Olga N. Pakhomova. The role of ESCRT-III and Annexin V in the repair of cell membrane permeabilization by the nanosecond pulsed electric field. *Bioelectrochemistry*, 140:107837, 2021.
70. Elizabeth Murphy, Hossein Ardehali, Robert S. Balaban, Fabio DiLisa, Gerald W. Dorn, Richard N. Kitsis, Kinya Otsu, Peipei Ping, Rosario Rizzuto, Michael N. Sack, Douglas Wallace, and Richard J. Youle. Mitochondrial function, biology, and role in disease: A scientific statement from the american heart association. *Circulation Research*, 118(12):1960–1991, 2016.
71. Tina Batista Napotnik, Yu-Hsuan Wu, Martin A. Gundersen, Damijan Miklavčič, and P. Thomas Vernier. Nanosecond electric pulses cause mitochondrial membrane permeabilization in jurkat cells. *Bioelectromagnetics*, 33(3):257–264, 2012.
72. Mohammad Naraghi and Erwin Neher. Linearized buffered Ca^{2+} diffusion in microdomains and its implications for calculation of $[\text{Ca}^{2+}]$ at the mouth of a calcium channel. *Journal of Neuroscience*, 17(18):6961–6973, 1997.
73. D. Needham and R. M. Hochmuth. Electro-mechanical permeabilization of lipid vesicles: Role of membrane tension and compressibility. *Biophysical Journal*, 55(5):1001–1009, 1989.
74. V. Nesin, A. M. Bowman, S. Xiao, and A. G. Pakhomov. Cell permeabilization and inhibition of voltage-gated Ca^{2+} and Na^{+} channel currents by nanosecond pulsed electric field. *Bioelectromagnetics*, 33(5):394–404, 2012.
75. Richard Nuccitelli, Kaying Lui, Mark Kreis, Brian Athos, and Pamela Nuccitelli. Nanosecond pulsed electric field stimulation of reactive oxygen species in human pancreatic cancer cells is Ca^{2+} -dependent. *Biochemical and Biophysical Research Communications*, 435(4):580–585, 2013.
76. Rosa Orlacchio, Jelena Kolosnjaj-Tabi, Nicolas Mattei, Philippe Lévêque, Marie Pierre Rols, Delia Arnaud-Cormos, and Muriel Golzio. Effects of nanosecond pulsed electric field (nsPEF) on a multicellular spheroid tumor model: Influence of pulse duration, pulse repetition rate, absorbed energy, and temperature. *International Journal of Molecular Sciences*, 24(19):14999, 2023.
77. Daniel P. Ott, Samit Desai, Jachen A. Solinger, Andres Kaeck, and Anne Spang. Coordination between escrt function and rab conversion during endosome maturation. *The EMBO Journal*, 2025.
78. Aysun Ozdemir et al. A novel proteolytic cleavage of rock 1 in cell death: Not only by caspases 3 and 7 but also by caspase 2. *Biochemical and Biophysical Research Communications*, 2021.
79. Cristina Paganini, Ugo Capasso Palmiero, Gabriella Pocsfalvi, Nicolas Touzet, Antonella Bongiovanni, and Paolo Arosio. Scalable production and isolation of extracellular vesicles: available sources and lessons from current industrial bioprocesses. *Biotechnology Journal*, 14(10):1800528, 2019.
80. Andrei G. Pakhomov, Juergen F. Kolb, Jody A. White, Ravindra P. Joshi, Shu Xiao, and Karl H. Schoenbach. Long-lasting plasma membrane permeabilization in mammalian cells by nanosecond pulsed electric field. *Bioelectromagnetics*, 28(8):655–663, 2007.
81. Andrei G. Pakhomov, Roman Shevin, Jody A. White, Juergen F. Kolb, Olga N. Pakhomova, Ravindra P. Joshi, and Karl H. Schoenbach. Membrane permeabilization and cell damage by ultrashort electric field shocks. *Archives of Biochemistry and Biophysics*, 465(1):109–118, 2007.
82. Andrei G. Pakhomov, Shu Xiao, Olga N. Pakhomova, Iurii Semenov, Marjorie A. Kuipers, and Bennett L. Ibey. Disassembly of actin structures by nanosecond pulsed electric field is a downstream effect of cell swelling. *Bioelectrochemistry*, 100:88–95, 2014.
83. Olga N. Pakhomova, Betsy W. Gregory, Iurii Semenov, Andrei G. Pakhomov, and Maria Rosaria Scarfi. Two modes of cell death caused by exposure to nanosecond pulsed electric field. *PLoS ONE*, 8(7):e70278, 2013.
84. Olga N. Pakhomova, Vera A. Khorokhorina, Angela M. Bowman, Raminta Rodaite-Riseviciene, Gintautas Saulis, Shu Xiao, and Andrei G. Pakhomov. Oxidative effects of nanosecond pulsed electric field exposure in cells and cell-free media. *Archives of Biochemistry and Biophysics*, 527(1):55–64, 2012.
85. Xiaoyun Pang et al. Structural elucidation of how arf small gtpases induce membrane tubulation for vesicle fission. *Proceedings of the National Academy of Sciences*, 122(12):e2417820122, 2025.
86. Michael Paulaitis, Kitty Agarwal, and Patrick Nana-Sinkam. Dynamic scaling of exosome sizes. *Langmuir*, 34(32):9387–9393, 2018.
87. Anna-Katharina Pfitzner et al. An escrt-iii polymerization sequence drives membrane deformation and fission. *Cell*, 182(5):1140–1155, 2020.
88. Marta Radulovic, Koen O. Schink, Eva M. Wenzel, Viola Nahse, Alberto Bongiovanni, Frank Lafont, and Harald Stenmark. ESCRT-mediated lysosome repair precedes lysophagy and promotes cell survival. *The EMBO Journal*, 37(21):e99753, 2018.
89. Eivina Radzeviciute-Valciuke, Veronika Malysko-Ptasinske, Egle Mickeviciute, Julita Kulbacka, Nina Rembialska, Aukse Zinkeviciene, Jurij Novickij, and Vitalij Novickij. Calcium electroporation causes ATP depletion in cells and is effective both in microsecond and nanosecond pulse range as a modality of electrochemotherapy. *Bioelectrochemistry*, 155:108574, 2024.
90. Maksim A. Rassokhin and Andrei G. Pakhomov. Cellular regulation of extension and retraction of pseudopod-like blebs produced by nanosecond pulsed electric field (nsPEF). *Cell Biochemistry and Biophysics*, 69(3):555–566, 2014.
91. Lea Rems, Marko Ušaj, Maša Kandušer, Matej Reberšek, and Damijan Miklavčič. Cell electrofusion using nanosecond electric pulses. *Scientific Reports*, 3:3385, 2013.
92. Rhiannon E. Roberts, Marianne Martin, Sabrina Marion, Geetha L. Elumalai, Kimberly Lewis, Maurice B. Hallett, and Ana-Maria Lennon-Dumenil. Ca^{2+} -activated cleavage of ezrin visualised dynamically in living myeloid cells during cell surface area expansion. *Journal of Cell Science*, 133(5):jcs236968, 2020.

93. Eric Rubinstein, Clotilde Thery, and Pascale Zimmermann. Tetraspanins affect membrane structures and the trafficking of molecular partners: what impact on extracellular vesicles? *Biochemical Society Transactions*, 2025.
94. Kazuhiro Sakamaki et al. Caspase-mediated cleavage of a scaffold protein, mrip, yields a truncated form that is involved in repetitive bleb formation. *The FEBS Journal*, 2025.
95. Laura Santangelo, Giovanna Giurato, Claudia Cicchini, Claudio Montaldo, Carmela Mancone, Roberta Tarallo, et al. The rna-binding protein syncrip is a component of the hepatocyte exosomal machinery controlling microrna sorting. *Cell Reports*, 17(3):799–808, 2016.
96. Louis L. Scheffer, S. C. Sreetama, Nidhi Sharma, Shruti Medikayala, Kevin J. Brown, Aurelie Defour, and Jyoti K. Jaiswal. Mechanism of Ca^{2+} -triggered ESCRT assembly and regulation of cell membrane repair. *Nature Communications*, 5:5646, 2014.
97. Alessandra Scotto Rosato et al. Two-pore and TRPML cation channels: Regulators of phagocytosis, autophagy and lysosomal exocytosis. *Pharmacology and Therapeutics*, 220:107713, 2021.
98. Katsumori Segawa, Shuntaro Kurata, and Shigekazu Nagata. Human type IV P-type ATPases that work as plasma membrane phospholipid flippases and their regulation by caspase and calcium. *Journal of Biological Chemistry*, 291(2):762–772, 2016.
99. Iurii Semenov, Shu Xiao, and Andrei G. Pakhomov. Primary pathways of intracellular Ca^{2+} mobilization by nanosecond pulsed electric field. *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1828(3):981–989, 2013.
100. Iurii Semenov, Shu Xiao, Olga N. Pakhomova, and Andrei G. Pakhomov. Recruitment of the intracellular Ca^{2+} by ultrashort electric stimuli: The impact of pulse duration. *Cell Calcium*, 54(3):145–150, 2013.
101. Thomas R. Shannon, Fei Wang, Jose Puglisi, Christopher Weber, and Donald M. Bers. A mathematical treatment of integrated Ca^{2+} dynamics within the ventricular myocyte. *Biophysical Journal*, 87(5):3351–3371, 2004.
102. Saumya Shukla, Kevin P. Larsen, Congcong Ou, Kelly Rose, and James H. Hurley. In vitro reconstitution of calcium-dependent recruitment of the human ESCRT machinery in lysosomal membrane repair. *Proceedings of the National Academy of Sciences of the United States of America*, 119(35):e2205590119, 2022.
103. Matthew J. Shurtleff, Margaret M. Temoche-Diaz, Katherine V. Karfilis, Shun Ri, and Randy Schekman. Y-box protein 1 is required to sort micrornas into exosomes in cells and in a cell-free reaction. *eLife*, 5:e19276, 2016.
104. M. Silkunas, A. Enos, I. Semenov, G. Silkuniene, E. Zivla, O. N. Pakhomova, and A. G. Pakhomov. Peculiarities of phosphatidylserine externalization by nano- and microsecond electric pulses. *Journal of Membrane Biology*, 259(1):19, 2026.
105. Giedre Silkuniene, Uma M. Mangalanathan, Andrei G. Pakhomov, and Olga N. Pakhomova. Silencing of ATP1A1 attenuates cell membrane disruption by nanosecond electric pulses. *Biochemical and Biophysical Research Communications*, 677:93–97, 2023.
106. M. Singh, G. Mazaheri-Tehrani, I. Martin-Fabiani, and O. G. Davies. Electroporation induced changes in extracellular vesicle profile. *Drug Delivery*, 32(1):2562224, 2025.
107. Tore Skotland, Alicia Llorente, and Kirsten Sandvig. Lipids in extracellular vesicles: what can be learned about membrane structure and function? *Cold Spring Harbor Perspectives in Biology*, 15(8):a041415, 2023.
108. Kyle C. Smith and James C. Weaver. A model of electroporation-induced cellular uptake and membrane conductance changes. *Biophysical Journal*, 87(4):2219–2231, 2004.
109. Kyle C. Smith and James C. Weaver. Transmembrane molecular transport during versus after extremely large, nanosecond electric pulses. *Biochemical and Biophysical Research Communications*, 412(1):8–12, 2011.
110. Jachen A. Solinger, Daniel P. Ott, and Anne Spang. Escrting the rabs through conversion. *Biochemical Society Transactions*, 2025.
111. Stine L. Sonder, Theresa L. Boye, Rebecca Tolle, Joern Dengjel, Kenji Maeda, Marja Jaattela, Adam C. Simonsen, Jyoti K. Jaiswal, and Jesper Nylandsted. Annexin A7 is required for ESCRT III-mediated plasma membrane repair. *Scientific Reports*, 9(1):6726, 2019.
112. Jiayang Song, Allen L. Garner, and Ravindra P. Joshi. Effect of thermal gradients created by electromagnetic fields on cell-membrane electroporation probed by molecular-dynamics simulations. *Physical Review Applied*, 7:024003, 2017.
113. Jiayang Song, Ravindra P. Joshi, and Karl H. Schoenbach. Synergistic effects of local temperature enhancements on cellular responses in the context of high-intensity, ultrashort electric pulses. *Medical & Biological Engineering & Computing*, 49(6):713–718, 2011.
114. Wojciech Szlasa, Natalia Sauer, Dominika Baczynska, Michal Zietek, Katarzyna Haczkiwicz-Lesniak, Piotr Karpinski, Marcin Fleszar, Filip Fortuna, Marta J. Kulus, Aleksandra Piotrowska, Aleksandra Kmiecik, Agata Baranska, Olivier Michel, Vitalij Novickij, Mounir Tarek, Paulina Kasperkiewicz, Piotr Dziegiel, Marzena Podhorska-Okolow, Jolanta Saczko, and Julita Kulbacka. Pulsed electric field induces exocytosis and overexpression of mage antigens in melanoma. *Scientific Reports*, 14:12546, 2024.
115. U. Szwedowicz, R. Dębicka, K. J. Pawlik, D. Przystupski, and A. Choromańska. Reversible electroporation triggers the release of microvesicles affecting viability, migration, and adhesion of melanoma cells. *Biomedicine & Pharmacotherapy*, 198:119283, 2026.
116. Hirohide Takahashi, Jonathan R. Mayers, Lei Wang, J. Michael Edwardson, and Anjon Audhya. Hrs and stam function synergistically to bind ubiquitin-modified cargoes in vitro. *Biophysical Journal*, 2015.
117. Wei Tang et al. Rab27 molecular switch of tumor exosome secretion. *International Journal of Molecular Medicine*, 58:182, 2026.
118. Mounir Tarek. Membrane electroporation: A molecular dynamics simulation. *Biophysical Journal*, 88(6):4045–4053, 2005.
119. Amit Thakur and Devendra Rai. Global requirements for manufacturing and validation of clinical grade extracellular vesicles. *The Journal of Liquid Biopsy*, 6:100278, 2024.
120. D. Peter Tieleman. The molecular basis of electroporation. *BMC Biochemistry*, 5:10, 2004.
121. Rochelle Tixeira et al. Rock1 but not limk1 or pak2 is a key regulator of apoptotic membrane blebbing and cell

- disassembly. *Cell Death and Differentiation*, 27:102–116, 2020.
122. Meriem Tkach, Joanna Kowal, Anna E. Zucchetti, Lara Enserink, Mickael Jouve, Daniel Lankar, Michael Saitakis, Lorena Martin-Jaular, and Clotilde Thery. Qualitative differences in t-cell activation by dendritic cell-derived extracellular vesicle subtypes. *The EMBO Journal*, 36(20):3012–3028, 2017.
 123. Gleb P. Tolstikh, Hope T. Beier, Caleb C. Roth, Gary L. Thompson, and Bennett L. Ibey. 600 ns pulse electric field-induced phosphatidylinositol 4,5-bisphosphate depletion. *Bioelectrochemistry*, 100:80–87, 2014.
 124. Gleb P. Tolstikh, Hope T. Beier, Caleb C. Roth, Gary L. Thompson, Jason A. Payne, Marjorie A. Kuipers, and Bennett L. Ibey. Activation of intracellular phosphoinositide signaling after a single 600 nanosecond electric pulse. *Bioelectrochemistry*, 94:23–29, 2013.
 125. Edwin van der Pol, Frank A. W. Coumans, Andre E. Grootemaat, Chris Gardiner, Ian L. Sargent, Paul Harrison, Auguste Sturk, Ton G. van Leeuwen, and Rienk Nieuwland. Particle size distribution of exosomes and microvesicles determined by transmission electron microscopy, flow cytometry, nanoparticle tracking analysis, and resistive pulse sensing. *Journal of Thrombosis and Haemostasis*, 12(7):1182–1192, 2014.
 126. Els J. van der Vlist, Ger J. A. Arkesteijn, Chris H. A. van de Lest, Willem Stoorvogel, Esther N. M. Nolte-’t Hoen, and Marca H. M. Wauben. CD4⁺ T cell activation promotes the differential release of distinct populations of nanosized vesicles. *Journal of Extracellular Vesicles*, 1:18364, 2012.
 127. Zlatko Vasilkoski, Annegret T. Esser, T. R. Gowrishankar, and James C. Weaver. Membrane electroporation: The absolute rate equation and nanosecond time scale pore creation. *Physical Review E*, 74(2):021904, 2006.
 128. Claudia Verderio, Michela Gabrielli, and Paola Giussani. Role of sphingolipids in the biogenesis and biological activity of extracellular vesicles. *Journal of Lipid Research*, 59(8):1325–1340, 2018.
 129. P. Thomas Vernier, Yinghua Sun, Laura Marcu, Soussan Salemi, Cheryl M. Craft, and Martin A. Gundersen. Calcium bursts induced by nanosecond electric pulses. *Biochemical and Biophysical Research Communications*, 310(2):286–295, 2003.
 130. P. Thomas Vernier, Matthew J. Ziegler, Yinghua Sun, Wenji V. Chang, Martin A. Gundersen, and D. Peter Tieleman. Nanopore formation and phosphatidylserine externalization in a phospholipid bilayer at high transmembrane potential. *Journal of the American Chemical Society*, 128(19):6288–6289, 2006.
 131. P. Thomas Vernier, Matthew J. Ziegler, Yinghua Sun, Martin A. Gundersen, and D. Peter Tieleman. Nanopore-facilitated, voltage-driven phosphatidylserine translocation in lipid bilayers: In cells and in silico. *Physical Biology*, 3(4):233–247, 2006.
 132. Carolina Villarroja-Beltri et al. Isgylation controls exosome secretion by promoting lysosomal degradation of mvb proteins. *Nature Communications*, 7:13588, 2016.
 133. Ariana K. von Lersner, Fabiane Fernandes, Patricia Midori Murobushi Ozawa, Marques Jackson, Matthieu Masurel, Hoangdung Ho, Sierra M. Lima, Tatyana Vagner, Bong Hwan Sung, Mohamed Wehbe, Kai Franze, Heather Pua, John T. Wilson, Jonathan M. Irish, Alissa M. Weaver, Dolores Di Vizio, and Andries Zijlstra. Multiparametric single-vesicle flow cytometry resolves extracellular vesicle heterogeneity and reveals selective regulation of biogenesis and cargo distribution. *ACS Nano*, 18, 2024.
 134. Hsin-Yi Wang, Alexandre Rumin, Milka Doktorova, Daniel Sputay, Sze H. Chan, Ann M. Wehman, Kandice R. Levental, and Ilya Levental. Loss of lipid asymmetry facilitates plasma membrane blebbing by decreasing membrane lipid packing. *Proceedings of the National Academy of Sciences of the United States of America*, 122(19):e2417145122, 2025.
 135. Sufen Wang, Jiexiao Chen, Meng-Tse Chen, P. Thomas Vernier, Martin A. Gundersen, and Miguel Valderrábano. Cardiac myocyte excitation by ultrashort high-field pulses. *Biophysical Journal*, 96(4):1640–1648, 2009.
 136. Douglas C. Watson, Benjamin C. Yung, Cristina Bergamaschi, Biswajit Chowdhury, Joseph Bear, Dimitrios Stellas, et al. Scalable, cgmp-compatible purification of extracellular vesicles carrying bioactive human heterodimeric il-15/lactadherin complexes. *Journal of Extracellular Vesicles*, 7(1):1442088, 2018.
 137. James Webber and Aled Clayton. How pure are your vesicles? *Journal of Extracellular Vesicles*, 2(1):19861, 2013.
 138. Joshua A. Welsh, Deborah C. I. Goberdhan, Lorraine O’Driscoll, Edit I. Buzas, Cherie Blenkinsop, et al. Minimal information for studies of extracellular vesicles (MISEV2023): From basic to advanced approaches. *Journal of Extracellular Vesicles*, 13(2):e12404, 2024.
 139. Daniel Wiczew, Natalia Szulc, and Mounir Tarek. Molecular dynamics simulations of the effects of lipid oxidation on the permeability of cell membranes. *Bioelectrochemistry*, 141:107869, 2021.
 140. Justin Krish Williams, Jordan Matthew Ngo, Isabelle Madeleine Lehman, and Randy Schekman. Annexin a6 mediates calcium-dependent exosome secretion during plasma membrane repair. *eLife*, 12:e86556, 2023.
 141. Justin Krish Williams, Jordan Matthew Ngo, Anand Murugupandian, Dorothy E. Croall, H. Criss Hartzell, and Randy Schekman. Calpains orchestrate secretion of annexin-containing microvesicles during membrane repair. *Journal of Cell Biology*, 224(7):e202408159, 2025.
 142. Thomas Wollert and James H. Hurley. Molecular mechanism of multivesicular body biogenesis by escrt complexes. *Nature*, 464(7290):864–869, 2010.
 143. Sang Su Woo et al. Munc13-4 functions as a ca2+ sensor for homotypic secretory granule fusion to generate endosomal exocytic vacuoles. *Molecular Biology of the Cell*, 2017.
 144. Zeyao Yan, Li Yin, Chunjing Hao, Kefu Liu, and Jian Qiu. Synergistic effect of pulsed electric fields and temperature on the inactivation of microorganisms. *AMB Express*, 11(1):47, 2021.
 145. Qi Yang, Shinji Kajimoto, Yuki Kobayashi, Hirotsugu Hiramatsu, and Takakazu Nakabayashi. Regulation of cell volume by nanosecond pulsed electric fields. *The Journal of Physical Chemistry B*, 125(38):10692–10700, 2021.
 146. Sung Hae Yun, Vasilii Mansurov, Lisha Yang, Jihwan Yoon, Normand Leblanc, Gale L. Craviso, and Josette Zaklit. Modulating Ca²⁺ influx into adrenal chromaffin cells with short-duration nanosecond electric pulses. *Biophysical Journal*, 123(16):2537–2556, 2024.

147. Andrea Zandrini, Giorgia Guerra, Krizia Sagini, Tatyana Vagner, Dolores Di Vizio, and Paolo Bergese. On the surface-to-bulk partition of proteins in extracellular vesicles. *Colloids and Surfaces B: Biointerfaces*, 218:112728, 2022.
148. Wenjun Zhang, Rong Liu, Yuhua Chen, Minghua Wang, and Juan Du. Crosstalk between oxidative stress and exosomes. *Oxidative Medicine and Cellular Longevity*, 2022:3553617, 2022.