

Irreducibility and convergence of nonlinear state-space models of Markov chains

Application: CMA-ES

Armand Gissler

Tuesday 3rd October, 2023

CMA-ES, École polytechnique & Inria
(Supervisors: Anne Auger & Nikolaus Hansen)



Inria

Control model [Meyn & Caines 1991, Meyn & Tweedie 2012]

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\text{CM}(\mathbf{F}))$$

Control model [Meyn & Caines 1991, Meyn & Tweedie 2012]

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\text{CM}(F))$$

- $F: X \times V \rightarrow X$ **smooth** (\mathcal{C}^∞)

Control model [Meyn & Caines 1991, Meyn & Tweedie 2012]

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\text{CM}(F))$$

- $F: X \times V \rightarrow X$ **smooth** (\mathcal{C}^∞)
- $\{v_{k+1}\}$ i.i.d. with a l.s.c. density p

Control model [Meyn & Caines 1991, Meyn & Tweedie 2012]

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\text{CM}(F))$$

- $F: X \times V \rightarrow X$ **smooth** (\mathcal{C}^∞)
- $\{v_{k+1}\}$ i.i.d. with a l.s.c. density p
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\text{CM}(F))$$

- $F: X \times V \rightarrow X$ **smooth** (\mathcal{C}^∞)
- $\{v_{k+1}\}$ i.i.d. with a l.s.c. density p
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\mathbf{CM}(\mathbf{F}))$$

- $F: X \times V \rightarrow X$ **smooth** (\mathcal{C}^∞)
- $\{v_{k+1}\}$ i.i.d. with a l.s.c. density p
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

- (i) *If $\mathbf{CM}(\mathbf{F})$ is forward accessible, then $\{\theta_k\}_{k \in \mathbb{N}}$ is a T -chain ;*

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\mathbf{CM}(\mathbf{F}))$$

- $F: X \times V \rightarrow X$ **smooth** (\mathcal{C}^∞)
- $\{v_{k+1}\}$ i.i.d. with a l.s.c. density p
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

- (i) If $\mathbf{CM}(\mathbf{F})$ is **forward accessible**, then $\{\theta_k\}_{k \in \mathbb{N}}$ is a *T-chain* ;
- (ii) then, $\{\theta_k\}_{k \in \mathbb{N}}$ is **irreducible** \Leftrightarrow there exists a **globally attracting state** ;

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\mathbf{CM}(\mathbf{F}))$$


- $F: X \times V \rightarrow X$ **smooth** (\mathcal{C}^∞)
- $\{v_{k+1}\}$ i.i.d. with a l.s.c. density p
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

- (i) *If $\mathbf{CM}(\mathbf{F})$ is forward accessible, then $\{\theta_k\}_{k \in \mathbb{N}}$ is a T -chain ;*
- (ii) *then, $\{\theta_k\}_{k \in \mathbb{N}}$ is irreducible \Leftrightarrow there exists a **globally attracting state** ;*
- (iii) *the (a)periodicity of $\mathbf{CM}(\mathbf{F})$ is equivalent to the (a)periodicity of $\{\theta_k\}_{k \in \mathbb{N}}$.*


Forward accessibility

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\mathbf{CM}(\mathbf{F}))$$



Forward accessibility


$$\begin{aligned}\theta_{k+1} &= F(\theta_k, v_{k+1}) && (\mathbf{CM}(\mathbf{F})) \\ &= F_{k+1}(\theta_0, v_1, \dots, v_{k+1})\end{aligned}$$



Forward accessibility

$$\begin{aligned}\theta_{k+1} &= F(\theta_k, v_{k+1}) && (\mathbf{CM}(\mathbf{F})) \\ &= F_{k+1}(\theta_0, v_1, \dots, v_{k+1})\end{aligned}$$

for $v_1, \dots, v_{k+1} \in \mathcal{O} = \text{supp } p$.




Forward accessibility

$$\begin{aligned}\theta_{k+1} &= F(\theta_k, v_{k+1}) && (\mathbf{CM}(\mathbf{F})) \\ &= F_{k+1}(\theta_0, v_1, \dots, v_{k+1})\end{aligned}$$

for $v_1, \dots, v_{k+1} \in \mathcal{O} = \text{supp } p$.

$$A_+(\theta) = \{F_k(\theta, v_1, \dots, v_k) \mid k \in \mathbb{N}, v_1, \dots, v_k \in \mathcal{O}\}$$

$\mathbf{CM}(\mathbf{F})$ is *forward accessible* when $A_+(\theta)$ has nonempty interior



Forward accessibility


$$\begin{aligned}\theta_{k+1} &= F(\theta_k, v_{k+1}) && (\mathbf{CM}(\mathbf{F})) \\ &= F_{k+1}(\theta_0, v_1, \dots, v_{k+1})\end{aligned}$$

for $v_1, \dots, v_{k+1} \in \mathcal{O} = \text{supp } p$.

$$A_+(\theta) = \{F_k(\theta, v_1, \dots, v_k) \mid k \in \mathbb{N}, v_1, \dots, v_k \in \mathcal{O}\}$$

$\mathbf{CM}(\mathbf{F})$ is *forward accessible* when $A_+(\theta)$ has nonempty interior

$$C_k = \text{Jac}_{v_{1..k}} F_k(\theta, v_1, \dots, v_k)$$



Forward accessibility

$$\begin{aligned}\theta_{k+1} &= F(\theta_k, v_{k+1}) && (\mathbf{CM}(\mathbf{F})) \\ &= F_{k+1}(\theta_0, v_1, \dots, v_{k+1})\end{aligned}$$

for $v_1, \dots, v_{k+1} \in \mathcal{O} = \text{supp } p$.

$$A_+(\theta) = \{F_k(\theta, v_1, \dots, v_k) \mid k \in \mathbb{N}, v_1, \dots, v_k \in \mathcal{O}\}$$


$\mathbf{CM}(\mathbf{F})$ is *forward accessible* when $A_+(\theta)$ has nonempty interior

$$C_k = \text{Jac}_{v_{1..k}} F_k(\theta, v_1, \dots, v_k)$$

Then:

$\mathbf{CM}(\mathbf{F})$ is forward accessible

$$\Leftrightarrow \forall \theta, \exists v_1, \dots, v_k \in \mathcal{O}, \text{rank } C_k = n.$$




Globally attracting state

x^* is globally attracting when

$\forall \theta_0, \exists v_1, \dots, v_k \in \mathcal{O} = \text{supp } p,$

$F_k(\theta_0, v_1, \dots, v_k) \in \text{Neighborhood}(x^*)$




Globally attracting state

x^* is globally attracting when

$$\forall \theta_0, \exists v_1, \dots, v_k \in \mathcal{O} = \text{supp } p,$$

$$F_k(\theta_0, v_1, \dots, v_k) \in \text{Neighborhood}(x^*)$$



If

$$C_k(x^*, v_1, \dots, v_k) = \text{Jac}_{v_{1..k}} F_k(x^*, v_1, \dots, v_k)$$


is of rank n

Globally attracting state

x^* is globally attracting when

$\forall \theta_0, \exists v_1, \dots, v_k \in \mathcal{O} = \text{supp } p,$

$$F_k(\theta_0, v_1, \dots, v_k) \in \text{Neighborhood}(x^*)$$



If

$$C_k(x^*, v_1, \dots, v_k) = \text{Jac}_{v_{1..k}} F_k(x^*, v_1, \dots, v_k)$$

is of rank n ,

then $\forall \theta, \exists u_1, \dots, u_j \in \mathcal{O},$

$$\text{rank } C_k(\theta, u_1, \dots, u_j) = n$$

Aperiodicity of $(\mathbf{CM}(\mathbf{F}))$

$(\mathbf{CM}(\mathbf{F}))$ is aperiodic $\Leftrightarrow \exists$ a steadily attracting state


Aperiodicity of $(\mathbf{CM}(\mathbf{F}))$

$(\mathbf{CM}(\mathbf{F}))$ is aperiodic $\Leftrightarrow \exists$ a steadily attracting state

θ^* is steadily attracting when for any $\theta_0 \in X$,

$\exists v_1, v_2, \dots \in \mathcal{O} = \text{supp } p$,

$$\lim_{k \rightarrow \infty} F_k(\theta_0, v_1, \dots, v_k) = \theta^*$$



$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\mathbf{CM}(\mathbf{F}))$$

- $F: X \times V \rightarrow X$ **smooth** (\mathcal{C}^∞)
- $\{v_{k+1}\}$ i.i.d. with a l.s.c. density p
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

- (i) *If $\mathbf{CM}(\mathbf{F})$ is forward accessible, then $\{\theta_k\}_{k \in \mathbb{N}}$ is a T -chain ;*
- (ii) *then, $\{\theta_k\}_{k \in \mathbb{N}}$ is irreducible \Leftrightarrow there exists a **globally attracting state** ;*
- (iii) *the (a)periodicity of $\mathbf{CM}(\mathbf{F})$ is equivalent to the (a)periodicity of $\{\theta_k\}_{k \in \mathbb{N}}$.*

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\text{CM(F)})$$

- $F: X \times V \rightarrow X$ **smooth** (\mathcal{C}^∞)
- $\{v_{k+1}\}$ i.i.d. with a l.s.c. density p
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

Suppose that θ^* is steadily attracting, i.e., for $\theta_0 \in X$,

$$\exists v_1, v_2, \dots \in \mathcal{O} = \text{supp } p, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists v_1, \dots, v_k \in \mathcal{O}$, such that

$$C_k = \text{Jac}_{v_1..k} F_j(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T-chain.

Example

$$\theta_{k+1} = \theta_k + v_{k+1}$$

with $v_{k+1} \sim \mathcal{N}(0, I_d)$.

Example

$$\theta_{k+1} = \theta_k + v_{k+1}$$

with $v_{k+1} \sim \mathcal{N}(0, I_d)$.

Then: 0 is globally attracting.

Example

$$\theta_{k+1} = \theta_k + v_{k+1}$$

with $v_{k+1} \sim \mathcal{N}(0, I_d)$.

Then: 0 is globally attracting.

Proof.

Set $v_1 = -\theta_0 \in \text{supp } p_{\mathcal{N}}$.

□

Example

$$\theta_{k+1} = \theta_k + v_{k+1}$$

with $v_{k+1} \sim \mathcal{N}(0, I_d)$.

Then: 0 is globally attracting.

Proof.

Set $v_1 = -\theta_0 \in \text{supp } p_{\mathcal{N}}$.

□

Then: $\text{rank Jac}_v F(0, v) = n$

Example

$$\theta_{k+1} = \theta_k + v_{k+1}$$

with $v_{k+1} \sim \mathcal{N}(0, I_d)$.

Then: 0 is globally attracting.

Proof.

Set $v_1 = -\theta_0 \in \text{supp } p_{\mathcal{N}}$.

□

Then: $\text{rank Jac}_v F(0, v) = n$

Proof.

$\text{Jac}_v F(0, v) = I_n$

□

Control model [Chotard & Auger 2019]

Consider

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\mathbf{CM(F)})$$

- $F: X \times V \rightarrow X$ **smooth** (\mathcal{C}^∞)
- $\{v_{k+1}\}$ i.i.d. with a l.s.c. density p
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

Suppose that θ^ is steadily attracting, i.e., for $\theta_0 \in X$,*

$$\exists v_1, v_2, \dots \in \mathcal{O}, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists v_1, \dots, v_k \in \mathcal{O}$, such that

$$C_k = \text{Jac}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T-chain.

Control model [Chotard & Auger 2019]

Consider

$$\theta_{k+1} = F(\theta_k, v_{k+1}) \quad (\mathbf{CM(F)})$$

- $F: X \times V \rightarrow X$ is \mathcal{C}^1
- $\{v_{k+1}\}$ i.i.d. with a l.s.c. density p
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

Suppose that θ^ is steadily attracting, i.e., for $\theta_0 \in X$,*

$$\exists v_1, v_2, \dots \in \mathcal{O}, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists v_1, \dots, v_k \in \mathcal{O}$, such that

$$C_k = \text{Jac}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T-chain.

Control model [Chotard & Auger 2019]

Consider $\theta_{k+1} = F(\theta_k, v_{k+1})$ **(CM(F))**

- $F: X \times V \rightarrow X$ is \mathcal{C}^1
- $\{v_{k+1}\}$ such that $v_{k+1} \sim p_{\theta_k}$ l.s.c.
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ open.

Theorem

Suppose that θ^* is steadily attracting, i.e., for $\theta_0 \in X$,

$$\exists v_1, v_2, \dots \in \mathcal{O}, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists v_1, \dots, v_k \in \mathcal{O}$, such that

$$C_k = \text{Jac}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T-chain.

Control model [Chotard & Auger 2019]

Consider

$$\theta_{k+1} = F(\theta_k, \alpha(\theta_k, u_{k+1})) \quad (\mathbf{CM(F)})$$

- $F: X \times V \rightarrow X$ is C^1
- $\{v_{k+1}\}$ such that $v_{k+1} = \alpha(\theta_k, u_{k+1}) \sim p_{\theta_k}$ l.s.c.
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ open.

Theorem

Suppose that θ^* is steadily attracting, i.e., for $\theta_0 \in X$,

$$\exists v_1, v_2, \dots \in \mathcal{O}, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists v_1, \dots, v_k \in \mathcal{O}$, such that

$$C_k = \text{Jac}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T-chain.

Control model [Chotard & Auger 2019]

Consider

$$\theta_{k+1} = F(\theta_k, \alpha(\theta_k, u_{k+1})) \quad (\mathbf{CM(F)})$$

- $F: X \times V \rightarrow X$ is C^1
- $\{v_{k+1}\}$ such that $v_{k+1} = \alpha(\theta_k, u_{k+1}) \sim p_{\theta_k}$ l.s.c.
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ open.

Theorem

Suppose that θ^* is steadily attracting, i.e., for $\theta_0 \in X$,

$$\exists (v_1, v_2, \dots) \in \mathcal{O}_{\theta_0}^\infty, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists (v_1, \dots, v_k) \in \mathcal{O}_{\theta^*}^k$, such that

$$C_k = \text{Jac}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T-chain.

Application: Evolution Strategies (ES)

$$\text{Find } x^* \in \arg \min_{x \in \mathbb{R}^d} f(x) \quad (\text{P})$$

Application: Evolution Strategies (ES)

$$\text{Find } x^* \in \arg \min_{x \in \mathbb{R}^d} f(x) \quad (\text{P})$$

ES approximate x^* by $\mathcal{N}(m_k, \sigma_k^2 I_d)$

Application: Evolution Strategies (ES)


$$\text{Find } x^* \in \arg \min_{x \in \mathbb{R}^d} f(x) \quad (\text{P})$$

ES approximate x^* by $\mathcal{N}(m_k, \sigma_k^2 I_d)$ by updating $\theta_k = (m_k, \sigma_k) \in \mathbb{R}^d \times \mathbb{R}_{++}$.

Algorithm: ES with stepsize adaptation

Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$




Algorithm: ES with stepsize adaptation

Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$



Algorithm: ES with stepsize adaptation


Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k u_{k+1}^i \sim \mathcal{N}(m_k, \sigma_k^2 I_d)$



Algorithm: ES with stepsize adaptation


Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k u_{k+1}^i \sim \mathcal{N}(m_k, \sigma_k^2 I_d)$
2. sort $f(x_{k+1}^i)$:



Algorithm: ES with stepsize adaptation


Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k u_{k+1}^i \sim \mathcal{N}(m_k, \sigma_k^2 I_d)$
2. sort $f(x_{k+1}^i)$:
 $f(x_{k+1}^{1:\lambda}) \leq \dots \leq f(x_{k+1}^{\lambda:\lambda})$



Algorithm: ES with stepsize adaptation


Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k u_{k+1}^i \sim \mathcal{N}(m_k, \sigma_k^2 I_d)$
2. sort $f(x_{k+1}^i)$:
 $f(x_{k+1}^{1:\lambda}) \leq \dots \leq f(x_{k+1}^{\lambda:\lambda})$
3. $m_{k+1} = \sum w_i x_{k+1}^{i:\lambda}$



Algorithm: ES with stepsize adaptation


Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k u_{k+1}^i \sim \mathcal{N}(m_k, \sigma_k^2 I_d)$
2. sort $f(x_{k+1}^i)$:
 $f(x_{k+1}^{1:\lambda}) \leq \dots \leq f(x_{k+1}^{\lambda:\lambda})$
3. $m_{k+1} = \sum w_i x_{k+1}^{i:\lambda} = x_k + \sigma_k \sum w_i u_{k+1}^{i:\lambda}$



Algorithm: ES with stepsize adaptation


Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k u_{k+1}^i \sim \mathcal{N}(m_k, \sigma_k^2 I_d)$
 2. sort $f(x_{k+1}^i)$:
 $f(x_{k+1}^{1:\lambda}) \leq \dots \leq f(x_{k+1}^{\lambda:\lambda})$
 3. $m_{k+1} = \sum w_i x_{k+1}^{i:\lambda} = x_k + \sigma_k \sum w_i u_{k+1}^{i:\lambda}$
 4. $\log \sigma_{k+1} = \log \sigma_k + \frac{\|\sum w_i u_{k+1}^{i:\lambda}\|}{\mathbb{E} \|\sum w_i u_{k+1}^i\|} - 1$
-



Algorithm: ES with stepsize adaptation


Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k u_{k+1}^i \sim \mathcal{N}(m_k, \sigma_k^2 I_d)$
 2. sort $f(x_{k+1}^i)$:
 $f(x_{k+1}^{1:\lambda}) \leq \dots \leq f(x_{k+1}^{\lambda:\lambda})$
 3. $m_{k+1} = \sum w_i x_{k+1}^{i:\lambda} = x_k + \sigma_k \sum w_i u_{k+1}^{i:\lambda}$
 4. $\log \sigma_{k+1} = \log \sigma_k + \frac{\|\sum w_i u_{k+1}^{i:\lambda}\|^2}{\mathbb{E} \|\sum w_i u_{k+1}^i\|^2} - 1$
-



Algorithm: ES with stepsize adaptation


Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k u_{k+1}^i \sim \mathcal{N}(m_k, \sigma_k^2 I_d)$
 2. sort $f(x_{k+1}^i)$:
 $f(x_{k+1}^{1:\lambda}) \leq \dots \leq f(x_{k+1}^{\lambda:\lambda})$
 3. $m_{k+1} = \sum w_i x_{k+1}^{i:\lambda} = x_k + \sigma_k \sum w_i u_{k+1}^{i:\lambda}$
 4. $\sigma_{k+1} = \sigma_k \times \exp\left(\frac{\|\sum w_i u_{k+1}^{i:\lambda}\|^2}{\mathbb{E}\|\sum w_i u_{k+1}^i\|^2} - 1\right)$
-



Convergence via analysis of CM(F)

$$\theta_k = (m_k, \sigma_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0)$$

Convergence via analysis of CM(F)

$$\theta_k = (m_k, \sigma_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0)$$

\Rightarrow no stationary measure

Convergence via analysis of CM(F)

$$\theta_k = (m_k, \sigma_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0)$$

\Rightarrow no stationary measure

$$z_k \stackrel{\text{def}}{=} \frac{m_k - x^*}{\sigma_k}$$

Convergence via analysis of CM(F)

$$\theta_k = (m_k, \sigma_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0)$$

\Rightarrow no stationary measure

$$z_k \stackrel{\text{def}}{=} \frac{m_k - x^*}{\sigma_k}$$

$$z_{k+1} = F(z_k, v_{k+1})$$

Convergence via analysis of CM(F)

$$\theta_k = (m_k, \sigma_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0)$$

\Rightarrow no stationary measure

$$z_k \stackrel{\text{def}}{=} \frac{m_k - x^*}{\sigma_k}$$

$$\begin{aligned} z_{k+1} &= F(z_k, v_{k+1}) \\ &= \exp\left(1 - \frac{\|w^T v_{k+1}\|^2}{\mathbb{E}\|w^T u_{k+1}\|^2}\right) \times (z_k + w^T v_{k+1}) \end{aligned}$$

Convergence via analysis of CM(F)

$$\theta_k = (m_k, \sigma_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0)$$

\Rightarrow no stationary measure

$$z_k \stackrel{\text{def}}{=} \frac{m_k - x^*}{\sigma_k}$$

$$\begin{aligned} z_{k+1} &= F(z_k, v_{k+1}) \\ &= \exp\left(1 - \frac{\|w^T v_{k+1}\|^2}{\mathbb{E}\|w^T u_{k+1}\|^2}\right) \times (z_k + w^T v_{k+1}) \end{aligned}$$

with $v_{k+1} = (u_{k+1}^{1:\lambda}, \dots, u_{k+1}^{\lambda:\lambda})$.

Control model [Chotard & Auger 2019]

Consider $\theta_{k+1} = F(\theta_k, \alpha(\theta_k, u_{k+1}))$ (CM(F))

- $F: X \times V \rightarrow X$ is C^1
- $\{v_{k+1}\}$ such that $v_{k+1} = \alpha(\theta_k, u_{k+1}) \sim p_{\theta_k}$ l.s.c.
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ open.

Theorem

Suppose that θ^* is steadily attracting, i.e., for $\theta_0 \in X$,

$$\exists (v_1, v_2, \dots) \in \mathcal{O}_{\theta_0}^\infty, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists (v_1, \dots, v_k) \in \mathcal{O}_{\theta^*}^k$, such that

$$C_k = \text{Jac}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T-chain.

$$\begin{aligned} z_{k+1} &= F(z_k, v_{k+1}) \\ &= \exp\left(1 - \frac{\|w^T v_{k+1}\|^2}{\mathbb{E}\|w^T u_{k+1}\|^2}\right) \times (z_k + w^T v_{k+1}) \end{aligned}$$

$$\begin{aligned} z_{k+1} &= F(z_k, v_{k+1}) \\ &= \exp\left(1 - \frac{\|w^T v_{k+1}\|^2}{\mathbb{E}\|w^T u_{k+1}\|^2}\right) \times (z_k + w^T v_{k+1}) \end{aligned}$$

Then: 0 is globally attracting.

$$\begin{aligned} z_{k+1} &= F(z_k, v_{k+1}) \\ &= \exp\left(1 - \frac{\|w^T v_{k+1}\|^2}{\mathbb{E}\|w^T u_{k+1}\|^2}\right) \times (z_k + w^T v_{k+1}) \end{aligned}$$

Then: 0 is globally attracting.

Proof.



$$\begin{aligned} z_{k+1} &= F(z_k, v_{k+1}) \\ &= \exp\left(1 - \frac{\|w^T v_{k+1}\|^2}{\mathbb{E}\|w^T u_{k+1}\|^2}\right) \times (z_k + w^T v_{k+1}) \end{aligned}$$

Then: 0 is globally attracting.

Proof.

If $v_1 = (-z_0, \dots, -z_0)$,



$$\begin{aligned} z_{k+1} &= F(z_k, v_{k+1}) \\ &= \exp\left(1 - \frac{\|w^T v_{k+1}\|^2}{\mathbb{E}\|w^T u_{k+1}\|^2}\right) \times (z_k + w^T v_{k+1}) \end{aligned}$$

Then: 0 is globally attracting.

Proof.

If $v_1 = (-z_0, \dots, -z_0)$,

then, $z_1 = \exp(\dots) \times (z_0 - \sum w_i z_0) = 0$. □

Irreducibility of ES

$$\begin{aligned} z_{k+1} &= F(z_k, v_{k+1}) \\ &= \exp\left(1 - \frac{\|w^T v_{k+1}\|^2}{\mathbb{E}\|w^T u_{k+1}\|^2}\right) \times (z_k + w^T v_{k+1}) \end{aligned}$$

Then: 0 is globally attracting.

Proof.

If $v_1 = (-z_0, \dots, -z_0)$,

then, $z_1 = \exp(\dots) \times (z_0 - \sum w_i z_0) = 0$. □

Then: $\text{rank Jac}_v F(0, 0) = n$.

Proof. □

Irreducibility of ES

$$\begin{aligned} z_{k+1} &= F(z_k, v_{k+1}) \\ &= \exp\left(1 - \frac{\|w^T v_{k+1}\|^2}{\mathbb{E}\|w^T u_{k+1}\|^2}\right) \times (z_k + w^T v_{k+1}) \end{aligned}$$

Then: 0 is globally attracting.

Proof.

If $v_1 = (-z_0, \dots, -z_0)$,

then, $z_1 = \exp(\dots) \times (z_0 - \sum w_i z_0) = 0$. □

Then: $\text{rank Jac}_v F(0, 0) = n$.

Proof.

$F(0, 0 + h) = \exp(1 + O(h)) \times w^T h$ □

Irreducibility of ES

$$\begin{aligned} z_{k+1} &= F(z_k, v_{k+1}) \\ &= \exp\left(1 - \frac{\|w^T v_{k+1}\|^2}{\mathbb{E}\|w^T u_{k+1}\|^2}\right) \times (z_k + w^T v_{k+1}) \end{aligned}$$

Then: 0 is globally attracting.

Proof.

If $v_1 = (-z_0, \dots, -z_0)$,

then, $z_1 = \exp(\dots) \times (z_0 - \sum w_i z_0) = 0$. □

Then: $\text{rank Jac}_v F(0, 0) = n$.

Proof.

$F(0, 0 + h) = \exp(1 + O(h)) \times w^T h = w^T h + o(h)$. □

Convergence

1. $\{z_k\}_{k \in \mathbb{N}}$ is **irreducible** and **aperiodic**

Convergence

1. $\{z_k\}_{k \in \mathbb{N}}$ is **irreducible** and **aperiodic**
2. we have a **drift** condition with

$$V(z) = \|z\|^\alpha$$

Convergence

1. $\{z_k\}_{k \in \mathbb{N}}$ is **irreducible** and **aperiodic**
2. we have a **drift** condition with

$$V(z) = \|z\|^\alpha$$

$\Rightarrow \{z_k\}_{k \in \mathbb{N}}$ admits a **unique invariant probability measure** π

Convergence

1. $\{z_k\}_{k \in \mathbb{N}}$ is **irreducible** and **aperiodic**
2. we have a **drift** condition with

$$V(z) = \|z\|^\alpha$$

$\Rightarrow \{z_k\}_{k \in \mathbb{N}}$ admits a **unique invariant probability measure** π and

$$\begin{aligned} & \lim_{t \rightarrow \infty} \frac{1}{t} \log \frac{\|m_t - x^*\|}{\|m_0 - x^*\|} \\ &= \lim_{t \rightarrow \infty} \frac{1}{t} \sum_{k=0}^{t-1} \log \frac{\|z_{k+1}\|}{\|z_k\|} - \log \exp \left(\frac{\|\sum w_i u_{k+1}^{i:\lambda}\|^2}{\mathbb{E} \|\sum w_i u_{k+1}^i\|^2} - 1 \right) \end{aligned}$$

Convergence

1. $\{z_k\}_{k \in \mathbb{N}}$ is **irreducible** and **aperiodic**
2. we have a **drift** condition with

$$V(z) = \|z\|^\alpha$$

$\Rightarrow \{z_k\}_{k \in \mathbb{N}}$ admits a **unique invariant probability measure** π and

$$\begin{aligned} & \lim_{t \rightarrow \infty} \frac{1}{t} \log \frac{\|m_t - x^*\|}{\|m_0 - x^*\|} \\ &= \lim_{t \rightarrow \infty} \frac{1}{t} \sum_{k=0}^{t-1} \log \frac{\|z_{k+1}\|}{\|z_k\|} - \log \exp \left(\frac{\|\sum w_i u_{k+1}^{i:\lambda}\|^2}{\mathbb{E} \|\sum w_i u_{k+1}^i\|^2} - 1 \right) \\ &= -\mathbb{E}_\pi \left[\frac{\|\sum w_i u^{i:\lambda}\|^2}{\mathbb{E} \|\sum w_i u^i\|^2} - 1 \mid u^1, \dots, u^\lambda \sim \mathcal{N}_d \right] \end{aligned}$$

Convergence

1. $\{z_k\}_{k \in \mathbb{N}}$ is **irreducible** and **aperiodic**
2. we have a **drift** condition with

$$V(z) = \|z\|^\alpha$$

$\Rightarrow \{z_k\}_{k \in \mathbb{N}}$ admits a **unique invariant probability measure** π and

$$\begin{aligned} & \lim_{t \rightarrow \infty} \frac{1}{t} \log \frac{\|m_t - x^*\|}{\|m_0 - x^*\|} \\ &= \lim_{t \rightarrow \infty} \frac{1}{t} \sum_{k=0}^{t-1} \log \frac{\|z_{k+1}\|}{\|z_k\|} - \log \exp \left(\frac{\|\sum w_i u_{k+1}^{i:\lambda}\|^2}{\mathbb{E} \|\sum w_i u_{k+1}^i\|^2} - 1 \right) \\ &= -\mathbb{E}_\pi \left[\frac{\|\sum w_i u^{i:\lambda}\|^2}{\mathbb{E} \|\sum w_i u^i\|^2} - 1 \mid u^1, \dots, u^\lambda \sim \mathcal{N}_d \right] \end{aligned}$$

Conclusion: ES converges *linearly* to x^*

$$\theta_{k+1} = F(\theta_k, \alpha(\theta_k, u_{k+1})) \quad (\mathbf{CM(F)})$$

- $F: X \times V \rightarrow X$ is C^1
- $\{v_{k+1}\}$ such that $v_{k+1} = \alpha(\theta_k, u_{k+1}) \sim p_{\theta_k}$ l.s.c.
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

Suppose that θ^ is steadily attracting, i.e., for $\theta_0 \in X$,*

$$\exists (v_1, v_2, \dots) \in \mathcal{O}_{\theta_0}^\infty, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists (v_1, \dots, v_k) \in \mathcal{O}_{\theta^}^k$, such that*

$$C_k = \text{Jac}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T-chain.

$$\theta_{k+1} = F(\theta_k, \alpha(\theta_k, u_{k+1})) \quad (\mathbf{CM(F)})$$

- $F: X \times V \rightarrow X$ is **locally Lipschitz**
- $\{v_{k+1}\}$ such that $v_{k+1} = \alpha(\theta_k, u_{k+1}) \sim p_{\theta_k}$ l.s.c.
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

Suppose that θ^ is steadily attracting, i.e., for $\theta_0 \in X$,*

$$\exists (v_1, v_2, \dots) \in \mathcal{O}_{\theta_0}^\infty, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists (v_1, \dots, v_k) \in \mathcal{O}_{\theta^}^k$, such that*

$$C_k = \text{Jac}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T-chain.

$$\theta_{k+1} = F(\theta_k, \alpha(\theta_k, u_{k+1})) \quad (\mathbf{CM(F)})$$

- $F: X \times V \rightarrow X$ is **locally Lipschitz**
- $\{v_{k+1}\}$ such that $v_{k+1} = \alpha(\theta_k, u_{k+1}) \sim p_{\theta_k}$ l.s.c.
- $X \subset \mathbb{R}^n$, $V \subset \mathbb{R}^p$ **open**.

Theorem

Suppose that θ^* is steadily attracting, i.e., for $\theta_0 \in X$,

$$\exists(v_1, v_2, \dots) \in \mathcal{O}_{\theta_0}^\infty, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists(v_1, \dots, v_k) \in \mathcal{O}_{\theta^*}^k$, such that **any**

$$C_k \in \text{Clarke}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T -chain.

$$\theta_{k+1} = F(\theta_k, \alpha(\theta_k, u_{k+1})) \quad (\text{CM(F)})$$

- $F: X \times V \rightarrow X$ is **locally Lipschitz**
- $\{v_{k+1}\}$ such that $v_{k+1} = \alpha(\theta_k, u_{k+1}) \sim p_{\theta_k}$ l.s.c.
- X and V are **manifolds**

Theorem

Suppose that θ^* is steadily attracting, i.e., for $\theta_0 \in X$,

$$\exists (v_1, v_2, \dots) \in \mathcal{O}_{\theta_0}^\infty, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists (v_1, \dots, v_k) \in \mathcal{O}_{\theta^*}^k$, such that **any**

$$C_k \in \text{Clarke}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T -chain.

Algorithm: ES with covariance matrix adaption


Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k u_{k+1}^i \sim \mathcal{N}(m_k, \sigma_k^2 I_d)$
2. sort $f(x_{k+1}^i)$:
 $f(x_{k+1}^{1:\lambda}) \leq \dots \leq f(x_{k+1}^{\lambda:\lambda})$
3. $m_{k+1} = \sum w_i x_{k+1}^{i:\lambda} = x_k + \sigma_k \sum w_i u_{k+1}^{i:\lambda}$
4. $\sigma_{k+1} = \sigma_k \times \exp\left(\frac{\|\sum w_i u_{k+1}^{i:\lambda}\|^2}{\mathbb{E}\|\sum w_i u_{k+1}^i\|^2} - 1\right)$



Algorithm: ES with covariance matrix adaption


Algorithm 1 ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k u_{k+1}^i \sim \mathcal{N}(m_k, \sigma_k^2 I_d)$
2. sort $f(x_{k+1}^i)$:
 $f(x_{k+1}^{1:\lambda}) \leq \dots \leq f(x_{k+1}^{\lambda:\lambda})$
3. $m_{k+1} = \sum w_i x_{k+1}^{i:\lambda} = x_k + \sigma_k \sum w_i u_{k+1}^{i:\lambda}$
4. $\sigma_{k+1} = \sigma_k \times \exp\left(\frac{\|\sum w_i u_{k+1}^{i:\lambda}\|}{\mathbb{E}\|\sum w_i u_{k+1}^i\|} - 1\right)$



Algorithm: ES with covariance matrix adaption


Algorithm 1 CMA-ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$, $C_0 \in \mathcal{S}_{++}^d$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,
 $x_{k+1}^i = x_k + \sigma_k \sqrt{C_k} u_{k+1}^i$
2. sort $f(x_{k+1}^i)$:
 $f(x_{k+1}^{1:\lambda}) \leq \dots \leq f(x_{k+1}^{\lambda:\lambda})$
3. $m_{k+1} = \sum w_i x_{k+1}^{i:\lambda} = x_k + \sigma_k \sqrt{C_k} \sum w_i u_{k+1}^{i:\lambda}$
4. $\sigma_{k+1} = \sigma_k \times \exp\left(\frac{\|\sum w_i u_{k+1}^{i:\lambda}\|}{\mathbb{E}\|\sum w_i u_{k+1}^i\|} - 1\right)$



Algorithm: ES with covariance matrix adaption

Algorithm 1 CMA-ES

Goal: $\min_{x \in \mathbb{R}^d} f(x)$

Given: $m_0 \in \mathbb{R}^d$, $\sigma_0 > 0$, $C_0 \in \mathcal{S}_{++}^d$

For $k = 0, 1, 2, \dots$:

1. $u_{k+1}^1, \dots, u_{k+1}^\lambda \sim \mathcal{N}(0, I_d)$,

$$x_{k+1}^i = x_k + \sigma_k \sqrt{C_k} u_{k+1}^i$$


2. sort $f(x_{k+1}^i)$:

$$f(x_{k+1}^{1:\lambda}) \leq \dots \leq f(x_{k+1}^{\lambda:\lambda})$$

3. $m_{k+1} = \sum w_i x_{k+1}^{i:\lambda} = x_k + \sigma_k \sqrt{C_k} \sum w_i u_{k+1}^{i:\lambda}$

4. $\sigma_{k+1} = \sigma_k \times \exp\left(\frac{\|\sum w_i u_{k+1}^{i:\lambda}\|}{\mathbb{E}\|\sum w_i u_{k+1}^i\|} - 1\right)$

5. $C_{k+1} = (1-c)C_k + c \sum w_i [\sqrt{C_k} u_{k+1}^{i:\lambda}] [\sqrt{C_k} u_{k+1}^{i:\lambda}]^T$



CMA-ES follows CM(F)

$$\theta_k = (m_k, \sigma_k, C_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0, 0)$$

CMA-ES follows CM(F)

$$\theta_k = (m_k, \sigma_k, C_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0, 0)$$

\Rightarrow no stationary measure

CMA-ES follows CM(F)

$$\theta_k = (m_k, \sigma_k, C_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0, 0)$$

\Rightarrow no stationary measure

$$z_k \stackrel{\text{def}}{=} \frac{m_k - x^*}{\sigma_k}$$

CMA-ES follows CM(F)

$$\theta_k = (m_k, \sigma_k, C_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0, 0)$$

\Rightarrow no stationary measure

$$z_k \stackrel{\text{def}}{=} \frac{m_k - x^*}{\sigma_k \det^{1/2d}(C_k)}$$

CMA-ES follows CM(F)

$$\theta_k = (m_k, \sigma_k, C_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0, 0)$$

\Rightarrow no stationary measure

$$z_k \stackrel{\text{def}}{=} \frac{m_k - x^*}{\sigma_k \det^{1/2d}(C_k)}$$

$$\Sigma_k \stackrel{\text{def}}{=} \frac{C_k}{\det^{1/d}(C_k)}$$

CMA-ES follows CM(F)

$$\theta_k = (m_k, \sigma_k, C_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0, 0)$$

\Rightarrow no stationary measure

$$z_k \stackrel{\text{def}}{=} \frac{m_k - x^*}{\sigma_k \det^{1/2d}(C_k)}$$

$$\Sigma_k \stackrel{\text{def}}{=} \frac{C_k}{\det^{1/d}(C_k)} \in \det^{-1}(\{1\})$$

CMA-ES follows CM(F)

$$\theta_k = (m_k, \sigma_k, C_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0, 0)$$

\Rightarrow no stationary measure

$$z_k \stackrel{\text{def}}{=} \frac{m_k - x^*}{\sigma_k \det^{1/2d}(C_k)}$$

$$\Sigma_k \stackrel{\text{def}}{=} \frac{C_k}{\det^{1/d}(C_k)} \in \det^{-1}(\{1\})$$

$$(z_{k+1}, \Sigma_{k+1}) = F((z_k, \Sigma_k), v_{k+1})$$

CMA-ES follows CM(F)

$$\theta_k = (m_k, \sigma_k, C_k) \xrightarrow[k \rightarrow \infty]{} (x^*, 0, 0)$$

\Rightarrow no stationary measure

$$z_k \stackrel{\text{def}}{=} \frac{m_k - x^*}{\sigma_k \det^{1/2d}(C_k)}$$

$$\Sigma_k \stackrel{\text{def}}{=} \frac{C_k}{\det^{1/d}(C_k)} \in \det^{-1}(\{1\})$$

$$(z_{k+1}, \Sigma_{k+1}) = F((z_k, \Sigma_k), v_{k+1})$$

with $v_{k+1} = (u_{k+1}^{1:\lambda}, \dots, u_{k+1}^{\lambda:\lambda})$.

$$\theta_{k+1} = F(\theta_k, \alpha(\theta_k, u_{k+1})) \quad (\mathbf{CM(F)})$$

- $F: X \times V \rightarrow X$ is **locally Lipschitz**
- $\{v_{k+1}\}$ such that $v_{k+1} = \alpha(\theta_k, u_{k+1}) \sim p_{\theta_k}$ l.s.c.
- X and V are **manifolds**

Theorem

Suppose that θ^* is steadily attracting, i.e., for $\theta_0 \in X$,

$$\exists (v_1, v_2, \dots) \in \mathcal{O}_{\theta_0}^\infty, \quad \lim F_k(\theta_0, v_1, \dots, v_k) = \theta^*.$$

If $\exists (v_1, \dots, v_k) \in \mathcal{O}_{\theta^*}^k$, such that **any**

$$C_k \in \text{Clarke}_{v_{1..k}} F_k(\theta^*, v_1, \dots, v_k)$$

is of rank n , then $\{\theta_k\}_k$ is an irreducible aperiodic T -chain.

Corollary

Under additional assumptions on the objective function f ,

$$(z_k, \Sigma_k)_{k \in \mathbb{N}}$$

is a irreducible, aperiodic T -chain.

Then (z_k, Σ_k) is positive recurrent and follows a LLN if there exists a drift $V: X \rightarrow [0, +\infty]$ such that

$$\mathbb{E}[V(z_1, \Sigma_1)] \leq (1 - \varepsilon) \times V(z_0, \Sigma_0) + b \times \mathbf{1}_{(z_0, \Sigma_0) \in K}$$

Theorem (Drift for the normalized chain)

When minimizing a **spherical** function $f: x \mapsto g(x^T x)$ ($g: \mathbb{R} \rightarrow \mathbb{R}$ increasing), then the irreducible, aperiodic Markov chain $(z_k, \Sigma_k)_{k \in \mathbb{N}}$ satisfies a Foster-Lyapunov condition with the potential defined by

$$V(z, \Sigma) = \alpha \times \frac{\|\Sigma z\|^2}{\lambda_{\max}(\Sigma)^2} + \beta \times \lambda_{\max}(\Sigma)$$


Theorem (Drift for the normalized chain)

When minimizing a **spherical** function $f: x \mapsto g(x^T x)$ ($g: \mathbb{R} \rightarrow \mathbb{R}$ increasing), then the irreducible, aperiodic Markov chain $(z_k, \Sigma_k)_{k \in \mathbb{N}}$ satisfies a Foster-Lyapunov condition with the potential defined by

$$V(z, \Sigma) = \alpha \times \frac{\|\Sigma z\|^2}{\lambda_{\max}(\Sigma)^2} + \beta \times \lambda_{\max}(\Sigma)$$

This can be generalized to when minimizing **ellipsoid** functions $f(x) = g(x^T H x)$ using the **affine-invariance** of CMA-ES.

Affine-Invariance



Thank you!

Scaling-invariant functions

