

Evolutionary Modes

Positive Selection
(Diversifying)

$$dS < dN \text{ or } \omega := dN/dS > 1$$

Negative Selection

$$dS > dN \text{ or } \omega < 1$$

Neutral Evolution

$$dS \approx dN \text{ or } \omega \approx 1$$

Codon-substitution models

- In 1994, first tractable mechanistic evolutionary models for codon sequences were proposed by **Muse and Gaut** (MG94), and, independently, by **Goldman and Yang** (GY94) [in the same issue of MBE, back to back]
- Markov models of codon substitution provide a powerful framework for **estimating substitution rates** from coding sequence data, as they
 - *encode our mechanistic understanding of the evolutionary process,*
 - *enable one to compute the phylogenetic likelihood,*
 - *permit hypothesis testing or Bayesian inference,*
 - *systematically account for confounding processes (unequal base frequencies, nucleotide substitution biases, etc.),*
 - *afford many opportunities for extension and refinement (still happening today).*

A likelihood approach for comparing synonymous and nonsynonymous nucleotide substitution rates, with application to the chloroplast genome

S. V. Muse and B. S. Gaut

Mol Biol Evol 11 715-724 (1994)

A codon-based model of nucleotide substitution for protein-coding DNA sequences.

N. Goldman and Z. Yang

Mol Biol Evol 11 725--736 (1994)