

Obtaining site-level dN/dS estimates with FEL

- dN/dS estimates at individual sites are not **precise**
- They are estimated from relatively small samples
- Precision improves with the number of sequences and divergence levels
- One approach to correct for this is to compute approximate site-level confidence intervals.

Codon	Partition	alpha	beta	LRT	Selection detected?	dN/dS with confidence intervals		
2	1	1.843	0.000	7.521	Neg. p = 0.0061	0.000(0.00-	0.09)
3	1	0.786	0.000	3.161	Neg. p = 0.0754	0.000(0.00-	0.16)
4	1	2.174	0.000	10.742	Neg. p = 0.0010	0.000(0.00-	0.07)
7	1	1.105	0.000	7.537	Neg. p = 0.0060	0.000(0.00-	0.11)
8	1	0.422	0.000	3.173	Neg. p = 0.0749	0.000(0.00-	0.28)
9	1	1.353	0.000	8.638	Neg. p = 0.0033	0.000(0.00-	0.08)
10	1	1.353	0.000	8.369	Neg. p = 0.0038	0.000(0.00-	0.09)
...								
247	1	1.353	0.000	8.088	Neg. p = 0.0045	0.000(0.00-	0.10)
248	1	0.451	0.000	3.496	Neg. p = 0.0615	0.000(0.00-	0.28)
249	1	0.000	2.700	7.881	Pos. p = 0.0050	10000.000(7599.84-10000.00)		
250	1	0.220	0.000	2.797	Neg. p = 0.0945	0.000(0.00-	0.61)
388	1	0.220	0.000	2.797	Neg. p = 0.0945	0.000(0.00-	0.61)

```
hyphy fel --alignment data/WestNileVirus_NS3.fna --ci Yes
```