


# Synonymous rate variation

- **dS** = constant for all sites (assumed by many models); this assumption appears to be nearly universally violated in biological data, due to e.g. secondary structure, localized codon usage bias, overlapping reading frames, etc.
- This can lead to, e.g. incorrect identification of relaxed constraint as selection and high false positive rates
- Most of HyPhy methods provide support for including dS



## Synonymous Site-to-Site Substitution Rate Variation Dramatically Inflates False Positive Rates of Selection Analyses: Ignore at Your Own Peril

Sadie R Wisotsky<sup>1 2</sup>, Sergei L Kosakovsky Pond<sup>2</sup>, Stephen D Shank<sup>2</sup>, Spencer V Muse<sup>1 3</sup>

Affiliations + expand

PMID: 32068869 PMCID: [PMC7403620](#) DOI: [10.1093/molbev/msaa037](#)

[Free PMC article](#)

# Allowing multi-nucleotide substitutions

- Some of the methods (e.g. BUSTED, aBSREL, RELAX) can extend substitution models to allow instantaneous double- and triple-“hits” (e.g. ACC to AGG)
- Sometimes multi-nucleotide changes along short branches at a single site can drive selection signal (possible false positives?)
- HyPhy includes a simple standard analysis for estimating alignment-wide multiple-hit rates.

Extra base hits: Widespread empirical support for instantaneous multiple-nucleotide changes

Alexander G. Lucaci , Sadie R. Wisotsky , Stephen D. Shank, Steven Weaver, Sergei L. Kosakovsky Pond 

Published: March 12, 2021 • <https://doi.org/10.1371/journal.pone.0248337>

[See the preprint](#)