

Polyhedral Computation

Jayant Apté
ASPITRG

October 3, 2012

Jayant Apté, ASPITRG

1

Part I


October 3, 2012

Jayant Apte, ASPITRG

2

The **Three Problems** of Polyhedral Computation

- Representation Conversion
 - H-representation of polyhedron
- V-Representation of polyhedron
- Projection
 - $\{P \mid P \in R^d\} \rightarrow \{P' \mid P' \in R^{(d-n)}\}$
- Redundancy Removal
 - Compute the minimal representation of a polyhedron



Outline

- Preliminaries
 - Representations of convex polyhedra, polyhedral cones
 - Homogenization - Converting a general polyhedron in \mathbb{R}^d to a cone in \mathbb{R}^{d+1}
 - Polar of a convex cone
- The three problems
- Representation Conversion
 - Review of LRS
 - Double Description Method
- Projection of polyhedral sets
 - Fourier-Motzkin Elimination
 - Block Elimination
 - Convex Hull Method (CHM)
- Redundancy removal
 - Redundancy removal using linear programming

Outline for Part 1

- Preliminaries
 - Representations of convex polyhedra, polyhedral cones
 - Homogenization – Converting a general polyhedron in \mathbb{R}^d to a cone in \mathbb{R}^{d+1}
- Polar of a convex cone
- The first problem
- Representation Conversion
 - Review of LRS
 - Double Description Method

Preliminaries

October 3, 2012


Jayant Apte, ASPITRG

Convex Polyhedron


- A subset P of R^d
- The set of solutions to a finite system of linear inequalities
- Called convex polytope if it is a convex polyhedron and bounded

Examples of polyhedra

Bounded- Polytope



Unbounded - polyhedron



October 3, 2012

Jayant Apte, ASPITRG

8

H-Representation of a Polyhedron

- The halfspace or inequality representation
- Polyhedron \mathcal{P} is the set $x \in \mathbb{R}^n$ obeying a system of linear inequalities i.e.
$$\mathcal{P} := \{x \in \mathbb{R}^n \mid Hx \leq h\}$$
- Can be written as $P(H, h)$
- $H \in \mathbb{R}^{m \times n}, h \in \mathbb{R}^m$ and the inequality is understood to hold elementwise
- Assumption: Polyhedron admits no lines.

October 3, 2012


Jayant Apte, ASPITRG

V-Representation of a Polyhedron

- $\mathcal{P} = conv\mathcal{S} + cone\mathcal{T}$
- \mathcal{S} is the finite set of extreme points, \mathcal{T} is the finite set of extreme directions(scaled to unit length).
- In other words, any point $x \in \mathcal{P}$ can be represented as,
$$x = \sum_{j=1}^J \beta_j s_j + \sum_{k=1}^K \gamma_k t_k$$
- J is the number of extreme points, K is the number of extreme directions, $\alpha_i \in \mathbb{R}, \beta_j \geq 0, \forall j, \sum_{j=1}^J \beta_j = 1, \gamma_k \geq 0.$

Example

$$\begin{aligned}x_1 &\leq 1 \\x_2 &\leq 1 \\-x_1 &\leq 0 \\-x_2 &\leq 0 \\-x_3 &\leq 0 \\x_2 + x_3 &\leq 2 \\-x_2 + x_3 &\leq 1 \\x_1 + x_3 &\leq 2 \\-x_1 + x_3 &\leq 1\end{aligned}$$



October 3, 2012

Jayant Apte, ASPITRG

11

Switching between the two representations : **Representation Conversion Problem**

- H-representation \longrightarrow V-representation: The Vertex Enumeration problem
- Methods:
 - Reverse Search, Lexicographic Reverse Search
 - Double-description method
- V-representation \longrightarrow H-representation
 - The Facet Enumeration Problem
- Facet enumeration can be accomplished by using polarity and doing Vertex Enumeration


Polyhedral Cone

- A special polyhedron
- Represented as:
- H representation:
 $C = \{x \mid \langle a_j, x \rangle \leq 0 \text{ for } j = 1, 2, \dots, m\}$ i.e.
 $C = C(A, 0)$
- V-representation
 $C = cone(r_1, \dots, r_n)$
 $= \{x \mid x = \sum_{i=1}^n \mu_i r_i, \mu_i \geq 0, i = 1, \dots, n\}$

October 3, 2012

Jayant Apte. ASPITRG

A cone in \mathbb{R}^3



October 3, 2012

Jayant Apté, ASPITRG

Homogenization

- Homogenization converts:
Any polyhedron in $R^d \rightarrow$ Pointed cone in $R^{(d+1)}$
- This way, we can consider polytopes/polyhedra
(bounded/unbounded) to be cones in +1 dimension
- Entire theory henceforth is developed for pointed
polyhedral cones

H-polyhedra

- If $P = P(A, z)$ is and \mathcal{H} -polyhedron, we define:
 - $C(P) := P \left(\begin{pmatrix} 0 & -1 \\ A & -z \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right)$ which is an \mathcal{H} -polyhedron(a cone)
in R^{d+1}
 - If P is defined as $a_i x \leq z_i$, $C(P)$ is defined by inequalities $a_i x - z_i x_{d+1} \leq 0$ and $x_{d+1} \geq 0$.
 - And $P = \{x \in R^d : \begin{pmatrix} x \\ 1 \end{pmatrix} \in C(P)\}$
 - Conversely, if $P = P(B, 0)$ is an \mathcal{H} -polyhedron in R^{d+1} ,
then $\{x \in R^d : \begin{pmatrix} x \\ 1 \end{pmatrix} \in P\}$ is an \mathcal{H} -polyhedron as well

October 3, 2012

Jayant Apte. ASPITRG

Example(d=2,d+1=3)

$$x_1 + x_2 \leq 18$$

$$x_1 - x_2 \leq 6$$

$$x_2 \leq 8$$

$$-x_1 + x_2 \leq 0$$

$$-x_1 - x_2 \leq -12$$

$$-x_2 \leq -4$$

↓

$$x_1 + x_2 - 18x_3 \leq 0$$

$$x_1 - x_2 - 6x_3 \leq 0$$


$$x_2 - 8x_3 \leq 0$$

$$-x_1 + x_2 \leq 0$$

$$-x_1 - x_2 + 12x_3 \leq 0$$

$$-x_2 + 4x_3 \leq 0$$


$$-x_3 \leq 0$$



October 3, 2012

Jayant Apte, ASPITRG

Example



October 3, 2012

Jayant Apté, ASPITRG

V -polyhedra

- If $P = conv(V) + cone(Y)$ is and \mathcal{V} -polyhedron, we define:

- $C(P) := cone \begin{pmatrix} V & Y \\ 1 & 0 \end{pmatrix}$

- which is a \mathcal{V} – polyhedron in R^{d+1}

- Conversely, if $C = cone(W)$ is any cone in R^{d+1} generated by vectors w_i with $w_{i(d+1)} \geq 0$, then $\{x \in R^d : \begin{pmatrix} x \\ 1 \end{pmatrix} \in C\}$ is a \mathcal{V} -polyhedron

Polar of a convex cone

- One notion of duality.
- If $\{r_i | i \in I\}$ are extreme rays of a closed convex cone C , then C consists of all non-negative combinations x of the r_i 's and,
 $C^\circ = \{y | \forall i \in I, \langle r_i, y \rangle \leq 0\}$ is called polar of C
- $C^{\circ\circ} = C$

Polar of a convex cone

- One notion of duality.
- If $\{r_i | i \in I\}$ are extreme rays of C , then C consists of all non-negative combinations x of the r_i 's then,
 $C^\circ = \{y | \forall i \in I, \langle r_i, y \rangle \leq 0\}$ ← Looks like an H-representation!!!
is called polar of C
- $C^{\circ\circ} = C$ ← Our ticket back to the original cone!!!

Polar: Intuition

Indicator Function

- For a convex set in R^n the indicator function

$\delta(\cdot|C)$ of C is given as:

$$\delta(x|C) = \begin{cases} 0, & \text{if } x \in C. \\ +\infty, & \text{if } x \notin C. \end{cases}$$


Positive Homogeneous Function

- A function f on R^n is said to be positive homogeneous(of degree 1) if for every x one has
 $f(\lambda x) = \lambda f(x) \quad 0 < \lambda < \infty$

Conjugate of a convex function defined over R^n


- It is defined as:
 $f^*(y) = \sup_{x \in R^n} \langle y, x \rangle - f(x)$
- For the indicator function of the convex set C defined above, it is given as,
 $\delta^*(y) = \sup_{x \in C} \langle y, x \rangle$

Polar: Intuition



- In first case if we find $\langle y, x \rangle > 0$ for some $x \in C$, we can always scale x to get higher value of the inner product because, for a cone, if $x \in C$, then, $\alpha x \in C, \alpha \geq 0$
- In second case 0 is the obvious supremum.

Polar: Intuition



The set $C^\circ = \{y | \forall x \in C, \langle y, x \rangle \leq 0\}$ so obtained is called the polar of C

Polar: Intuition


Every closed convex cone has a positive homogeneous indicator function associated with it

- ↓
- The conjugate of this indicator function is also a positive homogeneous indicator function
- ↓
- The set associated with it is also a cone and is called polar of C and denoted as C°

Use of Polar

- Representation Conversion
 - No need to have two different algorithms i.e. for
 $H \rightarrow V$ and $V \rightarrow H$.
- Redundancy removal
 - No need to have two different algorithms for removing redundancies from H-representation and V-representation
- Safely assume that input is always an H-representation for both these problems

Polarity: Intuition




$\langle r_i, y \rangle \leq 0$
 $x_1 + x_2 \leq 0$
 $2x_1 + x_2 \leq 0$
(Plotting in same space as the original cone)

October 3, 2012

Jayant Apte, ASPITRG

Polar: Intuition



October 3, 2012

Jayant Apte, ASPITRG

28

Problem #1

Representation Conversion

October 3, 2012

Jayant Apte, ASPITRG

Review of Lexicographic Reverse Search

October 3, 2012


Jayant Apte, ASPITRG

30

Reverse Search: High Level Idea

- Start with dictionary corresponding to the optimal vertex
- Ask yourself 'What pivot would have landed me at this dictionary if i was running simplex?'
- Go to that dictionary by applying reverse pivot to current dictionary
- Ask the same question again
- Generate the so-called 'reverse search tree'

Reverse Search on a Cube



Ref. David Avis, *Is: A Revised Implementation of the Reverse Search Vertex Enumeration Algorithm*

October 3, 2012

Jayant Apte, ASPITRG

Double Description Method

- First introduced in:
 - “ Motzkin, T. S.; Raiffa, H.; Thompson, G. L.; Thrall, R. M. (1953). "The double description method". *Contributions to the theory of games. Annals of Mathematics Studies*. Princeton, N. J.: Princeton University Press. pp. 51–73”
 - The primitive algorithm in this paper is very inefficient.
(How? We will see later)
 - Several authors came up with their own efficient implementation(viz. Fukuda, Padberg)
 - Fukuda's implementation is called *cdd*

Some Terminology

- A pair (A, R) is said to be a double description pair (DD pair) if the relationship:
 - $Ax \geq 0$ iff $x = R\lambda$ for some $\lambda \geq 0$ holds
 - Column size of A = Row Size of $R = d$
- Provides two different descriptions of the same object:
A *Polyhedral Cone*, formally defined as:
- A set $P(A)$ represented by A as: $P(A) = \{x \in \mathbb{R}^d : Ax \geq 0\}$
and is simultaneously represented by R as:
 $\{x \in \mathbb{R}^d : x = R\lambda \text{ for some } \lambda \geq 0\}$
- A is called the representation matrix, while R is called the generator matrix.

Double Description Method: The High Level Idea

- An *Incremental* Algorithm
- Starts with certain subset of rows of H-representation of a cone $Ax \geq 0$ to form initial H-representation
- Adds rest of the inequalities one by one constructing the corresponding V-representation every iteration
- Thus, constructing the V-representation *incrementally*.

How it works?

Initialization:

- Let $K \subset \{1, \dots, m\}$ i.e. the row indices of A
- Let A_K denote the submatrix of A consisting of rows indexed by K
- Suppose we have already found a generating matrix R of $P(A_K)$ i.e. (A_K, R) is a DD pair

Iteration:

- Given a DD pair (A_K, R) , select any row index $i \notin K$ and construct a DD pair (A_{K+i}, R') using the DD pair (A_K, R)

Termination:

- If $A = A_K$, we are done.

How it works?

Initialization:

- Let $K \subset \{1, \dots, m\}$ i.e. the row indices of A
- Let A_K denote the submatrix of A consisting of rows indexed by K
- Suppose we have already found a generating matrix R of $P(A_K)$ i.e. (A_K, R) is a DD pair

Iteration:

- Given a DD pair (A_K, R) , select any row index $i \notin K$ and construct a DD pair (A_{K+i}, R') using the DD pair (A_K, R)

Termination:

- If $A = A_K$, we are done.

Initialization

- method 1:
 - Find a DD pair (A_K, R) when $|K| = 1$.
- method 2:
 - Select a maximal submatrix A_K of A consisting of linearly independent rows of A .
 - The vectors r_j 's are obtained by solving the system:
$$A_K R = I \text{ where } I \text{ is } |K| \times |K|$$
$$A_K x \geq 0 \leftrightarrow x = A_K^{-1} \lambda, \lambda \geq 0$$

Initialization

 **Very trivial and inefficient**

- method 1:
Find a DD pair (A_K, R) when $|K| = 1$.
- method 2:
Select a maximal submatrix A_K of A consisting of linearly independent rows of A .
The vectors r_j 's are obtained by solving the system:
$$A_K R = I \text{ where } I \text{ is } |K| \times |K|$$
$$A_K x \geq 0 \leftrightarrow x = A_K^{-1} \lambda, \lambda \geq 0$$

Example: Initialization

Consider the problem of performing vertex enumeration on the polyhedron represented as follows:

$$\begin{array}{ll} -x_1 - x_2 + 18x_3 \geq 0 & (1) \\ -x_1 + x_2 + 6x_3 \geq 0 & (2) \\ x_1 - x_2 + 8x_3 \geq 0 & (3) \\ x_1 - x_2 \geq 0 & (4) \\ x_1 + x_2 - 12x_3 \geq 0 & (5) \\ x_2 - 4x_3 \geq 0 & (6) \end{array}$$

Example: Initialization

An Example:

Consider the problem of performing vertex enumeration on the polyhedron represented as follows:

$$\begin{array}{ll} -x_1 - x_2 + 18x_3 \geq 0 & (1) \\ -x_1 + x_2 + 6x_3 \geq 0 & (2) \\ x_1 - x_2 + 8x_3 \geq 0 & (3) \\ x_1 - x_2 \geq 0 & (4) \\ x_1 + x_2 - 12x_3 \geq 0 & (5) \\ x_2 - 4x_3 \geq 0 & (6) \end{array}$$

Example : Initialization

$$A_{\{1,4,6\}} = \begin{pmatrix} -1 & -1 & 18 \\ 1 & -1 & 0 \\ 0 & 1 & -4 \end{pmatrix}$$

$$A_{\{1,4,6\}}^{-1} = R = \begin{matrix} & \begin{matrix} A & B & C \end{matrix} \\ \begin{matrix} 4.0000 & 14.0000 & 9.0000 \\ 4.0000 & 4.0000 & 9.0000 \\ 1.0000 & 1.0000 & 1.0000 \end{matrix} & \end{matrix}$$

How it works?

Initialization:

- Let $K \subset \{1, \dots, m\}$ i.e. the row indices of A
- Let A_K denote the submatrix of A consisting of rows indexed by K

- Suppose we have already found a generating matrix R of $P(A_K)$ i.e. (A_K, R) is a DD pair

Iteration:

- Given a DD pair (A_K, R) , select any row index $i \notin K$ and construct a DD pair (A_{K+i}, R') using the DD pair (A_K, R)

Termination:

- If $A = A_K$, we are done.

Iteration: Insert a new constraint

- The newly inserted inequality $A_i x \geq 0$ partitions the space \mathbb{R}^d into three parts:

$$\begin{aligned}H_i^+ &= \{x \in \mathbb{R}^d : A_i x > 0\} \\H_i^0 &= \{x \in \mathbb{R}^d : A_i x = 0\} \\H_i^- &= \{x \in \mathbb{R}^d : A_i x < 0\}\end{aligned}$$


- Let J be the set of column indices of R . The rays $r_j (j \in J)$ are accordingly partitioned as:

$$\begin{aligned}J^+ &= \{j \in J : r_j \in H_i^+\} \\J^0 &= \{j \in J : r_j \in H_i^0\} \\J^- &= \{j \in J : r_j \in H_i^-\}\end{aligned}$$

October 3, 2012

Jayant Apte, ASPITRG


Iteration 1



October 3, 2012

Jayant Apte, ASPITRG

Iteration 1: Insert a new constraint




October 3, 2012

Jayant Apte, ASPITRG

46

Iteration 1: Insert a new constraint




October 3, 2012

Jayant Apte, ASPITRG

47

Iteration 1: Insert a new constraint



October 3, 2012

Jayant Apte, ASPITRG


Main Lemma for DD Method

Let (A_K, R) be a DD pair and let i be the new row index of A not in K . Then the pair (A_{K+i}, R') is a DD pair, where R' is the $d \times |J'|$ matrix with column vectors $r_j (j \in J')$ defined by,

$$J' = J^+ \cup J^0 \cup (J^+ \times J^-), \text{ and}$$

$$r_{jj'} = (A_i r_j) r_{j'} - (A_i r_{j'}) r_j \text{ for each } (j, j') \in J^+ \times J^-$$

Iteration 1: Insert a new constraint



October 3, 2012

Jayant Apte, ASPITRG

50


Iteration 1: Get the new DD pair

$$A_{\{1,2,4,6\}}$$

$$\begin{matrix} -1 & -1 & 18 \\ 1 & -1 & 0 \\ 0 & 1 & -4 \\ -1 & 1 & 6 \end{matrix}$$

	r_{AB}	r_{BC}	r_A	r_C
$R_{\{1,2,4,6\}}$	10.0000	12.0000	4.0000	9.0000
	4.0000	6.0000	4.0000	9.0000
	1.0000	1.0000	1.0000	1.0000


Iteration 2



October 3, 2012

Jayant Apte, ASPITRG

Iteration 2: Insert new constraint




October 3, 2012

Jayant Apte, ASPITRG

53

Iteration 2: Insert new constraint




October 3, 2012

Jayant Apté, ASPITRG

54

Iteration 2: Insert new constraint




October 3, 2012

Jayant Apte, ASPITRG

Get the new DD pair

$$A_{\{1,2,3,4,6\}} = \begin{pmatrix} -1 & -1 & 18 \\ 1 & -1 & 0 \\ 0 & 1 & -4 \\ -1 & 1 & 6 \\ 0 & -1 & 8 \end{pmatrix}$$

$$R_{\{1,2,3,4,6\}} = \begin{pmatrix} r_{AD} & r_{BD} & r_{CD} & r_A & r_B & r_C \\ 9.2000 & 10.0000 & 8.0000 & 10.0000 & 12.0000 & 4.0000 \\ 8.0000 & 8.0000 & 8.0000 & 4.0000 & 6.0000 & 4.0000 \\ 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 & 1.0000 \end{pmatrix}$$



Get the new DD pair

$$A_{\{1,2,3,4,5,6\}}$$

$$\begin{matrix} -1 & -1 & 18 \\ 1 & -1 & 0 \\ 0 & 1 & -4 \\ -1 & 1 & 6 \\ 0 & -1 & 8 \\ 1 & 1 & -12 \end{matrix}$$


$R_{\{1,2,3,4,5,6\}}$						
A	B	C	D	E	F	G
6.2609	6.4000	6.0000	8.0000	7.2000	9.2000	10.0000
5.7391	5.6000	6.0000	4.0000	4.8000	8.0000	8.0000
1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
H	I	J				

October 3, 2012

Jayant Apte, ASPITRG

58

Termination



October 3, 2012

Jayant Apte, ASPITRG

Efficiency Issues


- Is the implementation I just described good enough?

Efficiency Issues

- Is the implementation I just described good enough?
- **Hell no!**
 - The implementation just described suffers from profusion of redundancy

Efficiency Issues

We can totally do without this one!!




October 3, 2012

Jayant Apte, ASPITRG

62

Efficiency Issues

And without
all these!!



October 3, 2012

Jayant Apte, ASPITRG

63

Efficiency Issues

- In worst case an iteration can start with n extreme rays and end with $(\frac{n}{2})^2$ extreme rays.
- Hence, the number extreme rays can very soon grow out of hand.
- A straightforward implementation is quite useless.
- Redundancy removal for n extreme rays is equivalent to solving n linear programs which is also not a very exciting prospect.
- Hence, we focus on **Not letting redundant extreme rays to be created** in first place
- Fukuda's main contributions are in that direction

The Primitive DD method

```
procedure DoubleDescriptionMethod( $\Lambda$ );  
begin  
    Obtain any initial DD pair ( $\Lambda_K, R$ )  
    while  $K \neq \{1, 2, \dots, m\}$  do  
        begin  
            Select any index  $i$  from  $\{1, 2, \dots, m\}$ ;  
            Construct a DD pair ( $\Lambda_{K+i}, R'$ ) from ( $\Lambda_K, R$ );  
             $R := R'$ ;  $K := K + i$   
        end  
        Output  $R$   
    begin
```

What to do?

- Add some more structure
- Strengthen the *Main Lemma*

Add some more structure

October 3, 2012

Jayant Apte, ASPITRG

67

Some definitions

ray of P

- r is said to be a *ray* of P if $r \neq 0$ and $\alpha r \in P \forall \alpha > 0$
- If r and r' are such that $r = \alpha r'$ for some positive number α , we say $r \simeq r'$

zero set/active set

- For any vector $x \in P$, we define the *zero set* or *active set* $Z(x)$ as the set of inequality indices i such that $A_i x = 0$

Proposition 4. (Fukuda)

- Let r be a ray of P , $\bar{F} := \{x : A_{Z(r)}x = 0\}$, $F := \bar{F} \cap P$ and $rank(A_{Z(r)}) = d - k$ then

- (a) $rank(A_{Z(r) \cup \{i\}}) = d - k + 1 \forall i \neq Z(r)$;
- (b) F contains k linearly independent rays;
- (c) If $k \geq 2$ then r is a non-negative combination of two distinct rays r_1 and r_2 with $rank(A(Z(r_i))) > d - k, i = 1, 2$

Proposition 7. (Fukuda)

- Let r and r' be distinct rays of P . Then the following statements are equivalent:
 - (a) r and r' are adjacent extreme rays;
 - (b) r and r' are extreme rays and rank of the matrix $A_{Z(r)(r')}$ is $d - 2$
 - (c) if r'' is a ray with $Z(r'') \supset Z(r) \cap Z(r')$ then either $r'' \succeq r$ or $r'' \succeq r'$;

Proposition 7. (Fukuda)

- Let r and r' be distinct rays of P . Then the following statements are equivalent:

- (a) r and r' are adjacent extreme rays;
- (b) r and r' are extreme rays and rank of the matrix $A_{Z(r)(r')}$ is $d - 2$
- (c) if r'' is a ray with $Z(r'') \supset Z(r) \cap Z(r')$ then either $r'' \simeq r$ or $r'' \simeq r'$;

Algebraic Characterization of adjacency

Proposition 7. (Fukuda)

- Let r and r' be distinct rays of P . Then the following statements are equivalent:
 - (a) r and r' are adjacent extreme rays;
 - (b) r and r' are extreme rays and rank of the matrix $A_{Z(r)(r')}$ is $d - 2$


(c) if r'' is a ray with $Z(r'') \supset Z(r) \cap Z(r')$ then either
 $r'' \simeq r$ or $r'' \simeq r'$;

Combinatorial Characterization of adjacency
(Combinatorial Oracle)

Example: Combinatorial Adjacency Oracle

$$J^- = \{D\}$$


$$J^+ = \{A, B, C\}$$



October 3, 2012

Jayant Apte, ASPITRG

Example: Combinatorial Adjacency Oracle



October 3, 2012

Jayant Apte, ASPITRG

Strengthened Main Lemma for DD Method

Let (A_K, R) be a DD pair and let i be the new row index of A not in K . Then the pair (A_{K+i}, R') is a DD pair, where R' is the $d \times |J'|$ matrix with column vectors $r_j (j \in J')$ defined by,

$$\begin{aligned} J' &= J^+ \cup J^0 \cup Adj, \\ Adj &= \{(j, j') \in J^+ \times J^- : r_j \text{ and } r_{j'} \text{ are adjacent} \\ &\text{in } P(A_K)\}, \text{ and} \\ r_{jj'} &= (A_i r_j) r_{j'} - (A_i r_{j'}) r_j \text{ for each } (j, j') \in Adj \end{aligned}$$

Furthermore, if R is minimal generating matrix for $P(A_K)$ then R' is a minimal generating matrix for $P(A_{K+i})$

Procedural Description

```
procedure DDMETHODStandard( $\Lambda$ );
begin
    Obtain any initial DD pair  $(\Lambda_K, R)$ 
    while  $K \neq \{1, 2, \dots, m\}$  do
        begin
            Select any index  $i$  from  $\{1, 2, \dots, m\}$ ;
            Construct a DD pair  $(\Lambda_{K+i}, R')$  from  $(\Lambda_K, R)$ ;
            /*by using Strengthened Main Lemma */
             $R := R'$ ;  $K := K + i$ 
        end
        Output  $R$ 
    begin
```

Part 2

October 3, 2012

Jayant Apte, ASPITRG

77

- Projection of polyhedral sets
 - Fourier-Motzkin Elimination
 - Block Elimination
 - Convex Hull Method (CHM)
- Redundancy removal
 - Redundancy removal using linear programming

Projection of a Polyhedra

- Consider an \mathcal{H} -polyhedron $P = P(A, z) \subseteq \underline{R}^d$
- We want to project to $\{x \in R^d : x_k = 0\} \equiv R^{d-1}$ along the x_k axis
- We define:

$$\begin{aligned} proj_k(P) &:= \{x - x_k e_k : x \in P\} & (1) \\ &= \{x \in R^d : x_k = 0, \exists y \in R : x + y e_k \in P\} & (2) \end{aligned}$$

- This is projection of P in the direction of e_k
- The set $proj_k(P)$ is contained in the hyperplane
 $H_k := \{x \in R^d : x_k = 0\}$

Fourier-Motzkin Elimination

- Named after Joseph Fourier and Theodore Motzkin

How it works?

- We start with an \mathcal{H} -polyhedron $P = P(A, z) \subseteq R^d$
- Suppose we want to eliminate the variable x_k
- Consider coefficients of x_k in our system of inequalities, and assume that $a_{ik} > 0$ and $a_{jk} < 0$
- Then the respective inequalities can be written as,
$$a_i x \leq z_i \rightarrow a_{ik}x_k \leq a_{ik}x_k - a_i x + z_i$$
and
$$a_j x \leq z_j \rightarrow (-a_{jk}x_k) \geq -a_{jk}x_k + a_j x - z_j$$

How it works? Contd...

- Multiply these equations by $-a_{jk}$ and a_{ik} respectively
 $-a_{jk}a_{ik}x_k \leq -a_{jk}a_{ik}x_k - a_i x - a_{jk}z_i$
and
 $-a_{jk}a_{ik}x_k \geq -a_{jk}a_{ik}x_k + a_j a_{ik}x - a_{ik}z_j$
- These equations form upper bound and lower bound respectively on $-a_{jk}a_{ik}x_k$
- Combining the two we get,
 $a_{ik}a_j + (-a_{jk}a_j)x \leq a_{ik}z_j + (-a_{jk})z_j$

Efficiency of FM algorithm

- The number of inequalities goes beyond tractable limits within few elimination steps
- IF A has m rows, then $A^{\setminus k}$ may have as many as $\lfloor \frac{m^2}{4} \rfloor$ rows
- FM-elimination creates $O(m^2)$ new inequalities
- Useful only as a simple and elegant method that is easy to understand
- There have been efforts to introduce heuristics, as discussed in the paper by Lassez et al.


October 3, 2012


Jayant Apte, ASPITRG


Convex Hull Method(CHM)

- First appears in “C. Lassez and J.-L. Lassez, Quantifier elimination for conjunctions of linear constraints via aconvex hull algorithm, *IBM Research Report*, T.J. Watson Research Center (1991)”
- Found to be better than most other existing algorithms when dimension of projection is small
- Cited by Weidong Xu, Jia Wang, Jun Sun in their ISIT 2008 paper “A Projection Method for Derivation of Non-Shannon-Type Information Inequalities”

CHM intuition








CHM intuition


Is $x_2 \geq 4$ one of the inequalities defining $\pi(P)$?



Moral of the story

- We can make decisions about $\pi(P)$ without actually having its H-representation.
- It suffices to have P , the original polyhedron
- We run linear programs on P to make these decisions

Convex Hull Method: Flowchart



October 3, 2012


Jayant Apte, ASPITRG

90

Example

$$\begin{array}{ll} -16 + 0x_1 + 2x_2 + 1x_3 \geq 0 & (1) \\ -72 + 4x_1 + 4x_2 + 3x_3 \geq 0 & (2) \\ 0 + 0x_1 + 2x_2 - 1x_3 \geq 0 & (3) \\ -24 + 4x_1 + 4x_2 - 3x_3 \geq 0 & (4) \\ \text{Project } P \text{ onto } x_1, x_2 & 0 - 4x_1 + 4x_2 + 3x_3 \geq 0 \\ 3D \rightarrow 2D & 48 - 4x_1 + 4x_2 - 3x_3 \geq 0 \\ & 48 - 4x_1 - 4x_2 + 3x_3 \geq 0 \\ & 8 + 0x_1 - 2x_2 + 1x_3 \geq 0 \\ & -24 + 4x_1 - 4x_2 + 3x_3 \geq 0 \\ & 24 + 0x_1 - 2x_2 - 1x_3 \geq 0 \\ & 24 + 4x_1 - 4x_2 - 3x_3 \geq 0 \\ & 96 - 4x_1 - 4x_2 - 3x_3 \geq 0 \end{array}$$

Example




October 3, 2012

Jayant Apte, ASPITRG

92


Example



October 3, 2012

Jayant Apte, ASPITRG


93




How to get the extreme points of initial approximation?

- We need $d + 1$ points to have full dimensional convex hull
- Get first two points by maximizing and minimizing x_1
- Get rest of the points by running linear programs on initial set of constraints

Example: Get the first two points



Example: Get the first two points




October 3, 2012

Jayant Apte, ASPITRG

97

Example: Get the first two points




October 3, 2012

Jayant Apte, ASPITRG

Get rest of the points so you have a total of $d+1$ points


$d = 2$ so we need $d + 1 = 3$ points



October 3, 2012

Jayant Apte, ASPITRG

Get the rest of the points so you
have a total of $d+1$ points




October 3, 2012

Jayant Apte. ASPITRG

100

Get the rest of the points so you
have a total $d+1$ points



October 3, 2012


Jayant Apte, ASPITRG

101

How to get the initial facets of the projection?

```
procedure initial_hull( $E$ );  
begin  
    Let CH= $\phi$   
    for each  $p \in E$  do  
        Compute  $\sum_{j=1}^d \alpha_j x_j = \alpha$  the equation of the  
        hyperplane defined by  $E - \{p\}$   
        Let  $h = \sum_{j=1}^d \alpha_j x_j$   
        If  $h(p) \geq \alpha$   
            then  $CH = CH \cup \{-h \leq -\alpha\}$   
        else  $CH = CH \cup \{h \leq \alpha\}$   
    end  
end
```

How to get the initial facets of the projection?




October 3, 2012

Jayant Apte, ASPITRG

103

Convex Hull Method: Flowchart




October 3, 2012

Jayant Apte, ASPITRG

104

Convex Hull Method: Flowchart




October 3, 2012

Jayant Apte, ASPITRG

105

Convex Hull Method: Flowchart



October 3, 2012

Jayant Apte, ASPITRG

Incremental refinement

- Find a facet in current approximation of that is not actually the facet of $\pi(P)$
- How to do that?

given $\{h_i \leq \alpha_i\}$,


Maximize h_i over P

If $\text{Max}(h_i) = \alpha_i$

Then Label $\{h_i \leq \alpha_i\}$ as *terminal*

else Move it outward to find a new extreme point

Incremental refinement




October 3, 2012

Jayant Apte, ASPITRG

108

Incremental refinement




Maximization over P gives a point $(10, 8, 6)$ of P
This point corresponds to the point $(10,8)$ of $\pi(P)$

109

Jayant Apte, ASPITRG

October 3, 2012

Convex Hull Method: Flowchart



How to update the H-representation to accommodate new point ?


```
procedure update_convex_hull(new-pt);  
begin  
    for each  $C \in CH$  s.t.  $p \notin C$  do  
        Generate all subsets  $SE$  of  $d - 1$  extreme points in  $C$   
        for each  $SE$  do  
            If  $SE \cup \{p\}$  determines a unique hyperplane  $\sum_i \alpha_i x_i = 0$   
            then let  $h = \sum_i \alpha_i x_i$   
                If  $\forall q \in E, h(q) \leq \alpha_0$   
                    then  $C = \sum_i \alpha_i x_i \leq \alpha_0$   
                else if  $\forall q \in E, h(q) \geq \alpha_0$   
                    then  $C = -\sum_i \alpha_i x_i \leq -\alpha_0$   
                else  $C = \emptyset$   
                Let  $CH = CH \cup \{C\}$   
            end  
        end  
    end  
end
```

Jayant Apte, ASPITRG

111

October 3, 2012

Example




October 3, 2012

Jayant Apte, ASPITRG

112

Example




October 3, 2012

Jayant Apte, ASPITRG

113

Example: New candidate facets



October 3, 2012

Jayant Apte, ASPITRG


114

How to update the H-representation to accommodate new point ?

```
procedure update_convex_hull(new-pt);
begin
    for each  $C \in CH$  s.t.  $p \notin C$  do
        Generate all subsets  $SE$  of  $d - 1$  extreme points in  $C$ 
        for each  $SE$  do
            If  $SE \cup \{p\}$  determines a unique hyperplane  $\sum_i \alpha_i x_i = 0$ 
            then let  $h = \sum_i \alpha_i x_i$ 
                If  $\forall q \in E, h(q) \leq \alpha_0$ 
                    then  $C = \sum_i \alpha_i x_i \leq \alpha_0$ 
                else if  $\forall q \in E, h(q) \geq \alpha_0$ 
                    then  $C = -\sum_i \alpha_i x_i \leq -\alpha_0$ 
                else  $C = \emptyset$ 
                Let  $CH = CH \cup \{C\}$ 
            end
             $\forall C \in CH$  If  $p \notin C$  then  $CH = CH - \{C\}$ 
        end
    end
```

Example

$$x_2 \leq 8$$




October 3, 2012

Jayant Apte, ASPITRG

116

Example



$$CH = \{ -x_1 + x_2 \leq 0, x_2 \geq 6, x_1 + 2x_2 \leq 24, x_2 \leq 8 \} \cup \{ x_1 + x_2 \leq 18 \}$$

October 3, 2012

Jayant Apte, ASPITRG

117

How to update the H-representation to accommodate new point ?

```
procedure update_convex_hull(new-pt);
begin
    for each  $C \in CH$  s.t.  $p \notin C$  do
        Generate all subsets  $SE$  of  $d - 1$  extreme points in  $C$ 
        for each  $SE$  do
            If  $SE \cup \{p\}$  determines a unique hyperplane  $\sum_i \alpha_i x_i = 0$ 
            then let  $h = \sum_i \alpha_i x_i$ 
            If  $\forall q \in E, h(q) \leq \alpha_0$ 
            then  $C = \sum_i \alpha_i x_i \leq \alpha_0$ 
            else if  $\forall q \in E, h(q) \geq \alpha_0$ 
            then  $C = -\sum_i \alpha_i x_i \leq -\alpha_0$ 
            else  $C = \phi$ 
            Let  $CH = CH \cup \{C\}$ 
        end
    end
end
```


$\forall C \in CH$ If $p \notin C$ then $CH = CH - \{C\}$

October 3, 2012

Jayant Apte, ASPITRG

118

Example




October 3, 2012

Jayant Apte, ASPITRG

119

Example




October 3, 2012


Jayant Apte, ASPITRG

120


Example



2nd iteration



3rd iteration




October 3, 2012

Jayant Apte, ASPITRG

123

4th iteration: Done!!!



Comparison of Projection Algorithms

- Fourier Motzkin Elimination and Block Elimination doesn't work well when used on big problems
- CHM works very well when the dimension of projection is relatively small as compared to the dimension of original polyhedron.
- Weidong Xu, Jia Wang, Jun Sun have already used CHM to get non-Shannon inequalities.

Computation of minimal representation/Redundancy Removal

October 3, 2012

Jayant Apte, ASPITRG

126

Definitions

Definition A linear inequality $A_i x \leq b_i$ (for some $i \in \{1, \dots, m\}$) of a polyhedron P is *redundant* if it is implied by the other inequalities of P

Definition An extreme point v of polytope P is said to be redundant if it can be represented as convex combination of any other extreme points in the polytope

References

- K. Fukuda and A. Prodon. Double description method revisited. Technical report, Department of Mathematics, Swiss Federal Institute of Technology, Lausanne, Switzerland, 1995
- G. Ziegler, *Lectures on Polytopes*, Springer, 1994, revised 1998. Graduate Texts in Mathematics.
- Tien Huynh, Catherine Lassez, and Jean-Louis Lassez. *Practical issues on the projection of polyhedral sets*. Annals of mathematics and artificial intelligence, November 1992
- Catherine Lassez and Jean-Louis Lassez. *Quantifier elimination for conjunctions of linear constraints via a convex hull algorithm*. In Bruce Donald, Deepak Kapur, and Joseph Mundy, editors, Symbolic and Numerical Computation for Artificial Intelligence. Academic Press, 1992.
- R. T. Rockafellar. *Convex Analysis* (Princeton Mathematical Series). Princeton Univ Pr.
- W. Xu, J. Wang, J. Sun. A projection method for derivation of non-Shannon-type information inequalities. In IEEE International Symposium on Information Theory (ISIT), pp. 2116 – 2120, 2008.

Minkowski's Theorem for Polyhedral Cones

- For any $m \times d$ real matrix A , \exists some $d \times n$ real matrix R s.t. (A, R) is a DD pair, or in other words, the cone $P(A)$ is generated by R .

- Emphasis on finiteness of columns of R

Weyl's Theorem for Polyhedral Cones

- For any $d \times n$ real matrix R , \exists some $m \times d$ matrix A s.t. (A, R) is a DD pair, or in other words, the set generated by R is the cone $P(A)$
- It is the converse of Minkowski's Theorem
- Together they form the *Representation theorem* of Polyhedral Cones