

AHA/ACC/ESC/WHF CONSENSUS DOCUMENT

Fifth Universal Definition of Myocardial Infarction (2026): On Behalf of the Joint European Society of Cardiology (ESC)/American College of Cardiology (ACC)/American Heart Association (AHA)/World Heart Federation (WHF) Task Force for the Universal Definition of Myocardial Infarction

Endorsed by the European Association for Cardio-Thoracic Surgery (EACTS) and the Society of Thoracic Surgeons (STS)

Affirmation of Value by the Society for Cardiovascular Angiography and Interventions (SCAI)



Authors/Task Force Members

Nicholas L. Mills (ESC Chairperson) (United Kingdom)*; L. Kristin Newby (ACC/AHA Co-Chairperson) (United States of America)*; Sarah Zaman (WHF Co-Chairperson) (Australia)*; Wael A. Almahmeed (UAE); Chiara Bucciarelli-Ducci (United Kingdom); Monica M. Colvin (United States of America); George D. Dangas (United States of America); Holli A. DeVon (United States of America); Rohan Dharmakumar (United States of America); Thomas A. Gaziano (United States of America); Derek J. Hausenloy (Singapore); Mohamed Jeilan (Kenya); Konstantinos C. Koskinas (Switzerland); Bertil Lindahl (Sweden); Zuzana Motovska (Czech Republic); Maria Rubini Gimenez (Spain); Ana María Salvati (Argentina); Jacqueline E. Tamis-Holland (United States of America); ESC/ACC/AHA/WHF Universal Definition Document Advisory Group

Document Reviewers

Borja Ibanez, (ESC Review Co-Ordinator) (Spain); Elliott Antman, (ACC/AHA Review Co-ordinator) (United States of America); Junbo Ge, (WHF Review Co-ordinator) (China); Tayo Addo (United States of America); Junya Ako (Japan); Mirvat Alasnag (Saudi Arabia); Rasha Al-Lamee (United Kingdom); Paulo R.A. Caramori (Brazil); Mauricio G. Cohen (United States of America); Victoria Delgado (Spain); Bernard Dennis (United States of America); Sigrun Halvorsen (Norway); Julie Harris (United Kingdom); Milosz J. Jaguszewski (Poland); Hani Jneid (United States of America); Hugo Katus (Germany); Milan Milojevic (Serbia/Switzerland)‡; Michelle O'Donoghue (United States of America); Julie Redfern (Australia); Joseph F. Sabik III (United States of America)§; Clara Saldarriaga (Colombia); Nizal Sarrafzadegan (Iran/Canada)

Correspondence to: Nicholas L. Mills, British Heart Foundation Centre of Research Excellence, The University of Edinburgh, Royal Infirmary of Edinburgh, Edinburgh EH16 4SU, UK, Email nick.mills@ed.ac.uk; L. Kristin Newby, Division of Cardiology, Department of Medicine and Duke Clinical Research Institute, Duke Health, Durham, NC 27710, Email kristin.newby@duke.edu; or Sarah Zaman, Westmead Applied Research Centre, Faculty of Medicine and Health, The University of Sydney, Westmead, New South Wales, Australia, and Department of Cardiology, Westmead Hospital, Westmead, New South Wales, Australia, Email sarah.zaman@sydney.edu.au

*The chairperson and the two co-chairpersons contributed equally to the document and are joint corresponding authors.

‡Representing the European Association for Cardio-Thoracic Surgery (EACTS)

§Representing the Society of Thoracic Surgeons (STS)

Author/Task Force Member affiliations are listed in author information.

This article has been co-published in *European Heart Journal*, *Journal of the American College of Cardiology*, *Circulation*, and *Global Heart*. All rights reserved.

© 2026 European Society of Cardiology, American College of Cardiology, American Heart Association, Inc., and World Heart Foundation. The articles are identical except for minor stylistic and spelling differences in keeping with each journal's style. Any citation can be used when citing this article.

Circulation is available at www.ahajournals.org/journal/circ

Key Words: acute myocardial injury ■ cardiac troponin ■ chronic myocardial injury ■ coronary and cardiac imaging ■ expert consensus document ■ high-sensitivity cardiac troponin ■ International Classification of Diseases 11th Revision (ICD-11) ■ myocardial infarction ■ myocardial injury with non-obstructive coronary arteries (MINOCA) ■ primary myocardial infarction ■ procedure-related myocardial infarction ■ recurrent myocardial infarction ■ secondary myocardial infarction ■ Takotsubo syndrome ■ universal definition ■ unrecognized myocardial infarction ■ unstable angina

TABLE OF CONTENTS

1. Executive Summary.....	eXXX
2. Context.....	eXXX
3. Objective.....	eXXX
4. What Is New.....	eXXX
5. Definition of Myocardial Infarction.....	eXXX
6. Acute and Chronic Myocardial Injury.....	eXXX
6.1. Acute Myocardial Injury.....	eXXX
6.2. Chronic Myocardial Injury.....	eXXX
7. Classification of Myocardial Infarction.....	eXXX
7.1. Approaches to the Classification of Myocardial Infarction.....	eXXX
7.1.1. Classification Based on Etiology.....	eXXX
7.1.2. Classification Based on the Electrocardiogram.....	eXXX
7.2. Clinical Classification of Myocardial Infarction.....	eXXX
7.2.1. Primary Myocardial Infarction.....	eXXX
7.2.2. Secondary Myocardial Infarction.....	eXXX
7.2.3. Procedure-Related Myocardial Infarction.....	eXXX
8. Diagnosis of Myocardial Infarction Following Sudden Death.....	eXXX
9. Silent or Unrecognized Myocardial Infarction.....	eXXX
10. Unstable Angina.....	eXXX
11. Myocardial Injury With Non-Obstructive Coronary Arteries.....	eXXX
12. Symptoms of Myocardial Infarction.....	eXXX
13. The Electrocardiogram.....	eXXX
14. Cardiac Biomarkers.....	eXXX
14.1. Cardiac Troponin.....	eXXX
14.1.1. Defining the 99th Percentile Upper Reference Limit.....	eXXX
14.1.2. Analytical Considerations.....	eXXX
14.1.3. Serial Measurements.....	eXXX
14.1.4. Defining a Significant Rise and Fall.....	eXXX
14.1.5. Accelerated Diagnostic Pathways.....	eXXX
14.1.6. Interpretation of Cardiac Troponin for Diagnosis.....	eXXX
14.2. Alternative Biomarkers of Myocardial Injury.....	eXXX
15. Coronary and Cardiac Imaging.....	eXXX
15.1. Invasive Imaging.....	eXXX
15.1.1. Invasive Coronary Angiography.....	eXXX
15.1.2. Intravascular Imaging.....	eXXX
15.1.3. Invasive Functional Testing.....	eXXX
15.2. Non-Invasive Imaging.....	eXXX
15.2.1. Echocardiography.....	eXXX
15.2.2. Cardiac Magnetic Resonance Imaging.....	eXXX
15.2.3. Coronary Computed Tomography Angiography.....	eXXX
15.2.4. Nuclear Medicine.....	eXXX
16. Special Situations.....	eXXX
16.1. Recurrent Myocardial Infarction.....	eXXX
16.2. Heart Failure.....	eXXX
16.3. Chronic Kidney Disease.....	eXXX
16.4. Critical Illness.....	eXXX
16.5. Non-Cardiac Surgery.....	eXXX
16.6. Structural Cardiac Intervention.....	eXXX
16.7. Cardio-Oncology.....	eXXX
17. Implications for Patients, Secondary Prevention, and Cardiac Rehabilitation.....	eXXX
18. Implications for Public Health and Classification of Disease.....	eXXX
19. Implications and Adaptions for Low-Resource Settings.....	eXXX
20. Implications of the Definition on Research.....	eXXX
References.....	eXXX

1. EXECUTIVE SUMMARY

The Fifth Universal Definition of Myocardial Infarction (UDMI) updates the classification to better reflect underlying pathophysiology, align with clinical evaluation of patients, and incorporate objective diagnostic criteria. This classification recognizes that myocardial infarction occurs in three clinical settings. First, myocardial infarction can arise spontaneously due to an acute coronary pathology (**primary myocardial infarction**). While most presentations are due to atherothrombosis, alternative coronary pathologies can be identified using coronary imaging and functional testing. Second, myocardial infarction may result from another acute condition causing oxygen supply–demand imbalance (**secondary myocardial infarction**). Here, objective criteria are proposed to guide the use of cardiac and invasive or non-invasive coronary imaging, promoting consistent diagnosis and informing treatment. Third, myocardial infarction may occur as a complication of percutaneous or surgical cardiac procedures (**procedure-related**

Nonstandard Abbreviations and Acronyms	
ACC	American College of Cardiology
ACS	Acute coronary syndrome
AHA	American Heart Association
AI	Artificial intelligence
CAD	Coronary artery disease
CABG	Coronary artery bypass graft(ing)
CCTA	Coronary computed tomography angiography
CK-MB	Creatine kinase-muscle/brain isoenzyme
CKD	Chronic kidney disease
CMR	Cardiac magnetic resonance
cMyC	Cardiac myosin-binding protein
CCT	Computed tomography
ECG	Electrocardiogram
GFR	Estimated glomerular filtration rate
ESC	European Society of Cardiology
FFR	Fractional flow reserve
ICD	International Classification of Diseases
FR	Instantaneous wave-free ratio
IVUS	Intravascular ultrasound
LBBB	Left bundle branch block
LGE	Late gadolinium enhancement
LMICs	Low- and middle-income countries
MINOCA	Myocardial injury with non-obstructive coronary arteries
MINS	Myocardial injury after non-cardiac surgery
NSTEMI	Non-ST-segment elevation myocardial infarction
OCT	Optical coherence tomography
PCI	Percutaneous coronary intervention
PET	Positron emission tomography
PMI	Peri-operative myocardial injury
RFR	Resting full-cycle ratio
SCAD	Spontaneous coronary artery dissection
SPECT	Single photon emission computed tomography
STEMI	ST-segment elevation myocardial infarction
TAVI	Transcatheter aortic valve implantation
UDMI	Universal definition of myocardial infarction
URL	Upper reference limit
WHF	World Heart Federation

myocardial infarction). The diagnosis of procedure-related myocardial infarction does not rely on cardiac biomarker thresholds and is comparable for percutaneous and surgical procedures. In each of these three settings, guidance is provided on the interpretation of

symptoms and signs of myocardial ischemia, cardiac biomarkers, and imaging. Alignment of the classification of myocardial infarction with International Disease Classification (ICD)-11 codes, its distinction from myocardial injury, and implications for patients, public health, and research are discussed. It is anticipated that adoption of the classification in practice and research will ensure the diagnosis of myocardial infarction is meaningful for patients and clinicians with clear implications for care across healthcare settings.

2. CONTEXT

The Fifth UDMI is a joint statement from the European Society of Cardiology (ESC), the American College of Cardiology (ACC), the American Heart Association (AHA), and the World Heart Federation (WHF). This is a consensus statement and should be applied in conjunction with national and regional clinical practice guidelines.^{1–3} The statement was written by a global task force consisting of a multidisciplinary group of experts with representation from 12 countries across 5 continents. It has undergone independent peer review by a committee comprising 18 members selected by the above societies as well as by a panel of patients and representatives selected from the European Association for Cardio-Thoracic Surgery (EACTS), the Society of Thoracic Surgeons (STS), Clinical Practice Guidelines Committee and National Cardiac Societies of the ESC. Following this writing and peer review process the document was approved for publication.

The work of this task force builds on the success of prior consensus statements, which have improved the standardization of the diagnosis of myocardial infarction worldwide. The first international task force to define the diagnosis of myocardial infarction was established by the World Health Organization (WHO) in 1979, when the diagnosis was based on symptoms, electrocardiographic criteria, and cardiac enzymes, but excluded coronary angiography.⁴ This was updated in 2000, when the consensus statement was led by the ESC and ACC. This document was subsequently considered to be the first universal definition.⁵ Revision was necessary with the introduction of cardiac troponin testing and increasing use of coronary angiography in practice, and subsequent iterations in 2007,⁶ 2012,⁷ and 2018⁸ addressed the opportunities and challenges introduced by the development of high-sensitivity cardiac troponin assays and further advances in cardiac imaging. With these iterations, the UDMI has encouraged clinicians to consider the underlying mechanism of myocardial infarction and myocardial injury, stimulated research, and promoted the use of sex-specific diagnostic criteria. This important international scientific effort will continue to require regular revision as new evidence emerges.

Making the diagnosis of myocardial infarction promptly and accurately is important as treatment improves survival. Myocardial infarction can have a substantial impact on future health, well-being, and livelihood. Disruption of an atherosclerotic plaque resulting in thrombosis (atherothrombosis) is the predominant cause of myocardial infarction. However, due to greater access to coronary angiography, intravascular imaging, and functional testing, other acute coronary pathologies, such as spontaneous coronary artery dissection, coronary embolism, and vasospasm are being increasingly identified. Treatment varies depending on the underlying mechanism, but the initial approach to evaluation of a patient with suspected myocardial infarction is similar. In parallel, cardiac troponin testing has become widely available in both ambulatory and hospital settings, and with this, our recognition of alternative causes of acute and chronic myocardial injury.

The classification of disease should be clinically relevant, widely applicable, and inform the approach to management and prognosis. To achieve these goals the classification of myocardial infarction needs to reflect the underlying pathophysiology with objective and reproducible diagnostic criteria that can be consistently applied in practice. This is necessary to establish trust and ensure the implications of the diagnosis are clearly understood by both clinicians and the person whom it affects. To support adoption in practice, ICD codes that align with the classification are required. It also needs to recognize that not all healthcare systems have access to the coronary or cardiac imaging modalities that are required to identify the underlying mechanism and to confirm the diagnosis of myocardial infarction.⁹

3. OBJECTIVE

The Fifth Universal Definition of Myocardial Infarction provides an updated, evidence-based definition and classification of myocardial infarction to support consistent diagnosis in both clinical practice and research. A simplified classification based on the underlying pathophysiology and clinical context is proposed, with objective criteria to guide treatment.

The Universal Definition was developed jointly by an ESC/ACC/AHA/WHF Task Force, under the auspices of the ESC Clinical Practice Guidelines Committee, in accordance with established ESC processes agreed by the participating societies, and is endorsed by the EACTS and STS, with affirmation of value by SCAI. The Task Force worked collaboratively with the World Health Organization (WHO) on the clinical taxonomy and proposed ICD-11 codes which were under review by the Classification and Statistics Advisory Committee of the WHO Family of International Classifications at the time of publication. A summary of studies referenced in support of the statement are included in the Supplementary data online *Evidence Tables*.

4. WHAT IS NEW

The classification of myocardial infarction in the Third UDMI and the expansion in the Fourth UDMI were major advances, though some aspects remained difficult to apply consistently in practice. This is particularly true when myocardial infarction results from another acute illness or a coronary process other than atherothrombosis (previously type 2) or arises following coronary intervention (previously type 4a–c) or cardiac surgery (previously type 5).

A new clinical classification of myocardial infarction is proposed (Figure 1) based on pathophysiology, aligned with clinical evaluation of patients and supported by objective diagnostic criteria. Key updates from the Fourth UDMI are outlined in Table 1.

New or updated concepts:

- The criteria for acute and chronic myocardial injury are refined and the underlying mechanisms discussed (see *Section 6*). Sex-specific 99th percentile upper reference limits for cardiac troponin define myocardial injury to avoid systematic bias and the under-recognition of both myocardial infarction and other cardiac conditions associated with myocardial injury in female patients.
- The previous numerical classification of myocardial infarction has been replaced. Myocardial infarction is now classified into one of three clinical types:
- **Primary myocardial infarction:** spontaneous presentation due to a primary acute coronary pathology
- **Secondary myocardial infarction:** resulting from myocardial oxygen supply–demand imbalance due to another acute condition
- **Procedure-related myocardial infarction:** occurring as a complication of a percutaneous or surgical cardiac procedure.
- Primary myocardial infarction is likely when symptoms or signs of acute myocardial ischemia and injury arise spontaneously, and no alternative acute condition or cardiac procedure has triggered the presentation. The diagnosis is confirmed and etiology defined if atherothrombosis or an alternative acute coronary pathology is identified (see *Section 7.2.1*).
- Secondary myocardial infarction is considered when acute myocardial ischemia and injury is secondary to an alternative acute condition resulting in a myocardial oxygen supply–demand imbalance. The diagnosis is confirmed if ischemia is a consequence of obstructive coronary artery disease without acute coronary pathology, or myocardial injury results in a new or presumed new regional wall motion abnormality or absence of viable myocardium (see *Section 7.2.2*).
- Procedure-related myocardial infarction is considered when acute myocardial ischemia and injury occur due to a complication of any cardiac procedure (percutaneous or surgical) within 30 days. The




Figure 1. Central Illustration.

Fifth Universal Definition of Myocardial Infarction. Myocardial infarction can occur in three clinical settings: spontaneously due to a primary acute coronary pathology, secondary to an acute condition, or following a cardiac procedure. For image description, please refer to the figure legend and surrounding text.

diagnosis is confirmed if a coronary complication is identified **or** myocardial injury results in a new or presumed new regional wall motion abnormality or absence of viable myocardium. However, both a complication **and** a new or presumed new regional wall motion abnormality or absence of viable myocardium are required when the complication arises during the index procedure, or the procedure is performed in the management of acute myocardial infarction (see *Section 7.2.3*).

- The diagnosis of myocardial infarction following sudden death is discussed (see *Section 8*) and new, objective criteria are provided for silent or unrecognized myocardial infarction (see *Section 9*).
- The definition of MINOCA is updated to 'myocardial injury with non-obstructive coronary arteries' in recognition that this is a working rather than final diagnosis (see *Section 11*).
- Accelerated diagnostic pathways using high-sensitivity cardiac troponin assays are introduced and guidance is provided for the areas of practice in which the interpretation of cardiac troponin is challenging (see *Section 14*).
- New guidance is provided for the diagnosis of myocardial infarction in low-resource settings (see *Section 19*).

- International Classification of Disease 11th Revision (ICD-11) codes for use in hospital episode statistics and public health monitoring are proposed that align with the updated classification of myocardial infarction (see *Section 18*) to encourage implementation and further research (see *Section 20*).

5. DEFINITION OF MYOCARDIAL INFARCTION

Myocardial infarction is defined pathologically as myocardial cell death due to prolonged periods of ischemia caused by insufficient or absent coronary blood flow and reperfusion injury.^{10,11} A histopathological diagnosis requires autopsy, but a clinical diagnosis of myocardial infarction can be confirmed through the joint evaluation of clinical features, biomarkers, electrocardiography, and coronary or cardiac imaging. A clinical diagnosis requires indirect evidence of irreversible cell death and confirmation that cell death was caused by myocardial ischemia. No single criterion can confirm the diagnosis of acute myocardial infarction; therefore, several criteria are necessary that collectively increase the likelihood that myocardial infarction has occurred.

The clinical diagnosis of myocardial infarction requires evidence of acute myocardial injury (see *Section 6*) **and**

Table 1. Comparison of the Fourth and Fifth Universal Definitions of Myocardial Infarction

Fourth UDMI	Fifth UDMI—what's new	Rationale for change
Numerical classification of myocardial infarction: types 1, 2, 3, 4a, 4b, 4c, 5.	Clinical classification of myocardial infarction; primary, secondary, or procedure related.	To reflect underlying pathophysiology, align with clinical evaluation, and enable consistent application.
Type 1 myocardial infarction Restricted to atherothrombosis.	Primary myocardial infarction Includes all acute coronary pathologies: atherothrombosis; spontaneous coronary artery dissection; coronary embolism; vasospasm; and restenosis, stent thrombosis, or graft failure >30 days from procedure.	Prioritizes sensitivity to avoid missing any primary acute coronary pathology. Promotes coronary angiography and adjunctive testing to confirm diagnosis and identify underlying acute coronary pathology with new diagnostic codes for each mechanism. Reclassifies late stent/graft failure as <i>de novo</i> disease, not a procedure-related complication.
Type 2 myocardial infarction Myocardial oxygen supply–demand imbalance due to either: • Acute coronary pathology other than atherothrombosis, including spontaneous coronary artery dissection, coronary embolism, and vasospasm <i>or</i> • An alternative acute condition with or without coronary artery disease.	Secondary myocardial infarction Myocardial oxygen supply–demand imbalance due to an alternative acute condition: <i>and</i> • obstructive coronary artery disease without acute coronary pathology <i>and/or</i> • new or presumed new regional wall motion abnormality or absence of viable myocardium.	Prioritize specificity to differentiate myocardial infarction from acute myocardial injury in conditions resulting in oxygen supply–demand imbalance. Objective diagnostic criteria to allow consistent application in practice and identify patients in whom the diagnosis has treatment implications.
Type 3 myocardial infarction Cardiac death where myocardial infarction is the likely cause, but death occurs before diagnostic testing is performed.	Term removed. Where myocardial infarction is the likely cause of death, clinical classification (primary, secondary, or procedure-related) should be applied, based on setting or post-mortem findings.	Limited use in clinical practice.
Type 4 and 5 myocardial infarction Type 4a myocardial infarction: cardiac troponin concentration >5 times the 99th percentile with myocardial ischemia, imaging evidence of new loss of viable myocardium, or angiographic evidence of a procedural complication within 48 h. Type 4b due to stent thrombosis and type 4c due to stent restenosis, at any time after the procedure. Type 5 myocardial infarction: coronary artery bypass graft-related; cardiac troponin >10 times the 99th percentile, with new pathological Q waves, imaging evidence of new loss of viable myocardium, or angiographic evidence of a graft occlusion within 48 h.	Procedure-related myocardial infarction Coronary complication within 30 days of a cardiac procedure resulting in acute myocardial injury with one or more features. Both features are required when the complication arises during the procedure or in the setting of acute myocardial infarction. • Evidence of a coronary complication of percutaneous intervention or cardiac surgery • New or presumed new regional wall motion abnormality or absence of viable myocardium.	Prioritizes specificity to ensure the diagnosis is clinically meaningful, with objective cardiac imaging criteria. Diagnostic criteria the same for all cardiac procedures to enable meaningful comparisons. Excludes late stent/graft failure as a procedural complication as it typically reflects <i>de novo</i> disease.

UDMI indicates universal definition of myocardial infarction.

one or more of the following: symptoms or other evidence of acute myocardial ischemia that include new ischemic changes or pathological Q waves on the electrocardiogram (ECG); imaging evidence of an acute coronary pathology; or new loss of viable myocardium or a new regional wall motion abnormality in a pattern consistent with an ischemic etiology.⁸

These criteria are easier to apply when the clinical presentation is spontaneous acute onset of chest discomfort in the absence of other precipitating factors, and the initial evaluation should follow clinical practice guidelines.^{1,2} The diagnosis is often straightforward when the presentation is with central and radiating chest pain or discomfort (see *Section 12*) accompanied by ST-segment elevation on the ECG (see *Section 13*). However, many people do not have classical symptoms, and the ECG is often normal or not diagnostic of myocardial ischemia.¹² Thus, in a substantial proportion of cases, the working diagnosis of myocardial infarction is suggested by cardiac biomarker testing (see *Section 14*). Among those with acute myocardial injury but without ST-segment elevation, subsequent coronary or cardiac imaging identifies

an alternative diagnosis in around 1 in 20 cases, excluding the diagnosis of myocardial infarction.^{13,14} The clinical diagnosis is even more challenging in settings in which myocardial injury is common—acute illness or following a cardiac procedure. In these settings, where the cause of myocardial injury can be ischemic or non-ischemic, imaging should be encouraged to confirm or refute the diagnosis of myocardial infarction (see *Section 15*).

6. ACUTE AND CHRONIC MYOCARDIAL INJURY

The term *myocardial injury* was defined in the Third UDMI as an elevated cardiac troponin I or T level with at least one value above the 99th percentile upper reference limit (URL) regardless of the underlying cause.⁷ The Fourth UDMI added the terms *acute myocardial injury* and *chronic myocardial injury*.⁸ Both acute and chronic myocardial injury are common, but their prevalence varies depending on the approach to testing¹⁵ and the assay,¹⁶ with chronic myocardial injury up to five times more frequent with cardiac troponin T than cardiac troponin I.¹⁷

6.1. Acute Myocardial Injury

Acute myocardial injury is defined as a rise and/or fall in cardiac troponin with at least one value above the sex-specific 99th percentile URL (*Box 1*). Sex-specific thresholds are necessary to avoid a systematic bias and the under-recognition of myocardial injury in female patients (see *Section 14.1.1*).

Box 1: Criteria for Acute Myocardial Injury

Acute myocardial injury is defined as a rise and/or fall in cardiac troponin I or T with at least one value above the sex-specific 99th percentile URL.

Acute myocardial injury can be caused by ischemic and non-ischemic mechanisms. When demonstrated to be due to ischemia, acute myocardial injury is the hallmark of myocardial infarction, and it should be classified as such when the diagnostic criteria for myocardial infarction are met. However, many other cardiac and non-cardiac conditions are commonly associated with acute myocardial injury without myocardial ischemia.¹⁸ Non-ischemic mechanisms for acute myocardial injury include inflammation, physiological stress response, catecholamine stress, cardiotoxic agents, and trauma (*Table 2*).

In some cases, there may be multiple mechanisms responsible for acute myocardial injury, for example in sepsis, where both inflammation and the physiological stress response may contribute.¹⁹

Whether acute myocardial injury results in transient or permanent effects on myocardial function varies depending on the severity of the injury and underlying condition. In several conditions associated with significant increases in cardiac troponin, e.g. Takotsubo syndrome, severe left ventricular dysfunction and regional wall motion abnormalities may resolve without evidence of permanent damage, suggesting the underlying mechanism is due to a functional abnormality rather than myocardial ischemia or necrosis.^{20,21} In myocarditis, myocardial injury-related cardiac dysfunction may be fully reversible.²² Elevated cardiac troponin may not always reflect necrosis,^{23–25} as seen in athletes after strenuous exercise, where troponin release likely reflects reversible injury,²⁶ though some irreversible damage cannot be excluded.²⁷

Acute myocardial injury requires clinical evaluation to determine its etiology, assess functional impact, and identify ischemia, in which case, coronary and cardiac imaging is required to confirm or exclude myocardial infarction.

6.2. Chronic Myocardial Injury

Chronic myocardial injury is a term used to describe persistently elevated cardiac troponin due to an underlying

Table 2. Mechanisms of Acute Myocardial injury

Mechanism	Condition
Ischemia	Myocardial infarction
Inflammation	Myocarditis (autoimmune, infectious, toxic)
	Heart transplant rejection
	Cytokine mediated in sepsis
Hemodynamic stress	Supply–demand imbalance*
	Tachy- or brady-arrhythmia
	Acute heart failure
	Acute pulmonary embolism
	Malignant hypertension
Physiological stress	Strenuous exercise
Catecholamine stress	Takotsubo syndrome
	Subarachnoid hemorrhage
	Stroke
	Epileptic seizures
	Pheochromocytoma
Toxicity	Anthracyclines
	ErbB (erythroblastic oncogene B)-targeted therapies
	Tyrosine kinase inhibitors
	Immune checkpoint inhibitors
	Radiation induced
Trauma	Electrical cardioversion
	Percutaneous coronary or structural intervention
	Catheter ablation or device implantation
	Cardiac surgery
	Cardiac contusion

*Any acute condition that results in a myocardial oxygen imbalance with reduced supply (hypoxia, anemia, hypotension) and/or increased demand (tachycardia, hypertension).

cardiac condition or a non-cardiac condition associated with cardiac remodeling (*Box 2*). Cardiac troponin can be elevated chronically due to three mechanisms: increased release from the myocardium due to a cardiac condition; reduced clearance from the circulation; or analytical interference that can mimic either (see *Section 14*).

Box 2: Criteria for Chronic Myocardial Injury

Chronic myocardial injury is **considered** if two or more cardiac troponin I or T values are above the sex-specific 99th percentile URL when testing is performed in a stable clinical setting. Chronic myocardial injury is **confirmed** when a cardiac or non-cardiac condition associated with cardiac remodeling is identified and is excluded when the elevation in cardiac troponin is explained by analytical interference or reduced clearance.

At levels within the normal reference range cardiac troponin is cleared by the kidneys, whereas at higher concentrations elimination is thought to be predominantly by the liver.^{28,29} A halving of the estimated glomerular filtration rate (eGFR) will lead to a doubling of the basal level of cardiac troponin.²⁸ Given that the majority of patients with chronic elevations in cardiac troponin only have minor increases above the 99th percentile, it is important to consider the effect of reduced renal function when interpreting chronically elevated cardiac troponin levels (see *Section 16.3*).

Atherosclerotic coronary artery disease (CAD) is a chronic, progressive lifelong condition³⁰ and chronic myocardial injury can arise as a consequence of low-grade recurrent myocardial ischemia in patients with chronic coronary syndromes or ischemic cardiomyopathy, or may arise due to hemodynamic stress in chronic congestive cardiac failure or severe hypertension (Table 3). It may also occur due to left ventricular hypertrophy and in other cardiomyopathies or structural conditions, such as valvular and congenital heart disease.^{31–33} In many of these conditions the presence of chronic myocardial injury is an adverse prognostic feature.³⁴ However, persistently elevated cardiac troponin concentrations are also observed in older people without detectable heart disease. Furthermore, elevated cardiac troponin may not be due to myocardial injury at all but can arise due to *in vivo* or *ex vivo* antibody-mediated assay interference (see *Section 14.1.2*). The detection of a persistently elevated cardiac troponin concentration merits further investigation to determine whether this is due to chronic myocardial injury, in which case cardiac imaging may help to determine whether this is a consequence of structural or coronary heart disease.

Table 3. Mechanisms of Chronic Myocardial Injury

Etiology	Condition
Ischemia	Chronic coronary syndrome
	Ischemic cardiomyopathy
Other cardiomyopathies	Hypertrophic cardiomyopathy
	Hypertensive cardiomyopathy
	Dilated cardiomyopathy
	Chronic inflammatory cardiomyopathy
	Restrictive cardiomyopathy
	Infiltrative cardiomyopathy
	Uremic cardiomyopathy in chronic kidney disease
Hemodynamic stress	Chronic heart failure
	Systemic hypertension
	Pulmonary hypertension
Structural	Valvular heart disease
	Congenital heart disease
	Cardiac tumour

In an acute setting, it can be difficult to determine whether elevated cardiac troponin is due to acute or chronic myocardial injury, especially if serial measurements are taken at short intervals.³⁵ The priority in the acute setting is to exclude acute myocardial injury or infarction using a validated accelerated diagnostic pathway (see *Section 14.1.5*). In those patients where acute myocardial injury or infarction is excluded, troponin testing can be repeated in an outpatient setting or values compared with those from a previous assessment.^{17,36} Injury is more likely to be chronic and stable than progressive if the change is less than 20% between troponin measurements, but clinicians should be aware that this criterion is simply based on the analytical imprecision of cardiac troponin assays. If cardiac troponin levels remain elevated, then further investigation could be considered to confirm whether this is a result of chronic myocardial injury due to an underlying cardiac condition.

7. CLASSIFICATION OF MYOCARDIAL INFARCTION

7.1. Approaches to the Classification of Myocardial Infarction



Several pathophysiological mechanisms can lead to myocardial infarction. A logical classification of myocardial infarction should reflect the underlying pathophysiology, use objective diagnostic criteria, and be intuitive for consistent application and clear understanding by clinicians and patients.

7.1.1. Classification Based on Etiology

The Third UDMI introduced a classification that was based on the underlying cause, identifying five subtypes of myocardial infarction.⁷ This was expanded to seven subtypes in the fourth iteration.⁸ In recognition that cardiac troponin levels are elevated in many cardiac and non-cardiac conditions, two further diagnostic classifications were introduced: acute non-ischemic myocardial injury and chronic myocardial injury, defined by whether a rise and/or fall in cardiac troponin levels was observed on serial testing (see *Section 6*).

The introduction of different types of myocardial infarction represented a major conceptual step forward,³⁷ but has been challenging to apply consistently in practice.^{38,39} This is particularly true when myocardial infarction results from another acute illness or an acute coronary pathology other than atherothrombosis (previously type 2) or arises following coronary intervention (previously type 4a–c) or cardiac surgery (previously type 5).

Grouping non-atherosclerotic coronary causes (e.g. coronary dissection, embolism, or vasospasm) with myocardial oxygen supply–demand imbalance as type 2

myocardial infarction has limited its adoption in practice.⁴⁰ These patients follow very different diagnostic pathways and require different treatments. Under the Fourth UDMI, type 2 myocardial infarction was diagnosed two ways: retrospectively, when patients initially presumed to have type 1 myocardial infarction underwent coronary angiography that identified no acute coronary pathology; and prospectively, when supply–demand imbalance from another acute condition was evident at presentation. As many of these patients were not managed by cardiologists, the diagnosis was frequently made based on symptoms or signs of myocardial ischemia alone.⁴¹ Without coronary or cardiac imaging to confirm the cause or consequences of myocardial injury, a diagnosis of type 2 myocardial infarction had no direct treatment implications for many patients.

The previous classifications of type 4a and type 5 myocardial infarction following coronary intervention and cardiac surgery have been equally challenging to apply in practice as myocardial injury is ubiquitous in these settings and the classifications did not address complications following procedures other than revascularization. Thresholds that were 5 and 10 times greater than the 99th percentile URL of the cardiac troponin assay were arbitrary and applied along with additional criteria for the diagnosis of type 4a and type 5 myocardial infarction, respectively.⁸ However, for type 5 myocardial infarction, a large prospective international study showed that 97.5% of patients undergoing cardiac surgery had cardiac troponin levels more than 10 times the URL of the assay, and that thresholds more than 218 times the URL (95% confidence interval 40 to 318) were required to identify those at increased risk of peri-procedural mortality.⁴² Following coronary artery bypass graft (CABG) surgery several alternative criteria have been proposed⁴³ using cardiac troponin thresholds up to 170 times the URL or using less sensitive and specific biomarkers, such as creatine kinase-muscle/brain isoenzyme (CK-MB), which may no longer be widely available.⁴⁴ Likewise, definitions using alternative thresholds led to markedly different frequencies of the diagnosis of procedural myocardial infarction (type 4a) following percutaneous coronary intervention (PCI) and also had different prognostic significance.⁴⁵ Recent consensus statements have focused on the identification of prognostically relevant myocardial injury after coronary surgery and percutaneous coronary procedures.^{46,47} However, prognostic thresholds are not a substitute for clinical diagnosis and as the URLs between assays are not biologically equivalent the use of multiples of the URL to confirm the diagnosis is problematic. This variation in the diagnostic criteria has made it challenging to interpret myocardial infarction as an endpoint in clinical trials,⁴⁸ and to advise patients on the risks and benefits of revascularization using either approach.

Finally, while a classification that used alpha-numerical terms has encouraged research, these terms did not directly convey information about the underlying mechanism of myocardial infarction. Therefore, they were of less value when communicating with patients and with many clinicians.

7.1.2. Classification Based on the Electrocardiogram

The most widely adopted classification in practice stratifies myocardial infarction into two groups based on the presenting ECG, according to the presence or absence of ST-segment elevation (see *Section 1.3*). The recently released ICD-11 reflects the widespread use of this classification by introducing distinct codes for ST-segment elevation myocardial infarction (STEMI) (BA41.0) and non-ST-segment elevation myocardial infarction (NSTEMI) (BA41.1).^{1,2} Regional ST-segment elevation generally reflects an acute coronary occlusion requiring immediate restoration of blood flow to reduce infarct size.⁴⁹ However, not all patients with acute coronary occlusion demonstrate ST-segment elevation on the 12-lead ECG and other confounders (such as bundle branch block) can obscure interpretation of the ST segment (see *Section 1.3*).^{50–53} While imperfect, this classification is simple and useful to identify those patients likely to benefit from immediate coronary intervention or fibrinolysis.^{49,54} The terms in isolation offer limited insight into the underlying pathophysiological mechanism of myocardial infarction, however, and are therefore less useful at guiding management beyond the acute presentation.

7.2. Clinical Classification of Myocardial Infarction

Building on the existing clinical definition and classifications of myocardial infarction, while addressing some of the limitations that have prevented consistent application in practice, a simplified new clinical classification of myocardial infarction is proposed. It is aligned with the usual pathway for evaluation of patients with suspected myocardial infarction, considers the underlying pathophysiology, and incorporates more objective diagnostic criteria to enable more consistent application in practice, research, and comparative clinical trials.

This simplified clinical classification recognizes that myocardial infarction can arise in three settings: (i) spontaneously due to a primary acute coronary pathology (**primary myocardial infarction**); (ii) secondary to another acute condition resulting in myocardial oxygen supply–demand imbalance (**secondary myocardial infarction**); or (iii) as a complication following a cardiac procedure (**procedure-related myocardial infarction**). This revised classification aims to increase adoption and consistent application in practice (see Central illustration Figure 1).

7.2.1. Primary Myocardial Infarction

Primary myocardial infarction is considered in a patient with spontaneous onset of symptoms or signs of acute myocardial ischemia and acute myocardial injury where no alternative acute condition or cardiac procedure has triggered the presentation. While the majority are due to atherothrombosis, it includes all primary acute coronary pathologies. The diagnostic criteria for primary myocardial infarction prioritize sensitivity to minimize missing an acute coronary pathology and encourages coronary angiography to confirm the diagnosis and define the underlying mechanism (Box 3).

Box 3: Diagnostic Criteria for Primary Myocardial Infarction

The diagnosis is **likely** in patients with acute *myocardial injury* identified through the measurement of cardiac troponin when one or more clinical or electrocardiographic features are present:

- Symptoms consistent with acute myocardial ischemia
- New or presumed new ischemic changes on the electrocardiogram
- Development of pathological Q waves.

The diagnosis is **confirmed** with one or more additional features on imaging:

- Identification of an acute coronary pathology comprising atherothrombosis, spontaneous coronary artery dissection, vasospasm, or embolism; or restenosis, stent thrombosis, or bypass graft failure more than 30 days from revascularization
- Development of a new or presumed new regional wall motion abnormality and/or absence of viable myocardium in a pattern consistent with an ischemic etiology.

The diagnostic pathway for patients with possible primary myocardial infarction is illustrated in Figure 2. The diagnosis is likely in those with acute myocardial injury presenting with the spontaneous onset of symptoms suggestive of myocardial ischemia (see *Section 12*) or changes on the ECG consistent with acute myocardial ischemia (see *Section 13*). When more clinical features are present, the likelihood of myocardial infarction is greater. Coronary angiography and/or cardiac imaging are required to confirm the diagnosis, and invasive assessment with angiography or intracoronary imaging is essential to identify the underlying acute coronary pathology for further classification (Figure 3). However, where coronary angiography and cardiac imaging are not available or are not considered appropriate, a final diagnosis of primary myocardial infarction can be made based on an evaluation that considers biomarkers, electrocardiography, and clinical criteria alone.

Although the term primary myocardial infarction is used in this classification to differentiate from secondary and procedure-related myocardial infarction, it is synonymous with spontaneous myocardial infarction. In practice, when this diagnosis is made, clinicians may simply refer to the diagnosis as myocardial infarction.

In primary myocardial infarction an acute coronary pathology can cause a total or sub-total coronary artery occlusion, presenting with or without ST-segment elevation. Atherothrombosis (thrombus associated with plaque rupture or erosion) is the most common acute coronary pathology in primary myocardial infarction.^{55–57} In addition to causing acute coronary occlusion, thrombus formation can also cause distal embolization.⁵⁸ The degree of atherosclerosis and thrombus in the culprit lesion varies greatly and non-flow-limiting stenoses are also common causes of myocardial infarction.⁵⁹ Other less common but equally important acute coronary pathologies in primary myocardial infarction, with or without ST-segment elevation, include spontaneous coronary artery dissection (SCAD), epicardial or microcirculatory vasospasm (hence forth 'vasospasm'), or embolism from a non-coronary source; as well as restenosis, stent thrombosis, or graft failure arising more than 30 days from a revascularization procedure (see *Section 7.2.3* for events within 30 days). SCAD should be considered particularly in female patients under 50 years of age and in pregnancy or the post-partum period.^{60,61}

Coronary angiography does not always identify the underlying acute coronary pathology that is the cause of the primary myocardial infarction. Here, adjunctive intravascular imaging, such as optical coherence tomography (OCT) and intravascular ultrasound (IVUS) may be helpful (see *Section 15*). In some, both coronary and cardiac imaging are necessary to confirm or exclude the diagnosis of primary myocardial infarction. This is particularly relevant in patients with non-obstructive coronary arteries (see *Section 11*). In these patients, cardiac magnetic resonance (CMR) imaging is the modality of choice to confirm the diagnosis of myocardial infarction,^{1,62} but more often identifies an alternative cause of acute myocardial injury such as myocarditis or Takotsubo syndrome.⁶³

When coronary angiography is not performed, the diagnosis of STEMI or NSTEMI can be applied with primary myocardial infarction of unknown etiology (Table 4). However, when coronary angiography is performed, and the underlying acute coronary pathology identified, the final diagnosis should reflect this. This information will provide important additional insight into future risk of coronary events and treatment.

7.2.2. Secondary Myocardial Infarction

The diagnosis of secondary myocardial infarction is considered in patients with acute myocardial injury secondary to an alternative acute condition, causing




Figure 2. Diagnostic pathway for possible primary myocardial infarction.

NSTEMI, non-ST-segment elevation myocardial infarction; STEMI, ST-segment elevation myocardial infarction. ^aOr other electrocardiographic changes indicative of possible acute coronary occlusion (Figure 9). ^bFor patients with a working diagnosis of STEMI, acute myocardial injury is necessary to confirm the final diagnosis. ^cRise and/or fall of cardiac troponin I or T values and with at least one value above the sex-specific 99th percentile upper reference limit on serial testing. ^dWhere coronary angiography and cardiac imaging are not available or are not considered appropriate, a final diagnosis of primary myocardial infarction can be assigned based on an evaluation that considers biomarkers, electrocardiography, and clinical criteria alone without additional confirmatory features and the codes BA41.09 or BA41.19 applied as the etiology is unknown. ^eThis includes patients with a working diagnosis of myocardial injury with non-obstructive coronary arteries (MINOCA). For image description, please refer to the figure legend and surrounding text.

a mismatch between myocardial oxygen supply and demand when this results in symptoms or signs of myocardial ischemia.^{64–66} The diagnosis of secondary myocardial infarction is confirmed if the myocardial ischemia resulting from supply–demand mismatch is a *consequence* of obstructive CAD without evidence of an acute coronary pathology, or the resulting myocardial injury is sufficient to cause a new regional wall

motion abnormality or loss of myocardial viability. The diagnostic criteria prioritize specificity and encourage coronary angiography and cardiac imaging to reduce uncertainty, increasing the likelihood that the diagnosis has treatment implications for the patient (Box 4). If imaging provided evidence of an acute coronary pathology (Figure 3) then a diagnosis of primary myocardial infarction would be confirmed.




Figure 3. Primary myocardial infarction due to acute coronary pathology. For image description, please refer to the figure legend and surrounding text.

Box 4: Diagnostic Criteria for Secondary Myocardial Infarction

The diagnosis is **considered** in patients with acute myocardial injury identified through the measurement of cardiac troponin who present with an *alternative acute condition resulting in myocardial oxygen supply–demand mismatch*, when one or more clinical or electrocardiographic features are present:

- Symptoms consistent with acute myocardial ischemia
- New or presumed new ischemic changes on the electrocardiogram
- Development of pathological Q waves.

The diagnosis is **likely** with one or more additional features:

- Known coronary artery disease
- A strong clinical suspicion due to the extent of myocardial ischemia on the electrocardiogram or cardiac troponin elevation.

For patients in whom coronary and/or cardiac imaging is feasible and appropriate, the diagnosis is **confirmed** if one or more of the following features are present:

- Obstructive coronary artery disease, defined as $\geq 70\%$ stenosis in an epicardial vessel by angiography (or $\geq 50\%$ stenosis in an epicardial vessel that is flow-limiting on physiological assessment) **without** an acute coronary pathology
- Development of a new or presumed new regional wall motion abnormality or absence of viable myocardium in a pattern consistent with an ischemic etiology.

The pathway for patients in whom the diagnosis of secondary myocardial infarction is being considered is illustrated in Figure 4. Myocardial oxygen supply–demand imbalance can occur as a result of increased demand (e.g. tachycardia or severe hypertension) or reduced supply (e.g. hypotension, hypoxia, or anemia) (Figure 5). Therefore, acute myocardial ischemia can arise secondary to oxygen supply–demand imbalance in many acute conditions, the most common being tachyarrhythmia.^{41,67} The duration and severity of the trigger for supply–demand imbalance and whether the individual has underlying obstructive CAD or structural cardiac disease will influence the extent and severity of ischemia.⁶⁴ Furthermore, two or more triggers often coexist.⁶⁸ Therefore, it is not possible to define thresholds for any of the triggers of supply–demand imbalance that could be reliably applied to all.

Table 4. ICD-11 Codes for the Classification of Myocardial Infarction

Final diagnosis	ICD-11	Sixth digit stem code/extension code
Acute ST-segment elevation myocardial infarction (STEMI)*	BA41.0	Primary myocardial infarction BA41._1 atherothrombosis BA41._2 spontaneous coronary artery dissection BA41._3 coronary embolism BA41._4 coronary vasospasm BA41._5 restenosis >30 days from procedure BA41._6 stent thrombosis >30 days from procedure BA41._7 graft failure >30 days from procedure BA41._8 undetermined etiology BA41._9 unknown etiology (no coronary imaging) Secondary myocardial infarction BA41._A/ICD-11 code for alternative acute condition Procedure-related myocardial infarction BA41._B due to a complication of any percutaneous cardiac procedure within 30 days BA41._C due to a complication of any open cardiac surgical procedure within 30 days
Acute non-ST-segment elevation myocardial infarction (NSTEMI)	BA41.1	
Other diagnoses		
Unstable angina	BA40	
Unspecified myocardial infarction	BA41.Z	Sudden death without evaluation†
Unrecognized myocardial infarction	BA50	

ICD-11, 11th edition of the International Classification of Diseases.
*Including electrocardiographic findings of acute coronary occlusion (Figure 9).
†See Figure 8.

In this setting, symptoms of myocardial ischemia are challenging to interpret as patients are less likely to report chest discomfort and are more likely to have other symptoms due to the underlying acute condition.^{69,70} While myocardial ischemia secondary to supply–demand imbalance can manifest as regional ST-segment depression or T-wave inversion, these changes are often global. Acute myocardial injury is common in many conditions that result in supply–demand imbalance, but current cardiac troponin assays cannot differentiate ischemic from non-ischemic causes.⁷¹ Given that the diagnosis of myocardial infarction in this setting cannot be reliably established using symptoms or signs of myocardial ischemia and cardiac troponin levels alone,^{72,73} additional clinical assessment and cardiac imaging are needed, when it is feasible and appropriate based on the clinical status, to confirm or exclude the diagnosis of secondary myocardial infarction. Most patients in this setting will not have obstructive CAD or imaging evidence of new loss of viable myocardium, and the diagnosis of secondary myocardial infarction can be excluded.^{74,75} It is important to note that acute illness, particularly sepsis or bleeding, can also trigger atherothrombosis; therefore, primary myocardial infarction is an important differential diagnosis. If myocardial infarction is excluded, the underlying etiology of acute myocardial injury should be determined when possible (Table 2).

If there is persistent or recurrent myocardial ischemia and primary myocardial infarction is considered as a differential diagnosis, invasive coronary angiography is indi-

cated.^{1,2} Otherwise, imaging is often deferred until after the acute condition has been treated and non-invasive approaches can be applied, such as the use of coronary computed tomography angiography (CCTA).⁷⁵ For patients with comorbidities, advanced frailty, or limited life expectancy due to the underlying acute condition, additional cardiac imaging is sometimes not appropriate. Hence, it may not be possible to definitively confirm the diagnosis of secondary myocardial infarction and clinical judgment is required. Secondary myocardial infarction is more likely when patients have known CAD or significant myocardial injury that affects cardiac function.⁷⁴ Those with obstructive coronary disease are also more likely to have myocardial ischemia on the ECG.⁷² These features can support a clinical diagnosis when imaging is not feasible or appropriate.

One study suggests that application of these diagnostic criteria for secondary myocardial infarction would reduce the frequency of the diagnosis of myocardial infarction in acute illness and identify those patients at highest risk.⁷⁴ A diagnosis of secondary myocardial infarction confirmed by coronary and/or cardiac imaging would have treatment implications for those with CAD³ and/or left ventricular systolic dysfunction.^{1,2} The application of objective criteria to confirm the diagnosis of secondary myocardial infarction should encourage cardiac imaging in this setting. Imaging may also have additional benefits through the identification of non-obstructive but clinically relevant CAD or other previously unrecognized cardiac conditions, such as non-ischemic cardiomyopathy or valvular heart disease.⁷²






Figure 4. Diagnostic pathway for possible secondary myocardial infarction.

^aMyocardial oxygen supply–demand imbalance can occur as a result of increased demand (e.g. tachycardia or severe hypertension) and/or reduced supply (e.g. hypotension, hypoxia, or anemia). ^bRise and/or fall of cardiac troponin values and with at least one value above the sex-specific 99th percentile upper reference limit on serial testing. ^cIf primary myocardial infarction is an unlikely differential diagnosis, then non-invasive computed tomography (CT) coronary angiography following recovery from acute illness may be considered. ^dObstructive coronary artery disease is defined as $\geq 70\%$ stenosis in an epicardial vessel by angiography or $\geq 50\%$ stenosis that is flow-limiting on physiological assessment. For image description, please refer to the figure legend and surrounding text.

7.2.3. Procedure-Related Myocardial Infarction

The diagnosis of procedure-related myocardial infarction is made when myocardial infarction arises as a complication of a cardiac procedure within 30 days of the procedure (Box 5). While 30 days is an arbitrary period, this is aligned with most studies of post-procedure complications.⁷⁶ A cardiac procedure is defined as coronary angi-

ography with or without intervention, structural cardiac intervention, or catheter ablation, and any open or minimally invasive cardiac surgical procedure, including CABG, valve replacement, or other procedures for structural heart conditions. The risk of procedure-related myocardial infarction is higher following PCI or CABG but is recognized following many other cardiac procedures. The new diagnostic




Figure 5. Secondary myocardial infarction due to supply-demand imbalance.

criteria for procedure-related myocardial infarction are the same whether the complication arises from a percutaneous interventional or open surgical approach.

Box 5: Diagnostic Criteria for Procedure-Related Myocardial Infarction

The diagnosis is **considered** in patients with a suspected coronary complication arising within 30 days of a cardiac procedure and acute *myocardial injury* identified through the measurement of cardiac troponin when one or more clinical or electrocardiographic features are present:

- Symptoms consistent with acute myocardial ischemia
- New or presumed new ischemic changes on the electrocardiogram
- Development of pathological Q waves
- Unexplained sudden clinical deterioration if under sedation.

Immediately following a procedure, the diagnosis is **likely** with higher cardiac biomarker levels that have increased from pre-procedural levels and continue to rise at 6 h (e.g. >5 times URL for intervention) and 24 h (e.g. >35 times URL for surgery).

The diagnosis is **confirmed** with one or more additional features, except when the complication arises during the procedure or the procedure is performed in the setting of an acute myocardial infarction, when both are required:

- Angiographic evidence of a coronary complication of percutaneous intervention (e.g. side branch occlusion, iatrogenic dissection, no or slow flow, recoil or early restenosis, stent thrombosis), or surgery (e.g. bypass graft failure, impingement of a coronary artery), or either (e.g. embolism, occlusion of coronary ostium during valve intervention)
- Cardiac imaging showing a new or presumed new regional wall motion abnormality or absence of viable myocardium in a pattern consistent with an ischemic etiology within the territory subtended by the coronary artery in which the complication arose.

The pathway for patients in whom the diagnosis of procedure-related myocardial infarction is being considered is illustrated in Figure 6. Symptoms can be difficult to interpret immediately after cardiac procedures. Many patients experience some chest discomfort,⁷⁷ but if the symptoms are persistent, increase in severity, or develop suddenly after the patient being symptom-free




Figure 6. Diagnostic pathway for possible procedure-related myocardial infarction.

^aRising cardiac troponin values with at least one value above the sex-specific 99th percentile AND pre-procedural value (if known) on serial testing. ^bImmediately following a procedure, the diagnosis is more likely with higher cardiac biomarker levels that have increased from pre-procedural levels and continue to rise at 6 h (e.g. >5 times upper reference limit [URL] for intervention) and 24 h (e.g. >35 times URL for surgery). For image description, please refer to the figure legend and surrounding text.

at the end of the procedure, they are more likely due to a complication. Adding to the challenge, patients often receive sedation for cardiac procedures and receive general anesthesia for cardiac surgery. Although it may be impossible to determine whether chest discomfort is present, a sudden deterioration in clinical condition (e.g. pulmonary edema, unexpected hypotension, tachycardia, or ventricular arrhythmia) may be a sign of acute myocardial ischemia.

Electrocardiographic changes following coronary intervention are helpful, but ST-segment deviation and T-wave inversion are common after cardiac surgery due to other conditions (e.g. pericarditis, pneumopericardium, and electrolyte disturbance).^{78,79} Furthermore, what may appear as pathological Q waves can be transient early after CABG.⁸⁰ Thus, the presence of electrocardiographic changes requires further evaluation, but is not sufficient to diagnose procedure-related myocardial infarction.

Routine cardiac troponin measurement before and after coronary intervention or cardiac surgery is not supported by some clinical practice guidelines^{3,81} or the European Association for Cardio-Thoracic Surgery,⁴⁷ but is encouraged by the European Association of Percutaneous Cardiovascular Interventions⁴⁶; practice varies across the world. When cardiac troponin is measured, some degree of acute myocardial injury is ubiquitous.^{82–84} The extent of injury differs markedly depending on the duration and complexity of the procedure.^{42,85} For these reasons, it is not possible to use a single diagnostic threshold for cardiac troponin in this setting and cardiac imaging is required to confirm or exclude the diagnosis of procedure-related myocardial infarction.

Cardiac troponin testing is, however, essential in patients with clinical features of a suspected or evident coronary complication immediately following a procedure to guide the use of cardiac imaging. In this setting cardiac troponin thresholds may be helpful (and are recommended in some consensus statements)^{8,86} to identify those in whom cardiac imaging is needed to confirm or exclude the diagnosis. Whilst thresholds of >5 times the URL following coronary intervention and >35 times the URL following cardiac surgery are arbitrary, patients with cardiac troponin concentrations below these thresholds may be less likely to have a new or presumed new regional abnormality due to a procedure-related complication. Further research is required to validate this assumption (see *Section 20*). The sensitivity and negative predictive value of these thresholds to identify patients in whom cardiac imaging is not required to exclude procedure-related myocardial infarction is unknown. Different thresholds may be required for cardiac troponin I and T, for males and females, and for complex procedures, such as percutaneous intervention for a chronic total occlusion or combined valve and coronary surgery. If pre-procedural cardiac troponin levels are available, comparison with post-procedural levels is informative.⁸⁷ The timing of measurement is also important and can help differentiate transient myocardial injury due to instrumentation from injury due to myocardial ischemia and a procedural complication.^{88–90} For example, after cardiac surgery, increasing cardiac troponin concentrations between 6 and 24 h are more likely in those with new or presumed new absence of viable myocardium on CMR imaging.⁸⁸

Immediately following a procedure, or when the procedure is performed in the setting of acute myocardial infarction, both cardiac and coronary imaging are required to confirm the diagnosis of procedure-related myocardial infarction. The diagnosis is confirmed if angiographic evidence of a coronary complication is identified **AND** there is a new regional wall motion abnormality or loss of viable myocardium in a pattern consistent with an ischemic etiology on cardiac imag-

ing. When previous imaging is available, comparison with post-procedural imaging can help to differentiate new abnormalities. Because some complications arising during a procedure are transient and can be addressed promptly, the requirement for both features increases the specificity of the diagnosis of procedure-related myocardial infarction and ensures it is clinically meaningful. Examples of coronary complications that may arise during percutaneous intervention and cardiac surgery are given in *Box 5*.

When new symptoms or signs of myocardial ischemia and acute myocardial injury occur subsequent to an initially uncomplicated cardiac procedure and within 30 days, the diagnosis of procedure-related myocardial infarction is confirmed if either early restenosis or recoil, stent thrombosis, or graft failure **OR** a new regional wall motion abnormality or loss of viable myocardium is identified. To be considered a procedure-related myocardial infarction after an initially uncomplicated cardiac procedure, the acute coronary pathology needs to have arisen in the same vessel or territory treated in the index procedure. The criteria for, and causes of, procedure-related myocardial infarction are shown in *Figure 7*.

Patients may also have a myocardial infarction unrelated to the index procedure during this 30-day period, in which case the diagnosis of primary or secondary myocardial infarction would be considered depending on the setting (see *Section 7.2.1* and *Section 7.2.2*). When an acute coronary event related to a prior cardiac procedure occurs beyond 30 days, the diagnosis of primary myocardial infarction should be considered.

8. DIAGNOSIS OF MYOCARDIAL INFARCTION FOLLOWING SUDDEN DEATH

Patients with symptoms or signs of myocardial ischemia may die suddenly before clinical evaluation or investigations can be performed to confirm a diagnosis of myocardial infarction. An accurate diagnosis in this setting is important as it can carry implications for family members and insurance, death certification, and as an endpoint for clinical trials.

In cases of sudden death, clinical history and corroborative findings can be helpful and autopsy examinations are encouraged (*Figure 8*). Following a sudden death without any evaluation, the diagnosis of primary myocardial infarction is often considered as a possible cause of death. Clinical features such as a history of CAD, chest pain prior to death, or documented ventricular fibrillation in the absence of a known cardiomyopathy or channelopathy make the diagnosis more likely, but in the absence of an ECG or post-mortem investigation, the diagnosis remains uncertain.

When there is uncertainty, post-mortem imaging or an autopsy can confirm the diagnosis and identify the




Figure 7. Procedure-related myocardial infarction due to a complication of cardiac surgery or intervention. For image description, please refer to the figure legend and surrounding text.

underlying etiology for accurate classification into primary, secondary, or procedure-related myocardial infarction. Unfortunately, autopsies are performed in a minority of patients presumed to have died from myocardial infarction.⁹¹ Autopsy may exclude myocardial infarction by identifying non-cardiac causes of death or non-coronary causes of sudden cardiac death such as cardiomyopathy, including hypertrophic or arrhythmogenic cardiomyopathy, or cardiac sarcoidosis, while channelopathies are often associated with a structurally normal heart. Autopsy identification of coronary atherothrombosis, embolism, SCAD, or restenosis/thrombosis within a stent or bypass graft when it occurs more than 30 days from the index procedure, together with myocardial necrosis in the corresponding coronary artery region, confirms the diagnosis of primary myocardial infarction. However, it should be noted that histopathological features of myocardial necrosis are time dependent.⁹² Identification of myocardial necrosis and infarction in the absence of an acute coronary pathology, when there was clinical or autopsy evidence of an alternative acute condition that could increase myocardial oxygen demand or reduce supply, would support a diagnosis of secondary myocardial infarction. When the death arises within 30 days of a cardiac procedure with evidence

of an acute coronary or graft occlusion on autopsy, the diagnosis of procedure-related myocardial infarction can be confirmed.

Contemporary alternatives to autopsy, including post-mortem computed tomography (CT) with or without coronary angiography or CMR imaging with biopsy, can inform the diagnosis of myocardial infarction.^{92,93} However, these approaches may not be widely available in practice. CT can identify obstructive CAD and prior myocardial infarction or suggest an alternative diagnosis, but the spatial resolution of conventional CCTA cannot reliably detect coronary plaque rupture or thrombus, or necrotic myocardium.^{93,94} Emerging data suggest that more contemporary CT technologies, including photon-counting CT, may have the ability to detect plaque characteristics indicative of acute rupture.^{95,96} Post-mortem CMR imaging has shown good correlation with histological specimens in identifying patients with myocardial infarction although its sensitivity is limited.⁹⁷ Although these approaches are not routinely available in clinical practice and imaging protocols have not been standardized, the combination of CT-guided biopsy and CMR imaging has a sensitivity and specificity of >95% for the post-mortem diagnosis of acute myocardial infarction.⁹²

9. SILENT OR UNRECOGNIZED MYOCARDIAL INFARCTION

Individuals with no known history of myocardial infarction may have pathological Q waves identified on a routine ECG. The specificity of pathological Q waves is limited⁹⁸; therefore, when available, the diagnosis of unrecognized (also termed clinically “silent” or ‘old’) myocardial infarction should be confirmed by imaging, preferably with CMR with late gadolinium enhancement (see *Section 15*).^{99,100} In some cases, regional wall motion abnormalities are detected on echocardiography, or myocardial infarction is observed on CMR imaging when these investigations have been performed for another indication and also may represent a silent or unrecognized myocardial infarction.^{101,102}

Although some cases may be truly “silent,” with no symptoms or signs even on careful consideration by the patient, the term “silent” myocardial infarction can in some cases be misleading. Many patients in retrospect recall a history of symptoms but may not have considered them serious enough to seek acute care or recognize them as symptoms of a myocardial infarction, or they did seek assessment, but the diagnosis of myocardial infarction was not made. Unrecognized myocardial infarction is present in approximately 1 in 5 persons over the age of 70 years in the general population if CMR imaging is performed systematically, although most are small infarcts.¹⁰³ Furthermore, in a recent study, 3% of asymptomatic middle-aged individuals (mean age 50 years) had CMR imaging evidence of unrecognized myocardial infarction.¹⁰⁴ Among patients with chronic coronary syndromes, the frequency of unrecognized myocardial infarction is even higher and is associated with CAD in the infarct territory.¹⁰⁵ Unrecognized myocardial infarction is relevant because it is associated with adverse prognosis.^{102,103,106} Furthermore, the identification of an unrecognized myocardial infarction may have important implications for medical treatment, particularly in patients without prior known CAD.

10. UNSTABLE ANGINA

In patients with new symptoms of myocardial ischemia at rest or on minimal exertion (class 3 or 4 angina), or a sudden intensification of symptoms in those with chronic coronary syndrome, a diagnosis of unstable angina should be considered if myocardial infarction is excluded through serial cardiac troponin measurements. The diagnosis is more likely when symptoms are associated with signs of myocardial ischemia on the ECG, and the diagnosis is confirmed if atherothrombosis or an alternative acute coronary pathology is identified on coronary angiography with or without functional testing or intravascular imaging.¹

Unstable angina and primary myocardial infarction share similar underlying pathophysiological mechanisms,

and acute coronary syndrome (ACS) is still commonly used as an umbrella term to describe these diagnoses.² Historically, unstable angina was a frequent diagnosis in practice and was associated with adverse outcomes.¹⁰⁷ However, by measuring cardiac troponin in stored samples, studies demonstrated that a significant proportion of patients previously diagnosed with unstable angina would be reclassified as having myocardial infarction due to the lower sensitivity of the then utilized biomarker assays.¹⁰⁸ Those with unstable angina without an elevation in cardiac troponin had a more favourable prognosis.^{109–111} In other words, the diagnosis of unstable angina has become progressively less common over time due to the introduction of high-sensitivity cardiac troponin assays and lower diagnostic thresholds for myocardial infarction.¹¹²

While unstable angina by definition is not associated with acute myocardial injury at the time of assessment, it is not uncommon for cardiac troponin values to be *chronically* elevated in these patients, and this is associated with an adverse prognosis.¹¹³ A systematic assessment for causes of chronic myocardial injury should be performed, but differentiating this from late presentation of myocardial infarction can be difficult (see *Section 14.1.4*). If there is diagnostic uncertainty, cardiac imaging or repeating the cardiac troponin measurement on a subsequent visit can help to differentiate recent myocardial infarction from other types of injury.

Despite these concerns and limitations, unstable angina has not disappeared from clinical practice and is still considered part of the spectrum of ACS.¹

11. MYOCARDIAL INJURY WITH NON-OBSTRUCTIVE CORONARY ARTERIES

The definition of MINOCA is updated to “myocardial *injury* with non-obstructive coronary arteries.” The term can be applied as a working diagnosis in patients who present with clinical features of possible myocardial infarction who are subsequently found to have non-obstructive coronary arteries (no stenosis $\geq 50\%$) on coronary angiography.^{114,115} The previous term referred to “myocardial infarction” rather than “myocardial injury,” but this was problematic as most patients are subsequently found to have a non-coronary cardiac (e.g. myocarditis, Takotsubo syndrome, cardiomyopathy) or non-cardiac cause (e.g. pulmonary embolism).^{116,117} Updating the definition of MINOCA helps to resolve this issue, emphasizing its role as a working diagnosis for those that require further investigation to establish the underlying cause and final diagnosis. For patients in whom myocardial infarction is subsequently confirmed after further invasive or non-invasive testing, the final diagnosis should be classified as primary, secondary, or procedure-related myocardial infarction depending on the setting (see *Section 7.2*).

In patients with MINOCA, intravascular imaging can identify subtle abnormalities of an acute coronary pathology of primary myocardial infarction, such as plaque erosion/rupture, mural thrombus without plaque rupture, or SCAD.^{1,117} Similarly, invasive functional testing may identify microvascular dysfunction and/or epicardial coronary spasm that may have led to a primary myocardial infarction.¹ Clinical practice guidelines further recommend CMR imaging to aid in establishing the final diagnosis, ideally performed within 2 weeks after presentation, but thereafter if necessary.^{1,62} When CMR has been performed, myocardial infarction is the final diagnosis in 22%–27% of patients,^{118,119} while more often imaging identifies an alternative non-coronary cardiac diagnosis, such as acute myocarditis or Takotsubo syndrome, resulting in a change in management.¹²⁰ The diagnostic yield of CMR is higher if performed within 2 weeks of presentation¹²¹ as after this time there may be resolution of reversible myocardial changes, particularly in Takotsubo syndrome or acute myocarditis,¹²² while CMR findings will persist in myocardial infarction or in those with cardiomyopathy.¹²³

12. SYMPTOMS OF MYOCARDIAL INFARCTION

Chest pain is the most commonly reported presenting symptom of acute myocardial infarction.^{124–126} The sensitivity of chest pain for a diagnosis of myocardial infarction is high and is similar among women and men,^{127,128} but chest pain is not specific as it can be associated with multiple other conditions.⁶² Additionally, some patients report epigastric pain and/or referred pain to the jaw, neck, arm, or back, and some patients do not report chest pain at all¹²⁹ and present only with associated symptoms, such as breathlessness, palpitations, nausea, or vomiting.¹³⁰ Patients describe the quality of ischemic chest pain as pressure, tightness, or squeezing, and symptom severity does not associate with a diagnosis of myocardial infarction¹³¹; thus, chest discomfort may be a more appropriate term to describe the main symptom of myocardial ischemia.⁶² Use of the terms “typical” and “atypical” are discouraged as studies have found that symptoms labelled as atypical are more common among women evaluated for myocardial infarction and may contribute to delayed diagnoses.¹³² Chest discomfort is considered acute when there is an abrupt onset or change in intensity, duration, or overall pattern. Chest discomfort is considered stable when symptoms are chronic with reliable precipitating factors of exertion or emotional stress.¹³⁰ Patients with STEMI are more likely to present with persistent chest discomfort and to experience vomiting, dizziness, and diaphoresis compared with patients with NSTEMI, who are more likely to present with intermittent symptoms.^{125,133} The diagnosis of myocardial infarction is more often delayed among patients without chest pain, in

whom the risk of death in hospital is higher.^{129,130} Patients presenting with secondary myocardial infarction are less likely to present with chest pain and more likely to report breathlessness or other symptoms associated with the primary diagnosis.¹⁸

Studies of symptoms in myocardial infarction have included analyses by sex, age, race, ethnicity, and comorbidities. While chest pain is the most common symptom in both women and men, women more often present with additional symptoms that are less often recognized as cardiac by either patients or clinicians, which can contribute to delays in care.^{126,134} Compared with men with ACS, women had higher odds of presenting with pain between the shoulder blades, nausea or vomiting, and shortness of breath, and lower odds of chest pain and diaphoresis.¹³⁵ Myocardial infarction presenting without chest pain or discomfort is more common among older persons¹³⁶ and those with diabetes mellitus.^{137–139} Therefore, it is important to systematically inquire about additional symptoms, particularly in women, older persons, and people with diabetes, while noting that the majority of these patients still present with chest pain or discomfort. There are few studies examining the effects of pain perception in myocardial infarction. Persons with an unrecognized myocardial infarction had reduced pain sensitivity to cold pressor testing compared with persons who received a diagnosis of acute myocardial infarction.^{126,135,140} Whilst differences in symptoms have been observed by race or ethnicity,^{141,142} no consistent patterns have emerged to inform diagnosis.^{137,138}

13. THE ELECTROCARDIOGRAM

The 12-lead ECG remains a cornerstone in the diagnosis of myocardial infarction, particularly for rapid identification of patients with symptoms of possible myocardial infarction due to acute coronary occlusion who require immediate reperfusion (Figure 9). The working diagnosis of STEMI requires new ST-segment elevation at the J-point in at least two contiguous leads: ≥ 2.5 mm in men < 40 years, ≥ 2 mm in men ≥ 40 years, or ≥ 1.5 mm in women regardless of age in leads V_2 – V_3 ; and/or ≥ 1 mm in all other leads, in the absence of left ventricular hypertrophy or left bundle branch block (LBBB) (Table 5).

A posterior myocardial infarction is suggested by the presence of ST-segment depression ≥ 1 mm in lead V_1 , V_2 , and/or V_3 , particularly when the R-wave amplitude is greater than the S-wave (dominant) in V_1 or V_2 , confirmed by ST-segment elevation in the posterior leads V_7 to V_9 .¹⁴³ A right ventricular myocardial infarction is indicated by ST-segment elevation in the right precordial leads (V_3R – V_6R), especially when accompanied by ST-segment elevation in lead aVR.

Previous studies have shown that up to 1 in 4 patients managed as a NSTEMI without classical ST-segment elevation on conventional 12-lead ECG have an acute

Table 5. Electrocardiographic Findings of Myocardial Ischemia or Infarction

Feature	Description of electrocardiographic findings of myocardial ischemia or infarction
ST-segment elevation	New ST-segment elevation at the J-point in two contiguous leads: ≥1 mm in all leads other than leads V ₂ –V ₃ where the following apply: ≥2.5 mm in males <40 years, ≥2 mm in males ≥40 years, or ≥1.5 mm in females regardless of age, in the absence of bundle branch block and/or left ventricular hypertrophy
ST-segment depression	New horizontal or down sloping ST-segment depression ≥0.5 mm at the J-point in two or more contiguous leads
Hyperacute T waves	Symmetrical, broad T waves disproportionately large to the preceding QRS complex in two contiguous leads
de Winter T waves	Tall, prominent, symmetrical T waves with upsloping ST segment depression in the precordial leads
Biphasic T waves	Initial positive deflection followed by a negative deflection
T-wave inversion	New or dynamic T-wave inversion ≥1 mm in two contiguous leads
Wellens syndrome	Biphasic or deeply inverted T waves in leads V ₂ and V ₃
Pathologic Q waves	Q-wave duration ≥40 ms and/or a depth of ≥25% of the R wave in two contiguous leads
Sgarbossa criteria	In the presence of left bundle branch block or ventricular pacing; ST-segment elevation ≥1 mm concordant (in the same direction) with the QRS complex; ST-segment depression ≥1 mm in lead V ₁ , V ₂ , or V ₃ ; ST-segment elevation ≥5 mm not concordant with the QRS complex
Ventricular arrhythmia	Ventricular fibrillation or ventricular tachycardia

occlusion of the culprit artery.^{50,51,53} In the right clinical setting, other electrocardiographic patterns may be considered equivalent to STEMI.⁵² These include new or presumed new left and/or right bundle branch block when accompanied by other signs of myocardial ischemia.¹⁴⁴ Among patients with pre-existing LBBB or ventricular pacing the presence of ST-segment elevation ≥1 mm concordant (in the same direction) with the QRS complex; ST-segment depression ≥1 mm in lead V₁, V₂, or V₃; and ST-segment elevation ≥5 mm not concordant (in the opposite direction) with the QRS complex suggest acute coronary occlusion (Sgarbossa criteria).¹⁴⁵ The modified Sgarbossa criteria may improve the detection of acute coronary occlusion with LBBB or ventricular pacing by replacing the absolute discordant ST-segment elevation threshold with the proportional rule of an ST/S-wave ratio ≤0.25.¹⁴⁶ Further electrocardiographic patterns that suggest acute coronary occlusion include marked ST-segment depression or hyperacute T waves in V₁–V₂ with reciprocal changes elsewhere, de Winter pattern (upsloping ST-segment depression with tall, symmetric T waves in V₂–V₅),¹⁴⁷ Wellens syndrome (biphasic or deeply inverted T waves in V₂–V₃ during pain-free intervals),¹⁴⁸ Aslanger pattern (ST-segment elevation isolated to lead III with ST-segment depression in any of leads V₄–V₆ with a positive T wave), and the “South African flag” sign (ST-segment elevation in leads I, aVL, and V₂ and ST-segment depression in lead III).⁵²

Patients with previous coronary bypass grafts and/or complex CAD (e.g. chronic total occlusions with collateral supply) may present with acute coronary or graft occlusion without typical ST-segment elevation, and a high index of suspicion for myocardial infarction is required.¹⁴⁴

ST-segment elevation is not specific to myocardial infarction and can also be found in early repolarization or other cardiac (e.g. myocarditis, pericarditis, Brugada syndrome, and Takotsubo syndrome or cardiomyopathies) and non-cardiac (e.g. pulmonary embolism, hyperkale-

mia, hypothermia, elevated intracranial pressure) conditions. With an acute coronary pathology, ST-segment elevation is typically regional and often accompanied by reciprocal ST-segment depression. In contrast, in other conditions such as pericarditis or early repolarization, ST-segment elevation is often diffuse or global, not confined to a single coronary territory, and occurs without reciprocal changes.¹⁴⁹

In addition to ST-segment elevation, electrocardiographic findings associated with acute myocardial ischemia include ST-segment depression, T-wave inversion, hyperacute or biphasic T waves, pathological Q waves, and ventricular arrhythmias (Table 5). New horizontal or down-sloping ST-segment depression ≥0.5 mm and/or T-wave inversion >1 mm in two contiguous leads is suggestive of myocardial ischemia. Global ischemic changes (diffuse ST-segment depression along with ST-segment elevation in lead aVR) may reflect left main stem or balanced three vessel CAD. However, these electrocardiographic findings are not specific to myocardial infarction and can be observed in other cardiac and non-cardiac conditions.

Patients who present late following myocardial infarction due to an acute coronary occlusion are more likely to have pathological Q waves in two or more contiguous leads and biphasic T waves or T-wave inversions rather than ST-segment elevation or depression—although some patients presenting late may not necessarily have developed Q waves despite an acute coronary occlusion.¹⁵⁰ Q waves are usually permanent but may regress and disappear over months to years. The classical definition of a pathological Q wave requires a Q-wave duration ≥40 ms and/or a depth of ≥25% of the R wave in the same lead.¹⁵¹ While an alternative pathological Q wave definition was suggested in prior versions of the UDMI, the classical definition has a higher correlation with transmural myocardial infarction in CMR imaging studies.⁹⁸




Figure 8. Diagnostic pathway for possible primary myocardial infarction following sudden death.

VF, ventricular fibrillation. Following a sudden death in the community without clinical evaluation, the diagnosis of primary myocardial infarction can be considered. Clinical features such as a history of coronary artery disease, chest pain prior to death, or documented VF in the absence of known cardiomyopathy or channelopathies make the diagnosis more likely, but in the absence of an electrocardiogram or post-mortem investigations the diagnosis is uncertain, and the ICD-11 code of unspecified myocardial infarction (BA41.Z) should be applied to recognize this. Where investigations are incomplete with no biomarker testing or coronary imaging, the diagnosis of myocardial infarction of uncertain etiology is applied, but where there is uncertainty post-mortem imaging or an autopsy is required to confirm the diagnosis and identify the underlying acute coronary pathology. For image description, please refer to the figure legend and surrounding text.

In the setting of secondary myocardial infarction global ischemic changes are more common.⁶⁴ More marked electrocardiographic changes, particularly ST-segment elevation and depression, that do not resolve rapidly with treatment of the underlying cause (e.g. resolution of widespread ST-segment depression after cardioversion for tachyarrhythmia) should lead to reconsideration as to whether the diagnosis is primary myocardial infarction.

Procedure-related myocardial infarction presents with similar electrocardiographic findings as primary myocardial infarction. Early stent thrombosis often presents with classical ST-segment elevation with reciprocal ST-segment depression, while smaller procedural myocardial infarcts, such as side branch occlusion, may have more subtle or no electrocardiographic changes. Comparison with the pre-procedural ECG can be helpful. New electrocardiographic changes, such as ST-segment elevation or depression and T-wave inversion, are common after cardiac surgery and

do not always indicate acute graft failure.⁴³ For example, following cardiac surgery, post-pericardiotomy syndrome (an autoimmune inflammatory response to pericardial and pleural trauma), pneumopericardium (air in the pericardial space), and metabolic disturbances (due to hemodilution, renal dysfunction, blood products, and/or diuretics) can cause ST-segment elevation, depression, and/or T-wave changes. Furthermore, Q waves that meet the definition for a pathological Q wave can be transient early after cardiac surgery.⁸⁰ The early use of cardiac imaging and invasive coronary angiography is required to confirm whether these electrocardiographic changes are due to procedure-related myocardial infarction.

14. CARDIAC BIOMARKERS

Acute myocardial injury is integral to the diagnosis of myocardial infarction whether primary, secondary, or




Figure 9. Electrocardiographic features of acute coronary occlusion.

The electrocardiographic changes associated with primary myocardial infarction due to acute coronary occlusion depend on the time of presentation from the onset of persistent symptoms. The diagnosis of ST-segment elevation myocardial infarction can be made in patients with alternative electrocardiographic changes that are suggestive of acute coronary occlusion or in those who present late where the ST-segment changes have evolved. For image description, please refer to the figure legend and surrounding text.

procedure related. Since 2007, the UDMI has recommended cardiac troponin I or T (henceforth referred to as “cardiac troponin”) as the preferred biomarkers for the diagnosis of myocardial infarction due to their greater specificity for myocardial injury than other biomarkers.⁶ The definition of acute myocardial injury as a rise and/or fall in cardiac troponin with at least one value above the 99th percentile URL has remained consistent, with the use of sex-specific 99th percentile URLs recommended by the Fourth UDMI in 2018 when using high-sensitivity assays.⁸ Early diagnostic pathways incorporating high-sensitivity cardiac troponin assays, with some available at the point of care, have created opportunities to accelerate the diagnosis of patients with possible myocardial infarction. A diagnosis of myocardial infarction requires an understanding of cardiac troponin assay performance, decision threshold limitations, potential analytical interference, and when serial testing or alternative biomarkers are needed to differentiate acute from chronic elevation. Alternative biomarkers for myocardial infarction are covered in *Section 14.2* and implications for diagnosis of myocardial infarction in low-resource settings are discussed in *Section 19*.

14.1. Cardiac Troponin

Cardiac troponin plays a key role in myocardial contraction, and immunoassays that quantify cardiac troponins in the circulation play an important role in the diagnosis of myocardial infarction and recognition of both acute and chronic cardiac conditions.¹⁵² In the sarcomere, cardiac troponin I (inhibitory), T (tropomyosin-binding), and C (calcium-binding) subunits exist in a complex, which is released into the circulation as an intact ternary complex and as subunit fragments in acute myocardial infarction.¹⁵³ Cardiac troponin I and T have a half-life of 2 h and between 2 and 4 h in the circulation, respectively.¹⁵⁴ Immunoassays are available that target amino acid sequences that are specific to the cardiac isoforms of troponin I and T, which ensures these assays provide high specificity for myocardial injury.

High-sensitivity assays have greater analytical precision to enable detection of cardiac troponin at very low concentrations and are now available on multiple central laboratory and point-of care platforms.¹⁵⁵ For classification as a high-sensitivity assay, cardiac troponin assays should achieve a coefficient of variation of $\leq 10\%$ at the

sex-specific 99th percentile URL and be able to detect cardiac troponin in more than 50% of healthy females and males in the reference population.¹⁵⁶ Greater precision at the diagnostic threshold for myocardial infarction improves the reliability of cardiac troponin testing in practice. As such, high-sensitivity assays are recommended by clinical practice guidelines^{1,62} and are now used in many countries around the world.¹⁵⁷ These assays have enabled implementation of accelerated diagnostic pathways and more precise estimates of the 99th percentile URL for the diagnosis of myocardial infarction. As cardiac troponin assays are not standardized by the manufacturers, evidence from clinical studies may not apply to all assays and clinicians should refer to the International Federation of Clinical Chemistry and Laboratory Medicine Biomarkers Reference Tables¹⁵⁵ and apply assay-specific thresholds in practice.

14.1.1. Defining the 99th Percentile Upper Reference Limit

Since there is no gold standard for the diagnosis of myocardial infarction that is independent of cardiac troponin, it is not possible to define a decision threshold based on diagnostic performance studies. Therefore, the decision threshold is based instead on the distribution of cardiac troponin in reference populations without known cardiac conditions or cardiovascular risk factors. The 99th percentile URL is recommended to standardize the approach to recognizing those with myocardial injury across different assays, between sites, and across healthcare systems. It is conventional to use the 97.5th percentile as the URL for most biomarkers, but a higher threshold (the 99th percentile) is applied along with a rise and/or fall in cardiac troponin to increase the specificity for a diagnosis of myocardial infarction, as many other conditions also increase cardiac troponin concentrations. Given that the identification of those with myocardial injury is integral to the diagnosis of myocardial infarction, it is important that the 99th percentile is determined in sufficiently large and representative studies.¹⁵⁸

Cardiac troponin concentrations vary considerably by sex.^{159,160} For high-sensitivity cardiac troponin I and T assays, the 99th percentile URL in females is lower than the uniform URL.^{161,162} In contrast, the 99th percentile URL in males is above the uniform URL. As such, sex-specific 99th percentile URLs are necessary to define myocardial injury and avoid systematic bias that could contribute to the under-recognition of myocardial infarction and other cardiac conditions associated with myocardial injury in female patients.¹⁶³ Historically, when using a cardiac troponin I assay the use of sex-specific URLs compared with a uniform URL reclassified a greater proportion of females as having myocardial injury than the same comparison using a cardiac troponin T assay.^{163,164} However, the impact of sex-specific URLs is likely to be similar for cardiac troponin I and T when using the latest

generation high-sensitivity cardiac troponin T assay.¹⁶⁵ Enhanced precision at low concentrations has increased the proportion of healthy females in whom cardiac troponin T is quantifiable, enabling the accurate determination of URLs for both sexes: the URL in females is half the URL in males for cardiac troponin T in a large and representative global healthy reference population.¹⁶⁵

Whilst differences between female-specific and male-specific 99th percentile thresholds are similar in magnitude across all age ranges for both cardiac troponin I and T assays,¹⁵⁷ the age of the healthy reference population used to define the 99th percentile is particularly important as cardiac troponin increases significantly with age.^{159,166} Differences in cardiac troponin by age are as important as the differences observed by sex; however, unlike sex, age is a continuous measure, therefore defining appropriate age-adjusted 99th percentiles is more challenging. Most regulatory studies that define the normal reference range for cardiac troponin assays do not include a large enough sample of younger and older persons to reliably determine age-adjusted URLs, and no cardiac troponin assays have sought regulatory approval for the use of age-adjusted thresholds in clinical practice. Clinicians should be aware that 1 in 3 patients over 75 years with possible myocardial infarction have an elevated cardiac troponin concentration compared with 1 in 20 patients under 50 years old.¹⁶⁷ This is partly because older persons are more likely to have comorbid conditions that influence cardiac troponin, such as pre-existing or subclinical cardiac conditions or kidney disease, hypertension, and diabetes, and partly because the incidence of spontaneous/primary or secondary myocardial infarction is highest in this age group. Serial measurements are helpful to differentiate acute from chronic causes of elevated cardiac troponin values when there is uncertainty (see *Section 6*). Further research is required to define age-adjusted 99th percentile URLs for females and males in sufficiently large reference range populations, and to determine whether their application would improve the recognition of younger and older patients with myocardial injury due to myocardial infarction or other cardiac conditions (see *Section 20*). In the future, the use of statistical models or machine learning tools that incorporate variables beyond age and sex to include a measure of renal function or knowledge of pre-existing conditions could improve how normal cardiac troponin values are defined.¹⁶⁸

14.1.2. Analytical Considerations

Immunoassays for cardiac troponin are no different from immunoassays for other biomarkers in that they are susceptible to pre-analytical and analytical factors that can result in false positives and negatives.¹⁶⁹ For example, hemolysis and biotin can reduce, and fibrin micro-clots can increase, cardiac troponin concentrations.^{170,171} Ana-

lytical performance of assays at the point of care may be more variable in whole blood compared with plasma or serum on a central laboratory platform.¹⁷²

When the assay result is unexpected, the measurement should be repeated with a new sample. Heterophilic antibodies may interfere with immunoassays *ex vivo* resulting in analytical false positive values of cardiac troponin when in fact there is no elevation *in vivo*. However, with most cardiac troponin assays, heterophilic antibodies are an uncommon cause for false positive results.¹⁷³ Similarly, immunoglobulin binding with cardiac troponin *in vivo* can result in macrotroponin complexes¹⁷⁴ that are slowly cleared from the circulation with a half-life of several weeks, resulting in high measurable cardiac troponin concentrations even in the absence of acute or chronic myocardial injury.¹⁷⁵

Interference should be suspected when elevated cardiac troponin is not consistent with the clinical findings,¹⁷⁶ and is more likely when values increase and decrease with no apparent pattern on serial testing. Interference can be readily identified by measuring cardiac troponin using an alternative immunoassay, with discordance between cardiac troponin I and T or between cardiac troponin I assays being strongly suggestive of interference.¹⁷⁵ Macro-troponin complexes can be identified using immunoglobulin depletion or sucrose gradient ultracentrifugation in specialist laboratories.¹⁶⁹ The frequency of antibody-mediated interference and macrotroponin complexes in patients with possible myocardial infarction is unknown but may be responsible for as many as 1 in 2 elevated cardiac troponin I values in asymptomatic populations.¹⁷⁷

These analytical issues can also influence the accurate determination of the 99th percentile, which is particularly susceptible to outliers. Recent recommendations on sample processing, the size of the population, and the inclusion and exclusion criteria to define a truly healthy reference population, and the application of statistical methods to exclude outliers, should improve the reliability and consistency of studies defining the 99th percentile URL.¹⁵⁸ Larger sample sizes are likely to improve precision.

14.1.3. Serial Measurements

Serial cardiac troponin testing is generally needed to differentiate acute myocardial infarction from other causes of myocardial injury in patients with possible primary, secondary, and procedure-related myocardial infarction. The timing of serial measurements should be guided by a validated diagnostic pathway (see *Section 14.1.5*).

When using a high-sensitivity assay, very low cardiac troponin values below the limit of detection, limit of quantification, or a defined risk stratification threshold can be used to safely exclude myocardial infarction with a single measurement in many patients.^{66,178} In contrast, serial testing is required if higher thresholds are used to exclude myocardial infarction as the sensitivity of the 99th per-

centile URL at presentation is around 90% overall and as low as 70% among patients presenting within 3 h of symptom onset.^{179,180} The release kinetics of cardiac troponin follow a well-defined pattern in primary myocardial infarction, with increasing levels detectable within 2 to 3 h of symptom onset, peaking between 12 and 24 h, and remaining elevated for days or weeks (Figure 10). The release kinetics differ depending on whether there is acute coronary occlusion at presentation, and whether spontaneous or therapeutic epicardial and tissue level reperfusion occurs. One cannot rely on cardiac troponin results in the first few hours in patients with ST-segment elevation as levels can be below the 99th percentile URL in around 1 in 6 patients due to lack of perfusion of the infarcted territory.¹⁸¹ Among those with microvascular obstruction the peak cardiac troponin concentration is lower and delayed if intramyocardial hemorrhage is not present, whereas it is higher and occurs earlier in those with intramyocardial hemorrhage.^{182,183}

Less is known about the role of serial measurements for the diagnosis of secondary myocardial infarction due to supply-demand imbalance. In secondary myocardial infarction the presentation and maximum value of cardiac troponin may be lower than in primary myocardial infarction, but there is substantial overlap in values between these settings.⁷¹ On serial testing the absolute and relative changes in cardiac troponin concentrations are similar in both primary and secondary myocardial infarction and among those with other causes of acute myocardial injury.⁷¹ Interestingly, the initial cardiac troponin values are more similar (less than two-fold difference) than the peak values (up to five-fold difference) when comparing primary and secondary myocardial infarction. This could be due to delays in cardiac troponin release in those with acute coronary occlusion, differences in the amount of myocardium at jeopardy, or because ischemic injury is more likely to persist in primary than in secondary myocardial infarction, where the trigger of supply-demand imbalance may be rapidly corrected (e.g. with cardioversion).

Following cardiac surgery, cardiac troponin is elevated in all patients, and in 97.5% this exceeds the previously recommended diagnostic threshold of >10 times the URL.⁴² Interestingly, peak cardiac troponin concentrations occur earlier after surgery (3 to 12 h)⁴² than after primary myocardial infarction (12 to 24 h). This may reflect differences in cardiac troponin release following reperfusion or differences in the mechanism of myocardial injury, which in most patients undergoing cardiac surgery is due to surgical manipulation or limitations of cardioplegia without ongoing injury thereafter. Those who sustain a procedure-related myocardial infarction after cardiac surgery as identified by new late enhancement on CMR imaging are more likely to have a later peak.⁸⁸ A peak in cardiac troponin 12 to 24 h after car-




Figure 10. Rise and fall in cardiac troponin over time from symptom onset.

Following the onset of acute myocardial infarction, changes in cardiac troponin vary over time: (A) Small absolute changes in cardiac troponin occur early after symptom onset (<3 h). (B) Larger absolute changes in cardiac troponin occur later after symptom onset. (C) Minimal rise and fall in cardiac troponin may occur around 12 to 24 h from symptom onset. (D) Larger fall in cardiac troponin may occur after it has peaked (12 to 24 h from symptom onset). (E) Pattern may mimic chronic myocardial injury if presentation days after symptom onset. (F) Cardiac troponin may take weeks to return to normal after myocardial infarction and some patients may develop chronic myocardial injury following myocardial infarction. For image description, please refer to the figure legend and surrounding text.

diac surgery may indicate a procedure-related complication, given that values typically peak within 3 to 12 h.⁸⁹

Similarly, after PCI in patients with a chronic coronary syndrome, an increase in cardiac troponin concentration above the 99th percentile URL occurs in 1 in 2 patients,¹⁸⁴ with the majority having a peak value 12–24 h following intervention.¹⁸⁵ The timing of peak values differs between those with and without elevated values at baseline who are undergoing intervention for acute and chronic coronary syndromes, respectively.¹⁸⁶ New late enhancement on CMR imaging is more likely among those with elevated cardiac troponin concentrations 24 h following PCI than those without.¹⁸⁷ Among patients with myocardial ischemia or an overt complication of PCI, serial cardiac troponin measurements at baseline and 12–24 h after the event may be needed to recognize those with acute myocardial injury.

14.1.4. Defining a Significant Rise and Fall

No single absolute or relative value can be applied to define a significant rise or fall in cardiac troponin that applies to all clinical presentations and assays. Whilst accelerated diagnostic pathways use assay-specific thresholds combined with delta criteria to increase the

likelihood that myocardial injury is due to myocardial infarction (see *Section 14.1.5*), clinical judgment must be used to determine whether a particular change is sufficient to confirm the diagnosis of myocardial infarction, considering several relevant factors.

The influence that the timing of testing has on the rise and fall in cardiac troponin is illustrated in Figure 10. First, the **time from symptom onset** at which samples are obtained affects the magnitude of change in cardiac troponin concentrations on serial testing.^{8,188,189} Smaller absolute changes are observed within 3 h of symptom onset, with a more substantial increase observed between 3 and 12 h. When samples are obtained around the time of the peak concentration (12–24 h), there may be minimal changes on serial testing. Similarly, among those who present many days after the onset of symptoms, the pattern may mimic chronic myocardial injury. Differentiating residual acute myocardial injury following a recent myocardial infarction from chronic myocardial injury is particularly important. Second, the **time interval between samples** will substantially influence the absolute value of the rise or fall in cardiac troponin concentrations. For this reason, fixed time intervals are recommended at 1 or 2 h for accelerated diagnostic

algorithms to improve the reliability of these criteria.¹⁹⁰ Third, the **baseline concentration** is important. For cardiac troponin concentrations below or around the 99th percentile URL, absolute delta values rather than relative values improve precision.¹⁹⁰ It is important to note that absolute delta values are assay-specific, and a cut-off value cannot be transferred from one assay to another.¹⁵⁶ For clearly elevated baseline values, relative changes are more intuitive.⁸ However, for very high cardiac troponin values, it may not be necessary to demonstrate a rise and/or fall, as such values are rarely observed in those with chronic myocardial injury.¹⁵⁶ Please see *Section 6* for a more in-depth discussion of the criteria used to define acute and chronic myocardial injury.

14.1.5. Accelerated Diagnostic Pathways

One of the advantages of high-sensitivity cardiac troponin assays has been their use in accelerated diagnostic pathways. Multiple pathways have been validated for different assays, with a similar goal of improving the safety and efficiency of the assessment of patients with possible myocardial infarction without ST-segment elevation on the ECG.^{191,192} These pathways are illustrated in clinical practice guidelines^{1,2,193} and can be broadly classified into three types: (i) single-sample rule-out pathways that classify patients into two groups (low- or high-risk); (ii) serial testing pathways with repeated measurements at 1 or 2 h that use multiple thresholds and delta values to stratify patients into three groups (low-risk, observe, or high-risk); and (iii) conventional pathways that combine a clinical risk score with the 99th percentile at 0 and 3 h. Single-sample rule-out pathways use assay-specific optimized thresholds below the 99th percentile to identify the greatest proportion of low-risk patients as potentially suitable for discharge from the Emergency Department with a high sensitivity and negative predictive value for myocardial infarction.⁶³ A similar approach is applied in 0/1- and 0/2-hour serial testing pathways that also incorporate assay-specific optimized thresholds above the 99th percentile to identify high-risk patients with greater specificity and positive predictive value.¹⁹⁴

The thresholds used to triage patients with possible myocardial infarction in accelerated diagnostic pathways are often not based on the 99th percentile of the cardiac troponin assay and here uniform rather than sex-specific thresholds have been applied. For example, in the 0/1- and 0/2-hour serial testing pathways recommended by the ESC the same thresholds are applied in both male and female patients to differentiate those at low risk and suitable for discharge from those at increased risk who require further investigation or hospital admission.¹ It should be noted that these thresholds are used for risk stratification, rather than to define myocardial injury and confirm a final diagnosis of myocardial infarction, which is based on demonstrating a rise and/or fall in cardiac

troponin where at least one value is above the 99th percentile URL and sex-specific thresholds apply (see *Section 6*).

14.1.6. Interpretation of Cardiac Troponin for Diagnosis

Although the introduction of high-sensitivity assays and accelerated diagnostic pathways has contributed considerably to the early diagnosis of myocardial infarction, some additional considerations are needed when interpreting cardiac troponin values in patients with possible myocardial infarction (*Box 6*).

Box 6: Interpretation of Cardiac Troponin for Diagnosis of Myocardial Infarction

- In patients with possible myocardial infarction, cardiac troponin needs to be interpreted in conjunction with clinical assessment including a history, physical examination, and 12-lead electrocardiogram.
- In patients who present early after the onset of symptoms, increases in cardiac troponin concentration may not be detected even with high-sensitivity assays, and serial cardiac troponin testing is necessary.
- In patients with recurrent or persistent symptoms of myocardial ischemia, serial cardiac troponin testing is needed prior to excluding myocardial infarction.
- In unselected patients without a clinical indication for diagnostic testing, elevated cardiac troponin concentrations are common in multiple conditions, such as sepsis, kidney disease, or stroke.
- The 99th percentile for cardiac troponin is lower for females than males; the use of high-sensitivity assays with sex-specific thresholds will minimize the risk of underdiagnosis of myocardial injury and infarction in females compared with using a uniform threshold.
- Age is an important consideration when interpreting cardiac troponin as the prevalence of myocardial infarction increases with age, as does the prevalence of persistently elevated cardiac troponin values.
- When cardiac troponin concentrations are elevated, unless the diagnosis of myocardial infarction is confirmed, serial measurements are typically required to differentiate acute from chronic causes of myocardial injury.
- Analytical interference should be suspected when an elevated cardiac troponin value is not consistent with the clinical findings and is more likely when cardiac troponin values vary with no apparent pattern on serial testing.

14.2. Alternative Biomarkers of Myocardial Injury

The most sensitive and specific biomarker for the diagnosis of myocardial infarction is cardiac troponin measured using a high-sensitivity assay. In a recent survey of practice from 663 laboratories in 76 countries from Europe, Africa, South and North America, and Australasia, cardiac troponin testing was available in 98% of hospitals, and in 92% of these hospitals, testing was with a high-sensitivity assay.¹⁵⁷ If a high-sensitivity assay is not available, then a cardiac troponin assay that is not high-sensitivity is preferred over CK-MB, which has lower specificity and sensitivity.⁴⁴ If neither type of cardiac troponin assay is available, then an alternative would be to use CK-MB measured by a mass assay.^{44,83} A number of alternative biomarkers of myocardial injury are in development, such as cardiac myosin-binding protein C (cMyC). cMyC is a sarcomeric protein that is loosely associated with both myosin and actin, is more abundant than cardiac troponin, and rises more rapidly after iatrogenic myocardial injury, suggesting a role in early evaluation.¹⁹⁵ Prototype cMyC assays have comparable performance to cardiac troponin assays in accelerated diagnostic pathways for myocardial infarction,^{196,197} but whether testing of cMyC in combination with cardiac troponin will improve the diagnosis of myocardial infarction in patients for whom the interpretation of cardiac troponin is challenging due to analytical interference or kidney disease remains uncertain. And for these reasons, biomarkers other than cardiac troponin are not used routinely for the diagnosis of myocardial infarction.

15. CORONARY AND CARDIAC IMAGING

While a likely diagnosis of myocardial infarction can be made without imaging, confirmation of the diagnosis and determination of the underlying mechanism requires coronary angiography and/or cardiac imaging.

15.1. Invasive Imaging

15.1.1. Invasive Coronary Angiography

Invasive coronary angiography is the gold-standard test for the evaluation of patients with possible myocardial infarction, with the timing of coronary angiography defined in clinical practice guidelines.^{1,2,62} Not only can invasive coronary angiography confirm the diagnosis, but it can also determine the underlying acute coronary pathology in primary myocardial infarction. The prevalence of acute coronary pathologies other than atherothrombosis varies widely: SCAD (0.1%–4.0%), coronary embolism (0.2%–13%), epicardial coronary vasospasm (1%–20%), in-stent restenosis (1.0%–6.5%), stent thrombosis (0.6%–0.8%), and graft failure (unknown).^{61,198–201} This variation is likely due to differences in the populations studied, en-

rollment criteria (systematic or selected), lack of central adjudication, and variation in the use of adjuvant coronary or cardiac imaging, which can help to identify the culprit lesion and underlying pathology.^{1,2}

Historically, many patients with myocardial ischemia secondary to oxygen supply–demand imbalance have not undergone coronary angiography, with the diagnosis made based on clinical assessment.⁶⁴ Due to minimal use of coronary angiography in this setting, patients with primary myocardial infarction can be missed, and some patients with non-ischemic acute myocardial injury will be misdiagnosed as having secondary myocardial infarction. In a prospective study of patients with type 2 myocardial infarction defined using the previous classification who underwent systematic coronary angiography, 1 in 20 patients were found to have a primary myocardial infarction due to atherothrombosis.⁷² The use of more objective imaging criteria to define secondary myocardial infarction in this cohort reduced the diagnosis of myocardial infarction as most had neither obstructive CAD nor new left ventricular impairment.⁷⁴

In many parts of the world, particularly in low-resource settings, there is limited or no access to invasive coronary angiography.⁹ In some patients (e.g. those with a relative contraindication, limited life expectancy, or cognitive impairment), or based on patient values and preferences, invasive coronary angiography may not be performed. Consequently, many individuals with possible myocardial infarction do not undergo this key diagnostic test. In these situations, cardiac imaging can confirm the diagnosis, or when imaging is also not possible or not diagnostic, a clinical diagnosis of myocardial infarction can be made based on clinical presentation, the ECG, and cardiac biomarkers. Additionally, in some situations, such as in low-risk individuals, possible secondary myocardial infarction, or where the likelihood of revascularization is low, CCTA may be a useful alternative to aid the diagnosis and/or to guide the need for invasive coronary angiography, where available.¹

15.1.2. Intravascular Imaging

Intravascular imaging, comprising IVUS and OCT, if available, can be used to identify plaque rupture and abnormal coronary vessel wall morphology in patients with possible myocardial infarction.¹ This is important as approximately 14% of patients presenting with NSTEMI lack a clearly identifiable culprit lesion on invasive angiography.²⁰² Moreover, earlier studies reported that in non-Q wave myocardial infarction, more than a third of patients had no culprit lesion detected on angiography.²⁰³ Diagnostic uncertainty is more common among those without obstructive CAD (see *Section 11*) or with an acute coronary mechanism of primary myocardial infarction other than atherothrombosis.^{204–206} Intravascular imaging can confirm atherothrombosis and the type of plaque morphology (rupture, erosion, intraplaque cavity, mural thrombus) and/or identify an

alternative acute coronary pathology, such as intramural hematoma due to SCAD, isolated thrombus in coronary embolism, in-stent restenosis, stent thrombosis, or intimal bumping consistent with coronary artery spasm.^{117,207}

15.1.3. Invasive Functional Testing

Additional invasive functional testing may enhance evaluation in patients with possible myocardial infarction. In patients with spontaneous onset of symptoms and possible primary myocardial infarction, provocative testing (e.g. acetylcholine or ergonovine challenge) may aid in the recognition of epicardial or microvascular coronary spasm when coronary arteries appear angiographically non-obstructive.³ In patients with possible secondary myocardial infarction, particularly where the severity of coronary stenosis is unclear, invasive functional assessment using fractional flow reserve (FFR) or resting indices (e.g. resting full-cycle ratio [RFR] or instantaneous wave-free ratio [iFR]) may be helpful in detecting functionally significant stenosis that could cause myocardial ischemia in the context of supply–demand mismatch.^{3,208,209} Quantitative coronary angiography, in combination with computationally derived FFR, may also play a role in the detection of functionally significant lesions and may be helpful in supporting a diagnosis of secondary myocardial infarction.²¹⁰

15.2. Non-Invasive Imaging

Non-invasive cardiac imaging with echocardiography, CMR imaging, cardiac CT, single-photon emission computed tomography (SPECT), or positron emission tomography (PET) is valuable to help confirm the diagnosis of myocardial infarction or to identify an alternative mechanism for acute myocardial injury. In patients with confirmed myocardial infarction based on coronary angiography, the presence of regional wall motion abnormalities, loss of viable myocardium, and assessment of infarct size can corroborate the diagnosis and inform prognosis and treatment.^{211–214} Additionally, non-invasive imaging can identify complications of myocardial infarction, such as mitral regurgitation due to papillary muscle rupture, left ventricular aneurysm or pseudoaneurysm, ventricular septal defect, ventricular thrombus, right ventricular infarction, or pericardial effusion. Cardiac imaging can identify alternative mechanisms of acute myocardial injury, such as myocarditis, Takotsubo syndrome, or other cardiomyopathic processes.^{213,215–217}

15.2.1. Echocardiography

Echocardiography is a key bedside imaging tool in myocardial infarction, allowing assessment of regional wall motion abnormalities, infarct extent, ventricular thrombus, pseudoaneurysm, pericardial effusion/hematoma, and valvular dysfunction. It can identify alternative causes of the presentation, such as acute pericarditis, severe aortic

stenosis, aortic dissection, or massive pulmonary embolism.^{213,215–217} Point-of-care, hand-held echocardiography, when performed by trained personnel, can detect regional wall motion abnormalities induced by myocardial ischemia almost immediately.²¹⁸ When accompanied by evidence of acute myocardial injury, echocardiography can confirm the diagnosis of myocardial infarction.

Echocardiography has similar utility in the diagnosis of primary, secondary, and procedure-related myocardial infarction. In primary myocardial infarction, detection of regional wall motion abnormalities that correspond to the territory of the affected coronary artery can resolve uncertainty.^{211–213} In secondary myocardial infarction, differentiating between a global reduction in left ventricular function and regional wall motion abnormalities is important and can inform the diagnosis.²¹⁴ In procedure-related myocardial infarction, echocardiography can identify new regional wall motion abnormalities, assess left ventricular ejection fraction, and identify mechanical complications, and microvascular obstruction can be visualized using contrast echocardiography.^{213,219,220} It should be noted that immediately after cardiac surgery, adequate echo windows are not always obtainable. Further, not all myocardial infarction results in regional wall motion abnormalities, and where there is doubt, contrast echocardiography and CMR imaging can be helpful.^{221–224}



15.2.2. Cardiac Magnetic Resonance Imaging

CMR imaging provides detailed assessment of both reversible and irreversible myocardial injury. It can detect myocardial infarction, edema/inflammation, microvascular obstruction, and intramyocardial hemorrhage.^{225,226} Comprehensive CMR protocols include cine imaging (function), T2 (myocardial edema and inflammation, intramyocardial hemorrhage), T2* images (intramyocardial hemorrhage), first-pass perfusion and early gadolinium enhancement (microvascular obstruction), and late gadolinium enhancement (LGE) for myocardial necrosis, microvascular obstruction, and intramyocardial hemorrhage.²²⁷ Combining T2 with LGE helps discriminate between acute and chronic myocardial injury, though mild edema may persist for weeks in some infarcts.^{228–230} In patients with MINOCA the diagnostic yield is highest when CMR is performed within 2 weeks of myocardial infarction,²²⁴ whilst in patients with a large primary myocardial infarction, edema, microvascular obstruction and intramyocardial hemorrhage can persist for up to 6 months before resolving and evolving into fibrosis identifiable through LGE.

In primary myocardial infarction, CMR identifies infarct extent as well as microvascular obstruction and intramyocardial hemorrhage (Figure 11).^{231–235} CMR can identify complications, such as ventricular thrombus, papillary muscle infarction, right ventricular infarction, pseudoaneurysm formation (contained rup-




Figure 11. Stages of primary myocardial infarction by cardiac magnetic resonance imaging.

LGE, late gadolinium enhancement. Each stage of myocardial injury is progressive and is discernible from aggregate evidence derived from T2 maps (colour bar provided), LGE, and T2*-weighted (T2*w) short-axis cardiac magnetic resonance images along each row. *Stage 1* is marked by edema only (high signal on T2 map), absence of myocardial necrosis (lack of signal enhancement on LGE image), or myocardial hemorrhage (absence of low signal on T2*w images). *Stage 2* demonstrates both myocardial necrosis (high signal on LGE image) and edema (high signal on T2 map) but absence of persistent microvascular obstruction (absence of low signal within the zone of LGE image) or myocardial hemorrhage. *Stage 3* highlights the presence of microvascular obstruction (presence of low signal within the zone of high signal on LGE image) along with myocardial necrosis and edema. Note that myocardial hemorrhage is also absent at Stage 3 as evidenced by lack of low signal within the zone of infarction on T2*w image. *Stage 4* is characterized by myocardial hemorrhage (low signal within the zone of high signal in the T2*w image), along with microvascular obstruction, myocardial necrosis, and edema. High signal (brighter than surrounding tissue) and low signal (darker than surrounding tissue). Black arrowheads point to high signal on T2 maps, white arrow heads point to high signal on LGE, and blue arrow heads point to low signal on T2*w image. Note that low signal zones on Stage 3 and Stage 4 within LGE are demarcated by contoured zones within high signal zones in LGE. Red and green contours on LGE and T2w images are endocardial and epicardial contours segmenting the left ventricular myocardium. Adapted from Kumar *et al.* with permission.²³² For image description, please refer to the figure legend and surrounding text.

ture), and pericardial inflammation, as well as assess viability for revascularization (Figure 12). CMR also aids diagnosis when coronary angiography does not identify an acute coronary pathology (see *Section 6*), helping to differentiate myocardial infarction from alternative diagnoses such as acute myocarditis or Takotsubo syndrome. CMR can complement echocardiography in detecting new loss of viable myocardium or regional wall motion abnormality in a pattern consistent with ischemia.²³⁶ Similarly, in secondary or procedure-related myocardial infarction, CMR can identify new loss of viable myocardium or regional wall motion abnormality^{4,72,74,187}; however, image quality may be limited early after surgery due to artifact from chest drains or epicardial pacing wires.^{187,237}

15.2.3. Coronary Computed Tomography Angiography

CCTA is emerging as a valuable tool in the assessment of patients presenting with suspected ACS, offering sub-millimetre resolution to assess for presence of coronary atherosclerosis and plaque composition.^{238,239} Although current clinical guidelines do not recommend routine CCTA for suspected myocardial infarction in intermediate- and high-risk patients,^{1,2} it may be used to aid the diagnosis and has been found to offer prognostic information comparable to invasive coronary angiography in low-risk NSTEMI patients.²⁴⁰ CCTA may aid in differentiating primary from secondary myocardial infarction through plaque phenotyping^{241,242} and help identify non-atherothrom-




Figure 12. Role of cardiac magnetic resonance imaging in possible or confirmed primary myocardial infarction.

Illustration of common cardiac magnetic resonance imaging findings in patients with possible or confirmed primary myocardial infarction. (Top, left) Acute myocarditis with myocardial edema of the basal inferior and inferolateral wall (white arrow, left panel) with corresponding epicardial late enhancement (white arrow, right panel). (Top, middle) Focal myocardial infarction with transmural edema of the mid anterolateral wall (white arrow, left panel) and corresponding transmural late enhancement (white arrow, right panel). (Top, right) Takotsubo syndrome with transmural edema of the apical segments (white arrows, left panel), corresponding to the area of akinesia and in the absence of any late myocardial enhancement (right panel). (Bottom, left) Transmural myocardial infarction in the left circumflex territory with myocardial edema of the lateral wall (white arrow, left panel) and corresponding transmural late enhancement with a large dark or low signal core representing microvascular obstruction (white arrow, right panel). (Bottom, middle) Transmural myocardial infarction in the left anterior descending artery territory with left ventricular thrombus (white arrow, both panels). (Bottom, right) Transmural myocardial infarction with contained rupture of the basal inferoseptal wall with myocardial edema of the acutely injured wall (white arrow, left panel) and corresponding transmural late myocardial enhancement (white arrow, right panel). For image description, please refer to the figure legend and surrounding text.

botic causes for primary myocardial infarction, such as coronary artery dissection (although with limited resolution in smaller or more distal arteries).²⁴³ In secondary myocardial infarction, CCTA may detect obstructive or flow-limiting coronary artery disease,^{74,75} and in procedure-related myocardial infarction, assess graft patency.^{244,245} Emerging applications include photon-counting CT or late contrast enhancement CT may identify loss of viable myocardium or detect alternative causes of myocardial injury, such as myocarditis, Takotsubo syndrome, or cardiomyopathy.²⁴⁶

15.2.4. Nuclear Medicine

Nuclear medicine has advanced understanding of acute myocardial infarction but has limitations in clinical use. While SPECT and PET can assess myocardial blood flow

and viability, their spatial resolution restricts detection to transmural, not subendocardial, infarcts.^{247–249} Co-registration with CT or CMR enhances resolution and enables molecular imaging. Experimental PET tracers combined with CCTA may identify atherothrombosis²⁵⁰ and the source of thrombus due to coronary embolus in primary myocardial infarction.²⁵¹

16. SPECIAL SITUATIONS

Diagnosing acute myocardial infarction can be challenging when symptoms, ECG, cardiac biomarkers, or wall motion abnormalities are difficult to interpret—such as in early recurrent myocardial infarction, critical illness, recent surgery, or structural interventions. Elevated cardiac

biomarkers in heart failure, kidney disease, or cancer therapy may reflect non-ischemic injury, often requiring careful evaluation and multimodality imaging to confirm the diagnosis.

16.1. Recurrent Myocardial Infarction

Diagnosing a new myocardial infarction within the first month of the index event may be challenging, as electrocardiographic changes, cardiac biomarkers, and imaging findings may reflect residual abnormalities.^{252–255} Recurrent symptoms are common,²⁵⁶ but may stem from complications of the index myocardial infarction, such as pericarditis, chest trauma after cardiopulmonary resuscitation, or heart failure, rather than recurrent myocardial infarction. Symptoms may differ from the index event, with ~10% of males and ~16% of females reporting different symptoms from their first versus their recurrent presentation.²⁵⁷ Persistent ST-segment elevation or T-wave inversion may be caused by ventricular aneurysm or pericarditis.²⁵⁸ Further, cardiac troponin levels often remain elevated for weeks, so serial measurements are key, as rising values suggest recurrence.^{258,259} The use of CK-MB adds cost without providing diagnostic clarity.^{44,260} Coronary angiography is often required to confirm recurrent myocardial infarction, and to differentiate primary from procedure-related myocardial infarction within 30 days (see *Section 7*). CMR may help to distinguish multiple infarcts, suggesting the presence of more than one culprit lesion or that the culprit was not correctly identified during the index presentation.²⁶¹ Early recurrent myocardial infarction should still be classified according to the mechanism, rather than using a separate diagnostic code for recurrent myocardial infarction (Table 4).

16.2. Heart Failure

Myocardial injury without infarction is common in acute and chronic heart failure due to systolic and diastolic dysfunction.²⁶² Up to 90% of patients with acute heart failure have cardiac troponin T values above the URL,²⁶³ which may be misinterpreted as myocardial infarction due to overlapping symptoms.^{262,263} Cardiac troponin elevation can result from myocardial ischemia, supply–demand mismatch due to hypoxia, hypotension or tachycardia, apoptosis and autophagy, and direct cellular injury due to infiltrative processes, inflammation, or neurohormones.^{264,265} Experimental transient increases in left ventricular preload have resulted in cardiac troponin elevation, apoptosis, and myocardial stunning in the absence of ischemia,²⁶⁵ supporting preload-induced mechanical stretch as a possible mechanism for cardiac troponin release. Elevated cardiac troponins are of prognostic significance in acute and chronic heart failure^{266,267} and are indicative of the severity of heart failure.²⁶⁸

The spectrum of presentations with acute myocardial injury in heart failure includes primary or secondary myocardial infarction or non-ischemic causes. In patients presenting with heart failure, chest discomfort, or equivalent symptoms with signs of myocardial ischemia on the ECG, and/or known coronary artery disease, acute myocardial injury should raise the suspicion for primary myocardial infarction as the trigger for acute heart failure. Conditions that cause supply–demand imbalance, such as arrhythmias, hypoxia, hypotension, and anemia, increase the likelihood of secondary myocardial infarction. Clinical practice guidelines recommend that patients presenting with new-onset or acute heart failure should have cardiac troponin measured and an ECG performed on admission to exclude acute myocardial infarction.²⁶⁹ Serial measurements can improve discrimination between acute and chronic myocardial injury.⁷¹ Imaging may be required to exclude the diagnosis of primary or secondary myocardial infarction. Both diagnoses require careful interpretation and comparison with prior imaging as CAD and regional wall motion abnormalities frequently exist in heart failure.

16.3. Chronic Kidney Disease

Chronic kidney disease (CKD) is common among patients with myocardial infarction, and outcomes are worse among those with both conditions.^{270,271} Patients with CKD have a higher incidence of myocardial infarction due to shared cardiovascular risk factors and a greater risk of secondary myocardial infarction from infection, anemia, and hemodynamic stress associated with hemodialysis.

Cardiac troponin concentrations are frequently elevated but stable in CKD.²⁷² Whether this reflects chronic myocardial injury due to coexisting cardiac conditions or reduced clearance due to impaired glomerular filtration, or both, is not always clear.^{273,274} CKD is often a consequence of diabetes mellitus or hypertension, which also increase the risk of coronary heart disease.²⁷⁵ The term cardiovascular–kidney–metabolic syndrome has emerged to reflect this shared pathophysiology.²⁷⁶ As such, elevated cardiac troponin concentrations in patients with CKD should not be dismissed as being a consequence of reduced clearance alone.^{277,278}

Diagnosing myocardial infarction in advanced CKD is challenging due to the frequent presence of elevated cardiac troponin concentrations. In consecutive patients with possible myocardial infarction, cardiac troponin was three times more likely to be elevated in those with CKD compared with those without for both cardiac troponin T (70% vs 24%) and cardiac troponin I (47% vs 16%).^{272,279} While the final diagnosis is more likely to be primary myocardial infarction in those with CKD compared with those without (24% vs 11%), half of all patients with CKD and an elevated cardiac troponin concentration have

myocardial injury without infarction.²⁷² Therefore, serial cardiac troponin testing is particularly important to differentiate acute from chronic cardiac troponin elevations in CKD. The specificity and positive predictive value of the 99th percentile and high-risk thresholds are lower than among those without CKD.^{280,281} The use of higher thresholds for the diagnosis of myocardial infarction or greater absolute delta change values among those with CKD could improve the positive predictive value.²⁷⁹ However, these would be challenging to implement as the frequency of elevated cardiac troponin levels increases six-fold from 10% when the eGFR is 90 mL/min/1.73 m² or higher, to 66% when it falls below 30 mL/min/1.73 m².²⁷² Different thresholds would be required for each stage of CKD and for each assay. Adding to the challenge, patients with possible myocardial infarction and CKD are on average 20 years older than those without,²⁷⁹ which can also influence troponin thresholds. In the future, statistical models embedded into clinical decision support tools could address this complexity by integrating cardiac troponin and creatinine concentrations as continuous measures along with age, sex, and other comorbidities that influence cardiac troponin to estimate the probability of myocardial infarction.¹⁶⁸ However, the pathways for diagnosis of myocardial infarction (whether primary, secondary, or procedure-related) in patients with CKD are the same as for those without (see *Section 7.2*).

Due to greater diagnostic uncertainty and risk in patients with CKD, imaging is important to confirm the diagnosis of myocardial infarction. Concerns about the nephrotoxic effects of contrast agents have contributed to underinvestigation and worse outcomes.²⁸² Though the risk of contrast-induced nephropathy following coronary angiography in patients with CKD is often overstated,²⁸³ this risk, particularly in patients with advanced CKD, must be balanced against its potential benefits in guiding treatment.

16.4. Critical Illness

Critical illness, characterized by vital organ dysfunction, places a significant burden on the cardiovascular system and often requires artificial organ function replacement, sedation, and vasopressors.²⁸⁴ Systemic stress responses can lead to destabilization of atherosclerotic plaques, resulting in primary myocardial infarction. More often, critical illness leads to acute myocardial injury or secondary myocardial infarction due to oxygen supply-demand imbalance, which can arise from impaired oxygenation, anemia, impaired oxygen delivery, or a combination of these.

In analyses of discrepancies between clinical and post-mortem diagnoses in critically ill patients, myocardial infarction is among the most frequently missed diagnoses.^{285–287} Based on diagnostic criteria from previous UDMI classifications, an unrecognized myocardial infar-

tion occurred in 1 in 4 critically ill patients with known cardiovascular disease and was associated with a lower likelihood of survival.²⁸⁸ Often it is not possible to determine whether patients are experiencing symptoms due to sedation. Even in those who can report chest pain or dyspnea, these symptoms may be attributed to non-cardiac causes. Almost 40% of consecutive critically ill patients have myocardial injury and evidence of ischemia on the ECG.²⁸⁹ ST-segment elevation is common in critically ill patients and less specific for primary myocardial infarction.²⁹⁰ Without troponin data, ECG interpretation in critically ill patients shows only moderate reliability.²⁹¹ In critically ill patients with elevated cardiac troponin, a characteristic rise and fall pattern on serial testing is important to consider in the diagnosis of myocardial infarction.^{288,290,291} As with the diagnosis of myocardial infarction in other settings, confirmation (or exclusion) of the diagnosis requires coronary angiography and/or cardiac imaging (see *Section 7.2.1* and *Section 7.2.2*). Left ventricular dysfunction occurs in ~25% of patients in intensive care²⁹² and is associated with increased mortality, but half of cases stem from non-cardiac causes and most abnormalities are reversible.²⁹³ While the overall incidence of Takotsubo syndrome in critical illness is low (around 1.5%),²⁹⁴ there is some evidence that this is increasing in frequency,²⁹⁵ perhaps due to greater awareness. Coronary angiography may be needed to exclude myocardial infarction as a cause of regional left ventricular dysfunction.

Due to challenges in diagnosis and the high risk of primary and secondary myocardial infarction in critical illness, multidisciplinary care with cardiology input is essential. The risk of adverse outcomes from transportation of critically ill patients for coronary angiography and bleeding due to antithrombotic therapy needs careful consideration.^{296,297} Urgent coronary angiography may be necessary if primary myocardial infarction is likely or diagnosis will alter management. Otherwise, coronary and cardiac imaging to confirm or exclude a diagnosis of myocardial infarction may be deferred to prioritize stabilization of a critically ill patient.

16.5. Non-Cardiac Surgery

In patients undergoing non-cardiac surgery, myocardial infarction can result from a primary acute coronary pathology or be secondary to supply-demand mismatch. Any surgical procedure may increase the risk of myocardial infarction; however, the risk is greater for moderate- and high-risk surgeries, defined according to current guidelines,²⁹⁸ due to the greater chance for hemodynamic disturbance, blood loss, changes in body core temperature, and pro-inflammatory and prothrombotic changes.²⁹⁹

Diagnosing myocardial infarction after non-cardiac surgery is challenging due to symptom masking by

sedation and narcotics; most events occur within 2 days post-operatively, and up to two-thirds of patients have no ischemic symptoms.²⁹⁹ For this reason, when clinical deterioration or acute myocardial injury is identified, an ECG is essential to aid the diagnosis, with coronary angiography and/or cardiac imaging (see *Section 7.2.1* and *Section 7.2.2*) required to confirm the diagnosis of primary myocardial infarction; where ST-segment elevation is present, urgent coronary angiography is needed. In suspected secondary myocardial infarction, coronary or cardiac imaging is required to confirm the diagnosis, with timing based on its potential to impact peri-operative management. Post surgery, risks of anticoagulation and transport for coronary angiography require careful consideration. If active surveillance with cardiac troponin testing is performed, peri-operative myocardial injury (PMI) is identified in around 1 in 8 patients undergoing high-risk non-cardiac surgery.³⁰⁰ While the majority have peri-operative acute myocardial injury or secondary myocardial infarction due to supply-demand imbalance, a diagnosis of primary myocardial infarction occurs in 1 in 20 patients with myocardial injury in the post-operative setting (1 in 100 undergoing high-risk surgery overall).

The terms PMI and myocardial injury after non-cardiac surgery (MINS) have been used in this setting. PMI has been defined as acute myocardial injury with or without accompanying symptoms, or electrocardiographic or imaging evidence, of acute myocardial ischemia.²⁹⁸ MINS has been defined as a subset of PMI, in patients in whom myocardial injury was deemed most likely due to myocardial ischemia, in the absence of unexpected physiological stress from surgery or evidence of a cardiac non-coronary etiology.^{76,298} MINS may occur with or without concomitant symptoms or signs of ischemia and is associated with an eight-fold higher adjusted 30-day mortality.^{301,302} Identification of PMI or MINS should prompt an ECG, with further coronary and cardiac imaging required to confirm or exclude a diagnosis of primary or secondary myocardial infarction.

While clinical practice guidelines support routine post-operative cardiac troponin testing in high-risk patients or patients undergoing high-risk non-cardiac surgery,^{76,298,303} it is important to recognize that up to 40% of patients with cardiovascular disease or risk factors have baseline elevated cardiac troponin concentrations.³⁰⁴ For this reason, if assessment of cardiac troponin after surgery is planned, then a pre-operative value is essential in order to distinguish chronic myocardial injury from PMI or MINS.

16.6. Structural Cardiac Intervention

Structural cardiac interventions include a wide range of valve procedures, such as transcatheter aortic valve implantation (TAVI) and mitral transcatheter edge-to-edge repair (M-TEER), as well as septal/shunt closure and

appendage occlusion devices. Myocardial infarction can result from several mechanisms, including supply-demand imbalance in the setting of procedure-related hemodynamic changes, obstruction of coronary ostia due to displacement of valve leaflets, and coronary embolism of device material, air, or thrombus.

Myocardial infarction complicates up to 5.6% of TAVI, though pooled estimates suggest 1.1% based on previous definitions.³⁰⁵ Distinguishing procedure-related myocardial infarction from acute myocardial injury is challenging. Myocardial infarction from leaflet-related coronary ostium obstruction is uncommon, occurring in 0.7% of patients undergoing TAVI, but mortality is high.³⁰⁶ Myocardial infarction due to embolism from the valve—often presenting with ST-segment elevation and hypotension—is infrequent, occurring in 1% of patients undergoing TAVI.³⁰⁷ New LGE in an ischemic pattern was reported in as many as 18% of patients undergoing TAVI in whom CMR imaging was systematically performed.³⁰⁸ These lesions were typically small, multi-territorial, and unrelated to the presence of CAD, suggesting that procedure-related myocardial infarction due to embolism may be underrecognized.

Myocardial injury occurs in between 66% and 100% of patients following TAVI depending on the assay and definition used and whether transapical access was used.^{309–312} Myocardial injury was associated with short- and long-term mortality. Serial cardiac biomarker testing is important in patients with symptoms or signs of myocardial ischemia or clinical deterioration following structural intervention, but cardiac biomarkers should not be used in isolation to define a procedure-related myocardial infarction. The diagnosis of procedure-related myocardial infarction following structural cardiac interventions should follow the criteria outlined above (see *Section 7.2.3*), with the diagnosis requiring imaging evidence of new loss of viable myocardium in an ischemic pattern as a consequence of a procedural complication. This will facilitate comparisons between transcatheter and surgical approaches for the management of structural heart disease.

16.7. Cardio-Oncology

Patients receiving chemotherapy, particularly with anthracyclines or immune checkpoint inhibitors, are at increased risk of cardiac complications, including myocarditis, cancer therapy-related cardiac dysfunction, and thromboembolism, which can mimic or trigger myocardial infarction.³¹³ Cardiac troponin may rise from baseline or reflect new myocardial injury,^{314,315} so clinical guidelines recommend serial testing.³¹⁶ Therefore, when assessing a patient who is receiving cancer-related therapies, cardiac troponin values are more likely to be elevated, and in the appropriate clinical setting (see *Section 7.2.1* and *Section 7.2.2*) cardiac imaging is required to confirm

or exclude myocardial infarction. Given that both cancer and CAD are common, atherothrombosis remains the most common cause of primary myocardial infarction, but coronary vasospasm and embolism are more common in cancer due to the pro-inflammatory and hypercoagulability state of some cancers and the effects of chemotherapeutic agents.³¹⁷

17. IMPLICATIONS FOR PATIENTS, SECONDARY PREVENTION, AND CARDIAC REHABILITATION

Accurate diagnosis of myocardial infarction is important due to its significant prognostic, psychological, social, emotional, financial, insurance, and occupational impacts on patients and their families.³¹⁸ Following myocardial infarction, patients often report persistent symptoms such as fatigue, chest pain, low physical or mental energy, and reduced motivation that impact quality of life and well-being.^{318–320} Anxiety and depression are common, particularly among those with a prior history of depression, poor health, low socioeconomic status, and younger age at the time of diagnosis.^{320,321} Not all individuals perceive their recovery from myocardial infarction negatively; however, a small number of studies indicate that a myocardial infarction resulted in gratitude for one's life,³²² hope for the future,³²³ and having a second chance at life.³²⁴ Shared decision-making involving personalized discussions with patients, families, and clinicians is essential to incorporate values and concerns into diagnostic and treatment choices.

A diagnosis of myocardial infarction has lifelong treatment implications³²⁵ and may lead to financial burdens, including impacts on insurance and health and travel coverage. For patients living in high-income countries who are employed prior to myocardial infarction, most will return to work within one year, particularly if they are well-educated with a higher salary.³²⁶ Those with heart failure, diabetes, or depression are less likely to return to work, and regulations for occupations that require driving or flying may preclude some. Patients who return to work have a better psychological status following a diagnosis of myocardial infarction,³²⁷ with evidence that encouragement from clinicians increases the likelihood.^{326–328} Conversations with patients could include probing the consequences of a diagnosis of myocardial infarction on them personally with regards to their employment, relationships, or finances.

When there is diagnostic uncertainty between clinicians, patient outcomes have been found to be worse.³²⁹ By simplifying the classification into primary, secondary, and procedure-related myocardial infarction, we aim to make the diagnosis clearer and more understandable for both patients and clinicians. This in turn may facilitate tailoring of educational materials to patients who experience myocardial infarction. Cardiac rehabilitation

has demonstrated efficacy for people with myocardial infarction in clinical trials and is recommended by clinical practice guidelines. Cardiac rehabilitation is not currently tailored to the setting in which myocardial infarction arises and is less often offered to those with secondary myocardial infarction, where arguably the role for education is more important as patients are often managed outside specialist cardiac units.³³⁰

Survey data from Europe suggest that 90% of patients feel well informed about the role of secondary prevention following a diagnosis of primary myocardial infarction, but only 1 in 3 and 1 in 10 were aware of the targets for blood pressure and cholesterol, respectively.³³¹ Similar data from low- and middle-income countries (LMICs) suggest that less than half of patients in parts of Southern Africa are aware of secondary prevention targets.³³² No data are available for people with secondary myocardial infarction, perhaps because the role of CAD in the pathophysiology of the previous classification was uncertain, and therefore, the importance of risk factor modification was less clear. The use of diagnostic criteria for secondary myocardial infarction that include CAD may encourage inclusion of these patients in cardiac rehabilitation and education programmes. This is important as patient education improves engagement in cardiac rehabilitation, medication use, lifestyle choices, and understanding of symptoms,^{333,334} which could reduce delays in presentation among those with recurrent events.^{335,336} Clear diagnostic criteria for myocardial infarction may also help with strategies to improve referrals for rehabilitation, such as automatic referral systems, which in turn could improve participation.

18. IMPLICATIONS FOR PUBLIC HEALTH AND CLASSIFICATION OF DISEASE

Use of high-sensitivity cardiac troponin assays has increased the recognition of myocardial injury and, together with adoption of the Fourth UDMI, has led to an increase in the diagnoses of type 1 and type 2 myocardial infarction by 11% and 22%, respectively.³³⁷ However, as the Fourth UDMI was not aligned with the ICD-10 edition codes, this increase has not been reflected in morbidity and mortality data.³³⁸ For example, across two countries in Europe, fewer than 20% of patients who met the diagnostic criteria for type 2 myocardial infarction according to the Fourth UDMI received an ICD-10 code for myocardial infarction.⁴⁰ The increased recognition of myocardial injury has further presented health system challenges when myocardial infarction has been incorrectly assigned to patients with non-ischemic causes of acute myocardial injury.⁷³

Working together with the World Health Organization, the Fifth UDMI aims to address these challenges by including distinct ICD-11 codes for primary, secondary, and procedure-related myocardial infarction (Table 4).

For the first time, the ICD-11 code for acute myocardial infarction differentiates STEMI (BA41.0) from NSTEMI (BA41.1) and additional sixth digit stem codes have been incorporated to identify the underlying acute coronary pathology in primary myocardial infarction (1 to 9) and to differentiate primary from secondary (A) and procedure-related (B or C) myocardial infarction.

When primary myocardial infarction is considered to be the most likely diagnosis, but the acute coronary pathology remains uncertain following coronary angiography or cardiac imaging, or these investigations are not available or considered appropriate, specific stem codes are applied. In this scenario, BA41.0 or BA41.1 differentiates ST-segment elevation from non-ST-segment elevation myocardial infarction, respectively, with sixth digit stem codes applied for primary myocardial infarction of undetermined etiology following coronary imaging or for unknown etiology (no coronary imaging).

For secondary myocardial infarction, the structure of ICD-11 enables incorporation of associated diagnostic codes to indicate the acute alternative condition responsible for supply–demand mismatch to enable a better understanding of the risk of secondary myocardial infarction associated with common presentations. For procedure-related myocardial infarction, associated diagnostic codes differentiate percutaneous interventional (BA41.0B and BA41.1B) and open cardiac surgical (BA41.0C and BA41.1C) procedures. Finally, the ICD-11 code for unrecognized myocardial infarction (BA50) can be applied to better understand the frequency of this diagnosis and implications for patients.

While the implementation of ICD-11 coding across all countries will take time, it will enable epidemiological research, healthcare system planning, and public health monitoring of each type of myocardial infarction. The use of ICD-11 codes for STEMI would aid health service planning for delivery of immediate reperfusion therapy, while implementation of sixth digit stem codes to identify the acute coronary pathology will provide insights into the incidence of the less common mechanisms of primary myocardial infarction: SCAD, vasospasm, embolism, restenosis, stent thrombosis, or graft failure. In turn this will facilitate national and international comparisons of care and outcomes for these distinct conditions using routinely collected healthcare data. For successful implementation, healthcare providers—including clinicians and hospital coders—will need training on the new clinical classification of the Fifth UDMI and the associated ICD-11 codes.

19. IMPLICATIONS AND ADAPTIONS FOR LOW-RESOURCE SETTINGS

In 2021, there were 31.9 million new cases of ischemic heart disease globally—31.8% more than in 2010,³³⁹ driven mainly by ageing populations, urbanization, and rising risk factors of smoking, physical inactivity, obesity, and

diabetes in LMICs.^{340,341} People in low-resource settings, within or outside of LMICs, face many challenges in the diagnosis of myocardial infarction. This can be related to socioeconomic status, education and awareness, race and ethnic differences, as well as lack of timely access to emergency services, diagnostic testing, and treatment.³⁴² While clinical practice guidelines include access to high-sensitivity cardiac troponin testing as one of the 21 quality measures,³⁴³ in many LMICs there is limited availability.⁹ Guidance on the use of alternative biomarkers of myocardial injury for settings where high-sensitivity cardiac troponin testing is not available is found in *Section 14.2*. This further extends to electrocardiography, coronary angiography, and cardiac imaging, impacting the ability to diagnose myocardial infarction. For example, a study from Uganda reported that only 43% of healthcare facilities had access to troponin testing and 55% had access to electrocardiography.^{339,341,342,344} In China, the proportion of hospitals able to measure cardiac troponin increased from 25% to 84% between 2001 and 2011,³⁴⁵ while in India, coordinated strategies have been made to improve emergency ambulance services with pre-hospital electrocardiography.³⁴⁶ In the Eastern Mediterranean region, lack of access to cardiac catheterization affected the diagnosis of myocardial infarction.³⁴⁷ However, even where diagnostic testing is available, the affordability of these tests, which are often paid out of pocket in many LMICs, can further limit access.^{348–350}

Solutions include improved access to affordable point-of-care troponin testing, use of telemedicine for interpretation of the ECG, and widespread education to raise awareness.⁹ In 2023, the United Nations high-level meeting on universal health coverage for non-communicable diseases focused on a framework to achieve this by 2030, with global commitments to translate these into concrete actions. Basic diagnostic and laboratory equipment for the diagnosis of acute myocardial infarction in urgent care and emergency care facilities should be available to all. However, until such time, clinicians working in resource-limited settings still need to be able to make a diagnosis of myocardial infarction in the absence of cardiac biomarker testing or cardiac imaging (*Box 7*).³⁵¹

Box 7: Diagnostic Criteria for Myocardial Infarction in Low-Resource Settings Where Biomarker Testing and Cardiac Imaging Are Not Available

The diagnosis is **presumed** in those with symptoms consistent with acute myocardial ischemia where one or more clinical features are present:

- New ischemic changes on the electrocardiogram
- Development of pathological Q waves.

20. IMPLICATIONS OF THE DEFINITION ON RESEARCH

The previous classification of myocardial infarction was not always consistently applied in research, particularly in the evaluation of accelerated diagnostic pathways and in clinical trials of medical therapy and revascularization.^{352–358} The inclusion of all acute coronary mechanisms in the classification of primary myocardial infarction aims to improve the standardization of studies evaluating the performance of diagnostic pathways. Emerging artificial intelligence (AI) tools show promise in automating the interpretation of cardiac biomarkers and ECGs, enhancing the detection of ACS and supporting early myocardial infarction diagnoses.^{359–361} Machine-learning approaches are also providing new insights into the pathophysiological features of coronary atherosclerosis, which may ultimately guide personalized treatment and patient selection for invasive coronary angiography.³⁶² Further research is needed to determine if these tools, when used with appropriate clinician oversight, can improve diagnostic sensitivity, expedite decision-making, reduce diagnostic errors, and improve patient outcomes.

Many trials of therapy or intervention included patients with presumed primary myocardial infarction without confirming the diagnosis by coronary angiography, potentially enrolling those without true myocardial infarction, reducing the observed effectiveness of the intervention.³⁵⁵ Routine classification of primary myocardial infarction according to the acute coronary pathology (e.g. SCAD or coronary vasospasm) will enable the study of outcomes and support trials of tailored treatment pathways, which is particularly important for those pathologies where management may differ from atherothrombosis.³⁶³ Serial cardiac troponin measurements and cardiac imaging following reperfusion may improve staging of the severity of tissue injury in primary myocardial infarction,²³² improving patient selection for trials targeting events post infarction.

Embedding the UDMI classification with the associated ICD-11 codes into registries and electronic health records will facilitate enrollment and follow-up of patients with primary, secondary, and procedure-related myocardial infarction in registry-based or routine data-enabled clinical trials. Despite being common and linked to poorer outcomes, only one adequately powered randomized controlled trial has been conducted in patients with myocardial infarction and anemia.³⁵² This study compared liberal vs restrictive transfusion strategies and found no difference in death or recurrent infarction at 30 days. Subgroup analyses were consistent with the overall trial for those with type 2 myocardial infarction, who were expected to have the most to gain from liberal transfusion.³⁶⁴ These results underscore the need for clearer, objective diagnostic criteria for secondary myocardial infarction to ensure the enrollment of patients who are likely to benefit from interventions. Such criteria will facilitate enrollment into future studies that can

more rigorously evaluate targeted therapies in this group of patients.

The lack of consensus between interventional and surgical groups on defining procedure-related myocardial infarction has hindered trial interpretation for decades. *Post hoc* analyses demonstrate that varying criteria to define procedure-related myocardial infarction significantly affect reported rates of myocardial infarction across studies.^{48,365,366} This update to the UDMI aims to reduce variation between the definition of procedure-related myocardial infarction following percutaneous intervention and cardiac surgery to enable more reliable comparisons. Further research is needed to validate this approach and determine if the prognostic implications of procedure-related myocardial infarction are similar in both settings, which in turn will guide patient-centered clinical decision-making.

One of the objectives of this update in the classification is to improve consistency in the application of the diagnosis of myocardial infarction in clinical practice and clinical research so that the findings from research can be readily adopted into practice, allowing for improved epidemiological research and between-country comparisons. However, in recognition that there are areas in which uncertainty remains, suggestions for further research are provided (Box 8).

Box 8: Recommendations for Further Research

Research to encourage adoption:

- Evaluate the effects of implementation of the UDMI on the incidence of myocardial infarction across different healthcare systems and in different parts of the world.
- Evaluate the impact of implementation on the use of cardiac imaging in possible primary, secondary, and procedure-related myocardial infarction.
- Compare treatment and outcomes among patients with primary myocardial infarction stratified according to the underlying coronary etiology.
- Evaluate the impact of applying new diagnostic criteria for secondary myocardial infarction on treatment, management, and outcomes.
- Understand the prognostic implications and impact on health-related quality of life following procedure-related myocardial infarction and compare outcomes of coronary intervention with cardiac surgery.

Research to generate new knowledge in electrocardiography and biomarkers for myocardial infarction diagnosis:

- Determine whether strategies incorporating AI and machine learning (e.g. for electrocardiography and cardiac biomarker interpretation) can improve the accuracy and timeliness of myocardial infarction diagnosis.

- Improve our understanding of sex differences in the pathophysiology of myocardial infarction and the impact of sex-specific thresholds for cardiac troponin on investigation, treatment, and outcomes.
- Develop new approaches to define URLs for cardiac troponin with manufacturers and regulators that consider age, sex, and renal function so those that improve the recognition of myocardial injury due to myocardial infarction and/or other cardiac conditions in younger and older patients could be applied in practice.
- Determine the clinical and cost effectiveness of near-patient testing with point-of-care high-sensitivity cardiac troponin assays in the community, ambulance, and hospital settings.
- Determine whether routine cardiac troponin measurement following cardiac procedures improves the recognition of procedure-related myocardial infarction and outcomes.
- Evaluate the performance of sex-specific and uniform cardiac troponin thresholds (multiples of assay-specific URLs) to identify patients with a complication following a cardiac procedure who are unlikely to have a procedure-related myocardial infarction.
- Evaluate the utility of relative and absolute, assay-specific criteria for the recognition of chronic myocardial injury due to a cardiac condition.

Research to generate new knowledge in imaging for myocardial infarction diagnosis:

- Compare the diagnostic performance of echocardiography and cardiac magnetic resonance imaging in patients with possible secondary and procedure-related myocardial infarction.
- Determine the utility of CT coronary angiography with and without plaque phenotyping in patients with possible primary, secondary, and procedure-related myocardial infarction.
- Determine the utility of photon-counting CT coronary angiography with late iodine enhancement in possible myocardial infarction to identify acute coronary pathology and loss of viable myocardium.
- Determine if invasive physiological testing can improve diagnostic accuracy or outcomes in possible secondary myocardial infarction.
- Determine the diagnostic utility of cardiac imaging to investigate chronic myocardial injury in those without established cardiovascular disease.

Evidence Tables

Evidence tables are available at *Circulation* online.

Data Availability

No new data were generated or analysed in support of this research.

Author Information

Author/Task Force Member Affiliations: Wael A. Almahmeed, Heart Vascular and Thoracic Institute, Cleveland Clinic, Abu Dhabi, United Arab Emirates; Chiara Bucciarelli-Ducci, Cardiology and Cardiac MRI Department, Royal Brompton and Harefield Hospitals, Guys and St Thomas NHS Foundation Trust, London, United Kingdom, National Heart and Lung Institute, Imperial College London, London, United Kingdom, and School of Biomedical Engineering and Imaging Sciences, Kings College London, London, United Kingdom; Monica M. Colvin, Cardiovascular Division, University of Michigan, Ann Arbor, MI, United States of America; George D. Dangas, Cardiology, Mount Sinai Medical Center, New York, NY, United States of America; Holli A. DeVon, School of Nursing, University of California Los Angeles, Los Angeles, United States of America; Rohan Dharmakumar, Cardiovascular Imaging Research Center, Medical Imaging Research Institute, Department of Radiology and Imaging Sciences, Indiana University School of Medicine, Indianapolis, IN, United States of America, and Division of Cardiovascular Medicine, Department of Medicine, Indiana University School of Medicine, Indianapolis, IN, United States of America; Thomas A. Gaziano, Cardiovascular Medicine, Brigham and Women's Hospital, Boston, MA, United States of America, Health Policy and Management, Harvard School of Public Health, Boston, MA, United States of America, and Medicine, Harvard Medical School, Boston, MA, United States of America; Derek J. Hausenloy, Cardiovascular and Metabolic Disorders Program, Duke-NUS Medical School, Singapore, Singapore, National Heart Research Institute Singapore, National Heart Centre Singapore, Singapore, and The Hatter Cardiovascular Institute, University College London, London, United Kingdom; Mohamed Jeilan, Department of Medicine, Aga Khan University Hospital, Nairobi, Kenya; Konstantinos C. Koskinas, Department of Cardiology, Bern University Hospital, Inselspital, Bern, Switzerland; Bertil Lindahl, Department of Medical Sciences, Uppsala University, Uppsala, Sweden; Zuzana Motovska, Cardiac Center, Third Faculty of Medicine, Charles University and University Hospital Kralovske Vinohrady, Prague, Czech Republic; Maria Rubini Gimenez, Cardiology Department, Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain, and Cardiology Department, Ascires Biomedical Group, Valencia, Spain; Ana María Salvati, Cardiology, Clínica Modelo de Morón, Buenos Aires, Argentina, and Argentine Society of Cardiology, Buenos Aires, Argentina; Jacqueline E. Tamis-Holland, Cardiovascular Medicine, Cleveland Clinic, Heart, Vascular and Thoracic Institute, Cleveland, OH, United States of America.

APPENDIX

Approved by the ESC Clinical Practice Guidelines Committee on behalf of the ESC Board.

Approved by the ACC Presidential Team.

Approved by the AHA Executive Committee.

Approved by the WHF Executive Committee on behalf of the WHF Board, with recommendation from the Science Committee.

ESC/ACC/AHA/WHF Universal Definition Document Advisory Group includes Document Reviewers, ESC/ACC/AHA/WHF Committee Members, and ESC National Cardiac Societies.

Document Reviewers: Borja Ibanez (ESC Review Co-ordinator) (Spain), Elliott Antman (ACC/AHA Review Co-ordinator) (United States of America), Junbo Ge (WHF Review Co-ordinator) (China), Tayo Addo (United States of America), Junya Ako (Japan), Mirvat Alasnag (Saudi Arabia), Rasha Al-Lamee (United Kingdom), Paulo R. A. Caramori (Brazil), Mauricio G. Cohen (United States of America), Victoria Delgado (Spain), Bernard Dennis (United States of America), Sigrun Halvorsen (Norway), Julie Harris (United Kingdom), Milosz J. Jaguszewski (Poland), Hani Jneid (United States of America), Hugo Katus (Germany), Milan Milojevic (Serbia/Switzerland), Michelle O'Donoghue (United States of America), Julie Redfern (Australia), Joseph F. Sabik III (United States of America), Clara Saldarriaga (Colombia), Nizal Sarrafzadegan (Iran/Canada).

ESC CPG Committee members and WHF representatives who participated in the review of the Fifth Universal Definition of Myocardial Infarction:

Marianna Adamo (Italy), Suleman Aktaa (United Kingdom), Folkert W. Asselbergs (Netherlands), Michael A. Borger (Germany), Giuseppe Boriani (Italy), Margaritha Brida (Croatia), Robert A. Byrne (Ireland), Estelle Gandjbakhch (France), Bettina Heidecker (Germany), Anja Hennemuth (Germany), Borja Ibanez (Spain), Stefan James (Sweden), Ulf Landmesser (Germany), Gregory Y.H. Lip (United Kingdom), John William McEvoy (Ireland), Borislava Mihaylova (United Kingdom), Inge Moelgaard (Denmark), Jagat Narula (United States of America), Lis Neubeck (United Kingdom), Eva Prescott (Denmark), Bianca Rocca (Italy), Xavier Rossello (Spain), Anna Sannino (Germany), Felix C. Tanner (Switzerland), Katja Zeppenfeld (Netherlands).

ESC National Cardiac Societies actively involved in the review process of the Fifth Universal Definition of Myocardial Infarction:

Albania: Albanian Society of Cardiology, Naltin Shuka; Algeria: Algerian Society of Cardiology, Mohammed Chettibi; Armenia: Armenian Cardiologists Association, Hamlet G. Hayrapetyan; Austria: Austrian Society of Cardiology, Georg Delle Karth; Azerbaijan: Azerbaijan Society of Cardiology, Abbasaliyev Abbasali; Belgium: Belgian Society of Cardiology, Bernhard Gerber; Bosnia and Herzegovina: Association of Cardiologists in Bosnia and Herzegovina, Azra Durak Nalbantici; Bulgaria: Bulgarian Society of Cardiology, Elina Georgieva Baltadzhieva-Trendafilova; Croatia: Croatian Cardiac Society, Bosko Skoric; Cyprus: Cyprus Society of Cardiology, Kimon Myrianthopoulos; Czechia: Czech Society of Cardiology, Petr Ostadal; Denmark: Danish Society of Cardiology, Christian Juhl Terkelsen; Egypt: Egyptian Society of Cardiology, Moustafa Mokarrab; Estonia: Estonian Society of Cardiology, Toomas Marandi; Finland: Finnish Cardiac Society, Pekka Porela; France: French Society of Cardiology, Eric Van Belle; Georgia: Georgian Society of Cardiology, Zviad Kereselidze; Germany: German Cardiac Society, Tau Hartikainen; Greece: Hellenic Society of Cardiology, Konstantinos Toutouzias; Hungary: Hungarian Society of Cardiology, Istvan Ferenc Edes; Iceland: Icelandic Society of Cardiology, Geir Hirtlekar; Ireland: Irish Cardiac Society, James O'Neill; Israel: Israel Heart Society, David Hasdai; Italy: Italian Federation of Cardiology, Giuseppe Patti; Kazakhstan: Association of Cardiologists of Kazakhstan, Salim Berkinbayev; Kosovo (Republic of): Kosovo Society of Cardiology, Arlind Batalli; Kyrgyzstan: Kyrgyz Society of Cardiology, Olga Lunegova; Latvia: Latvian Society of Cardiology, Sanda Jegere; Lebanon: Lebanese Society of Cardiology, Ahmad Nabih Serhal; Libya: Libyan Cardiac Society, Aiman Smer; Lithuania: Lithuanian Society of Cardiology, Olivija Dobiliene; Luxembourg: Luxembourg Society of Cardiology, Jean Beissel; Malta: Maltese Cardiac Society, Matthew Mercieca Balbi; Moldova (Republic of): Moldavian Society of Cardiology, Aurel Andrei Grosu; Morocco: Moroccan Society of Cardiology, Samir S. Zlot; Netherlands: Netherlands Society of Cardiology, Roberto Diletti; North Macedonia: National Society of Cardiology of North Macedonia, Danica Petkoska Spirova; Norway: Norwegian Society of Cardiology, Kristin Angel; Poland: Polish Cardiac Society, Mariusz; Tomaniak; Portugal: Portuguese Society of Cardiology, Rita Calé Theotónio; Romania: Romanian Society of Cardiology, Razvan Ilie Radu; San Marino: San Marino Society of Cardiology, Marco Zavatta; Slovakia: Slovak Society of Cardiology, Martin Studencan; Slovenia: Slovenian Society of Cardiology, Matjaz Bunc; Spain: Spanish Society of Cardiology, Rut Andrea Riba; Sweden: Swedish Society of Cardiology, Erik Östgård Thunström; Switzerland: Swiss Society of Cardiology, Juan F. Iglesias; Syrian Arab Republic: Syrian Cardiovascular Association, Bani Marjeh; Mohammed Yassin; Tunisia: Tunisian Society of Cardiology and Cardiovascular Surgery, Ben El Hadj Zied; Türkiye: Turkish Society of Cardiology, Ertugrul Okuyan; Turkmenistan: Turkmen Committee of Cardiologists, Bahram Kadyrov; Ukraine: Ukrainian Association of Cardiology, Alexander Parkhomenko; United Kingdom of Great Britain and Northern Ireland: British Cardiovascular Society, David Hildick-Smith; Uzbekistan: Association of Cardiologists of Uzbekistan, Saodat Abdullaeva. Contributor either withdrew or was engaged in only a part of the review process: Newton Sigrist (Australia).

ARTICLE INFORMATION

Disclosures

All experts involved in the development of this Universal Definition have submitted declarations of interest, which are reported in a supplementary document to the Universal Definition. See the *European Heart Journal* online or <https://www.escardio.org/Guidelines> for submitted declarations of interests and evidence tables.

Disclaimer: This ESC/ACC/AHA/WHF Universal Definition Document represents the views of the ESC, ACC, AHA, and WHF and was produced after careful consideration of the scientific and medical knowledge and the available evidence at the time of its development. The ESC, ACC, AHA, and WHF are not responsible in the event of any contradiction, discrepancy, and/or ambiguity between the ESC/ACC/AHA/WHF Universal Definition Document and any other official recommendations or Expert Consensus Document issued by the relevant public health authorities, in particular in relation to good use of healthcare or therapeutic strategies. Health professionals are encouraged to take the ESC/ACC/AHA/WHF Universal Definition Document fully into account when exercising their clinical judgment, as well as in the determination and the implementation of preventive, diagnostic, or therapeutic medical strategies; however, the ESC/ACC/AHA/WHF Universal Definition Document does not override, in any way whatsoever, the individual responsibility of health professionals to make appropriate and accurate decisions in consideration of each patient's health condition and in consultation with that patient and, where appropriate and/or necessary, the patient's caregiver. Nor does the ESC/ACC/AHA/WHF Universal Definition Document exempt health professionals from taking into full and careful consideration

the relevant official updated recommendations, Universal Definition Documents or Expert Consensus Documents issued by the competent public health authorities, in order to manage each patient's case in light of the scientifically accepted data pursuant to their respective ethical and professional obligations. It is also the health professional's responsibility to verify the applicable rules and regulations relating to drugs and medical devices prior to making any clinical decision and to check whether a more recent version of this document exists. The ESC, ACC, AHA, and WHF warn readers that the technical language may be misinterpreted and declines any responsibility in this respect.

Permissions: The content of this ESC/ACC/AHA/WHF Universal Definition Document has been published for personal and educational use only. No commercial use is authorized. No part of the ESC/ACC/AHA/WHF Universal Definition Document may be translated or reproduced in any form without written permission from the ESC, ACC, AHA, WHF. Permissions can be obtained upon submission of a written request to Oxford University Press, the publisher of the *European Heart Journal* and the party authorized to handle such permissions on behalf of the ESC, ACC, AHA, and the WHF (journals.permissions@oup.com).

REFERENCES

- Byrne RA, Rossello X, Coughlan JJ, Barbato E, Berry C, Chieffo A, Claeys MJ, Dan G-A, Dweck MR, Galbraith M, et al; ESC Scientific Document Group. 2023 ESC guidelines for the management of acute coronary syndromes. *Eur Heart J*. 2023;44:3720–3826. doi: 10.1093/eurheartj/ehad191
- Rao SV, O'Donoghue ML, Ruel M, Rab T, Tamis-Holland JE, Alexander JH, Baber U, Baker H, Cohen MG, Cruz-Ruiz M, et al. 2025 ACC/AHA/ACEP/NAEMSP/SCAI guideline for the management of patients with acute coronary syndromes: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2025;151:e771–e862. doi: 10.1161/CIR.0000000000001309
- Vrints C, Andreotti F, Koskinas KC, Rossello X, Adamo M, Ainslie J, Banning AP, Budaj A, Buechel RR, Chiariello GA, et al; ESC Scientific Document Group. 2024 ESC guidelines for the management of chronic coronary syndromes. *Eur Heart J*. 2024;45:3415–3537. doi: 10.1093/eurheartj/ehae177
- Nomenclature and criteria for diagnosis of ischemic heart disease. Report of the Joint International Society and Federation of Cardiology/World Health Organization Task Force on Standardization of Clinical Nomenclature. *Circulation*. 1979;59:607–609. doi: 10.1161/01.cir.59.3.607
- Alpert JS, Thygesen K, Antman E, Bassand JP. Myocardial infarction redefined: a consensus document of the Joint European Society of Cardiology/American College of Cardiology Committee for the redefinition of myocardial infarction. *J Am Coll Cardiol*. 2000;36:959–969. doi: 10.1016/s0735-1097(00)00804-4
- Thygesen K, Alpert JS, White HD; Joint ESC/ACC/AHA/WHF Task Force for the Redefinition of Myocardial Infarction. Universal definition of myocardial infarction. *Eur Heart J*. 2007;28:2525–2538. doi: 10.1093/eurheartj/ehm355
- Thygesen K, Alpert JS, Jaffe AS, Simoons ML, Chaitman BR, White HD, Thygesen K, Alpert JS, White HD, Jaffe AS, et al; Joint ESC/ACC/AHA/WHF Task Force for Universal Definition of Myocardial Infarction. Third universal definition of myocardial infarction. *J Am Coll Cardiol*. 2012;60:1581–1598. doi: 10.1016/j.jacc.2012.08.001
- Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, White HD; Executive Group on behalf of the Joint European Society of Cardiology (ESC)/American College of Cardiology (ACC)/American Heart Association (AHA)/World Heart Federation (WHF) Task Force for the Universal Definition of Myocardial Infarction. Fourth universal definition of myocardial infarction (2018). *Circulation*. 2018;138:e618–e651. doi: 10.1161/CIR.0000000000000617
- Byrne R, Coughlan JJ, Rossello X, Ibanez B; Members of the Task Force for the 2023 ESC Guidelines for the management of acute coronary syndromes. Key priorities for the implementation of the 2023 ESC guidelines for the management of acute coronary syndromes in low-resource settings. *Eur Heart J Qual Care Clin Outcomes*. 2025;11:766–772. doi: 10.1093/ehjqcco/qcae107
- Jennings RB, Ganote CE. Structural changes in myocardium during acute ischemia. *Circ Res*. 1974;35:156–172. doi: 10.1161/res.35.3_supplement.iii-156
- Rossello X, Lobo-Gonzalez M, Ibanez B. Editor's choice: pathophysiology and therapy of myocardial ischaemia/reperfusion syndrome. *Eur Heart J Acute Cardiovasc Care*. 2019;8:443–456. doi: 10.1177/2048872619845283
- Fanaroff AC, Rymer JA, Goldstein SA, Simel DL, Newby LK. Does this patient with chest pain have acute coronary syndrome? The rational clinical

- cal examination systematic review. *JAMA*. 2015;314:1955–1965. doi: 10.1001/jama.2015.12735
13. Pasupathy S, Air T, Dreyer RP, Tavella R, Beltrame JF. Systematic review of patients presenting with suspected myocardial infarction and nonobstructive coronary arteries. *Circulation*. 2015;131:861–870. doi: 10.1161/CIRCULATIONAHA.114.011201
 14. Shanmuganathan M, Nikolaidou C, Burrage MK, Borlotti A, Kotronias R, Scarsini R, Banerjee A, Terentes-Printzios D, Pitcher A, Gara E, et al; Oxford Acute Myocardial Infarction (OxAMI) Study Investigators. Cardiovascular magnetic resonance before invasive coronary angiography in suspected non-ST-segment elevation myocardial infarction. *JACC Cardiovasc Imaging*. 2024;17:1044–1058. doi: 10.1016/j.jcmg.2024.05.007
 15. Shah ASV, Sandoval Y, Noaman A, Sexter A, Vaswani A, Smith SW, Gibbins M, Griffiths M, Chapman AR, Strachan FE, et al. Patient selection for high sensitivity cardiac troponin testing and diagnosis of myocardial infarction: prospective cohort study. *BMJ*. 2017;359:j4788. doi: 10.1136/bmj.j4788
 16. Steiro OT, Langorgen J, Tjora HL, Bjørneklett RO, Skadberg O, Bonarjee VVS, Mjelva OR, Steinsvik T, Lindahl B, Omland T, et al. Prognostic significance of chronic myocardial injury diagnosed by three different cardiac troponin assays in patients admitted with suspected acute coronary syndrome. *Clin Chem Lab Med*. 2024;62:729–739. doi: 10.1515/cclm-2023-0336
 17. Saeed N, Steiro OT, Langorgen J, Tjora HL, Bjørneklett RO, Skadberg O, Bonarjee VVS, Mjelva OR, Norekvål TM, Steinsvik T, et al. Diagnosing myocardial injury in an acute chest pain cohort; long-term prognostic implications of cardiac troponin T and I. *Clin Chem*. 2024;70:1241–1255. doi: 10.1093/clinchem/hvae110
 18. DeFilippis AP, Chapman AR, Mills NL, de Lemos JA, Arbab-Zadeh A, Newby LK, Morrow DA. Assessment and treatment of patients with type 2 myocardial infarction and acute nonischemic myocardial injury. *Circulation*. 2019;140:1661–1678. doi: 10.1161/CIRCULATIONAHA.119.040631
 19. Landesberg G, Jaffe AS, Gilon D, Levin PD, Goodman S, Abu-Baih A, Beeri R, Weissman C, Sprung CL, Landesberg A. Troponin elevation in severe sepsis and septic shock: the role of left ventricular diastolic dysfunction and right ventricular dilatation. *Crit Care Med*. 2014;42:790–800. doi: 10.1097/CCM.0000000000000107
 20. Singh T, Khan H, Gamble DT, Scally C, Newby DE, Dawson D. Takotsubo syndrome: pathophysiology, emerging concepts, and clinical implications. *Circulation*. 2022;145:1002–1019. doi: 10.1161/CIRCULATIONAHA.121.055854
 21. Templin C, Ghadri JR, Diekmann J, Napp LC, Bataiosu DR, Jaguszewski M, Cammann VL, Sarcon A, Geyer V, Neumann CA, et al. Clinical features and outcomes of takotsubo (stress) cardiomyopathy. *N Engl J Med*. 2015;373:929–938. doi: 10.1056/NEJMoa1406761
 22. Eichhorn C, Greulich S, Bucciarelli-Ducci C, Sznitman R, Kwong RY, Gräni C. Multiparametric cardiovascular magnetic resonance approach in diagnosing, monitoring, and prognostication of myocarditis. *JACC Cardiovasc Imaging*. 2022;15:1325–1338. doi: 10.1016/j.jcmg.2021.11.017
 23. Arnadóttir A, Pedersen S, Bo Hasselbalch R, Goetze JP, Friis-Hansen LJ, Bloch-Munster AM, Jensen JS, Bundgaard H, Iversen K. Temporal release of high-sensitivity cardiac troponin T and I and copeptin after brief induced coronary artery balloon occlusion in humans. *Circulation*. 2021;143:1095–1104. doi: 10.1161/CIRCULATIONAHA.120.046574
 24. Canty JM Jr. Myocardial injury, troponin release, and cardiomyocyte death in brief ischemia, failure, and ventricular remodeling. *Am J Physiol Heart Circ Physiol*. 2022;323:H1–H15. doi: 10.1152/ajpheart.00093.2022
 25. Mair J, Lindahl B, Hammarsten O, Muller C, Giannitsis E, Huber K, Möckel M, Plebani M, Thygesen K, Jaffe AS. How is cardiac troponin released from injured myocardium? *Eur Heart J Acute Cardiovasc Care*. 2018;7:553–560. doi: 10.1177/2048872617748553
 26. Aengevaeren VL, Baggish AL, Chung EH, George K, Kleiven O, Mingels AMA, Ørn S, Shave RE, Thompson PD, Eijssvogels TMH. Exercise-induced cardiac troponin elevations: from underlying mechanisms to clinical relevance. *Circulation*. 2021;144:1955–1972. doi: 10.1161/CIRCULATIONAHA.121.056208
 27. Fridrich O, Peretz A, Fox-Fisher I, Pyanin S, Dadon Z, Shcolnik E, Sadeh R, Fialkoff G, Sharkia I, Moss J, et al. Elevated cfDNA after exercise is derived primarily from mature polymorphonuclear neutrophils, with a minor contribution of cardiomyocytes. *Cell Rep Med*. 2023;4:101074. doi: 10.1016/j.xcrm.2023.101074
 28. Friden V, Starnberg K, Muslimovic A, Ricksten SE, Bjurman C, Forsgard N, Wickman A, Hammarsten O. Clearance of cardiac troponin T with and without kidney function. *Clin Biochem*. 2017;50:468–474. doi: 10.1016/j.clinbiochem.2017.02.007
 29. Muslimovic A, Friden V, Tenstad O, Starnberg K, Nystrom S, Wesen E, Esbjörner EK, Granholm K, Lindahl B, Hammarsten O. The liver and kidneys mediate clearance of cardiac troponin in the rat. *Sci Rep*. 2020;10:6791. doi: 10.1038/s41598-020-63744-8
 30. Zaman S, Wasfy JH, Kapil V, Ziaieian B, Parsonage WA, Sriswasdi S, Chico TJA, Capodanno D, Collieran R, Sutton NR, et al. The Lancet Commission on rethinking coronary artery disease: moving from ischaemia to atheroma. *Lancet*. 2025;405:1264–1312. doi: 10.1016/S0140-6736(25)00055-8
 31. Chin CW, Shah AS, McAllister DA, Joanna Cowell S, Alam S, Langrish JP, Strachan FE, Hunter AL, Maria Choy A, Lang CC, et al. High-sensitivity troponin I concentrations are a marker of an advanced hypertrophic response and adverse outcomes in patients with aortic stenosis. *Eur Heart J*. 2014;35:2312–2321. doi: 10.1093/eurheartj/ehu189
 32. Nellessen U, Goder S, Schobor R, Abawi M, Hecker H, Tschöke S. Serial analysis of troponin I levels in patients with ischemic and nonischemic dilated cardiomyopathy. *Clin Cardiol*. 2006;29:219–224. doi: 10.1002/clc.4960290510
 33. Willinger L, Brudy L, Meyer M, Oberhoffer-Fritz R, Ewert P, Müller J. Prognostic value of non-acute high sensitive troponin-T for cardiovascular morbidity and mortality in adults with congenital heart disease: a systematic review. *J Cardiol*. 2021;78:206–212. doi: 10.1016/j.jicc.2021.02.008
 34. Eggers KM, Lindahl B. Application of cardiac troponin in cardiovascular diseases other than acute coronary syndrome. *Clin Chem*. 2017;63:223–235. doi: 10.1373/clinchem.2016.261495
 35. Lindahl B, Thurston AJ, Tew YY, McDermott M, Fujisawa T, Lynch S, Cooper JG, Gray AJ, Jernberg T, Mills NL; POC-ET Investigators. Change in cardiac troponin T to differentiate acute from chronic myocardial injury in the emergency department. *Clin Biochem*. 2026;141:111055. doi: 10.1016/j.clinbiochem.2025.111055
 36. Roos A, Edgren G. Using historical cardiac troponins to identify patients at a high risk of myocardial infarction. *Heart*. 2022;109:127–133. doi: 10.1136/heartjnl-2022-321198
 37. Boeddinghaus J, Lopez-Ayala P, Thurston AJF, Lee KK, Tew YY, Doudesis D, Kadesjö E, Roos A, Raphael CE, Sandoval Y, et al; MI-2 Collaborators. Cardiovascular risk across myocardial injury and infarction categories using the universal definition: an individual patient-level data meta-analysis. *J Am Coll Cardiol*. 2026;87:2282–2297. doi: 10.1016/j.jacc.2025.11.043
 38. de Lemos JA, Newby LK, Mills NL. A proposal for modest revision of the definition of type 1 and type 2 myocardial infarction. *Circulation*. 2019;140:1773–1775. doi: 10.1161/CIRCULATIONAHA.119.042157
 39. Lindahl B, Mills NL. A new clinical classification of acute myocardial infarction. *Nat Med*. 2023;29:2200–2205. doi: 10.1038/s41591-023-02513-2
 40. Taggart C, Roos A, Kadesjö E, Anand A, Li Z, Doudesis D, Lee KK, Bularga A, Wereski R, Lowry MTH, et al. Application of the universal definition of myocardial infarction in clinical practice in Scotland and Sweden. *JAMA Netw Open*. 2024;7:e245853. doi: 10.1001/jamanetworkopen.2024.5853
 41. Gard A, Lindahl B, Baron T. Impact of clinical diagnosis of myocardial infarction in patients with elevated cardiac troponin. *Heart*. 2023;109:1533–1541. doi: 10.1136/heartjnl-2022-322298
 42. Devereaux PJ, Lamy A, Chan MTV, Allard RV, Lomivorotov VV, Landoni G, Zheng H, Paparella D, McGillion MH, Belley-Côté EP, et al; VISION Cardiac Surgery Investigators. High-sensitivity troponin I after cardiac surgery and 30-day mortality. *N Engl J Med*. 2022;386:827–836. doi: 10.1056/NEJMoa2000803
 43. Gaudio M, Jaffe AS, Milojevic M, Sandoval Y, Devereaux PJ, Thygesen K, Myers PO, Kluin J. Great debate: myocardial infarction after cardiac surgery must be redefined. *Eur Heart J*. 2024;45:4170–4177. doi: 10.1093/eurheartj/ehae416
 44. Jaffe AS, Lindahl B, Giannitsis E, Mueller C, Cullen L, Hammarsten O, Mockel M, Mair J, Krychtiuk KA, Huber K, et al. ESC Study Group on Cardiac Biomarkers of the Association for CardioVascular Care: a fond farewell at the retirement of CKMB. *Eur Heart J*. 2021;42:2260–2264. doi: 10.1093/eurheartj/ehaa1079
 45. Ueki Y, Otsuka T, Bar S, Koskinas KC, Heg D, Haner J, Siontis GCM, Praz F, Hunziker L, Lanz J, et al. Frequency and outcomes of periprocedural MI in patients with chronic coronary syndromes undergoing PPCI. *J Am Coll Cardiol*. 2022;79:513–526. doi: 10.1016/j.jacc.2021.11.047
 46. Bulluck H, Paradies V, Barbato E, Baumbach A, Botker HE, Capodanno D, De Caterina R, Cavallini C, Davidson SM, Feldman DN, et al. Prognostically relevant periprocedural myocardial injury and infarction associated with percutaneous coronary interventions: a consensus document of the ESC Working Group on Cellular Biology of the Heart and European Association of Percutaneous Cardiovascular Interventions (EAPCI). *Eur Heart J*. 2021;42:2630–2642. doi: 10.1093/eurheartj/ehab271

47. Gaudino M, Flather M, Capodanno D, Milojevic M, Bhatt DL, Biondi Zoccai G, Boden WE, Devereaux RJ, Doenst T, Farkouh M, et al. European Association of Cardio-Thoracic Surgery (EACTS) expert consensus statement on perioperative myocardial infarction after cardiac surgery. *Eur J Cardiothorac Surg*. 2024;65:ezad415. doi: 10.1093/ejcts/ezad415
48. Hara H, Serruys PW, Takahashi K, Kawashima H, Ono M, Gao C, Wang R, Mohr FW, Holmes DR, Davierwala PM, et al; SYNTAX Extended Survival Investigators. Impact of peri-procedural myocardial infarction on outcomes after revascularization. *J Am Coll Cardiol*. 2020;76:1622–1639. doi: 10.1016/j.jacc.2020.08.009
49. DeWood MA, Spores J, Notske R, Mouser LT, Burroughs R, Golden MS, Lang HT. Prevalence of total coronary occlusion during the early hours of transmural myocardial infarction. *N Engl J Med*. 1980;303:897–902. doi: 10.1056/NEJM198010163031601
50. Khan AR, Golwala H, Tripathi A, Bin Abdulhak AA, Bavishi C, Riaz H, Mallipeddi V, Pandey A, Bhatt DL. Impact of total occlusion of culprit artery in acute non-ST elevation myocardial infarction: a systematic review and meta-analysis. *Eur Heart J*. 2017;38:3082–3089. doi: 10.1093/eurheartj/ehx418
51. McLaren J, de Alencar JN, Aslanger EK, Meyers HP, Smith SW. From ST-segment elevation MI to occlusion MI: the new paradigm shift in acute myocardial infarction. *JACC Adv*. 2024;3:101314. doi: 10.1016/j.jaccadv.2024.101314
52. Ricci F, Martini C, Scordo DM, Rossi D, Gallina S, Fedorowski A, Sciarra L, Chahal CAA, Meyers HP, Herman R, et al. ECG patterns of occlusion myocardial infarction: a narrative review. *Ann Emerg Med*. 2025;85:330–340. doi: 10.1016/j.annemergmed.2024.11.019
53. Spirito A, Vainora L, Papadis A, Iacovelli F, Sardu C, Selberg A, Bär S, Kavaliaskaite R, Temperli F, Asatryan B, et al. Acute coronary occlusion in patients with non-ST-segment elevation out-of-hospital cardiac arrest. *J Am Coll Cardiol*. 2023;81:446–456. doi: 10.1016/j.jacc.2022.10.039
54. Miranda DF, Lobo AS, Walsh B, Sandoval Y, Smith SW. New insights into the use of the 12-lead electrocardiogram for diagnosing acute myocardial infarction in the emergency department. *Can J Cardiol*. 2018;34:132–145. doi: 10.1016/j.cjca.2017.11.011
55. Bentzon JF, Otsuka F, Virmani R, Falk E. Mechanisms of plaque formation and rupture. *Circ Res*. 2014;114:1852–1866. doi: 10.1161/CIRCRESAHA.114.302721
56. Fahed AC, Jang IK. Plaque erosion and acute coronary syndromes: phenotype, molecular characteristics and future directions. *Nat Rev Cardiol*. 2021;18:724–734. doi: 10.1038/s41569-021-00542-3
57. Kubo T, Imanishi T, Takarada S, Kuroi A, Ueno S, Yamano T, Tanimoto T, Matsuo Y, Masho T, Kitabata H, et al. Assessment of culprit lesion morphology in acute myocardial infarction: ability of optical coherence tomography compared with intravascular ultrasound and coronary angiography. *J Am Coll Cardiol*. 2007;50:933–939. doi: 10.1016/j.jacc.2007.04.082
58. Falk E, Nakano M, Bentzon JF, Finn AV, Virmani R. Update on acute coronary syndromes: the pathologists' view. *Eur Heart J*. 2013;34:719–728. doi: 10.1093/eurheartj/ehs411
59. Ambrose JA, Tannenbaum MA, Alexopoulos D, Hjelm Dahl-Monsen CE, Leavy J, Weiss M, Borrico S, Gorlin R, Fuster V. Angiographic progression of coronary artery disease and the development of myocardial infarction. *J Am Coll Cardiol*. 1988;12:56–62. doi: 10.1016/0735-1097(88)90356-7
60. Adam D, Alfonso F, Maas A, Vrints C; Writing Committee. European Society of Cardiology, Acute Cardiovascular Care Association, SCAD Study Group: a position paper on spontaneous coronary artery dissection. *Eur Heart J*. 2018;39:3353–68. doi: 10.1093/eurheartj/ehy080
61. Hayes SN, Kim ESH, Saw J, Adam D, Arslanian-Engoren C, Economy KE, Ganesh SK, Gulati R, Lindsay ME, Mieres JH, et al; American Heart Association Council on Peripheral Vascular Disease; Council on Clinical Cardiology; Council on Cardiovascular and Stroke Nursing; Council on Genomic and Precision Medicine; and Stroke Council. Spontaneous coronary artery dissection: current state of the science: a scientific statement from the American Heart Association. *Circulation*. 2018;137:e523–e557. doi: 10.1161/CIR.0000000000000564
62. Gulati M, Levy PD, Mukherjee D, Amsterdam E, Bhatt DL, Birtcher KK, Blankstein R, Boyd J, Bullock-Palmer RP, Conejo T, et al. 2021 AHA/ACC/ASE/CHEST/SAEM/SCCT/SCMR guideline for the evaluation and diagnosis of chest pain: executive summary: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2021;144:e368–e454. doi: 10.1161/CIR.0000000000001030
63. Tamis-Holland JE, Jneid H, Reynolds HR, Agewall S, Brilakis ES, Brown TM, Lerman A, Cushman M, Kumbhani DJ, Arslanian-Engoren C, et al; American Heart Association Interventional Cardiovascular Care Committee of the Council on Clinical Cardiology; Council on Cardiovascular and Stroke Nursing; Council on Epidemiology and Prevention; and Council on Quality of Care and Outcomes Research. Contemporary diagnosis and management of patients with myocardial infarction in the absence of obstructive coronary artery disease: a scientific statement from the American Heart Association. *Circulation*. 2019;139:e891–e908. doi: 10.1161/CIR.0000000000000670
64. Chapman AR, Taggart C, Boedinghaus J, Mills NL, Fox KAA. Type 2 myocardial infarction: challenges in diagnosis and treatment. *Eur Heart J*. 2025;46:504–517. doi: 10.1093/eurheartj/ehae803
65. Sandoval Y, Jaffe AS. Type 2 myocardial infarction: JACC review topic of the week. *J Am Coll Cardiol*. 2019;73:1846–1860. doi: 10.1016/j.jacc.2019.02.018
66. Shah AS, Anand A, Sandoval Y, Lee KK, Smith SW, Adamson PD, Chapman AR, Langdon T, Sandeman D, Vaswani A, et al; High-STEACS investigators. High-sensitivity cardiac troponin I at presentation in patients with suspected acute coronary syndrome: a cohort study. *Lancet*. 2015;386:2481–2488. doi: 10.1016/S0140-6736(15)00391-8
67. Saaby L, Poulsen TS, Hosbond S, Larsen TB, Pyndt Diederichsen AC, Hallas J, Thygesen K, Mickley H. Classification of myocardial infarction: frequency and features of type 2 myocardial infarction. *Am J Med*. 2013;126:789–797. doi: 10.1016/j.amjmed.2013.02.029
68. Bularga A, Taggart C, Mendusic F, Kimenai DM, Wereski R, Lowry MTH, Lee KK, Ferry AV, Stewart SS, McAllister DA, et al; High-Sensitivity Troponin in the Evaluation of Patients with Suspected Acute Coronary Syndrome (High-STEACS) Investigators. Assessment of oxygen supply-demand imbalance and outcomes among patients with type 2 myocardial infarction: a secondary analysis of the High-STEACS cluster randomized clinical trial. *JAMA Netw Open*. 2022;5:e220162. doi: 10.1001/jamanetworkopen.2022.20162
69. Neumann JT, Sorensen NA, Rubsamen N, Ojeda F, Renne T, Qaderi V, Teltrop E, Kramer S, Quantius L, Zeller T, et al. Discrimination of patients with type 2 myocardial infarction. *Eur Heart J*. 2017;38:3514–3520. doi: 10.1093/eurheartj/ehx457
70. Sarkisian L, Saaby L, Poulsen TS, Gerke O, Jangaard N, Hosbond S, Diederichsen ACP, Thygesen K, Mickley H. Clinical characteristics and outcomes of patients with myocardial infarction, myocardial injury, and nonelevated troponins. *Am J Med*. 2016;129:446.e5–446.e21. doi: 10.1016/j.amjmed.2015.11.006
71. Wereski R, Kimenai DM, Taggart C, Doudesis D, Lee KK, Lowry MTH, Bularga A, Lowe DJ, Fujisawa T, Apple FS, et al. Cardiac troponin thresholds and kinetics to differentiate myocardial injury and myocardial infarction. *Circulation*. 2021;144:528–538. doi: 10.1161/CIRCULATIONAHA.121.054302
72. Bularga A, Hung J, Daghem M, Stewart S, Taggart C, Wereski R, Singh T, Meah MN, Fujisawa T, Ferry AV, et al. Coronary artery and cardiac disease in patients with type 2 myocardial infarction: a prospective cohort study. *Circulation*. 2022;145:1188–1200. doi: 10.1161/CIRCULATIONAHA.121.058542
73. McCarthy C, Murphy S, Cohen JA, Rehman S, Jones-O'Connor M, Olshan DS, Singh A, Vaduganathan M, Januzzi JL, Wasfy JH. Misclassification of myocardial injury as myocardial infarction: implications for assessing outcomes in value-based programs. *JAMA Cardiol*. 2019;4:460–464. doi: 10.1001/jamacardio.2019.0716
74. Boedinghaus J, Bularga A, Taggart C, Wereski R, McDermott M, Thurston AJF, Ferry AV, Williams MC, Baker AH, Dweck MR, et al; DEMAND-MI Investigators. Implications of a new clinical classification of acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care*. 2025;14:131–141. doi: 10.1093/ehjacc/zuaf002
75. McCarthy CP, Murphy SP, Amponsah DK, Rambarat PK, Lin C, Liu Y, Mohebi R, Levin A, Raghavan A, Miksenas H, et al. Coronary computed tomographic angiography with fractional flow reserve in patients with type 2 myocardial infarction. *J Am Coll Cardiol*. 2023;82:1676–1687. doi: 10.1016/j.jacc.2023.08.020
76. Ruetzler K, Smilowitz NR, Berger JS, Devereaux PJ, Maron BA, Newby LK, de Jesus Perez V, Sessler DI, Wijeyesundera DN. Diagnosis and management of patients with myocardial injury after noncardiac surgery: a scientific statement from the American Heart Association. *Circulation*. 2021;144:e287–e305. doi: 10.1161/CIR.0000000000001024
77. Kurz DJ, Naegeli B, Bertel O. A double-blind, randomized study of the effect of immediate intravenous nitroglycerin on the incidence of postprocedural chest pain and minor myocardial necrosis after elective coronary stenting. *Am Heart J*. 2000;139:35–43. doi: 10.1016/s0002-8703(00)90306-5
78. Belley-Cote EP, Whitlock RP, Lamy A, Alhussein MM, Lomivorotov V, Brady K, Chan MTV, Allard RV, Ajello S, Wang CY, et al. Association between electrocardiographic changes and myocardial injury or death after cardiac surgery. *Can J Cardiol*. 2025;41:2601–2610. doi: 10.1016/j.cjca.2025.09.035

79. Polzl L, Thielmann M, Cymorek S, Nagele F, Hirsch J, Graber M, Sappeler N, Eder J, Staggli S, Theurl F, et al. Impact of myocardial injury after coronary artery bypass grafting on long-term prognosis. *Eur Heart J*. 2022;43:2407–2417. doi: 10.1093/eurheartj/ehac054
80. Crescenzi G, Bove T, Pappalardo F, Scandroglio AM, Landoni G, Aletti G, Zangrillo A, Alfieri O. Clinical significance of a new Q wave after cardiac surgery. *Eur J Cardiothorac Surg*. 2004;25:1001–1005. doi: 10.1016/j.ejcts.2004.02.030
81. Virani SS, Newby LK, Arnold SV, Bittner V, Brewer LC, Demeter SH, Dixon DL, Fearon WF, Hess B, Johnson HM, et al; Peer Review Committee Members. 2023 AHA/ACC/ACCP/ASPC/NLA/PCNA guideline for the management of patients with chronic coronary disease: a report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. *Circulation*. 2023;148:e9–e119. doi: 10.1161/CIR.0000000000001168
82. Heuts S, Gollmann-Tepekoylu C, Denessen EJS, Olsthoorn JR, Romeo JLR, Maessen JG, van 't Hof AWJ, Bekers O, Hammarsten O, Pölzl L, et al. Cardiac troponin release following coronary artery bypass grafting: mechanisms and clinical implications. *Eur Heart J*. 2023;44:100–112. doi: 10.1093/eurheartj/ehac604
83. Moussa ID, Klein LW, Shah B, Mehran R, Mack MJ, Brilakis ES, Reilly JP, Zoghbi G, Holper E, Stone GW. Consideration of a new definition of clinically relevant myocardial infarction after coronary revascularization: an expert consensus document from the Society for Cardiovascular Angiography and Interventions (SCAI). *J Am Coll Cardiol*. 2013;62:1563–1570. doi: 10.1016/j.jacc.2013.08.720
84. Ndrepepa G, Collieran R, Braun S, Cassese S, Hieber J, Fusaro M, Kufner S, Ott I, Byrne RA, Huser O, et al. High-sensitivity troponin T and mortality after elective percutaneous coronary intervention. *J Am Coll Cardiol*. 2016;68:2259–2268. doi: 10.1016/j.jacc.2016.08.059
85. Kini A, Marmur JD, Kini S, Dargas G, Cocke TP, Wallenstein S, Brown E, Ambrose JA, Sharma SK. Creatine kinase-MB elevation after coronary intervention correlates with diffuse atherosclerosis, and low-to-medium level elevation has a benign clinical course: implications for early discharge after coronary intervention. *J Am Coll Cardiol*. 1999;34:663–671. doi: 10.1016/s0735-1097(99)00298-3
86. Garcia-Garcia HM, McFadden EP, Farb A, Mehran R, Stone GW, Spertus J, Onuma Y, Morel M-A, van Es G-A, Zuckerman B, et al; Academic Research Consortium. Standardized end point definitions for coronary intervention trials: the Academic Research Consortium-2 consensus document. *Eur Heart J*. 2018;39:2192–2207. doi: 10.1093/eurheartj/ehy223
87. Armillotta M, Bergamaschi L, Paolisso P, Belmonte M, Angeli F, Sansonetti A, Stefanizzi A, Bertolini D, Bodega F, Amicone S, et al. Prognostic relevance of type 4a myocardial infarction and periprocedural myocardial injury in patients with non-ST-segment-elevation myocardial infarction. *Circulation*. 2025;151:760–772. doi: 10.1161/CIRCULATIONAHA.124.070729
88. Alam SR, Stirrat C, Spath N, Zamvar V, Pessotto R, Dweck MR, Moore C, Semple S, El-Medany A, Manoharan D, et al. Myocardial inflammation, injury and infarction during on-pump coronary artery bypass graft surgery. *J Cardiothorac Surg*. 2017;12:115. doi: 10.1186/s13019-017-0681-6
89. Omran H, Deutsch MA, Groezinger E, Zittermann A, Renner A, Neumann JT, Westermann D, Myles P, Ramosaj B, Pauly M, et al. High-sensitivity cardiac troponin I after coronary artery bypass grafting for post-operative decision-making. *Eur Heart J*. 2022;43:2388–2403. doi: 10.1093/eurheartj/ehab918
90. Spirito A, Sartori S, Koshy AN, Feng Y, Vogel B, Baber U, Sweeny J, Khera S, Kini AS, Windecker S, et al. Mortality after procedural or spontaneous myocardial infarction. *J Am Coll Cardiol*. 2024;84:467–477. doi: 10.1016/j.jacc.2024.04.061
91. Jangaard N, Sarkisian L, Saaby L, Mikkelsen S, Lassen AM, Marcussen N, Thomsen JL, Diederichsen ACP, Thygesen K, Mickley H. Incidence, frequency, and clinical characteristics of type 3 myocardial infarction in clinical practice. *Am J Med*. 2017;130:862.e9–862.e14. doi: 10.1016/j.amjmed.2016.12.034
92. Wagenveld IM, Blokker GM, Pezzato A, Wielopolski PA, Renken NS, von der Thüsen JH, Krestin GP, Hunink MGM, Oosterhuis JW, Weustink AC. Diagnostic accuracy of postmortem computed tomography, magnetic resonance imaging, and computed tomography-guided biopsies for the detection of ischaemic heart disease in a hospital setting. *Eur Heart J Cardiovasc Imaging*. 2018;19:739–748. doi: 10.1093/ehjci/jeu015
93. Stassi C, Mondello C, Baldino G, Cardia L, Gualniera P, Calapai F, Sapienza D, Asmundo A, Ventura Spagnolo E. State of the art on the role of postmortem computed tomography angiography and magnetic resonance imaging in the diagnosis of cardiac causes of death: a narrative review. *Tomography*. 2022;8:961–973. doi: 10.3390/tomography8020077
94. Michaud K, Jacobsen C, Basso C, Banner J, Blokker BM, de Boer HH, Dedout F, O'Donnell C, Giordano C, Magnin V, et al. Application of postmortem imaging modalities in cases of sudden death due to cardiovascular diseases-current achievements and limitations from a pathology perspective. *Virchows Arch*. 2023;482:385–406. doi: 10.1007/s00428-022-03458-6
95. Cau R, Saba L, Balestrieri A, Meloni A, Mannelli L, La Grutta L, Bossone E, Mantini C, Politi C, Suri JS, et al. Photon-counting computed tomography in atherosclerotic plaque characterization. *Diagnostics*. 2024;14:1065. doi: 10.3390/diagnostics14111065
96. Kotronias RA, de Maria GL, Xie C, Thomas S, Chan K, Portolan L, Langrish JP, Walsh J, Cahill TJ, Lucking AJ, et al; OxAMI and ORFAN Investigators. Benchmarking photon-counting computed tomography angiography against invasive assessment of coronary stenosis: implications for severely calcified coronaries. *JACC Cardiovasc Imaging*. 2025;18:572–585. doi: 10.1016/j.jcmg.2024.11.005
97. Gascho D, von Allmen A, Landsmann A, Hunermond T, Tappero C, Thali MJ, Deininger-Czermak E. Diagnostic value of T1- and T2-weighted 3-Tesla MRI for postmortem detection and age stage classification of myocardial infarction. *Forensic Sci Med Pathol*. 2024;20:14–22. doi: 10.1007/s12024-023-00592-8
98. Delewi R, Ijff G, van de Hoef TP, Hirsch A, Robbers LF, Nijveldt R, van der Laan AM, van der Vleuten PA, Lucas C, Tijssen JGP, et al. Pathological Q waves in myocardial infarction in patients treated by primary PCI. *JACC Cardiovasc Imaging*. 2013;6:324–331. doi: 10.1016/j.jcmg.2012.08.018
99. Antiochos P, Ge Y, Steel K, Bingham S, Abdullah S, Mikolich JR, Arai AE, Bandettini WP, Patel AR, Farzaneh-Far A, et al; SPINS Study Investigators. Imaging of clinically unrecognized myocardial fibrosis in patients with suspected coronary artery disease. *J Am Coll Cardiol*. 2020;76:945–957. doi: 10.1016/j.jacc.2020.06.063
100. Boland LL, Folsom AR, Sorlie PD, Taylor HA, Rosamond WD, Chambless LE, Cooper LS. Occurrence of unrecognized myocardial infarction in subjects aged 45 to 65 years (the ARIC study). *Am J Cardiol*. 2002;90:927–931. doi: 10.1016/s0002-9149(02)02655-3
101. Kwong RY, Chan AK, Brown KA, Chan CW, Reynolds HG, Tsang S, Davis RB. Impact of unrecognized myocardial scar detected by cardiac magnetic resonance imaging on event-free survival in patients presenting with signs or symptoms of coronary artery disease. *Circulation*. 2006;113:2733–2743. doi: 10.1161/CIRCULATIONAHA.105.570648
102. Kwong RY, Sattar H, Wu H, Vorobiof G, Gandla V, Steel K, Siu S, Brown KA. Incidence and prognostic implication of unrecognized myocardial scar characterized by cardiac magnetic resonance in diabetic patients without clinical evidence of myocardial infarction. *Circulation*. 2008;118:1011–1020. doi: 10.1161/CIRCULATIONAHA.107.727826
103. Barbier CE, Bjerner T, Johansson L, Lind L, Ahlstrom H. Myocardial scars more frequent than expected: magnetic resonance imaging detects potential risk group. *J Am Coll Cardiol*. 2006;48:765–771. doi: 10.1016/j.jacc.2006.05.041
104. Fernandez-Friera L, Garcia-Alvarez A, Oliva B, Garcia-Lunar I, Garcia I, Moreno-Arciniegas A, Gómez-Talavera S, Pérez-Herreras C, Sánchez-González J, de Vega VM, et al. Association between subclinical atherosclerosis burden and unrecognized myocardial infarction detected by cardiac magnetic resonance in middle-aged low-risk adults. *Eur Heart J Cardiovasc Imaging*. 2024;25:968–975. doi: 10.1093/ehjci/jeae044
105. Hammar P, Nordenskjöld AM, Lindahl B, Duvernoy O, Ahlstrom H, Johansson L, Hadziosmanovic N, Bjerner T. Unrecognized myocardial infarctions assessed by cardiovascular magnetic resonance are associated with the severity of the stenosis in the supplying coronary artery. *J Cardiovasc Magn Reson*. 2015;17:98. doi: 10.1186/s12968-015-0202-5
106. Nordenskjöld AM, Hammar P, Ahlstrom H, Bjerner T, Duvernoy O, Lindahl B. Unrecognized myocardial infarction assessed by cardiac magnetic resonance imaging is associated with adverse long-term prognosis. *PLoS One*. 2018;13:e0200381. doi: 10.1371/journal.pone.0200381
107. Braunwald E. Unstable angina: a classification. *Circulation*. 1989;80:410–414. doi: 10.1161/01.cir.80.2.410
108. Gerhardt W, Katus H, Ravkilde J, Hamm C, Jorgensen FJ, Peheim E, Ljungdahl L, Löfdahl P. S-troponin T in suspected ischemic myocardial injury compared with mass and catalytic concentrations of S-creatinine kinase isoenzyme MB. *Clin Chem*. 1991;37:1405–1411. doi: 10.1093/clinchem/37.8.1405
109. Diderholm E, Andren B, Frostfeldt G, Genberg M, Jernberg T, Lagerqvist B, Lindahl B, Venge P, Wallentin L; Fast Revascularisation during InStability in Coronary artery disease (FRISC II) Investigators. The prognostic and therapeutic implications of increased troponin T levels and ST depression in unstable coronary artery disease: the FRISC II invasive troponin

- T electrocardiogram substudy. *Am Heart J*. 2002;143:760–767. doi: 10.1067/mhj.2002.121733
110. Lindahl B, Toss H, Siegbahn A, Venge P, Wallentin L; for the FRISC Study Group. Markers of myocardial damage and inflammation in relation to long-term mortality in unstable coronary artery disease: FRISC Study Group: Fragmin During Instability in Coronary Artery Disease. *N Engl J Med*. 2000;343:1139–1147. doi: 10.1056/NEJM200010193431602
 111. Morrow DA, Antman EM, Tanasijevic M, Rifai N, de Lemos JA, McCabe CH, Cannon CP, Braunwald E. Cardiac troponin I for stratification of early outcomes and the efficacy of enoxaparin in unstable angina: a TIMI-11B substudy. *J Am Coll Cardiol*. 2000;36:1812–1817. doi: 10.1016/s0735-1097(00)00942-6
 112. Braunwald E, Morrow DA. Unstable angina: is it time for a requiem? *Circulation*. 2013;127:2452–2457. doi: 10.1161/CIRCULATIONAHA.113.001258
 113. Eggers KM, Jernberg T, Lindahl B. Unstable angina in the era of cardiac troponin assays with improved sensitivity: a clinical dilemma. *Am J Med*. 2017;130:1423–1430.e5. doi: 10.1016/j.amjmed.2017.05.037
 114. Agewall S, Beltrame JF, Reynolds HR, Niessner A, Rosano G, Caforio AL, De Caterina R, Zimarino M, Roffi M, Kjeldsen S, et al; WG on Cardiovascular Pharmacotherapy. ESC working group position paper on myocardial infarction with non-obstructive coronary arteries. *Eur Heart J*. 2017;38:143–153. doi: 10.1093/eurheartj/ehw149
 115. Collet JP, Thiele H, Barabato E, Barthelémy O, Bauersachs J, Bhatt DL, Dendale P, Dorobantu M, Edvardsson T, Folliquet T, et al; ESC Scientific Document Group. 2020 ESC guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J*. 2021;42:1289–1367. doi: 10.1093/eurheartj/ehaa575
 116. Heitner JF, Senthikumar A, Harrison JK, Klem I, Sketch MH Jr, Ivanov A, Hamo C, Van Assche L, White J, Washam J, et al. Identifying the infarct-related artery in patients with non-ST-segment-elevation myocardial infarction. *Circ Cardiovasc Interv*. 2019;12:e007305. doi: 10.1161/CIRCINTERVENTIONS.118.007305
 117. Reynolds HR, Maehara A, Kwong RY, Sedlak T, Saw J, Smilowitz NR, Mahmud E, Wei J, Marzo K, Matsumura M, et al. Coronary optical coherence tomography and cardiac magnetic resonance imaging to determine underlying causes of myocardial infarction with nonobstructive coronary arteries in women. *Circulation*. 2021;143:624–640. doi: 10.1161/CIRCULATIONAHA.120.052008
 118. Liang K, Bisaccia G, Leo I, Williams MGL, Dastidar A, Strange JW, Sammut E, Johnson TW, Bucciarelli-Ducci C. CMR reclassifies the majority of patients with suspected MINOCA and non MINOCA. *Eur Heart J Cardiovasc Imaging*. 2023;25:8–15. doi: 10.1093/ehjci/jead182
 119. Mileva N, Paolisso P, Gallinoro E, Fabbicatore D, Munhoz D, Bergamaschi L, Belmonte M, Panayotov P, Pizzi C, Barabato E, et al. Diagnostic and prognostic role of cardiac magnetic resonance in MINOCA: systematic review and meta-analysis. *JACC Cardiovasc Imaging*. 2023;16:376–389. doi: 10.1016/j.jcmg.2022.10.029
 120. Rajwani A, Giudicatti L, Telyuk P, Maredia N, Ithayhid A, Chieng D, Pasupathy S, Beltrame J, McQuillan B, Spiro J, et al. Clinical impact of cardiac magnetic resonance imaging in myocardial infarction with non-obstructive coronary arteries: a prospective multicentre cohort study. *Heart*. 2025;112:95–102. doi: 10.1136/heartjnl-2024-325181
 121. Williams MGL, Liang K, De Garate E, Spagnoli L, Fiori E, Dastidar A, Benedetto U, Biglino G, Johnson TW, Luscher T, et al. Peak troponin and CMR to guide management in suspected ACS and nonobstructive coronary arteries. *JACC Cardiovasc Imaging*. 2022;15:1578–1587. doi: 10.1016/j.jcmg.2022.03.017
 122. Schulz-Menger J, Collini V, Groschel J, Adler Y, Brucato A, Christian V, Ferreira VM, Gandjbakhch E, Heidecker B, Kerneis M, et al; ESC Scientific Document Group. 2025 ESC guidelines for the management of myocarditis and pericarditis. *Eur Heart J*. 2025;46:3952–4041. doi: 10.1093/eurheartj/ehaf192
 123. Dastidar AG, Baritussio A, De Garate E, Drobni Z, Biglino G, Singhal P, Milano EG, Angelini GD, Dorman S, Strange J, et al. Prognostic role of CMR and conventional risk factors in myocardial infarction with nonobstructed coronary arteries. *JACC Cardiovasc Imaging*. 2019;12:1973–1982. doi: 10.1016/j.jcmg.2018.12.023
 124. Anderson HVS, Masri SC, Abdallah MS, Chang AM, Cohen MG, Elgendy IY, Gulati M, LaPoint K, Madan N, Mousa ID, et al. 2022 ACC/AHA key data elements and definitions for chest pain and acute myocardial infarction: a report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Data Standards. *Circ Cardiovasc Qual Outcomes*. 2022;15:e000112. doi: 10.1161/HCQ.0000000000000112
 125. Angerud KH, Sederholm Lawesson S, Isaksson RM, Thülen I, Swahn E. Differences in symptoms, first medical contact and pre-hospital delay times between patients with ST- and non-ST-elevation myocardial infarction. *Eur Heart J Acute Cardiovasc Care*. 2019;8:201–207. doi: 10.1177/2048872617741734
 126. Lichtman JH, Leifheit EC, Safdar B, Bao H, Krumholz HM, Lorenze NP, Daneshvar M, Sertus JA, D'Onofrio G. Sex differences in the presentation and perception of symptoms among young patients with myocardial infarction: evidence from the VIRGO study (Variation in Recovery: Role of Gender on Outcomes of Young AMI Patients). *Circulation*. 2018;137:781–790. doi: 10.1161/CIRCULATIONAHA.117.031650
 127. Devon HA, Rosenfeld A, Steffen AD, Daya M. Sensitivity, specificity, and sex differences in symptoms reported on the 13-item acute coronary syndrome checklist. *J Am Heart Assoc*. 2014;3:e000586. doi: 10.1161/JAHA.113.000586
 128. Ferry AV, Anand A, Strachan FE, Mooney L, Stewart SD, Marshall L, Chapman AR, Lee KK, Jones S, Orme K, et al. Presenting symptoms in men and women diagnosed with myocardial infarction using sex-specific criteria. *J Am Heart Assoc*. 2019;8:e012307. doi: 10.1161/JAHA.119.012307
 129. Brieger D, Eagle KA, Goodman SG, Steg PG, Budaj A, White K, Montalescot G; GRACE Investigators. Acute coronary syndromes without chest pain, an underdiagnosed and undertreated high-risk group: insights from the Global Registry of Acute Coronary Events. *Chest*. 2004;126:461–469. doi: 10.1378/chest.126.2.461
 130. Canto AJ, Kiefe CI, Goldberg RJ, Rogers WJ, Peterson ED, Wenger NK, Vaccarino V, Frederick PD, Sopko G, Zheng Z-J, et al. Differences in symptom presentation and hospital mortality according to type of acute myocardial infarction. *Am Heart J*. 2012;163:572–579. doi: 10.1016/j.ahj.2012.01.020
 131. Galinski M, Saget D, Ruscev M, Gonzalez G, Ameur L, Lapostolle F, Adnet F. Chest pain in an out-of-hospital emergency setting: no relationship between pain severity and diagnosis of acute myocardial infarction. *Pain Pract*. 2015;15:343–347. doi: 10.1111/papr.12178
 132. Khan IA, Karim HMR, Panda CK, Ahmed G, Nayak S. Atypical presentations of myocardial infarction: a systematic review of case reports. *Cureus*. 2023;15:e35492. doi: 10.7759/cureus.35492
 133. Kirchberger I, Meisinger C, Heier M, Kling B, Wende R, Greschik C, von Scheidt W, Kuch B. Patient-reported symptoms in acute myocardial infarction: differences related to ST-segment elevation: the MONICA/KORA Myocardial Infarction Registry. *J Intern Med*. 2011;270:58–64. doi: 10.1111/j.1365-2796.2011.02365.x
 134. Stehli J, Martin C, Brennan A, Dinh DT, Lefkowitz J, Zaman S. Sex differences persist in time to presentation, revascularization, and mortality in myocardial infarction treated with percutaneous coronary intervention. *J Am Heart Assoc*. 2019;8:e012161. doi: 10.1161/JAHA.119.012161
 135. van Oosterhout REM, de Boer AR, Maas A, Rutten FH, Bots ML, Peters SAE. Sex differences in symptom presentation in acute coronary syndromes: a systematic review and meta-analysis. *J Am Heart Assoc*. 2020;9:e014733. doi: 10.1161/JAHA.119.014733
 136. DeVon HA, Vuckovic K, Burke LA, Mirzaei S, Breen K, Robinson N, Zegre-Hemsey J. What's the risk? Older women report fewer symptoms for suspected acute coronary syndrome than younger women. *Biores Open Access*. 2018;7:131–138. doi: 10.1089/biores.2018.0020
 137. Bjorck L, Nielsen S, Jernberg T, Zverkova-Sandstrom T, Giang KW, Rosengren A. Absence of chest pain and long-term mortality in patients with acute myocardial infarction. *Open Heart*. 2018;5:e000909. doi: 10.1136/openhrt-2018-000909
 138. Kumar A, Sanghera A, Sanghera B, Mohamed T, Midgen A, Pattison S, Marston L, Jones MM. Chest pain symptoms during myocardial infarction in patients with and without diabetes: a systematic review and meta-analysis. *Heart*. 2023;109:1516–1524. doi: 10.1136/heartjnl-2022-322289
 139. Nanna MG, Hajduk AM, Krumholz HM, Murphy TE, Dreyer RP, Alexander KP, Geda M, Tsang S, Welty FK, Safdar B, et al. Sex-based differences in presentation, treatment, and complications among older adults hospitalized for acute myocardial infarction: the SILVER-AMI study. *Circ Cardiovasc Qual Outcomes*. 2019;12:e005691. doi: 10.1161/CIRCOUTCOMES.119.005691
 140. Øhrn AM, Nielsen CS, Schirmer H, Stubhaug A, Wilsaard T, Lindekleiv H. Pain tolerance in persons with recognized and unrecognized myocardial infarction: a population-based, cross-sectional study. *J Am Heart Assoc*. 2016;5:e003846. doi: 10.1161/JAHA.116.003846
 141. DeVon HA, Burke LA, Nelson H, Zerwic JJ, Riley B. Disparities in patients presenting to the emergency department with potential acute coronary

- syndrome: it matters if you are Black or White. *Heart Lung*. 2014;43:270–277. doi: 10.1016/j.hrtlung.2014.04.019
142. King-Shier K, Quan H, Kapral MK, Tsuyuki R, An L, Banerjee S, Southern DA, Khan N. Acute coronary syndromes presentations and care outcomes in white, South Asian and Chinese patients: a cohort study. *BMJ Open*. 2019;9:e022479. doi: 10.1136/bmjopen-2018-022479
 143. Jinatongthai P, Kongwatcharapong J, Foo CY, Phrommintikul A, Nathisuwan S, Thakkinian A, Reid CM, Chaiyakunapruk N. Comparative efficacy and safety of reperfusion therapy with fibrinolytic agents in patients with ST-segment elevation myocardial infarction: a systematic review and network meta-analysis. *Lancet*. 2017;390:747–759. doi: 10.1016/S0140-6736(17)31441-1
 144. Widimsky P, Rohac F, Stasek J, Kala P, Rokyta R, Kuzmanov B, Jakl M, Poloczek M, Kanovsky J, Bernat I, et al. Primary angioplasty in acute myocardial infarction with right bundle branch block: should new onset right bundle branch block be added to future guidelines as an indication for reperfusion therapy? *Eur Heart J*. 2012;33:86–95. doi: 10.1093/eurheartj/ehr291
 145. Sgarbossa EB, Pinski SL, Barbagelata A, Underwood DA, Gates KB, Topol EJ, Califf RM, Wagner GS. Electrocardiographic diagnosis of evolving acute myocardial infarction in the presence of left bundle-branch block. GUSTO-1 (Global Utilization of Streptokinase and Tissue Plasminogen Activator for Occluded Coronary Arteries) Investigators. *N Engl J Med*. 1996;334:481–487. doi: 10.1056/NEJM19960223340801
 146. Smith SW, Dodd KW, Henry TD, Dvorak DM, Pearce LA. Diagnosis of ST-elevation myocardial infarction in the presence of left bundle branch block with the ST-elevation to S-wave ratio in a modified Sgarbossa rule. *Ann Emerg Med*. 2012;60:766–776. doi: 10.1016/j.annemergmed.2012.07.119
 147. de Winter RJ, Verouden NJ, Wellens HJ, Wilde AA; Interventional Cardiology Group of the Academic Medical Center. A new ECG sign of proximal LAD occlusion. *N Engl J Med*. 2008;359:2071–2073. doi: 10.1056/NEJMc0804737
 148. de Zwaan C, Bär FW, Wellens HJ. Characteristic electrocardiographic pattern indicating a critical stenosis high in left anterior descending coronary artery in patients admitted because of impending myocardial infarction. *Am Heart J*. 1982;103:730–736. doi: 10.1016/0002-8703(82)90480-x
 149. Rossello X, Wiegnerinck RF, Alguesuari J, Bardaji A, Wörner F, Sutil M, Ferrero A, Cinca J. New electrocardiographic criteria to differentiate acute pericarditis and myocardial infarction. *Am J Med*. 2014;127:233–239. doi: 10.1016/j.amjmed.2013.11.006
 150. Raitt MH, Maynard C, Wagner GS, Cerqueira MD, Selvester RH, Weaver WD. Appearance of abnormal Q waves early in the course of acute myocardial infarction: implications for efficacy of thrombolytic therapy. *J Am Coll Cardiol*. 1995;25:1084–1088. doi: 10.1016/0735-1097(94)00514-q
 151. Krone RJ, Friedman E, Thanavaro S, Miller JP, Kleiger RE, Oliver GC. Long-term prognosis after first Q-wave (transmural) or non-Q-wave (non-transmural) myocardial infarction: analysis of 593 patients. *Am J Cardiol*. 1983;52:234–239. doi: 10.1016/0002-9149(83)90114-5
 152. Westermann D, Neumann JT, Sörensen NA, Blankenberg S. High-sensitivity assays for troponin in patients with cardiac disease. *Nat Rev Cardiol*. 2017;14:472–483. doi: 10.1038/nrcardio.2017.48
 153. Vylegzhanina AV, Kogan AE, Katrukha IA, Koshkina EV, Bereznikova AV, Filatov VL, Bloshchitsyna MN, Bogomolova AP, Katrukha AG. Full-size and partially truncated cardiac troponin complexes in the blood of patients with acute myocardial infarction. *Clin Chem*. 2019;65:882–892. doi: 10.1373/clinchem.2018.301127
 154. Kristensen JH, Hasselbalch RB, Strandkjaer N, Jorgensen N, Ostergaard M, Møller-Sørensen PH, Christian Nilsson J, Afzal S, Rørbæk Kamstrup P, Dahl M, et al. Half-life and clearance of cardiac troponin I and troponin T in humans. *Circulation*. 2024;150:1187–1198. doi: 10.1161/CIRCULATIONAHA.123.066565
 155. International Federation of Clinical Chemistry and Laboratory Medicine. *Biomarkers Reference Tables. High Sensitivity Cardiac Troponin I and T Assay Analytical Characteristics*. 2026. Accessed April 13, 2026. <https://ifcc.org/ifcc-education-division/emd-committees/committee-on-clinical-applications-of-cardiac-bio-markers-c-cb/biomarkers-reference-tables>
 156. Apple FS, Collinson PO; IFCC Task Force on Clinical Applications of Cardiac Biomarkers. Analytical characteristics of high-sensitivity cardiac troponin assays. *Clin Chem*. 2012;58:54–61. doi: 10.1373/clinchem.2011.165795
 157. Collinson P, Hammerer-Lercher A, Aakre KM, Gruson D, Suvisaari J, Pulkki K, Stankovic S, Baum H, Lowry MT, Mills NL, et al. Implementation of high sensitivity troponin into routine clinical practice: results of the extended CARDiac MARKers guideline uptake in Europe group (CARMAGUE) survey. *Clin Chim Acta*. 2024;558:117900. doi: 10.1016/j.cca.2024.117900
 158. Aakre KM, Saenger AK, Body R, Collinson P, Hammarsten O, Jaffe AS, Kavsak P, Omland T, Ordonez-Lianos J, Apple FS. Analytical considerations in deriving 99th percentile upper reference limits for high-sensitivity cardiac troponin assays: educational recommendations from the IFCC Committee on Clinical Application of Cardiac Bio-Markers. *Clin Chem*. 2022;68:1022–1030. doi: 10.1093/clinchem/hvac092
 159. McEvoy JW, Tang O, Wang D, Ndumele CE, Coresh J, Christenson RH, Selvin E. Myocardial injury thresholds for 4 high-sensitivity troponin assays in U.S. adults. *J Am Coll Cardiol*. 2023;81:2028–2039. doi: 10.1016/j.jacc.2023.03.403
 160. Welsh P, Preiss D, Shah ASV, McAllister D, Briggs A, Boachie C, McConnachie A, Hayward C, Padmanabhan S, Welsh C, et al. Comparison between high-sensitivity cardiac troponin T and cardiac troponin I in a large general population cohort. *Clin Chem*. 2018;64:1607–1616. doi: 10.1373/clinchem.2018.292086
 161. Cao M, Pierce AE, Norman MS, Thakur B, Diercks K, Hale C, Issioui Y, Diercks DB. Systematic review of sex-specific high sensitivity cardiac troponin I and T thresholds. *Clin Ther*. 2024;46:988–994. doi: 10.1016/j.clinthera.2024.09.025
 162. Kimenai DM, Janssen E, Eggers KM, Lindahl B, den Ruijter HM, Bekers O, Appelman Y, Meex SJR. Sex-specific versus overall clinical decision limits for cardiac troponin I and T for the diagnosis of acute myocardial infarction: a systematic review. *Clin Chem*. 2018;64:1034–1043. doi: 10.1373/clinchem.2018.286781
 163. Shah AS, Griffiths M, Lee KK, McAllister DA, Hunter AL, Ferry AV, Cruikshank A, Reid A, Stoddart M, Strachan F, et al. High sensitivity cardiac troponin and the under-diagnosis of myocardial infarction in women: prospective cohort study. *BMJ*. 2015;350:g7873. doi: 10.1136/bmj.g7873
 164. Rubini Gimenez M, Twerenbold R, Boeddinghaus J, Nestelberger T, Puelacher C, Hillinger P, Wildi K, Jaeger C, Grimm K, Heitzelmann K-F, et al. Clinical effect of sex-specific cutoff values of high-sensitivity cardiac troponin T in suspected myocardial infarction. *JAMA Cardiol*. 2016;1:912–920. doi: 10.1001/jamacardio.2016.2882
 165. Daniels LB, Mueller C, Giannitsis E, Meex SJR, Buehlmann D, Dilba P, Bendig G, Cole M, Body R, Christenson RH, et al; TSIX Investigators. Establishing reference values in healthy participants for the cardiac troponin T high-sensitivity Gen 6 Assay: REF-TSIX global reference study. *Clin Chem*. 2026;72:488–502. doi: 10.1093/clinchem/hvag011
 166. de Bakker M, Anand A, Shipley M, Fujisawa T, Shah ASV, Kardys I, Boersma E, Brunner EJ, Mills NL, Kimenai DM. Sex differences in cardiac troponin trajectories over the life course. *Circulation*. 2023;147:1798–1808. doi: 10.1161/CIRCULATIONAHA.123.064386
 167. Lowry MTH, Doudesis D, Wereski R, Kimenai DM, Tuck C, Ferry AV, Bularga A, Taggart C, Lee K, Chapman AR, et al; High-STEACS Investigators. Influence of age on the diagnosis of myocardial infarction. *Circulation*. 2022;146:1135–1148. doi: 10.1161/CIRCULATIONAHA.122.059994
 168. Doudesis D, Lee KK, Boeddinghaus J, Bularga A, Ferry AV, Tuck C, Lowry MTH, Lopez-Ayala P, Nestelberger T, Koehlin L, et al; CoDE-ACS Investigators. Machine learning for diagnosis of myocardial infarction using cardiac troponin concentrations. *Nat Med*. 2023;29:1201–1210. doi: 10.1038/s41591-023-02325-4
 169. Hammarsten O, Warner JV, Lam L, Kavsak P, Lindahl B, Aakre KM, Collinson P, Jaffe AS, Saenger AK, Body R, et al. Antibody-mediated interferences affecting cardiac troponin assays: recommendations from the IFCC Committee on Clinical Applications of Cardiac Biomarkers. *Clin Chem Lab Med*. 2023;61:1411–1419. doi: 10.1515/cclm-2023-0028
 170. Harley K, Bissonnette S, Inzitari R, Schulz K, Apple FS, Kavsak PA, Gunsolus IL. Independent and combined effects of biotin and hemolysis on high-sensitivity cardiac troponin assays. *Clin Chem Lab Med*. 2021;59:1431–1443. doi: 10.1515/cclm-2021-0124
 171. Herman DS, Kavsak PA, Greene DN. Variability and error in cardiac troponin testing: an ACLPS critical review. *Am J Clin Pathol*. 2017;148:281–295. doi: 10.1093/ajcp/aaq066
 172. Christenson RH, Frenk LDS, de Graaf HJ, van Domburg TSY, Wijnands FPG, Foolen HWJ, Kemper DWM, Bruinen AL, Meijering BDM, Fonville JM, et al. Point-of-care: roadmap for analytical characterization and validation of a high-sensitivity cardiac troponin I assay in plasma and whole blood matrices. *J Appl Lab Med*. 2022;7:971–988. doi: 10.1093/jalm/jfac028
 173. Chaulin AM. False-positive causes in serum cardiac troponin levels. *J Clin Med Res*. 2022;14:80–87. doi: 10.14740/jocmr4664
 174. Salaun E, Drory S, Cote MA, Tremblay V, Bedard E, Steinberg C, Paré D, O'Connor K, Cieza T, Coté N, et al. Role of antitroponin antibodies and macrotroponin in the clinical interpretation of cardiac troponin. *J Am Heart Assoc*. 2024;13:e035128. doi: 10.1161/JAHA.123.035128

175. Mair J, Giannitsis E, Mills NL, Mueller C; Study Group on Biomarkers of the European Society of Cardiology Association for Acute CardioVascular Care. How to deal with unexpected cardiac troponin results. *Eur Heart J Acute Cardiovasc Care*. 2022;11:e1–e3. doi: 10.1093/ehjacc/zuac023
176. Bularga A, Oskoui E, Fujisawa T, Jenks S, Sutherland R, Apple FS, Hammarsten O, Mills NL. Macrotrponin complex as a cause for cardiac troponin increase after COVID-19 vaccination and infection. *Clin Chem*. 2022;68:1015–1019. doi: 10.1093/clinchem/hvac100
177. Lam L, Tse R, Gladding P, Kyle C. Effect of macrotrponin in a cohort of community patients with elevated cardiac troponin. *Clin Chem*. 2022;68:1261–1271. doi: 10.1093/clinchem/hvac118
178. Jaffe AS, Body R, Mills NL, Aakre KM, Collinson PO, Saenger A, Hammarsten O, Wereski R, Omland T, Sandoval Y, et al; IFCC Committee on Cardiac Bio-Markers. Single troponin measurement to rule out myocardial infarction: JACC review topic of the week. *J Am Coll Cardiol*. 2023;82:60–69. doi: 10.1016/j.jacc.2023.04.040
179. Hoeller R, Rubini Gimenez M, Reichlin T, Twerenbold R, Zellweger C, Moehring B, Wildi K, Freese M, Stelzig C, Hartmann B, et al. Normal presenting levels of high-sensitivity troponin and myocardial infarction. *Heart*. 2013;99:1567–1572. doi: 10.1136/heartjnl-2013-303643
180. Lowry MTH, Doudesis D, Boeddinghaus J, Kimenai DM, Bularga A, Taggart C, Wereski R, Ferry AV, Stewart SD, Tuck C, et al; High-STEACS Investigators. Troponin in early presenters to rule out myocardial infarction. *Eur Heart J*. 2023;44:2846–2858. doi: 10.1093/eurheartj/ehad376
181. Wereski R, Chapman AR, Lee KK, Smith SW, Lowe DJ, Gray A, Mills NL. High-sensitivity cardiac troponin concentrations at presentation in patients with ST-segment elevation myocardial infarction. *JAMA Cardiol*. 2020;5:1302–1304. doi: 10.1001/jamacardio.2020.2867
182. Liu T, Howarth AG, Chen Y, Nair AR, Yang HJ, Ren D, Tang R, Sykes J, Kovacs MS, Dey D, et al. Intramyocardial hemorrhage and the "wave front" of reperfusion injury compromising myocardial salvage. *J Am Coll Cardiol*. 2022;79:35–48. doi: 10.1016/j.jacc.2021.10.034
183. Vora KP, Kalra A, Shah CD, Bhatt K, Kumar A, Pandya T, Poptani V, Chan SF, Singh D, Jambunathan N, et al. In-hospital mortality in hemorrhagic myocardial infarction. *NEJM Evid*. 2025;4:EVIDoa2400294. doi: 10.1056/EVIDoa2400294
184. Silvain J, Zeitouni M, Paradies V, Zheng HL, Ndrepepa G, Cavallini C, Feldman DN, Sharma SK, Mehili J, Gili S, et al. Procedural myocardial injury, infarction and mortality in patients undergoing elective PCI: a pooled analysis of patient-level data. *Eur Heart J*. 2021;42:323–334. doi: 10.1093/eurheartj/ehaa885
185. Miller WL, Garratt KN, Burritt MF, Reeder GS, Jaffe AS. Timing of peak troponin T and creatine kinase-MB elevations after percutaneous coronary intervention. *Chest*. 2004;125:275–280. doi: 10.1378/chest.125.1.275
186. Miller WL, Garratt KN, Burritt MF, Lennon RJ, Reeder GS, Jaffe AS. Baseline troponin level: key to understanding the importance of post-PCI troponin elevations. *Eur Heart J*. 2006;27:1061–1069. doi: 10.1093/eurheartj/ehi760
187. Selvanayagam JB, Porto I, Channon K, Petersen SE, Francis JM, Neubauer S, Banning AP. Troponin elevation after percutaneous coronary intervention directly represents the extent of irreversible myocardial injury: insights from cardiovascular magnetic resonance imaging. *Circulation*. 2005;111:1027–1032. doi: 10.1161/01.CIR.0000156328.28485.AD
188. Eggers KM, Batra G, Lindahl B, Ghukasyan Lakic T, Lindback J, Budaj A, Cornel JH, Giannitsis E, Katus HA, Storey RF, et al. Temporal biomarker concentration patterns during the early course of acute coronary syndrome. *Clin Chem Lab Med*. 2024;62:1167–1176. doi: 10.1515/cclm-2023-1253
189. Jaffe AS, Moeckel M, Giannitsis E, Huber K, Mair J, Mueller C, Plebani M, Thygesen K, Lindahl B. In search for the Holy Grail: suggestions for studies to define delta changes to diagnose or exclude acute myocardial infarction: a position paper from the Study Group on Biomarkers of the Acute Cardiovascular Care Association. *Eur Heart J Acute Cardiovasc Care*. 2014;3:313–316. doi: 10.1177/2048872614541906
190. Reichlin T, Irfan A, Twerenbold R, Reiter M, Hochholzer W, Burkhalter H, Bassetti S, Steuer S, Winkler K, Peter F, et al. Utility of absolute and relative changes in cardiac troponin concentrations in the early diagnosis of acute myocardial infarction. *Circulation*. 2011;124:136–145. doi: 10.1161/CIRCULATIONAHA.111.023937
191. Hollander JE, Than M, Mueller C. State-of-the-art evaluation of emergency department patients presenting with potential acute coronary syndromes. *Circulation*. 2016;134:547–564. doi: 10.1161/CIRCULATIONAHA.116.021886
192. Lowry MTH, Anand A, Mills NL. Implementing an early rule-out pathway for acute myocardial infarction in clinical practice. *Heart*. 2021;107:1912–1919. doi: 10.1136/heartjnl-2019-316242
193. Kontos MC, de Lemos JA, Deitelzweig SB, Diercks DB, Gore MO, Hess EP, McCarthy CP, McCord JK, Musey Jr PL, Villines TC, et al; Writing Committee. 2022 ACC expert consensus decision pathway on the evaluation and disposition of acute chest pain in the emergency department: a report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol*. 2022;80:1925–60. doi: 10.1016/j.jacc.2022.08.750
194. Reichlin T, Schindler C, Drexler B, Twerenbold R, Reiter M, Zellweger C, Moehring B, Ziller R, Hoeller R, Rubini Gimenez M, et al. One-hour rule-out and rule-in of acute myocardial infarction using high-sensitivity cardiac troponin T. *Arch Intern Med*. 2012;172:1211–1218. doi: 10.1001/archinternmed.2012.3698
195. Baker JO, Tyther R, Liebetrau C, Clark J, Howarth R, Patterson T, Möllmann H, Nef H, Sicard P, Kailey B, et al. Cardiac myosin-binding protein C: a potential early biomarker of myocardial injury. *Basic Res Cardiol*. 2015;110:23. doi: 10.1007/s00395-015-0478-5
196. Kaier TE, Twerenbold R, Lopez-Ayala P, Nestelberger T, Boeddinghaus J, Alaour B, Huber I-M, Zhi Y, Koechlin L, Wussler D, et al; ASPACE Investigators. A 0/1h-algorithm using cardiac myosin-binding protein C for early diagnosis of myocardial infarction. *Eur Heart J Acute Cardiovasc Care*. 2022;11:325–335. doi: 10.1093/ehjacc/zuac007
197. Kaier TE, Twerenbold R, Puelacher C, Marjot J, Imambaccus N, Boeddinghaus J, Nestelberger T, Badertscher P, Sabti Z, Giménez MR, et al. Direct comparison of cardiac myosin-binding protein C with cardiac troponins for the early diagnosis of acute myocardial infarction. *Circulation*. 2017;136:1495–1508. doi: 10.1161/CIRCULATIONAHA.117.028084
198. Iqbal J, Sumaya W, Tatman V, Parviz Y, Morton AC, Grech ED, Campbell S, Storey RF, Gunn J. Incidence and predictors of stent thrombosis: a single-centre study of 5,833 consecutive patients undergoing coronary artery stenting. *EuroIntervention*. 2013;9:62–69. doi: 10.4244/EIJV9I1A10
199. Moussa ID, Mohanane D, Saucedo J, Stone GW, Yeh RW, Kennedy KF, Waksman R, Teirstein P, Moses JW, Simonson C. Trends and outcomes of restenosis after coronary stent implantation in the United States. *J Am Coll Cardiol*. 2020;76:1521–1531. doi: 10.1016/j.jacc.2020.08.002
200. Popovic B, Agrinier N, Bouchahda N, Pinelli S, Maigrat CH, Metzendorf PA, Selton Suty C, Juillière Y, Camenzind E. Coronary embolism among ST-segment-elevation myocardial infarction patients: mechanisms and management. *Circ Cardiovasc Interv*. 2018;11:e005587. doi: 10.1161/CIRCINTERVENTIONS.117.005587
201. Wang J, Gao H, Xiao J, Gao M, Liu Y, Gao J. The clinical features and prognosis of type 4C myocardial infarction in patients with non-ST-segment elevation myocardial infarction. *Ann Transl Med*. 2021;9:1153. doi: 10.21037/atm-21-2587
202. Balbi MM, Scarparo P, Tovar MN, Masdjedi K, Daemen J, Den Dekker W, Ligthart J, Witberg K, Cummins P, Wilschut J, et al. Culprit lesion detection in patients presenting with non-ST elevation acute coronary syndrome and multivessel disease. *Cardiovasc Revasc Med*. 2022;35:110–118. doi: 10.1016/j.carrev.2021.03.019
203. Kerensky RA, Wade M, Deedwania P, Boden WE, Pepine CJ; Veterans Affairs Non-Q-Wave Infarction Strategies In-Hospital (VANQWISH) Trial Investigators. Revisiting the culprit lesion in non-Q-wave myocardial infarction: Results from the VANQWISH trial angiographic core laboratory. *J Am Coll Cardiol*. 2002;39:1456–1463. doi: 10.1016/s0735-1097(02)01770-9
204. Andersson HB, Pedersen F, Engstrom T, Helqvist S, Jensen MK, Jorgensen E, Kelbæk H, Råder SBEW, Saunamäki K, Bates E, et al. Long-term survival and causes of death in patients with ST-elevation acute coronary syndrome without obstructive coronary artery disease. *Eur Heart J*. 2018;39:102–110. doi: 10.1093/eurheartj/ehx491
205. Gehrie ER, Reynolds HR, Chen AY, Neelon BH, Roe MT, Gibler WB, Ohman EM, Newby LK, Peterson ED, Hochman JS. Characterization and outcomes of women and men with non-ST-segment elevation myocardial infarction and nonobstructive coronary artery disease: results from the Can Rapid Risk Stratification of Unstable Angina Patients Suppress Adverse Outcomes With Early Implementation of the ACC/AHA Guidelines (CRUSADE) quality improvement initiative. *Am Heart J*. 2009;158:688–694. doi: 10.1016/j.ahj.2009.08.004
206. Larsen AI, Nilsen DW, Yu J, Mehran R, Nikolsky E, Lansky AJ, Caixeta A, Parise H, Fahy M, Cristea E, et al. Long-term prognosis of patients presenting with ST-segment elevation myocardial infarction with no significant coronary artery disease (from the HORIZONS-AMI trial). *Am J Cardiol*. 2013;111:643–648. doi: 10.1016/j.amjcard.2012.11.011
207. Johnson TW, Raber L, di Mario C, Bourantas C, Jia H, Mattesini A, Gonzalo N, de la Torre Hernandez JM, Prati F, Koskinas K, et al. Clinical use of intracoronary imaging, part 2: acute coronary syndromes, ambiguous coronary

- angiography findings, and guiding interventional decision-making: an expert consensus document of the European Association of Percutaneous Cardiovascular Interventions. *Eur Heart J*. 2019;40:2566–2584. doi: 10.1093/eurheartj/ehz332
208. De Bruyne B, Pijls NH, Kalesan B, Barbato E, Tonino PA, Piroth Z, Jagic N, Möbius-Winkler S, Rioufol G, Witt N, et al; FAME 2 Trial Investigators. Fractional flow reserve-guided PCI versus medical therapy in stable coronary disease. *N Engl J Med*. 2012;367:991–1001. doi: 10.1056/NEJMoa1205361
 209. Eftekhari A, Holck EN, Westra J, Olsen NT, Bruun NH, Jensen LO, Engstrøm T, Christiansen EH. Instantaneous wave free ratio vs. fractional flow reserve and 5-year mortality: iFR SWEDEHEART and DEFINE FLAIR. *Eur Heart J*. 2023;44:4376–4384. doi: 10.1093/eurheartj/ehad582
 210. Masdjedi K, van Zandvoort LJC, Balbi MM, Gijzen FJH, Ligthart JMR, Rutten MCM, Lemmert ME, Wilschut JM, Diletti R, de Jaegere P, et al. Validation of a three-dimensional quantitative coronary angiography-based software to calculate fractional flow reserve: the FAST study. *EuroIntervention*. 2020;16:591–599. doi: 10.4244/EIJ-D-19-00466
 211. Kontos MC. Role of echocardiography in the emergency department for identifying patients with myocardial infarction and ischemia. *Echocardiography*. 1999;16:193–205. doi: 10.1111/j.1540-8175.1999.tb00804.x
 212. Rybicki FJ, Udelson JE, Peacock WF, Goldhaber SZ, Issebacher EM, Kazerooni E, Kontos MC, Litt H, Woodard PK. 2015 ACR/ACC/AHA/AATS/ACEP/ASNC/NACCI/SAEM/SCCT/SCMR/SCPC/SNMMI/STR/STS appropriate utilization of cardiovascular imaging in emergency department patients with chest pain: a joint document of the American College of Radiology Appropriateness Criteria Committee and the American College of Cardiology Appropriate Use Criteria Task Force. *J Am Coll Cardiol*. 2016;67:853–879. doi: 10.1016/j.jacc.2015.09.011
 213. Stillman AE, Oudkerk M, Bluemke D, Bremerich J, Esteves FP, Garcia EV, Gutberlet M, Hundley WG, Jerosch-Herold M, Kuipers D, et al; North American Society of Cardiovascular Imaging. Assessment of acute myocardial infarction: current status and recommendations from the North American Society for Cardiovascular Imaging and the European Society of Cardiac Radiology. *Int J Cardiovasc Imaging*. 2011;27:7–24. doi: 10.1007/s10554-010-9714-0
 214. Thune JJ, Kober L, Pfeffer MA, Skali H, Anavekar NS, Bourgoin M, Ghali JK, Arnold JMO, Velazquez EJ, Solomon SD. Comparison of regional versus global assessment of left ventricular function in patients with left ventricular dysfunction, heart failure, or both after myocardial infarction: the Valsartan in Acute Myocardial Infarction Echocardiographic Study. *J Am Soc Echocardiogr*. 2006;19:1462–1465. doi: 10.1016/j.echo.2006.05.028
 215. Kontos MC, Diercks DB, Kirk JD. Emergency department and office-based evaluation of patients with chest pain. *Mayo Clin Proc*. 2010;85:284–299. doi: 10.4065/mcp.2009.0560
 216. Lewis WR. Echocardiography in the evaluation of patients in chest pain units. *Cardiol Clin*. 2005;23:531–539, vii. doi: 10.1016/j.ccl.2005.08.009
 217. Scirica BM. Acute coronary syndrome: emerging tools for diagnosis and risk assessment. *J Am Coll Cardiol*. 2010;55:1403–1415. doi: 10.1016/j.jacc.2009.09.071
 218. Geers J, Balfour A, Molek P, Barron P, Botezatu S, Joshi SS, White A, Buchwald M, Everett R, McCarley J, et al. Systematic hand-held echocardiography in patients hospitalized with acute coronary syndrome. *Eur Heart J Cardiovasc Imaging*. 2024;25:1441–1450. doi: 10.1093/ehjci/jeae149
 219. Korosoglou G, Geiger B, Hansen A, Hardt SA, Giannitsis E, Selter C, Katus HA, Kuecherer H. Usefulness of real-time myocardial perfusion imaging to evaluate alterations of myocardial blood flow in patients with stable angina pectoris undergoing elective percutaneous coronary interventions. *Am J Cardiol*. 2005;96:885–891. doi: 10.1016/j.amjcard.2005.05.041
 220. Mair J, Jaffe A, Lindahl B, Mills N, Mockel M, Cullen L, Giannitsis E, Hammarsten O, Huber K, Krychtiuk K, et al. The clinical approach to diagnosing peri-procedural myocardial infarction after percutaneous coronary interventions according to the fourth universal definition of myocardial infarction: from the Study Group on Biomarkers of the European Society of Cardiology (ESC) Association for Acute CardioVascular Care (ACVC). *Biomarkers*. 2022;27:407–417. doi: 10.1080/1354750X.2022.2055792
 221. Doolub G, Khurshid S, Theriault-Lauzier P, Nolin Lapalme A, Tastet O, So D, Labrecque Langlais E, Cobin D, Avram R. Revolutionising acute cardiac care with artificial intelligence: opportunities and challenges. *Can J Cardiol*. 2024;40:1813–1827. doi: 10.1016/j.cjca.2024.06.011
 222. Kusunose K, Abe T, Haga A, Fukuda D, Yamada H, Harada M, Sata M. A deep learning approach for assessment of regional wall motion abnormality from echocardiographic images. *JACC Cardiovasc Imaging*. 2020;13:374–381. doi: 10.1016/j.jcmg.2019.02.024
 223. Laumer F, Di Vece D, Cammann VL, Wurdinger M, Petkova V, Schonberger M, Schönberger A, Mercier JC, Niederseer D, Seifert B, et al. Assessment of artificial intelligence in echocardiography diagnostics in differentiating takotsubo syndrome from myocardial infarction. *JAMA Cardiol*. 2022;7:494–503. doi: 10.1001/jamacardio.2022.0183
 224. Upton R, Mumith A, Beqiri A, Parker A, Hawkes W, Gao S, Porumb M, Sarwar R, Marques P, Markham D, et al. Automated echocardiographic detection of severe coronary artery disease using artificial intelligence. *JACC Cardiovasc Imaging*. 2022;15:715–727. doi: 10.1016/j.jcmg.2021.10.013
 225. Ibanez B, Aletras AH, Arai AE, Arheden H, Bax J, Berry C, Bucciarelli-Ducci C, Croisille P, Dall'Armellina E, Dharmakumar R, et al. Cardiac MRI endpoints in myocardial infarction: experimental and clinical trials: JACC scientific expert panel. *J Am Coll Cardiol*. 2019;74:238–256. doi: 10.1016/j.jacc.2019.05.024
 226. Vora KP, Kumar A, Krishnam MS, Prato FS, Raman SV, Dharmakumar R. Microvascular obstruction and intramyocardial hemorrhage in reperfused myocardial infarctions: pathophysiology and clinical insights from imaging. *JACC Cardiovasc Imaging*. 2024;17:795–810. doi: 10.1016/j.jcmg.2024.02.003
 227. Rossello X, Lopez-Ayala P, Fernandez-Jimenez R, Oliver E, Galan-Arriola C, de Molina-Iracheta A, Agüero J, López GJ, Lobo-Gonzalez M, Vilchez-Tschischke JP, et al. R2 prime (R2') magnetic resonance imaging for post-myocardial infarction intramyocardial haemorrhage quantification. *Eur Heart J Cardiovasc Imaging*. 2020;21:1031–1038. doi: 10.1093/ehjci/jez306
 228. Abdel-Aty H, Zagrosek A, Schulz-Menger J, Taylor AJ, Messroghli D, Kumar A, Gross M, Dietz R, Friedrich MG. Delayed enhancement and T2-weighted cardiovascular magnetic resonance imaging differentiate acute from chronic myocardial infarction. *Circulation*. 2004;109:2411–2416. doi: 10.1161/01.CIR.0000127428.10985.C6
 229. Bulluck H, Rosmini S, Abdel-Gadir A, White SK, Bhuvana AN, Treibel TA, Fontana M, Ramlall M, Hamarneh A, Sirkar A, et al. Residual myocardial iron following intramyocardial hemorrhage during the convalescent phase of reperfused ST-segment-elevation myocardial infarction and adverse left ventricular remodeling. *Circ Cardiovasc Imaging*. 2016;9:e004940. doi: 10.1161/CIRCIMAGING.116.004940
 230. Wang G, Yang HJ, Kali A, Cokic I, Tang R, Xie G, Yang Q, Francis J, Li S, Dharmakumar R. Influence of myocardial hemorrhage on staging of reperfused myocardial infarctions with T2 cardiac magnetic resonance imaging: insights into the dependence on infarction type with ex vivo validation. *JACC Cardiovasc Imaging*. 2019;12:693–703. doi: 10.1016/j.jcmg.2018.01.018
 231. Ganame J, Messalli G, Dymarkowski S, Rademakers FE, Desmet W, Van de Werf F, Bogaert J. Impact of myocardial haemorrhage on left ventricular function and remodelling in patients with reperfused acute myocardial infarction. *Eur Heart J*. 2009;30:1440–1449. doi: 10.1093/eurheartj/ehp093
 232. Kumar A, Connelly K, Vora K, Bainey KR, Howarth A, Leipsic J, Betteridge-LeBlanc S, Prato FS, Leong-Poi H, Main A, et al. The Canadian Cardiovascular Society classification of acute atherothrombotic myocardial infarction based on stages of tissue injury severity: an expert consensus statement. *Can J Cardiol*. 2024;40:1–14. doi: 10.1016/j.cjca.2023.09.020
 233. Lechner I, Reindl M, Stiermaier T, Tiller C, Holzknecht M, Oberhollenzer F, von der Emde S, Mayr A, Feistritz H-J, Carberry J, et al. Clinical outcomes associated with various microvascular injury patterns identified by CMR after STEMI. *J Am Coll Cardiol*. 2024;83:2052–2062. doi: 10.1016/j.jacc.2024.03.408
 234. Vyas R, Chahal KH, Bhuta S, Pasadyn V, Katterle K, Niedoba MJ, Vora K, Dharmakumar R, Gupta R. Impact of intramyocardial hemorrhage on clinical outcomes in ST-elevation myocardial infarction: a systematic review and meta-analysis. *J Soc Cardiovasc Angiogr Interv*. 2022;1:100444. doi: 10.1016/j.jscai.2022.100444
 235. Wu KC, Zerhouni EA, Judd RM, Lugo-Olivieri CH, Barouch LA, Schulman SP, Blumenthal RS, Lima JA. Prognostic significance of microvascular obstruction by magnetic resonance imaging in patients with acute myocardial infarction. *Circulation*. 1998;97:765–772. doi: 10.1161/01.cir.97.7.765
 236. Locca D, Bucciarelli-Ducci C, Ferrante G, La Manna A, Keenan NG, Grasso A, Barlis P, Del Furia F, Prasad SK, Kaski JC, et al. New universal definition of myocardial infarction applicable after complex percutaneous coronary interventions? *JACC Cardiovasc Interv*. 2010;3:950–958. doi: 10.1016/j.jcin.2010.06.015
 237. Hwang HY, Yeom SY, Park EA, Lee W, Jang MJ, Kim KB. Serial cardiac magnetic resonance imaging after surgical coronary revascularization for left ventricular dysfunction. *J Thorac Cardiovasc Surg*. 2020;159:1798–1805. doi: 10.1016/j.jtcvs.2019.04.109

238. Kolossvary M, Szilveszter B, Merkely B, Maurovich-Horvat P. Plaque imaging with CT: a comprehensive review on coronary CT angiography based risk assessment. *Cardiovasc Diagn Ther*. 2017;7:489–506. doi: 10.21037/cdt.2016.11.06
239. Puchner SB, Liu T, Mayrhofer T, Truong QA, Lee H, Fleg JL, Nagurney JT, Udelson JE, Hoffmann U, Ferencik M. High-risk plaque detected on coronary CT angiography predicts acute coronary syndromes independent of significant stenosis in acute chest pain: results from the ROMICAT-II trial. *J Am Coll Cardiol*. 2014;64:684–692. doi: 10.1016/j.jacc.2014.05.039
240. Kofoed KF, Engstrom T, Sigvardsen PE, Linde JJ, Torp-Pedersen C, de Knecht M, Hansen PR, Fritz-Hansen T, Bech J, Heitmann M, et al. Prognostic value of coronary CT angiography in patients with non-ST-segment elevation acute coronary syndromes. *J Am Coll Cardiol*. 2021;77:1044–1052. doi: 10.1016/j.jacc.2020.12.037
241. Gray AJ, Roobottom C, Smith JE, Goodacre S, Oatey K, O'Brien R, Storey RF, Curzen N, Keating L, Kardos A, et al; RAPID-CTCA Investigators. Early computed tomography coronary angiography in patients with suspected acute coronary syndrome: randomised controlled trial. *BMJ*. 2021;374:n2106. doi: 10.1136/bmj.n2106
242. Meah MN, Bularga A, Tzolos E, Chapman AR, Daghem M, Hung JD, Chiong J, Taggart C, Wereski R, Gray A, et al. Distinguishing type 1 from type 2 myocardial infarction by using CT coronary angiography. *Radiol Cardiothorac Imaging*. 2022;4:e220081. doi: 10.1148/ryct.220081
243. Pergola V, Continisio S, Mantovani F, Motta R, Mattesi G, Marrazzo G, Dellino CM, Montonati C, De Conti G, Galzerano D, et al. Spontaneous coronary artery dissection: the emerging role of coronary computed tomography. *Eur Heart J Cardiovasc Imaging*. 2023;24:839–850. doi: 10.1093/ehjci/jead060
244. Dai N, Chen Z, Zhou F, Zhou Y, Hu N, Duan S, Wang W, Zhang L, Qian J, Ge J. Coronary CT angiography-derived plaque characteristics and physiologic patterns for peri-procedural myocardial infarction and subsequent events. *Eur Heart J Cardiovasc Imaging*. 2023;24:897–908. doi: 10.1093/ehjci/jead025
245. Serruys PW, Kageyama S, Pompilio G, Andreini D, Pontone G, Mushtaq S, La Meir M, De Mey J, Tanaka K, Doenst T, et al. Coronary bypass surgery guided by computed tomography in a low-risk population. *Eur Heart J*. 2024;45:1804–1815. doi: 10.1093/eurheartj/ehae199
246. Palmisano A, Vignale D, Tadic M, Moroni F, De Stefano D, Gatti M, Boccia E, Faletti R, Oppizzi M, Peretto G, et al. Myocardial late contrast enhancement CT in troponin-positive acute chest pain syndrome. *Radiology*. 2022;302:545–553. doi: 10.1148/radiol.211288
247. Çitaku H, Miftari R, Stubljarić D, Krasniqi X. Size of acute myocardial infarction correlates with earlier time of initiation of reperfusion therapy with cardiac perfusion scintigraphy: a national single-center study. *Med Sci Monit Basic Res*. 2021;27:e933214. doi: 10.12659/MSMBR.933214
248. Wagner A, Mahrholdt H, Holly TA, Elliott MD, Regenfus M, Parker M, Klocke FJ, Bonow RO, Kim RJ, Judd RM. Contrast-enhanced MRI and routine single photon emission computed tomography (SPECT) perfusion imaging for detection of subendocardial myocardial infarction: an imaging study. *Lancet*. 2003;361:374–379. doi: 10.1016/S0140-6736(03)12389-6
249. Wells RG. Instrumentation in molecular imaging. *J Nucl Cardiol*. 2016;23:1343–1347. doi: 10.1007/s12350-016-0498-z
250. Tzolos E, Bing R, Andrews J, MacAskill MG, Tavares AAS, Macnaught G, Clark T, Mills NL, Fujisawa T, Nash J, et al. Noninvasive in vivo coronary artery thrombus imaging. *JACC Cardiovasc Imaging*. 2023;16:820–832. doi: 10.1016/j.jcmg.2022.10.002
251. Tzolos E, Bing R, Newby DE, Dweck MR. Categorising myocardial infarction with advanced cardiovascular imaging. *Lancet*. 2021;398:e9. doi: 10.1016/S0140-6736(21)01329-5
252. Gibson CM, Karha J, Murphy SA, James D, Morrow DA, Cannon CP, Giugliano RP, Antman EM, Braunwald E; TIMI Study Group. Early and long-term clinical outcomes associated with reinfarction following fibrinolytic administration in the Thrombolysis in Myocardial Infarction trials. *J Am Coll Cardiol*. 2003;42:7–16. doi: 10.1016/s0735-1097(03)00506-0
253. Kernis SJ, Harjai KJ, Stone GW, Grines LL, Boura JA, Yerkley MW, O'Neill W, Grines CL. The incidence, predictors, and outcomes of early reinfarction after primary angioplasty for acute myocardial infarction. *J Am Coll Cardiol*. 2003;42:1173–1177. doi: 10.1016/s0735-1097(03)00920-3
254. Nair R, Johnson M, Kravitz K, Huded C, Rajeswaran J, Anabila M, Blackstone E, Menon V, Lincoff AM, Kapadia S, et al. Characteristics and outcomes of early recurrent myocardial infarction after acute myocardial infarction. *J Am Heart Assoc*. 2021;10:e019270. doi: 10.1161/JAHA.120.019270
255. Song J, Murugiah K, Hu S, Gao Y, Li X, Krumholz HM, Zheng X; China PEACE Collaborative Group. Incidence, predictors, and prognostic impact of recurrent acute myocardial infarction in China. *Heart*. 2020;107:313–318. doi: 10.1136/heartjnl-2020-317165
256. Holzmann MJ, Andersson T, Doemland ML, Roux S. Recurrent myocardial infarction and emergency department visits: a retrospective study on the Stockholm Area Chest Pain Cohort. *Open Heart*. 2023;10:e002206. doi: 10.1136/openhrt-2022-002206
257. Stromback U, Engstrom A, Lundqvist R, Lundblad D, Vikman I. The second myocardial infarction: is there any difference in symptoms and prehospital delay compared to the first myocardial infarction? *Eur J Cardiovasc Nurs*. 2018;17:652–659. doi: 10.1177/1474515118777391
258. Kwok CS, Bennett S, Azam Z, Welsh V, Potluri R, Loke YK, Mallen CD. Misdiagnosis of acute myocardial infarction: a systematic review of the literature. *Crit Pathw Cardiol*. 2021;20:155–162. doi: 10.1097/HPC.0000000000000256
259. Adamson PD, McAllister D, Pilbrow A, Pickering JW, Poppe K, Shah A, Whalley G, Ellis C, Mills NL, Newby DE, et al. Convalescent troponin and cardiovascular death following acute coronary syndrome. *Heart*. 2019;105:1717–1724. doi: 10.1136/heartjnl-2019-315084
260. Apple FS, Murakami MM. Cardiac troponin and creatine kinase MB monitoring during in-hospital myocardial reinfarction. *Clin Chem*. 2005;51:460–463. doi: 10.1373/clinchem.2004.042887
261. Ekstrom K, Nepper-Christensen L, Ahtarovski KA, Kyhl K, Goransson C, Bertelsen L, Ghotbi AA, Kelbæk H, Helqvist S, Høfsten DE, et al. Impact of multiple myocardial scars detected by CMR in patients following STEMI. *JACC Cardiovasc Imaging*. 2019;12:2168–2178. doi: 10.1016/j.jcmg.2019.01.032
262. Kociol RD, Pang PS, Gheorghiade M, Fonarow GC, O'Connor CM, Felker GM. Troponin elevation in heart failure prevalence, mechanisms, and clinical implications. *J Am Coll Cardiol*. 2010;56:1071–1078. doi: 10.1016/j.jacc.2010.06.016
263. Felker GM, Mentz RJ, Teerlink JR, Voors AA, Pang PS, Ponikowski P, Greenberg BH, Filippatos G, Davison BA, Cotter G, et al. Serial high sensitivity cardiac troponin T measurement in acute heart failure: insights from the RELAX-AHF study. *Eur J Heart Fail*. 2015;17:1262–1270. doi: 10.1002/ejhf.341
264. Januzzi JL Jr, Filippatos G, Nieminen M, Gheorghiade M. Troponin elevation in patients with heart failure: on behalf of the Third Universal Definition of Myocardial Infarction Global Task Force: Heart Failure Section. *Eur Heart J*. 2012;33:2265–2271. doi: 10.1093/eurheartj/ehs191
265. Weil BR, Suzuki G, Young RF, Iyer V, Canty JM Jr. Troponin release and reversible left ventricular dysfunction after transient pressure overload. *J Am Coll Cardiol*. 2018;71:2906–2916. doi: 10.1016/j.jacc.2018.04.029
266. Aimo A, Januzzi JL Jr, Vergaro G, Ripoli A, Latini R, Masson S, Magnoli M, Anand IS, Cohn JN, Tavazzi L, et al. Prognostic value of high-sensitivity troponin T in chronic heart failure: an individual patient data meta-analysis. *Circulation*. 2018;137:286–297. doi: 10.1161/CIRCULATIONAHA.117.031560
267. Frank Peacock W 4th, De Marco T, Fonarow GC, Diercks D, Wynne J, Apple FS, Wu AHB; ADHERE Investigators. Cardiac troponin and outcome in acute heart failure. *N Engl J Med*. 2008;358:2117–2126. doi: 10.1056/NEJMoa0706824
268. Packer M, Januzzi JL, Ferreira JP, Anker SD, Butler J, Filippatos G, Pocock SJ, Brueckmann M, Jamal W, Cotton D, et al; EMPEROR-Reduced Trial Committees and Investigators. Concentration-dependent clinical and prognostic importance of high-sensitivity cardiac troponin T in heart failure and a reduced ejection fraction and the influence of empagliflozin: the EMPEROR-Reduced trial. *Eur J Heart Fail*. 2021;23:1529–1538. doi: 10.1002/ejhf.2256
269. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumach A, Bohm M, Burri H, Butler J, Čelutkienė J, Chioncel O, et al; ESC Scientific Document Group. 2021 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J*. 2021;42:3599–3726. doi: 10.1093/eurheartj/ehab368
270. Gallacher PJ, Yeung D, Bell S, Shah ASV, Mills NL, Dhaun N. Kidney replacement therapy: trends in incidence, treatment, and outcomes of myocardial infarction and stroke in a nationwide Scottish study. *Eur Heart J*. 2024;45:1339–1351. doi: 10.1093/eurheartj/ehae080
271. Szumner K, Lundman P, Jacobson SH, Schon S, Lindback J, Stenestrand U, Wallentin L, Jernberg T; SWEDEHEART. Relation between renal function, presentation, use of therapies and in-hospital complications in acute coronary syndrome: data from the SWEDEHEART register. *J Intern Med*. 2010;268:40–49. doi: 10.1111/j.1365-2796.2009.02204.x
272. Gallacher PJ, Miller-Hodges E, Shah ASV, Anand A, Dhaun N, Mills NL; High-STEACS Investigators. Use of high-sensitivity cardiac troponin in

- patients with kidney impairment: a randomized clinical trial. *JAMA Intern Med.* 2021;181:1237–1239. doi: 10.1001/jamainternmed.2021.1184
273. Keddis MT, El-Zoghby ZM, El Ters M, Rodrigo E, Pellikka PA, Jaffe AS, Cosio FG. Cardiac troponin T before and after kidney transplantation: determinants and implications for posttransplant survival. *Am J Transplant.* 2013;13:406–414. doi: 10.1111/j.1600-6143.2012.04317.x
 274. van der Linden N, Cornelis T, Kimenai DM, Klinkenberg LJ, Hilderink JM, Luck S, Litjens EJR, Peeters FECM, Streng AS, Breidhardt T, et al. Origin of cardiac troponin T elevations in chronic kidney disease. *Circulation.* 2017;136:1073–1075. doi: 10.1161/CIRCULATIONAHA.117.029986
 275. Matsushita K, Ballew SH, Wang AY, Kalyesubula R, Schaeffner E, Agarwal R. Epidemiology and risk of cardiovascular disease in populations with chronic kidney disease. *Nat Rev Nephrol.* 2022;18:696–707. doi: 10.1038/s41581-022-00616-6
 276. Ndumele CE, Neeland JJ, Tuttle KR, Chow SL, Mathew RO, Khan SS, Coresh J, Baker-Smith CM, Carnethon MR, Després J-P, et al; American Heart Association. A synopsis of the evidence for the science and clinical management of cardiovascular-kidney-metabolic (CKM) syndrome: a scientific statement from the American Heart Association. *Circulation.* 2023;148:1636–1664. doi: 10.1161/CIR.0000000000001186
 277. Brunner FJ, Kroger F, Blaum C, Gossling A, Lorenz T, van Erckelens E, Brätz J, Westermann D, Blankenberg S, Zeller T, et al. Association of high-sensitivity troponin T and I with the severity of stable coronary artery disease in patients with chronic kidney disease. *Atherosclerosis.* 2020;313:81–87. doi: 10.1016/j.atherosclerosis.2020.09.024
 278. deFilippi C, Seliger SL, Kelley W, Duh SH, Hise M, Christenson RH, Wolf M, Gaggin H, Januzzi J. Interpreting cardiac troponin results from high-sensitivity assays in chronic kidney disease without acute coronary syndrome. *Clin Chem.* 2012;58:1342–1351. doi: 10.1373/clinchem.2012.185322
 279. Knott JD, Ola O, De Micheli L, Akula A, Mehta RA, Dworak M, Crockford E, Lobo R, Slusser J, Rastas N, et al. Diagnosis of acute myocardial infarction in patients with renal failure using high-sensitivity cardiac troponin T. *Eur Heart J Acute Cardiovasc Care.* 2024;13:546–558. doi: 10.1093/ehjacc/zaue079
 280. Miller-Hodges E, Anand A, Shah ASV, Chapman AR, Gallacher P, Lee KK, Farrah T, Halbesma N, Blackmur JP, Newby DE, et al. High-sensitivity cardiac troponin and the risk stratification of patients with renal impairment presenting with suspected acute coronary syndrome. *Circulation.* 2018;137:425–435. doi: 10.1161/CIRCULATIONAHA.117.030320
 281. Twerenbold R, Badertscher P, Boeddinghaus J, Nestelberger T, Wildi K, Puelacher C, Sabti Z, Rubini Gimenez M, Tschirky S, du Fay de Lavallaz J, et al. 0/1-Hour triage algorithm for myocardial infarction in patients with renal dysfunction. *Circulation.* 2018;137:436–451. doi: 10.1161/CIRCULATIONAHA.117.028901
 282. Gallacher PJ, Miller-Hodges E, Shah ASV, Farrah TE, Halbesma N, Blackmur JP, Chapman AR, Adamson PD, Anand A, Strachan FE, et al; High-STEACS Investigators. High-sensitivity cardiac troponin and the diagnosis of myocardial infarction in patients with kidney impairment. *Kidney Int.* 2022;102:149–159. doi: 10.1016/j.kint.2022.02.019
 283. Davenport MS, Perazella MA, Yee J, Dillman JR, Fine D, McDonald RJ, Rodby RA, Wang CL, Weinreb JC. Use of intravenous iodinated contrast media in patients with kidney disease: consensus statements from the American College of Radiology and the National Kidney Foundation. *Radiology.* 2020;294:660–668. doi: 10.1148/radiol.2019192094
 284. Maslove DM, Tang B, Shankar-Hari M, Lawler PR, Angus DC, Baillie JK, Baron RM, Bauer M, Buchman TG, Calfee CS, et al. Redefining critical illness. *Nat Med.* 2022;28:1141–1148. doi: 10.1038/s41591-022-01843-x
 285. Pastores SM, Dulu A, Voigt L, Raoof N, Alicea M, Halpern NA. Premortem clinical diagnoses and postmortem autopsy findings: discrepancies in critically ill cancer patients. *Crit Care.* 2007;11:R48. doi: 10.1186/cc5782
 286. Perkins GD, McAuley DF, Davies S, Gao F. Discrepancies between clinical and postmortem diagnoses in critically ill patients: an observational study. *Crit Care.* 2003;7:R129–R132. doi: 10.1186/cc2359
 287. Roosen J, Frans E, Wilmer A, Knockaert DC, Bobbaers H. Comparison of premortem clinical diagnoses in critically ill patients and subsequent autopsy findings. *Mayo Clin Proc.* 2000;75:562–567. doi: 10.4065/75.6.562
 288. Docherty AB, Alam S, Shah AS, Moss A, Newby DE, Mills NL, Stanworth SJ, Lone NI, Walsh TS; TROPICAL Investigators. Unrecognised myocardial infarction and its relationship to outcome in critically ill patients with cardiovascular disease. *Intensive Care Med.* 2018;44:2059–2069. doi: 10.1007/s00134-018-5425-0
 289. Lim W, Holinski P, Devereaux PJ, Tkaczyk A, McDonald E, Clarke F, Qushmaq I, Terrenato I, Schunemann H, Crowther M, et al. Detecting myocardial infarction in critical illness using screening troponin measurements and ECG recordings. *Crit Care.* 2008;12:R36. doi: 10.1186/cc6815
 290. Rennyson SL, Hunt J, Haley MW, Norton HJ, Littmann L. Electrocardiographic ST-segment elevation myocardial infarction in critically ill patients: an observational cohort analysis. *Crit Care Med.* 2010;38:2304–2309. doi: 10.1097/CCM.0b013e3181fa02cd
 291. Lim W, Qushmaq I, Cook DJ, Devereaux PJ, Heels-Ansdell D, Crowther MA, Tkaczyk A, Meade MO, Cook RJ. Reliability of electrocardiogram interpretation in critically ill patients. *Crit Care Med.* 2006;34:1338–1343. doi: 10.1097/01.CCM.0000214679.23957.90
 292. Marcelino PA, Marum SM, Fernandes AP, Germano N, Lopes MG. Routine transthoracic echocardiography in a general intensive care unit: an 18 month survey in 704 patients. *Eur J Intern Med.* 2009;20:e37–e42. doi: 10.1016/j.iejim.2008.09.015
 293. Cavefors O, Holmqvist J, Bech-Hanssen O, Einarsson F, Norberg E, Lundin S, Omerovic E, Ricksten S-E, Redfors B, Oras J. Regional left ventricular systolic dysfunction associated with critical illness: incidence and effect on outcome. *ESC Heart Fail.* 2021;8:5415–5423. doi: 10.1002/ehf2.13633
 294. Champion S, Belcour D, Vandroux D, Drouet D, Gauzere BA, Bouchet B, Bossard G, Djouhri S, Jabot J, Champion M, et al. Stress (tako-tsubo) cardiomyopathy in critically-ill patients. *Eur Heart J Acute Cardiovasc Care.* 2015;4:189–196. doi: 10.1177/2048872614547686
 295. Vallabhajosyula S, Deshmukh AJ, Kashani K, Prasad A, Sakhuja A. Tako-tsubo cardiomyopathy in severe sepsis: national trends, predictors, and outcomes. *J Am Heart Assoc.* 2018;7:e009160. doi: 10.1161/JAHA.118.009160
 296. Murata M, Nakagawa N, Kawasaki T, Yasuo S, Yoshida T, Ando K, Okamori S, Okada Y. Adverse events during intrahospital transport of critically ill patients: a systematic review and meta-analysis. *Am J Emerg Med.* 2022;52:13–19. doi: 10.1016/j.ajem.2021.11.021
 297. Mjumentier-Decrucq E, Poissy J, Favory R, Nseir S, Onimus T, Guerry MJ, Duracher A, Mathieu D. Adverse events during intrahospital transport of critically ill patients: incidence and risk factors. *Ann Intensive Care.* 2013;3:10. doi: 10.1186/2110-5820-3-10
 298. Halvorsen S, Mehilli J, Cassese S, Hall TS, Abdelhamid M, Barbato E, De Hert S, de Laval I, Geisler T, Hinterbuchner L, et al; ESC Scientific Document Group. 2022 ESC guidelines on cardiovascular assessment and management of patients undergoing non-cardiac surgery. *Eur Heart J.* 2022;43:3826–3924. doi: 10.1093/eurheartj/ehac270
 299. Devereaux PJ, Xavier D, Pogue J, Guyatt G, Sigamani A, Garutti I, Leslie K, Rao-Melacini P, Chrolavicius S, Yang H, et al; POISE (Perioperative ISchemic Evaluation) Investigators. Characteristics and short-term prognosis of perioperative myocardial infarction in patients undergoing non-cardiac surgery: a cohort study. *Ann Intern Med.* 2011;154:523–528. doi: 10.7326/0003-4819-154-8-201104190-00003
 300. Puelacher C, Gualandro DM, Glarner N, Lurati Buse G, Lampart A, Bolliger D, Steiner LA, Grossenbacher M, Burri-Winkler K, Gerhard H, et al; BASEL-PMI Investigators. Long-term outcomes of perioperative myocardial infarction/injury after non-cardiac surgery. *Eur Heart J.* 2023;44:1690–1701. doi: 10.1093/eurheartj/ehac798
 301. Smilowitz NR, Redel-Traub G, Hausvater A, Armanious A, Nicholson J, Puelacher C, Berger JS. Myocardial injury after noncardiac surgery: a systematic review and meta-analysis. *Cardiol Rev.* 2019;27:267–273. doi: 10.1097/CRD.0000000000000254
 302. Devereaux PJ, Biccari BM, Sigamani A, Xavier D, Chan MTV, Srinathan SK, Walsh M, Abraham V, Pearse R, Wang CY, et al; Writing Committee for the VISION Study Investigators. Association of postoperative high-sensitivity troponin levels with myocardial injury and 30-day mortality among patients undergoing noncardiac surgery. *JAMA.* 2017;317:1642–1651. doi: 10.1001/jama.2017.4360
 303. Thompson A, Fleischmann KE, Smilowitz NR, de Las Fuentes L, Mukherjee D, Aggarwal NR, Ahmad FS, Allen RB, Altin SE, Auerbach A, et al; Peer Review Committee Members. 2024 AHA/ACC/ACS/ASNC/HRS/SCA/SCCT/SCMR/SVM guideline for perioperative cardiovascular management for noncardiac surgery: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation.* 2024;150:e351–e442. doi: 10.1161/CIR.0000000000001285
 304. Nagele P, Brown F, Gage BF, Gibson DW, Miller JP, Jaffe AS, Apple FS, Scott MG. High-sensitivity cardiac troponin T in prediction and diagnosis of myocardial infarction and long-term mortality after noncardiac surgery. *Am Heart J.* 2013;166:325–332.e1. doi: 10.1016/j.ahj.2013.04.018
 305. Genereux P, Head SJ, Van Mieghem NM, Kodali S, Kirtane AJ, Xu K, Smith C, Serruys PW, Kappetein AP, Leon MB. Clinical outcomes after transcatheter

- aortic valve replacement using valve academic research consortium definitions: a weighted meta-analysis of 3,519 patients from 16 studies. *J Am Coll Cardiol*. 2012;59:2317–2326. doi: 10.1016/j.jacc.2012.02.022
306. Ribeiro HB, Webb JG, Makkar RR, Cohen MG, Kapadia SR, Kodali S, Tamburino C, Barbanti M, Chakravarty T, Jilaihawi H, et al. Predictive factors, management, and clinical outcomes of coronary obstruction following transcatheter aortic valve implantation: insights from a large multicenter registry. *J Am Coll Cardiol*. 2013;62:1552–1562. doi: 10.1016/j.jacc.2013.07.040
 307. Akuka A, Landes U, Manevich L, Rubinshtein R, Danenberg HD. Coronary embolism after transcatheter aortic valve replacement—case series and review of literature. *Am J Cardiol*. 2023;205:234–240. doi: 10.1016/j.amjcard.2023.07.137
 308. Kim WK, Rolf A, Liebetrau C, Van Linden A, Blumenstein J, Kempfert J, Bachmann G, Nef H, Hamm C, Walther T, et al. Detection of myocardial injury by CMR after transcatheter aortic valve replacement according to VARC-3 criteria. *JACC Cardiovasc Interv*. 2023;16:1221–1232. doi: 10.1016/j.jcin.2023.03.022
 309. Ribeiro HB, Nombela-Franco L, Munoz-Garcia AJ, Lemos PA, Amat-Santos I, Serra V, de Brito FS, Abizaid A, Sarmiento-Leite R, Puri R, et al. Predictors and impact of myocardial injury after transcatheter aortic valve replacement: a multicenter registry. *J Am Coll Cardiol*. 2015;66:2075–2088. doi: 10.1016/j.jacc.2015.08.881
 311. Rodes-Cabau J, Gutierrez M, Bagur R, De Larochelliere R, Doyle C, Cote M, Villeneuve J, Bertrand OF, Larose E, Manazzoni J, et al. Incidence, predictive factors, and prognostic value of myocardial injury following uncomplicated transcatheter aortic valve implantation. *J Am Coll Cardiol*. 2011;57:1988–1999. doi: 10.1016/j.jacc.2010.11.060
 312. Sinning JM, Hammerstingl C, Schueler R, Neugebauer A, Keul S, Ghanem A, Mellert F, Schiller W, Müller C, Vasa-Nicotera M, et al. The prognostic value of acute and chronic troponin elevation after transcatheter aortic valve implantation. *EuroIntervention*. 2016;11:1522–1529. doi: 10.4244/EIJY15M02_02
 313. Tocchetti CG, Farmakis D, Koop Y, Andres MS, Couch LS, Formisano L, Ciardiello F, Pane F, Au L, Emmerich M, et al. Cardiovascular toxicities of immune therapies for cancer: a scientific statement of the Heart Failure Association (HFA) of the ESC and the ESC Council of Cardio-Oncology. *Eur J Heart Fail*. 2024;26:2055–2076. doi: 10.1002/ehf.3340
 314. Cardinale D, Sandri MT, Colombo A, Colombo N, Boeri M, Lamantia G, Civelli M, Peccatori F, Martinelli G, Fiorentini C, et al. Prognostic value of troponin I in cardiac risk stratification of cancer patients undergoing high-dose chemotherapy. *Circulation*. 2004;109:2749–2754. doi: 10.1161/01.CIR.0000130926.51766.CC
 315. Tzolos E, Adamson PD, Hall PS, Macpherson IR, Oikonomidou O, MacLean M, Lewis SC, McVicar H, Newby DE, Mills NL, et al. Dynamic changes in high-sensitivity cardiac troponin I in response to anthracycline-based chemotherapy. *Clin Oncol (R Coll Radiol)*. 2020;32:292–297. doi: 10.1016/j.clon.2019.11.008
 316. Lyon AR, Lopez-Fernandez T, Couch LS, Asteggiano R, Aznar MC, Bergler-Klein J, Boriani G, Cardinale D, Cordoba R, Cosyns B, et al; ESC Scientific Document Group. 2022 ESC guidelines on cardio-oncology developed in collaboration with the European Hematology Association (EHA), the European Society for Therapeutic Radiology and Oncology (ESTRO) and the International Cardio-Oncology Society (IC-OS). *Eur Heart J*. 2022;43:4229–4361. doi: 10.1093/eurheartj/ehac244
 317. Jiang C, Xu H, Wu Y. Effect of chemotherapy in tumor on coronary arteries: mechanisms and management. *Life Sci*. 2024;338:122377. doi: 10.1016/j.lfs.2023.122377
 318. Juneag L, Asplund K, Svedlund M. A qualitative study: perceptions of the psychosocial consequences and access to support after an acute myocardial infarction. *Intensive Crit Care Nurs*. 2014;30:22–30. doi: 10.1016/j.iccn.2013.07.002
 319. Gwaltney C, Reaney M, Krohe M, Martin MM, Falvey H, Mollon P. Symptoms and functional limitations in the first year following a myocardial infarction: a qualitative study. *Patient*. 2017;10:225–235. doi: 10.1007/s40271-016-0194-8
 320. Lichtman JH, Froelicher ES, Blumenthal JA, Carney RM, Doering LV, Frasure-Smith N, Freedland KE, Jaffe AS, Leitheit-Limson EC, Sheps DS, et al; American Heart Association Statistics Committee of the Council on Epidemiology and Prevention and the Council on Cardiovascular and Stroke Nursing. Depression as a risk factor for poor prognosis among patients with acute coronary syndrome: systematic review and recommendations: a scientific statement from the American Heart Association. *Circulation*. 2014;129:1350–1369. doi: 10.1161/CIR.0000000000000019
 321. Murphy B, Le Grande M, Alvarenga M, Worcester M, Jackson A. Anxiety and depression after a cardiac event: prevalence and predictors. *Front Psychol*. 2019;10:3010. doi: 10.3389/fpsyg.2019.03010
 322. Legler SR, Beale EE, Celano CM, Beach SR, Healy BC, Huffman JC. State gratitude for one's life and health after an acute coronary syndrome: prospective associations with physical activity, medical adherence and re-hospitalizations. *J Posit Psychol*. 2019;14:283–291. doi: 10.1080/17439760.2017.1414295
 323. Dreyer RP, Pavlo AJ, Horne A, Dunn R, Danvers K, Brush J, Slade M, Davidson L. Conceptual framework for personal recovery in patients with acute myocardial infarction. *J Am Heart Assoc*. 2021;10:e022354. doi: 10.1161/JAHA.121.022354
 324. Sjöström-Strand A, Ivarsson B, Sjöberg T. Women's experience of a myocardial infarction: 5 years later. *Scand J Caring Sci*. 2011;25:459–466. doi: 10.1111/j.1471-6712.2010.00849.x
 325. Rossello X, Pocock SJ, Julian DG. Long-term use of cardiovascular drugs: challenges for research and for patient care. *J Am Coll Cardiol*. 2015;66:1273–1285. doi: 10.1016/j.jacc.2015.07.018
 326. Smedegaard L, Numé AK, Charlot M, Kragholm G, Gislason G, Hansen PR. Return to work and risk of subsequent detachment from employment after myocardial infarction: insights from Danish nationwide registries. *J Am Heart Assoc*. 2017;6:e006486. doi: 10.1161/JAHA.117.006486
 327. Boudrez H, De Backer G. Recent findings on return to work after an acute myocardial infarction or coronary artery bypass grafting. *Acta Cardiol*. 2000;55:341–349. doi: 10.2143/AC.55.6.2005765
 328. Mittag O, Kolenda KD, Nordman KJ, Bernien J, Maurischat C. Return to work after myocardial infarction/coronary artery bypass grafting: patients' and physicians' initial viewpoints and outcome 12 months later. *Soc Sci Med*. 2001;52:1441–1450. doi: 10.1016/s0277-9536(00)00250-1
 329. Sepehrvand N, Zheng Y, Armstrong PW, Welsh R, Goodman SG, Tymchak W, Khadour F, Chan M, Weiss D, Ezekowitz JA, et al. Alignment of site versus adjudication committee-based diagnosis with patient outcomes: insights from the Providing Rapid Out of Hospital Acute Cardiovascular Treatment 3 trial. *Clin Trials*. 2016;13:140–148. doi: 10.1177/1740774515601437
 330. McCarthy CP, Murphy S, Cohen JA, Rehman S, Jones-O'Connor M, Olshan DS, Singh A, Vaduganathan M, Januzzi JL, Wasfy JH. Underutilization of cardiac rehabilitation for type 2 myocardial infarction. *J Am Coll Cardiol*. 2019;73:2005–2007. doi: 10.1016/j.jacc.2019.01.032
 331. Zeymer U, Goss F, Kunadt M, Oldenburg S, Hochadel M, Thiele H, Werdan K; GULLIVE-R Investigators. Patient knowledge about risk factors, achievement of target values, and guideline-adherent secondary prevention therapies 12 months after acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care*. 2024;13:537–545. doi: 10.1093/ehjacc/zaue066
 332. Hertz JT, Sakita FM, Manavalan P, Mmbaga BT, Thielman NM, Staton CA. Knowledge, attitudes, and preventative practices regarding ischemic heart disease among emergency department patients in northern Tanzania. *Public Health*. 2019;175:60–67. doi: 10.1016/j.puhe.2019.06.017
 333. Gallagher R, Roach K, Belshaw J, Kirkness A, Sadler L, Warrington D. A pre-test post-test study of a brief educational intervention demonstrates improved knowledge of potential acute myocardial infarction symptoms and appropriate responses in cardiac rehabilitation patients. *Aust Crit Care*. 2013;26:49–54. doi: 10.1016/j.aucc.2012.01.002
 334. Redfern J, Hyun K, Chew DP, Astley C, Chow C, Aliprandi-Costa B, Howell T, Carr B, Lintern K, Ranasinghe I, et al. Prescription of secondary prevention medications, lifestyle advice, and referral to rehabilitation among acute coronary syndrome inpatients: results from a large prospective audit in Australia and New Zealand. *Heart*. 2014;100:1281–1288. doi: 10.1136/heartjnl-2013-305296
 335. Albarqouni L, Smenes K, Meinertz T, Schunkert H, Fang X, Ronel J, Ladwig K-H. Patients' knowledge about symptoms and adequate behaviour during acute myocardial infarction and its impact on delay time: findings from the multicentre MEDEA Study. *Patient Educ Couns*. 2016;99:1845–1851. doi: 10.1016/j.pec.2016.06.007
 336. Grace SL, Turk-Adawi KI, Contractor A, Atrey A, Campbell NR, Derman W, Ghisi GLM, Sarkar BK, Yeo TJ, Lopez-Jimenez F, et al. Cardiac rehabilitation delivery model for low-resource settings: an international council of cardiovascular prevention and rehabilitation consensus statement. *Prog Cardiovasc Dis*. 2016;59:303–322. doi: 10.1016/j.pcad.2016.08.004
 337. Chapman AR, Adamson PD, Shah ASV, Anand A, Strachan FE, Ferry AV, Lee KK, Berry C, Findlay I, Cruikshank A, et al; High-STEACS

- Investigators. High-sensitivity cardiac troponin and the universal definition of myocardial infarction. *Circulation*. 2020;141:161–171. doi: 10.1161/CIRCULATIONAHA.119.042960
338. Thygesen K, Jaffe AS. Adjusting the MI codes into the framework of the universal definition of myocardial infarction. *J Am Coll Cardiol*. 2021;77:858–860. doi: 10.1016/j.jacc.2021.01.003
 339. GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. 2020;396:1204–1222. doi: 10.1016/S0140-6736(20)30925-9
 340. Makuvire TT, Latif Z, Kumar P, Samad Z, Warraich HJ. Trends in ischemic heart disease among females in low-and middle-income countries from 1990 to 2019. *Int J Cardiol*. 2023;388:131113. doi: 10.1016/j.ijcard.2023.06.004
 341. Yusuf S, Hawken S, Ounpuu S, Dans T, Avezum A, Lanas F, McQueen M, Budaj A, Pais P, Varigos J, et al; INTERHEART Study Investigators. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet*. 2004;364:937–952. doi: 10.1016/S0140-6736(04)17018-9
 342. Beza L, Leslie SL, Alemayehu B, Gary R. Acute coronary syndrome treatment delay in low to middle-income countries: a systematic review. *Int J Cardiol Heart Vasc*. 2021;35:100823. doi: 10.1016/j.ijcha.2021.100823
 343. Rossello X, Gonzalez-Del-Hoyo M, Aktaa S, Gale CP, Barbash I, Claeys MJ, Coughlan JJ, Ferreira JP, Galbraith M, Leosdottir M, et al. European Society of Cardiology quality indicators for the management of acute coronary syndromes: developed in collaboration with the Association for Acute CardioVascular Care and the European Association of Percutaneous Cardiovascular Interventions of the ESC. *Eur Heart J Acute Cardiovasc Care*. 2025;14:145–154. doi: 10.1093/ehjacc/zuaf014
 344. Kibirige D, Atuhe D, Kampiire L, Kiggundu DS, Donggo P, Nabbaale J, Mwebaze RM, Kalyesubula R, Lumu W. Access to medicines and diagnostic tests integral in the management of diabetes mellitus and cardiovascular diseases in Uganda: insights from the ACCODAD study. *Int J Equity Health*. 2017;16:154. doi: 10.1186/s12939-017-0651-6
 345. Zhan L, Masoudi FA, Li X, Hu S, Venkatesh AK, Spertus JA, Lin Z, Desai NR, Li J, Krumholz HM, et al; China PEACE Collaborative Group. Trends in cardiac biomarker testing in China for patients with acute myocardial infarction, 2001 to 2011: China PEACE-retrospective AMI study. *PLoS One*. 2015;10:e0122237. doi: 10.1371/journal.pone.0122237
 346. Mehta S, Granger C, Grines CL, Jacobs A, Henry TD, Rokos I, Lansky A, Baumbach A, Botelho R, Ferre A, et al. Confronting system barriers for ST-elevation MI in low and middle income countries with a focus on India. *Indian Heart J*. 2018;70:185–190. doi: 10.1016/j.ijhj.2017.06.020
 347. Turk-Adawi K, Sarrafzadegan N, Fadhil I, Taubert K, Sadeghi M, Wenger NK, Tan NS, Grace SL. Cardiovascular disease in the eastern Mediterranean region: epidemiology and risk factor burden. *Nat Rev Cardiol*. 2018;15:106–119. doi: 10.1038/nrcardio.2017.138
 348. Bueno H, Rossello X, Pocock S, Van de Werf F, Chin CT, Danchin N, Lee SW-L, Medina J, Vega A, Huo Y. Regional variations in hospital management and post-discharge mortality in patients with non-ST-segment elevation acute coronary syndrome. *Clin Res Cardiol*. 2018;107:836–844. doi: 10.1007/s00392-018-1254-y
 349. Rossello X, Huo Y, Pocock S, Van de Werf F, Chin CT, Danchin N, Lee SW-L, Medina J, Vega A, Bueno H. Global geographical variations in ST-segment elevation myocardial infarction management and post-discharge mortality. *Int J Cardiol*. 2017;245:27–34. doi: 10.1016/j.ijcard.2017.07.039
 350. Xavier D, Pais P, Devereaux PJ, Xie C, Prabhakaran D, Reddy KS, Gupta R, Joshi P, Kerkar P, Thanikachalam S, et al; CREATE registry investigators. Treatment and outcomes of acute coronary syndromes in India (CREATE): a prospective analysis of registry data. *Lancet*. 2008;371:1435–1442. doi: 10.1016/S0140-6736(08)60623-6
 351. Mendis S, Thygesen K, Kuulasmaa K, Giampaoli S, Mahonen M, Ngu Blackett K, Lisheng L; Writing Group on behalf of the participating experts of the WHO consultation for revision of WHO definition of myocardial infarction. World Health Organization definition of myocardial infarction: 2008–09 revision. *Int J Epidemiol*. 2011;40:139–146. doi: 10.1093/ije/dyq165
 352. Carson JL, Brooks MM, Hebert PC, Goodman SG, Bertolet M, Glynn SA, Chaitman BR, Simon T, Lopes RD, Goldsweig AM, et al; MINT Investigators. Restrictive or liberal transfusion strategy in myocardial infarction and anemia. *N Engl J Med*. 2023;389:2446–2456. doi: 10.1056/NEJMoa2307983
 353. Chapman AR, Anand A, Boeddinghaus J, Ferry AV, Sandeman D, Adamson PD, Andrews J, Tan S, Cheng SF, D'Souza M, et al. Comparison of the efficacy and safety of early rule-out pathways for acute myocardial infarction. *Circulation*. 2017;135:1586–1596. doi: 10.1161/CIRCULATIONAHA.116.025021
 354. Fearon WF, Zimmermann FM, De Bruyne B, Piroth Z, van Straten AHM, Szekeley L, Davidavičius G, Kalinauskas G, Mansour S, Kharbanda R, et al; FAME 3 Investigators. Fractional flow reserve-guided PCI as compared with coronary bypass surgery. *N Engl J Med*. 2022;386:128–137. doi: 10.1056/NEJMoa2112299
 355. Kunadian V, Mossop H, Shields C, Bardgett M, Watts P, Teare MD, Pritchard J, Adams-Hall J, Runnett C, Ripley DP, et al; British Heart Foundation SENIOR-RITA Trial Team and Investigators. Invasive treatment strategy for older patients with myocardial infarction. *N Engl J Med*. 2024;391:1673–1684. doi: 10.1056/NEJMoa2407791
 356. Mueller C, Giannitsis E, Christ M, Ordonez-Llanos J, deFilippi C, McCord J, Body R, Panteghini M, Jernberg T, Plebani M, et al; RAPID-AMI Investigators. Multicenter evaluation of a 0-hour/1-hour algorithm in the diagnosis of myocardial infarction with high-sensitivity cardiac troponin T. *Ann Emerg Med*. 2016;68:76–87.e4. doi: 10.1016/j.annemergmed.2015.11.013
 357. Spitzer E, McFadden E, Vranckx P, Garcia-Garcia HM, Seltzer JH, Held C, de Vries T, Menon V, Brown KJ, Soliman Oll, et al. Critical appraisal of contemporary clinical endpoint definitions in coronary intervention trials: a guidance document. *JACC Cardiovasc Interv*. 2019;12:805–819. doi: 10.1016/j.jcin.2018.12.031
 358. Stone GW, Sabik JF, Serruys PW, Simonton CA, Genereux P, Puskas J, Kandzari DE, Morice M-C, Lembo N, Brown WM, et al; EXCEL Trial Investigators. Everolimus-eluting stents or bypass surgery for left main coronary artery disease. *N Engl J Med*. 2016;375:2223–2235. doi: 10.1056/NEJMoa1610227
 359. Al-Zaiti SS, Martin-Gill C, Zegre-Hemsey JK, Bouzid Z, Faramand Z, Alrawashdeh MO, Gregg RE, Helman S, Riek NT, Kraevsky-Phillips K, et al. Machine learning for ECG diagnosis and risk stratification of occlusion myocardial infarction. *Nat Med*. 2023;29:1804–1813. doi: 10.1038/s41591-023-02396-3
 360. Cho KH, Ji YH, Joo S, Chang M, Oh S, Lim J, Kim JH, Lee SH, Hyun DY, Lee N, et al. Novel artificial intelligence model using electrocardiogram for detecting acute myocardial infarction needing revascularization. *Eur Heart J Digit Health*. 2025;6:608–618. doi: 10.1093/ehjdh/ztaf049
 361. Elias P, Jain SS, Poterucha T, Randazzo M, Lopez Jimenez F, Khara R, Perez M, Ouyang D, Pirruccello J, Salerno M, et al. Artificial intelligence for cardiovascular care, part 1: advances: JACC review topic of the week. *J Am Coll Cardiol*. 2024;83:2472–2486. doi: 10.1016/j.jacc.2024.03.400
 362. Alizadehsani R, Khosravi A, Roshanzamir M, Abdar M, Sarrafzadegan N, Shafie D, Khozeimeh F, Shoeibi A, Nahavandi S, Panahiazar M, et al. Coronary artery disease detection using artificial intelligence techniques: a survey of trends, geographical differences and diagnostic features 1991–2020. *Comput Biol Med*. 2021;128:104095. doi: 10.1016/j.combiomed.2020.104095
 363. Dang QM, Psaltis PJ, Burgess S, Chandrasekhar J, Mukherjee S, Kritharides L, Jepson N, Fairley S, Ildayhid A, Layland J, et al. The Australian-New Zealand spontaneous coronary artery dissection cohort study: predictors of major adverse cardiovascular events and recurrence. *Eur Heart J*. 2025;46:2012–2023. doi: 10.1093/eurheartj/ehaf097
 364. DeFilippis AP, Abbott JD, Herbert BM, Bertolet MH, Chaitman BR, White HD, Goldsweig AM, Polonsky TS, Gupta R, Alsweller C, et al; MINT Investigators. Restrictive versus liberal transfusion in patients with type 1 or type 2 myocardial infarction: a prespecified analysis of the MINT trial. *Circulation*. 2024;150:1826–1836. doi: 10.1161/CIRCULATIONAHA.124.071208
 365. Chaitman BR, Alexander KP, Cyr DD, Berger JS, Reynolds HR, Bangalore S, Boden WE, Lopes RD, Demkow M, Piero Perna G, et al; ISCHEMIA Research Group. Myocardial infarction in the ISCHEMIA trial: impact of different definitions on incidence, prognosis, and treatment comparisons. *Circulation*. 2021;143:790–804. doi: 10.1161/CIRCULATIONAHA.120.047987
 366. Gregson J, Stone GW, Ben-Yehuda O, Redfors B, Kandzari DE, Morice MC, Leon MB, Kosmidou I, Lembo NJ, Brown WM, et al. Implications of alternative definitions of peri-procedural myocardial infarction after coronary revascularization. *J Am Coll Cardiol*. 2020;76:1609–1621. doi: 10.1016/j.jacc.2020.08.016