

# The density of covering numbers

Nathan McNew  
Towson University

Joint work with Jai Setty (NYU)

Combinatorial and Additive Number Theory  
CUNY Graduate Center, New York  
May 23, 2025

# Covering Systems

A set of integer residue classes forms a **covering system** if their union contains every integer.

# Covering Systems

A set of integer residue classes forms a **covering system** if their union contains every integer. Denote this set  $\{a_i \pmod{m_i}\}$ ,  $m_i > 1$ . Every integer  $n$  satisfies  $n \equiv a_i \pmod{m_i}$  for some  $i$ .

# Covering Systems

A set of integer residue classes forms a **covering system** if their union contains every integer. Denote this set  $\{a_i \pmod{m_i}\}$ ,  $m_i > 1$ . Every integer  $n$  satisfies  $n \equiv a_i \pmod{m_i}$  for some  $i$ .

If the moduli  $m_i$  are distinct we call it a **distinct covering system**.

# Covering Systems

A set of integer residue classes forms a **covering system** if their union contains every integer. Denote this set  $\{a_i \pmod{m_i}\}$ ,  $m_i > 1$ . Every integer  $n$  satisfies  $n \equiv a_i \pmod{m_i}$  for some  $i$ .

If the moduli  $m_i$  are distinct we call it a **distinct covering system**.

For example, we can check that the set:

$$\{0 \pmod{2}, 0 \pmod{3}, 1 \pmod{4}, 1 \pmod{6}, 11 \pmod{12}\}$$

forms a distinct covering system, by considering the residues modulo 12:

0 1 2 3 4 5 6 7 8 9 10 11

# Covering Systems

A set of integer residue classes forms a **covering system** if their union contains every integer. Denote this set  $\{a_i \pmod{m_i}\}$ ,  $m_i > 1$ . Every integer  $n$  satisfies  $n \equiv a_i \pmod{m_i}$  for some  $i$ .

If the moduli  $m_i$  are distinct we call it a **distinct covering system**.

For example, we can check that the set:

$$\{0 \pmod{2}, 0 \pmod{3}, 1 \pmod{4}, 1 \pmod{6}, 11 \pmod{12}\}$$

forms a distinct covering system, by considering the residues modulo 12:

0 1 2 3 4 5 6 7 8 9 10 11

# Covering Systems

A set of integer residue classes forms a **covering system** if their union contains every integer. Denote this set  $\{a_i \pmod{m_i}\}$ ,  $m_i > 1$ . Every integer  $n$  satisfies  $n \equiv a_i \pmod{m_i}$  for some  $i$ .

If the moduli  $m_i$  are distinct we call it a **distinct covering system**.

For example, we can check that the set:

$$\{0 \pmod{2}, 0 \pmod{3}, 1 \pmod{4}, 1 \pmod{6}, 11 \pmod{12}\}$$

forms a distinct covering system, by considering the residues modulo 12:

0 1 2 3 4 5 6 7 8 9 10 11

# Covering Systems

A set of integer residue classes forms a **covering system** if their union contains every integer. Denote this set  $\{a_i \pmod{m_i}\}$ ,  $m_i > 1$ . Every integer  $n$  satisfies  $n \equiv a_i \pmod{m_i}$  for some  $i$ .

If the moduli  $m_i$  are distinct we call it a **distinct covering system**.

For example, we can check that the set:

$$\{0 \pmod{2}, 0 \pmod{3}, 1 \pmod{4}, 1 \pmod{6}, 11 \pmod{12}\}$$

forms a distinct covering system, by considering the residues modulo 12:

0 1 2 3 4 5 6 7 8 9 10 11



# Covering Systems

A set of integer residue classes forms a **covering system** if their union contains every integer. Denote this set  $\{a_i \pmod{m_i}\}$ ,  $m_i > 1$ . Every integer  $n$  satisfies  $n \equiv a_i \pmod{m_i}$  for some  $i$ .

If the moduli  $m_i$  are distinct we call it a **distinct covering system**.

For example, we can check that the set:

$$\{0 \pmod{2}, 0 \pmod{3}, 1 \pmod{4}, 1 \pmod{6}, 11 \pmod{12}\}$$

forms a distinct covering system, by considering the residues modulo 12:

0 1 2 3 4 5 6 7 8 9 10 11

# Erdős and some conjectures

- Erdős introduced covering systems in 1950, introduced the Minimum Modulus Problem: “Are there distinct covering systems whose smallest modulus exceeds  $M$  for any  $M$ ?”
- Many additional conjectures and problems of Erdős and others.
- Is there a distinct covering system with only odd moduli?  
(Erdős-Selfridge conjecture)

# Exciting Developments

- Minimum Modulus Problem

- [Hough](#) (2015):  $M$  cannot be arbitrarily large.
- [Owens](#) (2014), and [Balister](#), [Bollobás](#), [Morris](#), [Sahasrabudhe](#), [Tiba](#) (2018):

$$42 \leq m_1 < 616000.$$

# Exciting Developments

- Minimum Modulus Problem

- [Hough](#) (2015):  $M$  cannot be arbitrarily large.
- [Owens](#) (2014), and [Balister](#), [Bollobás](#), [Morris](#), [Sahasrabudhe](#), [Tiba](#) (2018):

$$42 \leq m_1 < 616000.$$

- The [Erdős-Selfridge](#) conjecture remains open.

- [Hough](#), [Nielsen](#) (2017) a modulus is divisible by either 2 or 3.
- [Balister](#) et al (2018): a modulus is divisible by 2, 9 or 15.

# Covering Numbers

[Haight](#) (1979), answering an [Erdős](#) problem, defines **covering numbers**: integers whose divisors form the moduli of a distinct covering system.

Since we found a covering system with moduli 2, 3, 4, 6, and 12, we see that 12 is a covering number. In fact it is the smallest covering number.

# Covering Numbers

[Haight](#) (1979), answering an [Erdős](#) problem, defines **covering numbers**: integers whose divisors form the moduli of a distinct covering system.

Since we found a covering system with moduli 2, 3, 4, 6, and 12, we see that 12 is a covering number. In fact it is the smallest covering number.

Any multiple of a covering number is also covering, so call  $n$  a **primitive covering number** if it is a covering number but no proper divisor of  $n$  is.

Primitive covering numbers (A160559): 12, 80, 90, 210, 280, 378, 448, 1386...

# Covering Numbers

[Haight](#) (1979), answering an [Erdős](#) problem, defines **covering numbers**: integers whose divisors form the moduli of a distinct covering system.

Since we found a covering system with moduli 2, 3, 4, 6, and 12, we see that 12 is a covering number. In fact it is the smallest covering number.

Any multiple of a covering number is also covering, so call  $n$  a **primitive covering number** if it is a covering number but no proper divisor of  $n$  is.

Primitive covering numbers (A160559): 12, 80, 90, 210, 280, 378, 448, 1386...

**Erdős-Selfridge Conjecture:** There are no odd covering numbers.

After [Bellobas](#) et al, every covering number is a multiple of 2, 9 or 15.

# Abundant Numbers

Define the “abundancy index”  $h(n) := \frac{\sigma(n)}{n} = \sum_{d|n} \frac{1}{d}$ .



# Abundant Numbers

Define the “abundancy index”  $h(n) := \frac{\sigma(n)}{n} = \sum_{d|n} \frac{1}{d}$ .

A number  $n$  is **abundant** if  $\sigma(n) := \sum_{d|n} d > 2n$

# Abundant Numbers

Define the “abundancy index”  $h(n) := \frac{\sigma(n)}{n} = \sum_{d|n} \frac{1}{d}$ .

A number  $n$  is **abundant** if  $\sigma(n) := \sum_{d|n} d > 2n \iff h(n) > 2$ .

# Abundant Numbers

Define the “abundancy index”  $h(n) := \frac{\sigma(n)}{n} = \sum_{d|n} \frac{1}{d}$ .

A number  $n$  is **abundant** if  $\sigma(n) := \sum_{d|n} d > 2n \iff h(n) > 2$ .

An arithmetic progression mod  $d$  “covers” a set of integers of density  $\frac{1}{d}$ , so a necessary (not sufficient) condition for  $n$  to be a covering number is

$$\sum_{d|n, d>1} \frac{1}{d} \geq 1$$

# Abundant Numbers

Define the “abundancy index”  $h(n) := \frac{\sigma(n)}{n} = \sum_{d|n} \frac{1}{d}$ .

A number  $n$  is **abundant** if  $\sigma(n) := \sum_{d|n} d > 2n \iff h(n) > 2$ .

An arithmetic progression mod  $d$  “covers” a set of integers of density  $\frac{1}{d}$ , so a necessary (not sufficient) condition for  $n$  to be a covering number is

$$\sum_{d|n, d>1} \frac{1}{d} > 1$$

# Abundant Numbers

Define the “abundancy index”  $h(n) := \frac{\sigma(n)}{n} = \sum_{d|n} \frac{1}{d}$ .

A number  $n$  is **abundant** if  $\sigma(n) := \sum_{d|n} d > 2n \iff h(n) > 2$ .

An arithmetic progression mod  $d$  “covers” a set of integers of density  $\frac{1}{d}$ , so a necessary (not sufficient) condition for  $n$  to be a covering number is

$$\sum_{d|n, d>1} \frac{1}{d} > 1 \implies h(n) - 1 > 1$$

# Abundant Numbers

Define the “abundancy index”  $h(n) := \frac{\sigma(n)}{n} = \sum_{d|n} \frac{1}{d}$ .

A number  $n$  is **abundant** if  $\sigma(n) := \sum_{d|n} d > 2n \iff h(n) > 2$ .

An arithmetic progression mod  $d$  “covers” a set of integers of density  $\frac{1}{d}$ , so a necessary (not sufficient) condition for  $n$  to be a covering number is

$$\sum_{d|n, d>1} \frac{1}{d} > 1 \implies h(n) - 1 > 1 \implies h(n) > 2.$$

# Abundant Numbers

Define the “abundancy index”  $h(n) := \frac{\sigma(n)}{n} = \sum_{d|n} \frac{1}{d}$ .

A number  $n$  is **abundant** if  $\sigma(n) := \sum_{d|n} d > 2n \iff h(n) > 2$ .

An arithmetic progression mod  $d$  “covers” a set of integers of density  $\frac{1}{d}$ , so a necessary (not sufficient) condition for  $n$  to be a covering number is

$$\sum_{d|n, d>1} \frac{1}{d} > 1 \implies h(n) - 1 > 1 \implies h(n) > 2.$$

So the covering numbers form a subset of the abundant numbers.

# Abundant Numbers

Define the “abundancy index”  $h(n) := \frac{\sigma(n)}{n} = \sum_{d|n} \frac{1}{d}$ .

A number  $n$  is **abundant** if  $\sigma(n) := \sum_{d|n} d > 2n \iff h(n) > 2$ .

An arithmetic progression mod  $d$  “covers” a set of integers of density  $\frac{1}{d}$ , so a necessary (not sufficient) condition for  $n$  to be a covering number is

$$\sum_{d|n, d>1} \frac{1}{d} > 1 \implies h(n) - 1 > 1 \implies h(n) > 2.$$

So the covering numbers form a subset of the abundant numbers.



# Primitive Covering Numbers

## Theorem (Sun, 2007)

*There are infinitely many primitive covering numbers  $n = 2^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ , where  $\alpha_k = 1$ ,  $\alpha_{k-1} = \left\lfloor \frac{p_k - 1}{p_{k-1} - 1} \right\rfloor$*

$$p_i = 1 + \tau(p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_{i-1}^{\alpha_{i-1}}) \quad (1 < i < k)$$

$$\text{and } p_k \leq \tau(p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_{k-1}^{\alpha_{k-1}}).$$

# Primitive Covering Numbers

## Theorem (Sun, 2007)

*There are infinitely many primitive covering numbers  $n = 2^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ , where  $\alpha_k = 1$ ,  $\alpha_{k-1} = \left\lfloor \frac{p_k - 1}{p_{k-1} - 1} \right\rfloor$*

$$p_i = 1 + \tau(p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_{i-1}^{\alpha_{i-1}}) \quad (1 < i < k)$$

$$\text{and } p_k \leq \tau(p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_{k-1}^{\alpha_{k-1}}).$$

**Conjecture (Sun):** Every primitive covering number has (roughly) this form.

# Primitive Covering Numbers

## Theorem (Sun, 2007)

*There are infinitely many primitive covering numbers  $n = 2^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ , where  $\alpha_k = 1$ ,  $\alpha_{k-1} = \left\lfloor \frac{p_k - 1}{p_{k-1} - 1} \right\rfloor$*

$$p_i = 1 + \tau(p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_{i-1}^{\alpha_{i-1}}) \quad (1 < i < k)$$

$$\text{and } p_k \leq \tau(p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_{k-1}^{\alpha_{k-1}}).$$

**Conjecture** (Sun): Every primitive covering number has (roughly) this form.

Jones, White (2016) find an infinite family of counterexamples  $2^\alpha pq$ .

# Primitive Covering Numbers

## Theorem (Sun, 2007)

*There are infinitely many primitive covering numbers  $n = 2^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$ , where  $\alpha_k = 1$ ,  $\alpha_{k-1} = \left\lfloor \frac{p_k - 1}{p_{k-1} - 1} \right\rfloor$*

$$p_i = 1 + \tau(p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_{i-1}^{\alpha_{i-1}}) \quad (1 < i < k)$$

$$\text{and } p_k \leq \tau(p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_{k-1}^{\alpha_{k-1}}).$$

**Conjecture** (Sun): Every primitive covering number has (roughly) this form.

Jones, White (2016) find an infinite family of counterexamples  $2^\alpha pq$ .

Harrington, Jones, Phillips (2017) find infinitely many more of form  $2^\alpha p^\beta q$ .

# Covering Numbers

## Question (Phillips)

Do the covering numbers have a density?

# Covering Numbers

## Question (Phillips)

Do the covering numbers have a density?

If so, can this density be estimated?

# Notation Summary

- $\tau(n)$ : number of divisors,
- $P^+(n), P^-(n)$ : largest, smallest prime factor
- $\mathcal{A}, \mathcal{C}$ , sets of abundant and covering numbers
- $\mathcal{P}_{\mathcal{A}}, \mathcal{P}_{\mathcal{C}}$  their primitive subsets
- $\mathcal{S}(x) := \#\{n \leq x \mid n \in \mathcal{S}\}$  the counting function of the set  $\mathcal{S}$ .
- $d(\mathcal{S}) := \lim_{x \rightarrow \infty} \frac{\mathcal{S}(x)}{x}$  denotes the natural density of  $\mathcal{S}$  if it exists.

# The Density of Abundant Numbers

It isn't immediately obvious the density of abundant numbers exists.



# The Density of Abundant Numbers

It isn't immediately obvious the density of abundant numbers exists.

- Behrend (1932) showed that  $\frac{\mathcal{A}(n)}{n} < 0.47$  for all integers  $n$ .

# The Density of Abundant Numbers

It isn't immediately obvious the density of abundant numbers exists.

- Behrend (1932) showed that  $\frac{A(n)}{n} < 0.47$  for all integers  $n$ .
- Davenport (1933) proved the abundant numbers have a density.

# The Density of Abundant Numbers

It isn't immediately obvious the density of abundant numbers exists.

- Behrend (1932) showed that  $\frac{A(n)}{n} < 0.47$  for all integers  $n$ .
- Davenport (1933) proved the abundant numbers have a density.
- Erdős (1934): The sum  $\sum_{a \in \mathcal{P}_A} \frac{1}{a}$  converges, giving an elementary proof of the existence of the density of abundant numbers.

# The Density of Abundant Numbers

It isn't immediately obvious the density of abundant numbers exists.

- [Behrend](#) (1932) showed that  $\frac{A(n)}{n} < 0.47$  for all integers  $n$ .
- [Davenport](#) (1933) proved the abundant numbers have a density.
- [Erdős](#) (1934): The sum  $\sum_{a \in \mathcal{P}_A} \frac{1}{a}$  converges, giving an elementary proof of the existence of the density of abundant numbers.

## Subsequent improvements

Year	Author	Lower Bound	Upper Bound
1933	<a href="#">Behrend</a>	0.241	0.314

# The Density of Abundant Numbers

It isn't immediately obvious the density of abundant numbers exists.

- Behrend (1932) showed that  $\frac{A(n)}{n} < 0.47$  for all integers  $n$ .
- Davenport (1933) proved the abundant numbers have a density.
- Erdős (1934): The sum  $\sum_{a \in \mathcal{P}_A} \frac{1}{a}$  converges, giving an elementary proof of the existence of the density of abundant numbers.

## Subsequent improvements

Year	Author	Lower Bound	Upper Bound
1933	Behrend	0.241	0.314
1955	Salié	0.246	

# The Density of Abundant Numbers

It isn't immediately obvious the density of abundant numbers exists.

- Behrend (1932) showed that  $\frac{A(n)}{n} < 0.47$  for all integers  $n$ .
- Davenport (1933) proved the abundant numbers have a density.
- Erdős (1934): The sum  $\sum_{a \in \mathcal{P}_A} \frac{1}{a}$  converges, giving an elementary proof of the existence of the density of abundant numbers.

## Subsequent improvements

Year	Author	Lower Bound	Upper Bound
1933	Behrend	0.241	0.314
1955	Salié	0.246	
1971-72	Wall et al.	0.2441	0.2909

# The Density of Abundant Numbers

It isn't immediately obvious the density of abundant numbers exists.

- Behrend (1932) showed that  $\frac{A(n)}{n} < 0.47$  for all integers  $n$ .
- Davenport (1933) proved the abundant numbers have a density.
- Erdős (1934): The sum  $\sum_{a \in \mathcal{P}_A} \frac{1}{a}$  converges, giving an elementary proof of the existence of the density of abundant numbers.

## Subsequent improvements

Year	Author	Lower Bound	Upper Bound
1933	Behrend	0.241	0.314
1955	Salié	0.246	
1971-72	Wall et al.	0.2441	0.2909
1998	Deléglise	0.2474	0.2480

# The Density of Abundant Numbers

It isn't immediately obvious the density of abundant numbers exists.

- Behrend (1932) showed that  $\frac{A(n)}{n} < 0.47$  for all integers  $n$ .
- Davenport (1933) proved the abundant numbers have a density.
- Erdős (1934): The sum  $\sum_{a \in \mathcal{P}_A} \frac{1}{a}$  converges, giving an elementary proof of the existence of the density of abundant numbers.

## Subsequent improvements

Year	Author	Lower Bound	Upper Bound
1933	Behrend	0.241	0.314
1955	Salié	0.246	
1971-72	Wall et al.	0.2441	0.2909
1998	Deléglise	0.2474	0.2480
2010	Kobayashi	0.2476171	0.2476475



# Covering Numbers

## Question (Phillips)

Do the covering numbers have a density?

If so, can this density be estimated?

# A key lemma

Lemma (M.)

*If  $n$  is a primitive covering number then  $P^+(n) < \tau(n)$ .*

# A key lemma

Lemma (M.)

*If  $n$  is a primitive covering number then  $P^+(n) < \tau(n)$ .*

This inequality,  $P^+(n) < \tau(n)$  doesn't happen too frequently.

# A key lemma

## Lemma (M.)

*If  $n$  is a primitive covering number then  $P^+(n) < \tau(n)$ .*

This inequality,  $P^+(n) < \tau(n)$  doesn't happen too frequently.

On average,  $\frac{1}{x} \sum_{n \leq x} \tau(n) \approx \log x$ , while  $\frac{1}{x} \sum_{n \leq x} P^+(n) \approx \frac{\pi^2}{12} \frac{x}{\log x}$ .

# Bounding Primitive Covering Numbers

Key idea: Numbers typically have large prime factors and few divisors.

# Bounding Primitive Covering Numbers

Key idea: Numbers typically have large prime factors and few divisors.

- **Smooth numbers:**  $\psi(x, y)$  counts integers  $\leq x$  with  $P^+(n) \leq y$

$$\psi(x, y) = x \exp \left( (-1 + o(1)) \frac{\log x}{\log y} \log \frac{\log x}{\log y} \right)$$

Valid when  $x \geq y \geq \exp((\log \log x)^{5/3+\epsilon})$  ([Hildebrand](#)).

# Bounding Primitive Covering Numbers

Key idea: Numbers typically have large prime factors and few divisors.

- **Smooth numbers:**  $\psi(x, y)$  counts integers  $\leq x$  with  $P^+(n) \leq y$

$$\psi(x, y) = x \exp \left( (-1 + o(1)) \frac{\log x}{\log y} \log \frac{\log x}{\log y} \right)$$

Valid when  $x \geq y \geq \exp((\log \log x)^{5/3+\epsilon})$  ([Hildebrand](#)).

- **Many divisors:**  $\Delta(x, y)$  counts integers  $\leq x$  with  $\tau(n) > y$

$$\Delta(x, y) = x \exp \left( (-1 + o(1)) \frac{\log y}{\log 2} \log \log y \right)$$

Valid when  $x \geq y > (\log x)^{2 \log 2 + \epsilon}$  ([Norton](#)).

# Primitive Covering Numbers

$$\text{So } \mathcal{P}_C(x) \leq \psi(x, y) + \Delta(x, y) \approx \frac{x}{\exp\left(\frac{\log x}{\log y} \log \frac{\log x}{\log y}\right)} + \frac{x}{\exp\left(\frac{\log y}{\log 2} \log \log y\right)}.$$



# Primitive Covering Numbers

$$\text{So } \mathcal{P}_C(x) \leq \psi(x, y) + \Delta(x, y) \approx \frac{x}{\exp\left(\frac{\log x}{\log y} \log \frac{\log x}{\log y}\right)} + \frac{x}{\exp\left(\frac{\log y}{\log 2} \log \log y\right)}.$$

Pick  $y = \exp(\sqrt{\log 2 \log x})$ .

# Primitive Covering Numbers

$$\text{So } \mathcal{P}_C(x) \leq \psi(x, y) + \Delta(x, y) \approx \frac{x}{\exp\left(\frac{\log x}{\log y} \log \frac{\log x}{\log y}\right)} + \frac{x}{\exp\left(\frac{\log y}{\log 2} \log \log y\right)}.$$

Pick  $y = \exp(\sqrt{\log 2 \log x})$ .

# Primitive Covering Numbers

$$\text{So } \mathcal{P}_C(x) \leq \psi(x, y) + \Delta(x, y) \approx \frac{x}{\exp\left(\frac{\log x}{\log y} \log \frac{\log x}{\log y}\right)} + \frac{x}{\exp\left(\frac{\log y}{\log 2} \log \log y\right)}.$$

Pick  $y = \exp(\sqrt{\log 2 \log x})$ .

## Theorem

$$\mathcal{P}_C(x) \ll x \exp\left(\left(-\frac{1}{2\sqrt{\log 2}} + \epsilon\right) \sqrt{\log x} \log \log x\right)$$

# Primitive Covering Numbers

$$\text{So } \mathcal{P}_C(x) \leq \psi(x, y) + \Delta(x, y) \approx \frac{x}{\exp\left(\frac{\log x}{\log y} \log \frac{\log x}{\log y}\right)} + \frac{x}{\exp\left(\frac{\log y}{\log 2} \log \log y\right)}.$$

Pick  $y = \exp(\sqrt{\log 2 \log x})$ .

## Theorem

$$\mathcal{P}_C(x) \ll x \exp\left(\left(-\frac{1}{2\sqrt{\log 2}} + \epsilon\right) \sqrt{\log x} \log \log x\right)$$

## Corollary

- *The reciprocal sum  $\sum_{c \in \mathcal{P}_C} \frac{1}{c}$  converges.*

# Primitive Covering Numbers

$$\text{So } \mathcal{P}_C(x) \leq \psi(x, y) + \Delta(x, y) \approx \frac{x}{\exp\left(\frac{\log x}{\log y} \log \frac{\log x}{\log y}\right)} + \frac{x}{\exp\left(\frac{\log y}{\log 2} \log \log y\right)}.$$

Pick  $y = \exp(\sqrt{\log 2 \log x})$ .

## Theorem

$$\mathcal{P}_C(x) \ll x \exp\left(\left(-\frac{1}{2\sqrt{\log 2}} + \epsilon\right) \sqrt{\log x} \log \log x\right)$$

## Corollary

- *The reciprocal sum  $\sum_{c \in \mathcal{P}_C} \frac{1}{c}$  converges.*
- *The density  $d(\mathcal{C})$  of covering numbers exists.*

# Bounding $d(\mathcal{A})$ - the Behrend–Deléglise Approach

Define  $A_y(x) = d(\{n : P^-(n) > y, h(n) > x\})$

# Bounding $d(\mathcal{A})$ - the Behrend–Deléglise Approach

Define  $A_y(x) = d(\{n : P^-(n) > y, h(n) > x\})$

Behrend's goal: Bound the density  $d(\mathcal{A}) = A_1(2)$ .

# Bounding $d(\mathcal{A})$ - the Behrend–Deléglise Approach

Define  $A_y(x) = d(\{n : P^-(n) > y, h(n) > x\})$

Behrend's goal: Bound the density  $d(\mathcal{A}) = A_1(2)$ .

Fix  $y$ , each  $n$  factors uniquely as  $n = sr$  with  $P^+(s) \leq y$  and  $P^-(r) > y$ .



# Bounding $d(\mathcal{A})$ - the Behrend–Deléglise Approach

Define  $A_y(x) = d(\{n : P^-(n) > y, h(n) > x\})$

Behrend's goal: Bound the density  $d(\mathcal{A}) = A_1(2)$ .

Fix  $y$ , each  $n$  factors uniquely as  $n = sr$  with  $P^+(s) \leq y$  and  $P^-(r) > y$ .

If  $h(n) \geq 2$  then (since  $h$  is multiplicative),  $h(s)h(r) \geq 2$ , so  $h(r) \geq \frac{2}{h(s)}$ .

$$d(\mathcal{A}) = \sum_{P^+(s) \leq y} \frac{1}{s} A_y\left(\frac{2}{h(s)}\right)$$

# Bounding $d(\mathcal{A})$ - the Behrend–Deléglise Approach

Define  $A_y(x) = d(\{n : P^-(n) > y, h(n) > x\})$

Behrend's goal: Bound the density  $d(\mathcal{A}) = A_1(2)$ .

Fix  $y$ , each  $n$  factors uniquely as  $n = sr$  with  $P^+(s) \leq y$  and  $P^-(r) > y$ .

If  $h(n) \geq 2$  then (since  $h$  is multiplicative),  $h(s)h(r) \geq 2$ , so  $h(r) \geq \frac{2}{h(s)}$ .

$$d(\mathcal{A}) = \sum_{P^+(s) \leq y} \frac{1}{s} A_y\left(\frac{2}{h(s)}\right) = \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y\left(\frac{2}{h(s)}\right) + \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s} A_y\left(\frac{2}{h(s)}\right)$$

# Bounding these sums

$$d(\mathcal{A}) = \sum_{\substack{s \leq z \\ P^+(\bar{s}) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right) + \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right)$$

# Bounding these sums

$$d(\mathcal{A}) = \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right) + \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right)$$

Bound the tail sum trivially:  $\sum_{\substack{n > z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right) \leq \left( \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \right) \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s}$

# Bounding these sums

$$d(\mathcal{A}) = \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right) + \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right)$$

Bound the tail sum trivially: 
$$\sum_{\substack{n > z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right) \leq \left( \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \right) \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s}$$

Bound the  $A_y(x)$  in the first sum using moments of  $h_y(n) := \sum_{d|n} \frac{1}{d}$ .

Set  $\mu_{y,r} := \lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} h_y(n)^r$  then  $A_y(x) \leq \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \frac{\mu_{y,r} - 1}{x^r - 1}$

# Bounding these sums

$$d(\mathcal{A}) = \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right) + \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right)$$

Bound the tail sum trivially: 
$$\sum_{\substack{n > z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right) \leq \left( \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \right) \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s}$$

Bound the  $A_y(x)$  in the first sum using moments of  $h_y(n) := \sum_{\substack{d|n \\ P^-(d) > y}} \frac{1}{d}$ .

Set  $\mu_{y,r} := \lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} h_y(n)^r$  then  $A_y(x) \leq \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \frac{\mu_{y,r} - 1}{x^r - 1}$  Pick  $r$  to give the best bound.

# Bounding these sums

$$d(\mathcal{A}) = \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right) + \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right)$$

Bound the tail sum trivially: 
$$\sum_{\substack{n > z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{h(s)} \right) \leq \left( \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \right) \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s}$$

Bound the  $A_y(x)$  in the first sum using moments of  $h_y(n) := \sum_{\substack{d|n \\ P^-(d) > y}} \frac{1}{d}$ .

Set  $\mu_{y,r} := \lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} h_y(n)^r$  then  $A_y(x) \leq \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \frac{\mu_{y,r} - 1}{x^r - 1}$  Pick  $r$  to give the best bound.

Deléglise takes  $y = 500$ ,  $z = 10^{14}$  to get  $d(\mathcal{A}) < 0.2480$ .

# Bounding $d(\mathcal{C})$

Since  $\mathcal{C} \subset \mathcal{A}$ ,  $d(\mathcal{C}) \leq d(\mathcal{A}) < 0.24765$ . ([Kobayashi](#))



# Bounding $d(\mathcal{C})$

Since  $\mathcal{C} \subset \mathcal{A}$ ,  $d(\mathcal{C}) \leq d(\mathcal{A}) < 0.24765$ . ([Kobayashi](#))

To improve this, we need an analogue of the multiplicative function  $h(n)$ .

# Bounding $d(\mathcal{C})$

Since  $\mathcal{C} \subset \mathcal{A}$ ,  $d(\mathcal{C}) \leq d(\mathcal{A}) < 0.24765$ . ([Kobayashi](#))

To improve this, we need an analogue of the multiplicative function  $h(n)$ .

Let  $r(n)$  count the maximum number of residues modulo  $n$  that can be covered by distinct congruences mod  $d$ , with  $d \mid n$ ,  $d > 1$ .

# Bounding $d(\mathcal{C})$

Since  $\mathcal{C} \subset \mathcal{A}$ ,  $d(\mathcal{C}) \leq d(\mathcal{A}) < 0.24765$ . ([Kobayashi](#))

To improve this, we need an analogue of the multiplicative function  $h(n)$ .

Let  $r(n)$  count the maximum number of residues modulo  $n$  that can be covered by distinct congruences mod  $d$ , with  $d \mid n$ ,  $d > 1$ .

Now, define the “covering index” 
$$c(n) := 1 + \frac{r(n)}{n}$$

# Bounding $d(\mathcal{C})$

Since  $\mathcal{C} \subset \mathcal{A}$ ,  $d(\mathcal{C}) \leq d(\mathcal{A}) < 0.24765$ . (Kobayashi)

To improve this, we need an analogue of the multiplicative function  $h(n)$ .

Let  $r(n)$  count the maximum number of residues modulo  $n$  that can be covered by distinct congruences mod  $d$ , with  $d \mid n$ ,  $d > 1$ .

Now, define the “covering index” 
$$c(n) := 1 + \frac{r(n)}{n}$$

The integer  $n$  is a covering number  $\iff r(n) = n \iff c(n) = 2$ .

# Bounding $d(\mathcal{C})$ with $c(n)$

**Lemma:** For all  $n$ ,  $c(n) \leq h(n)$ .

# Bounding $d(\mathcal{C})$ with $c(n)$

**Lemma:** For all  $n$ ,  $c(n) \leq h(n)$ .

Unlike  $h(n)$ ,  $c(n)$  is not multiplicative.

# Bounding $d(\mathcal{C})$ with $c(n)$

**Lemma:** For all  $n$ ,  $c(n) \leq h(n)$ .

Unlike  $h(n)$ ,  $c(n)$  is not multiplicative.

## Theorem

*If  $\gcd(m, n) = 1$ , then  $c(mn) \leq c(m)h(n)$ .*

# Bounding $d(\mathcal{C})$ with $c(n)$

**Lemma:** For all  $n$ ,  $c(n) \leq h(n)$ .

Unlike  $h(n)$ ,  $c(n)$  is not multiplicative.

## Theorem

If  $\gcd(m, n) = 1$ , then  $c(mn) \leq c(m)h(n)$ .

Since  $c(mn) = 2 \iff h(m) \geq \frac{2}{c(n)}$ ,

$$d(\mathcal{C}) < \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y\left(\frac{2}{c(s)}\right) + \left( \prod_{p \leq y} \left(1 - \frac{1}{p}\right) \right) \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s}.$$



# Bounding $d(\mathcal{C})$ with $c(n)$

**Lemma:** For all  $n$ ,  $c(n) \leq h(n)$ .

Unlike  $h(n)$ ,  $c(n)$  is not multiplicative.

## Theorem

If  $\gcd(m, n) = 1$ , then  $c(mn) \leq c(m)h(n)$ .

Since  $c(mn) = 2 \iff h(m) \geq \frac{2}{c(n)}$ ,

$$d(\mathcal{C}) < \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y\left(\frac{2}{c(s)}\right) + \left( \prod_{p \leq y} \left(1 - \frac{1}{p}\right) \right) \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s}.$$

Unfortunately, computing  $c(n)$  is **very** computationally intensive...

# An Inclusion/Exclusion

# An Inclusion/Exclusion

Let  $S_2(i, j)$  be Stirling numbers of the 2<sup>nd</sup> kind,  $B_{r,d} := -\sum_{k=1}^d (-r)^k S_2(d, k)$ .

# An Inclusion/Exclusion

Let  $S_2(i, j)$  be Stirling numbers of the 2<sup>nd</sup> kind,  $B_{r,d} := -\sum_{k=1}^d (-r)^k S_2(d, k)$ .  
( $B_{1,d}$  is the negative of the sequence of complementary Bell numbers.)

# An Inclusion/Exclusion

Let  $S_2(i, j)$  be Stirling numbers of the 2<sup>nd</sup> kind,  $B_{r,d} := -\sum_{k=1}^d (-r)^k S_2(d, k)$ .  
( $B_{1,d}$  is the negative of the sequence of complementary Bell numbers.)

## Theorem

*If  $n = \ell m$  with  $(\ell, m) = 1$  and  $r(\ell) = \ell - 1$  then*

$$c(n) \leq 1 + \frac{\ell - 1}{\ell} + \frac{1}{\ell} \sum_{d|m, d>1} \frac{B_{\tau(\ell), \omega(d)}}{d} =: c'(n).$$

# Using $c'(n)$ to bound the density

$$d(\mathcal{C}) < \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{c'(s)} \right) + \left( \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \right) \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s}$$

# Using $c'(n)$ to bound the density

$$d(\mathcal{C}) < \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{c'(s)} \right) + \left( \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \right) \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s}$$

$c'(s)$  is far more efficient to compute than  $c(s)$ , but not *that* efficient.

# Using $c'(n)$ to bound the density

$$d(\mathcal{C}) < \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{c'(s)} \right) + \left( \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \right) \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s}$$

$c'(s)$  is far more efficient to compute than  $c(s)$ , but not *that* efficient.  
No hope of computing this sum with  $y = 500$ ,  $z = 10^{14}$  like Deléglise.



# Using $c'(n)$ to bound the density

$$d(\mathcal{C}) < \sum_{\substack{s \leq z \\ P^+(s) \leq y}} \frac{1}{s} A_y \left( \frac{2}{c'(s)} \right) + \left( \prod_{p \leq y} \left( 1 - \frac{1}{p} \right) \right) \sum_{\substack{s > z \\ P^+(s) \leq y}} \frac{1}{s}$$

$c'(s)$  is far more efficient to compute than  $c(s)$ , but not *that* efficient. No hope of computing this sum with  $y = 500$ ,  $z = 10^{14}$  like Deléglise. With  $y = 200$ ,  $z = 10^9$  we found:

$$0.1032 \leq d(\mathcal{C}) \leq 0.1197$$

# A more flexible version of Deléglise's approach

Code to compute this sum spends a long time handling  $s$  coprime to 6.

# A more flexible version of Deléglise's approach

Code to compute this sum spends a long time handling  $s$  coprime to 6.

We know there are no such covering numbers.

# A more flexible version of Deléglise's approach

Code to compute this sum spends a long time handling  $s$  coprime to 6.

We know there are no such covering numbers.

It also spends a while on odd multiples of 3.

# A more flexible version of Deléglise's approach

Code to compute this sum spends a long time handling  $s$  coprime to 6.

We know there are no such covering numbers.

It also spends a while on odd multiples of 3.

The density of such covering numbers is tiny.

# A more flexible version of Deléglise's approach

Code to compute this sum spends a long time handling  $s$  coprime to 6.

We know there are no such covering numbers.

It also spends a while on odd multiples of 3.

The density of such covering numbers is tiny.

Do we need to treat all the numbers  $s$  in a “large box”  $P^+(s) < y$ ,  $s < z$ ?

# A more flexible version of Deléglise's approach

Code to compute this sum spends a long time handling  $s$  coprime to 6.

We know there are no such covering numbers.

It also spends a while on odd multiples of 3.

The density of such covering numbers is tiny.

Do we need to treat all the numbers  $s$  in a “large box”  $P^+(s) < y$ ,  $s < z$ ?

We'd rather focus on numbers divisible by large powers of 2...

# A more flexible version of Deléglise's approach

Code to compute this sum spends a long time handling  $s$  coprime to 6.

We know there are no such covering numbers.

It also spends a while on odd multiples of 3.

The density of such covering numbers is tiny.

Do we need to treat all the numbers  $s$  in a “large box”  $P^+(s) < y$ ,  $s < z$ ?

We'd rather focus on numbers divisible by large powers of 2...



# Smooth-Rough Divisor Partition

## Definition

For a pair  $(a, q)$  with  $P^+(a) \leq q$ , denote by  $M_{a,q} = \{ar : P^-(r) \geq q\}$ .

A set  $W$  of such pairs is a **smooth-rough divisor partition** of  $\mathbb{N}$  if:

- $M_{a,q}$  are disjoint,
- $\mathbb{N} = \bigsqcup_{(a,q) \in W} M_{a,q}$ ,

# Smooth-Rough Divisor Partition

## Definition

For a pair  $(a, q)$  with  $P^+(a) \leq q$ , denote by  $M_{a,q} = \{ar : P^-(r) \geq q\}$ .

A set  $W$  of such pairs is a **smooth-rough divisor partition** of  $\mathbb{N}$  if:

- $M_{a,q}$  are disjoint,
- $\mathbb{N} = \bigsqcup_{(a,q) \in W} M_{a,q}$ ,

Further partition  $W = W_{<} \sqcup W_{=}$

$$W_{<} := \{(a, q) \in W : P^+(a) < q\}$$

$$W_{=} := \{(a, q) \in W : P^+(a) = q\}$$

# Smooth-Rough-Divisor Partitions

Example:

$$W = \{(1, 5), (2, 3), (3, 3), (4, 2)\}$$

$$W_{<} = \{(1, 5), (2, 3)\},$$

$$W_{=} = \{(3, 3), (4, 2)\}$$

For any smooth-rough divisor partition  $W$ ,

$$d(\mathcal{C}) = \sum_{(a,q) \in W_{<}} \frac{1}{a} A_q \left( \frac{2}{c'(a)} \right) + \sum_{(a,q) \in W_{=}} \frac{1}{a(1 - 1/q)} A_{q+1} \left( \frac{2}{c''(a, q)} \right)$$

# Computation

Unclear how to pick the optimal smooth-rough divisor partition  $W$ .

# Computation

Unclear how to pick the optimal smooth-rough divisor partition  $W$ .

After much numerical experimentation and several hours of computation...

## Theorem

$$0.10324 < d(\mathcal{C}) < 0.10356$$

Lower bound from the primitive covering numbers.

# Computation

Unclear how to pick the optimal smooth-rough divisor partition  $W$ .

After much numerical experimentation and several hours of computation...

## Theorem

$$0.10324 < d(C) < 0.10356$$

Lower bound from the primitive covering numbers. First use the infinite families identified by [Sun](#), et al.

# Computation

Unclear how to pick the optimal smooth-rough divisor partition  $W$ .

After much numerical experimentation and several hours of computation...

## Theorem

$$0.10324 < d(C) < 0.10356$$

Lower bound from the primitive covering numbers. First use the infinite families identified by [Sun](#), et al. Then use integer programming (Gurobi) and tricks with almost-covering numbers to find “sporadic” examples.

# Computation

Unclear how to pick the optimal smooth-rough divisor partition  $W$ .

After much numerical experimentation and several hours of computation...

## Theorem

$$0.10324 < d(\mathcal{C}) < 0.10356$$

Lower bound from the primitive covering numbers. First use the infinite families identified by [Sun](#), et al. Then use integer programming (Gurobi) and tricks with almost-covering numbers to find “sporadic” examples.

Identify all primitive covering numbers  $< 773\,500 = 2^2 \times 5^3 \times 7 \times 13 \times 17$ .



# Abundant Numbers

The smooth-rough divisor partitions were useful for bounding  $d(\mathcal{C})$ .

# Abundant Numbers

The smooth-rough divisor partitions were useful for bounding  $d(\mathcal{C})$ .  
Can we use them to bound  $d(\mathcal{A})$ ?

# Abundant Numbers

The smooth-rough divisor partitions were useful for bounding  $d(\mathcal{C})$ .  
Can we use them to bound  $d(\mathcal{A})$ ? Yes!

# Abundant Numbers

The smooth-rough divisor partitions were useful for bounding  $d(\mathcal{C})$ .

Can we use them to bound  $d(\mathcal{A})$ ? Yes!

Recall, [Kobayashi](#) (2010):  $0.2476171 < d(\mathcal{A}) < 0.2476475$ .

# Abundant Numbers

The smooth-rough divisor partitions were useful for bounding  $d(\mathcal{C})$ .

Can we use them to bound  $d(\mathcal{A})$ ? Yes!


Recall, [Kobayashi](#) (2010):  $0.2476171 < d(\mathcal{A}) < 0.2476475$ .

Using smooth-rough divisor partitions and 3 minutes of runtime we find

$$0.2476127 < d(\mathcal{A}) < 0.2476379.$$


# Abundant density bounds

Lower	Upper	Time
0.2476127	0.2476379	3m07s
0.2476158	0.2476299	13m08s
0.2476175	0.2476253	56m56s
0.2476185	0.2476226	4h54m59s
0.2476190	0.2476214	26h38m12s
0.24761929	0.24762053	6d05h14m31s
0.24761940	0.24762022	



# Abundant density bounds

Lower	Upper	Time
0.2476127	0.2476379	3m07s
0.2476158	0.2476299	13m08s
0.2476175	0.2476253	56m56s
0.2476185	0.2476226	4h54m59s
0.2476190	0.2476214	26h38m12s
0.24761929	0.24762053	6d05h14m31s
0.24761940	0.24762022	



Further optimization and about two weeks of distributed computation gives:

Theorem

$$0.24761951 < d(\mathcal{A}) < 0.24761989$$

# THANK YOU!