

# The distribution of intermediate prime factors

Nathan McNew  
Towson University

Joint work with Paul Pollack and Akash Singha Roy (UGA)

33èmes Journées Arithmétiques  
University of Luxembourg  
July 3rd, 2025

# Extreme prime factors

The study of the largest and smallest prime factors of integers has a rich history, motivated by the analysis of factoring algorithm run times.

# Extreme prime factors

The study of the largest and smallest prime factors of integers has a rich history, motivated by the analysis of factoring algorithm run times.

Let  $P^+(n)$  and  $P^-(n)$  denote the largest and smallest prime factors of  $n$ .

# Extreme prime factors

The study of the largest and smallest prime factors of integers has a rich history, motivated by the analysis of factoring algorithm run times.

Let  $P^+(n)$  and  $P^-(n)$  denote the largest and smallest prime factors of  $n$ .

What can we say about the distribution of these functions for  $n \in [2, x]$ ?

# Extreme prime factors

The study of the largest and smallest prime factors of integers has a rich history, motivated by the analysis of factoring algorithm run times.

Let  $P^+(n)$  and  $P^-(n)$  denote the largest and smallest prime factors of  $n$ .

What can we say about the distribution of these functions for  $n \in [2, x]$ ?

These questions are closely related to smooth/rough numbers: An integer is  $y$ -smooth (rough) if all of its prime factors are no larger (smaller) than  $y$ .

# The distribution of the largest prime factor

**Mean value of  $P^+(n)$ ,  $n \in [2, x]$**

# The distribution of the largest prime factor

**Mean value of  $P^+(n)$ ,  $n \in [2, x]$**

$$\frac{1}{x} \sum_{2 \leq n \leq x} P^+(n) \sim \frac{\pi^2 x}{12 \log x} \quad (\text{Knuth, Pardo, 1976,})$$

# The distribution of the largest prime factor

**Mean value of  $P^+(n)$ ,  $n \in [2, x]$**

$$\frac{1}{x} \sum_{2 \leq n \leq x} P^+(n) \sim \frac{\pi^2 x}{12 \log x} \quad (\text{Knuth, Pardo, 1976, Erdős, Alladi, 1977, Kemeny, 1993})$$



# The distribution of the largest prime factor

**Mean value of  $P^+(n)$ ,  $n \in [2, x]$**

$$\frac{1}{x} \sum_{2 \leq n \leq x} P^+(n) \sim \frac{\pi^2 x}{12 \log x} \quad (\text{Knuth, Pardo, 1976, Erdős, Alladi, 1977, Kemeny, 1993})$$

In fact

$$\frac{\pi^2 x}{12 \log x} + \frac{d_2 x}{\log^2 x} + \dots + \frac{d_m x}{\log^m x} + O\left(\frac{x}{\log^{m+1} x}\right) \quad (\text{De Koninck and Ivić, 1984})$$

Uniformly for all  $m$ .

$$d_m = \frac{1}{2^{m+1}} \sum_{j=0}^m \frac{(-2)^j \zeta^{(j)}(2)}{j!}. \quad (\text{Naslund, 2013})$$

# Largest Prime Factor

**Mean value:**

$$\frac{\pi^2 x}{12 \log x} + \frac{d_2 x}{\log^2 x} + \dots + \frac{d_m x}{\log^m x} + O\left(\frac{x}{\log^{m+1} x}\right)$$

# Largest Prime Factor

**Mean value:**  $\frac{\pi^2 x}{12 \log x} + \frac{d_2 x}{\log^2 x} + \dots + \frac{d_m x}{\log^m x} + O\left(\frac{x}{\log^{m+1} x}\right)$

**Median value:**

# Largest Prime Factor

**Mean value:**

$$\frac{\pi^2 x}{12 \log x} + \frac{d_2 x}{\log^2 x} + \dots + \frac{d_m x}{\log^m x} + O\left(\frac{x}{\log^{m+1} x}\right)$$

**Median value:**

$$x^{\frac{1}{\sqrt{e}} + o(1)}$$

(Selfridge and Wunderlich, 1974)

# Largest Prime Factor

**Mean value:**

$$\frac{\pi^2 x}{12 \log x} + \frac{d_2 x}{\log^2 x} + \dots + \frac{d_m x}{\log^m x} + O\left(\frac{x}{\log^{m+1} x}\right)$$

**Median value:**

$$x^{\frac{1}{\sqrt{e}} + o(1)}$$

(Selfridge and Wunderlich, 1974)

$$e^{\frac{\gamma-1}{\sqrt{e}}} x^{\frac{1}{\sqrt{e}}} \left(1 + O\left(\frac{1}{\log x}\right)\right)$$

(Naslund, 2013)

# Largest Prime Factor

**Mean value:**

$$\frac{\pi^2 x}{12 \log x} + \frac{d_2 x}{\log^2 x} + \dots + \frac{d_m x}{\log^m x} + O\left(\frac{x}{\log^{m+1} x}\right)$$

**Median value:**

$$x^{\frac{1}{\sqrt{e}} + o(1)}$$

(Selfridge and Wunderlich, 1974)

$$e^{\frac{\gamma-1}{\sqrt{e}}} x^{\frac{1}{\sqrt{e}}} \left(1 + O\left(\frac{1}{\log x}\right)\right)$$

(Naslund, 2013)

**Mode:**

# Largest Prime Factor

**Mean value:**

$$\frac{\pi^2 x}{12 \log x} + \frac{d_2 x}{\log^2 x} + \dots + \frac{d_m x}{\log^m x} + O\left(\frac{x}{\log^{m+1} x}\right)$$

**Median value:**

$$x^{\frac{1}{\sqrt{e}} + o(1)}$$

(Selfridge and Wunderlich, 1974)

$$e^{\frac{\gamma-1}{\sqrt{e}}} x^{\frac{1}{\sqrt{e}}} \left(1 + O\left(\frac{1}{\log x}\right)\right)$$

(Naslund, 2013)

**Mode:**

$$e^{\sqrt{\frac{1}{2} \log x (\log_2 x + \log_3 x)} + O\left(\sqrt{\frac{\log x}{\log_2 x}}\right)}$$

(De Koninck, 1994)

# Largest Prime Factor

**Mean value:**  $\frac{\pi^2 x}{12 \log x} + \frac{d_2 x}{\log^2 x} + \dots + \frac{d_m x}{\log^m x} + O\left(\frac{x}{\log^{m+1} x}\right)$

**Median value:**  $x^{\frac{1}{\sqrt{e}} + o(1)}$  (Selfridge and Wunderlich, 1974)

$$e^{\frac{\gamma-1}{\sqrt{e}}} x^{\frac{1}{\sqrt{e}}} \left(1 + O\left(\frac{1}{\log x}\right)\right) \quad (\text{Naslund, 2013})$$

**Mode:**  $e^{\sqrt{\frac{1}{2} \log x (\log_2 x + \log_3 x)} + O\left(\sqrt{\frac{\log x}{\log_2 x}}\right)}$  (De Koninck, 1994)

$$\exp\left(\sqrt{\nu(x) \log x} + \frac{1}{4} - \frac{\nu(x)-3}{8\nu(x)^2-12\nu(x)+4}\right) \left(1 + O\left(\left(\frac{\log \log x}{\log x}\right)^{1/4}\right)\right) \quad (\text{M, 2015})$$

where  $\nu(x)$  is defined implicitly as the solution to  $e^{\nu(x)} = 1 + \sqrt{\nu(x) \log x} - \nu(x)$ .



# Smallest Prime Factor

Mean value:

$$\frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{x}{2 \log x}$$

# Smallest Prime Factor

Mean value:

$$\frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{x}{2 \log x} \sim \frac{1}{x} \sum_{p \leq x} p$$

# Smallest Prime Factor

**Mean value:**

$$\frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{x}{2 \log x} \sim \frac{1}{x} \sum_{p \leq x} p$$

**Median:**

# Smallest Prime Factor

**Mean value:**

$$\frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{x}{2 \log x} \sim \frac{1}{x} \sum_{p \leq x} p$$

**Median:** 2

# Smallest Prime Factor

**Mean value:**

$$\frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{x}{2 \log x} \sim \frac{1}{x} \sum_{p \leq x} p$$

**Median:** 2 (or 2.5)

# Smallest Prime Factor

**Mean value:**

$$\frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{x}{2 \log x} \sim \frac{1}{x} \sum_{p \leq x} p$$

**Median:** 2 (or 2.5)

**Mode:**

# Smallest Prime Factor

**Mean value:**

$$\frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{x}{2 \log x} \sim \frac{1}{x} \sum_{p \leq x} p$$

**Median:** 2 (or 2.5)

**Mode:** 2

# Average size

**Average size:** Knuth and Trabb Pardo (1976) consider the average size of the largest prime factors as part of an analysis of factoring algorithm runtimes. (Size of  $n$  is  $\log n$ .)



# Average size

**Average size:** Knuth and Trabb Pardo (1976) consider the average size of the largest prime factors as part of an analysis of factoring algorithm runtimes. (Size of  $n$  is  $\log n$ .)

$$\frac{1}{x} \sum_{2 \leq n \leq x} \log P^+(n) \sim \lambda \log x \quad \lambda = \int_0^1 e^{\text{Li}(t)} dt = 0.624329\dots \text{ (Golomb-Dickman constant)}$$

# Average size

**Average size:** Knuth and Trabb Pardo (1976) consider the average size of the largest prime factors as part of an analysis of factoring algorithm runtimes. (Size of  $n$  is  $\log n$ .)

$$\frac{1}{x} \sum_{2 \leq n \leq x} \log P^+(n) \sim \lambda \log x \quad \lambda = \int_0^1 e^{\text{Li}(t)} dt = 0.624329\dots \text{ (Golomb-Dickman constant)}$$

- On average  $P^+(n)$  has about  $5/8=0.625$  as many digits as  $n$ .

# Average size

**Average size:** Knuth and Trabb Pardo (1976) consider the average size of the largest prime factors as part of an analysis of factoring algorithm runtimes. (Size of  $n$  is  $\log n$ .)

$$\frac{1}{x} \sum_{2 \leq n \leq x} \log P^+(n) \sim \lambda \log x \quad \lambda = \int_0^1 e^{\text{Li}(t)} dt = 0.624329\dots \text{ (Golomb-Dickman constant)}$$

- On average  $P^+(n)$  has about  $5/8=0.625$  as many digits as  $n$ .
- The geometric mean of the largest prime factors is  $\left( \prod_{2 \leq n \leq x} P^+(n) \right)^{\frac{1}{x}} = x^{\lambda+o(1)}$ .

# Average size

**Average size:** Knuth and Trabb Pardo (1976) consider the average size of the largest prime factors as part of an analysis of factoring algorithm runtimes. (Size of  $n$  is  $\log n$ .)

$$\frac{1}{x} \sum_{2 \leq n \leq x} \log P^+(n) \sim \lambda \log x \quad \lambda = \int_0^1 e^{\text{Li}(t)} dt = 0.624329\dots \text{ (Golomb-Dickman constant)}$$

- On average  $P^+(n)$  has about  $5/8=0.625$  as many digits as  $n$ .
- The geometric mean of the largest prime factors is  $\left( \prod_{2 \leq n \leq x} P^+(n) \right)^{\frac{1}{x}} = x^{\lambda+o(1)}$ .

For smallest prime factors Tennenbaum shows:  $\frac{1}{x} \sum_{2 \leq n \leq x} \log P^-(n) \sim e^{-\gamma} \log_2 x$

# Average size

**Average size:** Knuth and Trabb Pardo (1976) consider the average size of the largest prime factors as part of an analysis of factoring algorithm runtimes. (Size of  $n$  is  $\log n$ .)

$$\frac{1}{x} \sum_{2 \leq n \leq x} \log P^+(n) \sim \lambda \log x \quad \lambda = \int_0^1 e^{\text{Li}(t)} dt = 0.624329\dots \text{ (Golomb-Dickman constant)}$$

- On average  $P^+(n)$  has about  $5/8=0.625$  as many digits as  $n$ .
- The geometric mean of the largest prime factors is  $\left( \prod_{2 \leq n \leq x} P^+(n) \right)^{\frac{1}{x}} = x^{\lambda+o(1)}$ .

For smallest prime factors Tennenbaum shows:  $\frac{1}{x} \sum_{2 \leq n \leq x} \log P^-(n) \sim e^{-\gamma} \log_2 x$

Connection to permutations: On average the longest cycle in a permutation of length  $n$  is  $\lambda n$  (Golomb, 1964) and the average shortest cycle is  $e^{-\gamma} \log n$  (Shepp, Lloyd, 1966).

# Other distributional properties of $P^+(n)$

# Other distributional properties of $P^+(n)$

**Equidistribution in residue classes:** If  $q \leq \log^A x$  (for fixed  $A$ ) and  $(a, q) = 1$  then

$$\#\{n \leq x : P^+(n) \equiv a \pmod{q}\} \sim \frac{x}{\varphi(q)} \quad (\text{Banks, Harmon, Shparlinski, 2005})$$

# Other distributional properties of $P^+(n)$

**Equidistribution in residue classes:** If  $q \leq \log^A x$  (for fixed  $A$ ) and  $(a, q) = 1$  then

$$\#\{n \leq x : P^+(n) \equiv a \pmod{q}\} \sim \frac{x}{\varphi(q)} \quad (\text{Banks, Harmon, Shparlinski, 2005})$$

**Benford's leading digit law:** Fix a base  $b$  and an initial string of digits  $D$ .



# Other distributional properties of $P^+(n)$

**Equidistribution in residue classes:** If  $q \leq \log^A x$  (for fixed  $A$ ) and  $(a, q) = 1$  then

$$\#\{n \leq x : P^+(n) \equiv a \pmod{q}\} \sim \frac{x}{\varphi(q)} \quad (\text{Banks, Harmon, Shparlinski, 2005})$$

**Benford's leading digit law:** Fix a base  $b$  and an initial string of digits  $D$ .

$$\#\{n \leq x : P^+(n) \text{ starts with } D \text{ in base } b\} \sim \frac{x \log(1 + D^{-1})}{\log b} \quad (\text{Pollack, Singha Roy, 2022})$$

# Other distributional properties of $P^+(n)$

**Equidistribution in residue classes:** If  $q \leq \log^A x$  (for fixed  $A$ ) and  $(a, q) = 1$  then

$$\#\{n \leq x : P^+(n) \equiv a \pmod{q}\} \sim \frac{x}{\varphi(q)} \quad (\text{Banks, Harmon, Shparlinski, 2005})$$

**Benford's leading digit law:** Fix a base  $b$  and an initial string of digits  $D$ .

$$\#\{n \leq x : P^+(n) \text{ starts with } D \text{ in base } b\} \sim \frac{x \log(1 + D^{-1})}{\log b} \quad (\text{Pollack, Singha Roy, 2022})$$

Both also hold for the  $k$ th largest prime factor ( $q \leq \frac{\log x}{(\log_2 x)^{k-1+\epsilon}}$ ).

# Other distributional properties of $P^+(n)$

**Equidistribution in residue classes:** If  $q \leq \log^A x$  (for fixed  $A$ ) and  $(a, q) = 1$  then

$$\#\{n \leq x : P^+(n) \equiv a \pmod{q}\} \sim \frac{x}{\varphi(q)} \quad (\text{Banks, Harmon, Shparlinski, 2005})$$

**Benford's leading digit law:** Fix a base  $b$  and an initial string of digits  $D$ .

$$\#\{n \leq x : P^+(n) \text{ starts with } D \text{ in base } b\} \sim \frac{x \log(1 + D^{-1})}{\log b} \quad (\text{Pollack, Singha Roy, 2022})$$

Both also hold for the  $k$ th largest prime factor ( $q \leq \frac{\log x}{(\log_2 x)^{k-1+\epsilon}}$ ). Neither holds for  $P^-(n)$ .

# Other distributional properties of $P^+(n)$

**Equidistribution in residue classes:** If  $q \leq \log^A x$  (for fixed  $A$ ) and  $(a, q) = 1$  then

$$\#\{n \leq x : P^+(n) \equiv a \pmod{q}\} \sim \frac{x}{\varphi(q)} \quad (\text{Banks, Harmon, Shparlinski, 2005})$$

**Benford's leading digit law:** Fix a base  $b$  and an initial string of digits  $D$ .

$$\#\{n \leq x : P^+(n) \text{ starts with } D \text{ in base } b\} \sim \frac{x \log(1 + D^{-1})}{\log b} \quad (\text{Pollack, Singha Roy, 2022})$$

Both also hold for the  $k$ th largest prime factor ( $q \leq \frac{\log x}{(\log_2 x)^{k-1+\epsilon}}$ ). Neither holds for  $P^-(n)$ .

What about “in between” these extremes?

# Other distributional properties of $P^+(n)$

**Equidistribution in residue classes:** If  $q \leq \log^A x$  (for fixed  $A$ ) and  $(a, q) = 1$  then

$$\#\{n \leq x : P^+(n) \equiv a \pmod{q}\} \sim \frac{x}{\varphi(q)} \quad (\text{Banks, Harmon, Shparlinski, 2005})$$

**Benford's leading digit law:** Fix a base  $b$  and an initial string of digits  $D$ .

$$\#\{n \leq x : P^+(n) \text{ starts with } D \text{ in base } b\} \sim \frac{x \log(1 + D^{-1})}{\log b} \quad (\text{Pollack, Singha Roy, 2022})$$

Both also hold for the  $k$ th largest prime factor ( $q \leq \frac{\log x}{(\log_2 x)^{k-1+\epsilon}}$ ). Neither holds for  $P^-(n)$ .

What about “in between” these extremes? ...the middle prime factor?

# The middle prime factor

De Koninck and Luca (2013), initiated the study of “middle” prime factors.

# The middle prime factor

De Koninck and Luca (2013), initiated the study of “middle” prime factors. Suppose

$$n = p_1 p_2 \cdots p_{\Omega(n)} \quad p_1 \leq p_2 \leq \cdots \leq p_{\Omega(n)}$$

# The middle prime factor

De Koninck and Luca (2013), initiated the study of “middle” prime factors. Suppose

$$n = p_1 p_2 \cdots p_{\Omega(n)} \quad p_1 \leq p_2 \leq \cdots \leq p_{\Omega(n)}$$

Define the middle prime factor  $P^{(\frac{1}{2})}(n) := p_{\lceil \Omega(n)/2 \rceil}$ .



# The middle prime factor

De Koninck and Luca (2013), initiated the study of “middle” prime factors. Suppose

$$n = p_1 p_2 \cdots p_{\Omega(n)} \quad p_1 \leq p_2 \leq \cdots \leq p_{\Omega(n)}$$

Define the middle prime factor  $P^{(\frac{1}{2})}(n) := p_{\lceil \Omega(n)/2 \rceil}$ .

Note that when an integer has an even number of prime factors we take the smaller of the two “middle” primes.

# The middle prime factor

De Koninck and Luca (2013), initiated the study of “middle” prime factors. Suppose

$$n = p_1 p_2 \cdots p_{\Omega(n)} \quad p_1 \leq p_2 \leq \cdots \leq p_{\Omega(n)}$$

Define the middle prime factor  $P^{(\frac{1}{2})}(n) := p_{\lceil \Omega(n)/2 \rceil}$ .

Note that when an integer has an even number of prime factors we take the smaller of the two “middle” primes.  $P^{(\frac{1}{2})}(140) = P^{(\frac{1}{2})}(2 \cdot 2 \cdot 5 \cdot 7) = 2$ .

# The middle prime factor

De Koninck and Luca (2013), initiated the study of “middle” prime factors. Suppose

$$n = p_1 p_2 \cdots p_{\Omega(n)} \quad p_1 \leq p_2 \leq \cdots \leq p_{\Omega(n)}$$

Define the middle prime factor  $P^{(\frac{1}{2})}(n) := p_{\lceil \Omega(n)/2 \rceil}$ .

Note that when an integer has an even number of prime factors we take the smaller of the two “middle” primes.  $P^{(\frac{1}{2})}(140) = P^{(\frac{1}{2})}(2 \cdot 2 \cdot 5 \cdot 7) = 2$ .

We can generalize this to any percentile. For  $\alpha \in (0, 1)$  define  $P^{(\alpha)} := p_{\lceil \alpha \Omega(n) \rceil}$

# The distribution of the intermediate prime factor

For any  $\alpha \in (0, 1)$ ,

- Mean:  $\frac{1}{x} \sum_{2 \leq n \leq x} P^{(\alpha)}(n) \sim \frac{x}{2 \log x}$

# The distribution of the intermediate prime factor

For any  $\alpha \in (0, 1)$ ,

- Mean:  $\frac{1}{x} \sum_{2 \leq n \leq x} P^{(\alpha)}(n) \sim \frac{x}{2 \log x} \sim \frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{1}{x} \sum_{p \leq x} p$

# The distribution of the intermediate prime factor

For any  $\alpha \in (0, 1)$ ,

- Mean:  $\frac{1}{x} \sum_{2 \leq n \leq x} P^{(\alpha)}(n) \sim \frac{x}{2 \log x} \sim \frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{1}{x} \sum_{p \leq x} p$
- Mode: 2

# The distribution of the intermediate prime factor

For any  $\alpha \in (0, 1)$ ,

- Mean:  $\frac{1}{x} \sum_{2 \leq n \leq x} P^{(\alpha)}(n) \sim \frac{x}{2 \log x} \sim \frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{1}{x} \sum_{p \leq x} p$
- Mode: 2
- $\log \log P^{(\alpha)}(n)$  has normal order  $\alpha \log \log n$ . (De Koninck, Katai, 2014)  $P^{(\frac{1}{2})}(n) \approx e^{\sqrt{\log n}}$

# The distribution of the intermediate prime factor

For any  $\alpha \in (0, 1)$ ,

- Mean:  $\frac{1}{x} \sum_{2 \leq n \leq x} P^{(\alpha)}(n) \sim \frac{x}{2 \log x} \sim \frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim \frac{1}{x} \sum_{p \leq x} p$
- Mode: 2
- $\log \log P^{(\alpha)}(n)$  has normal order  $\alpha \log \log n$ . (De Koninck, Katai, 2014)  $P^{(\frac{1}{2})}(n) \approx e^{\sqrt{\log n}}$
- $\log_2 P^{(\alpha)}(n)$  follows a normal distribution. (De Koninck, Doyon, Ouellet, 2019)

$$\frac{1}{x} \# \left\{ n \leq x : \frac{\log_2 P^{(\alpha)}(n) - \alpha \log_2 x}{\sqrt{\log_2 x}} < t \right\} = \Phi(2t) + O_\alpha \left( \frac{1}{\sqrt{\log_3 x}} \right)$$

for  $|t| \ll (\log_2 x)^{1/8-\epsilon}$ .  $\Phi(t) = \int_{-\infty}^t e^{-u^2/2} du$  is the Gaussian distribution function.



# The distribution of the middle prime factor

Does  $P^{(\alpha)}(n)$  follow Benford's law? Is it equidistributed in coprime residue classes?

# The distribution of the middle prime factor

Does  $P^{(\alpha)}(n)$  follow Benford's law? Is it equidistributed in coprime residue classes? **Yes!**

# The distribution of the middle prime factor

Does  $P^{(\alpha)}(n)$  follow Benford's law? Is it equidistributed in coprime residue classes? **Yes!**

Theorem (M., P., S.,  $P^{(\alpha)}(n)$  follows Benford's law)

*Fix an integer  $b \geq 2$ , and fix a positive integer  $D$ . The set of  $n$  for which  $P^{(\alpha)}(n)$  begins with  $D$  in base  $b$  has asymptotic density  $\log(1 + D^{-1})/\log b$ .*

# The distribution of the middle prime factor

Does  $P^{(\alpha)}(n)$  follow Benford's law? Is it equidistributed in coprime residue classes? **Yes!**

Theorem (M., P., S.,  $P^{(\alpha)}(n)$  follows Benford's law)

*Fix an integer  $b \geq 2$ , and fix a positive integer  $D$ . The set of  $n$  for which  $P^{(\alpha)}(n)$  begins with  $D$  in base  $b$  has asymptotic density  $\log(1 + D^{-1})/\log b$ .*

Theorem (M., P., S.,  $P^{(\alpha)}(n)$  is equidistributed in residue classes)

*The number of  $n \leq x$  for which  $P^{(\frac{1}{2})}(n) \equiv a \pmod{q}$  is  $(1 + o(1))x/\phi(q)$  as  $x \rightarrow \infty$ , uniformly in moduli  $q \leq (\log x)^{\frac{1}{2}-\epsilon}$  and in coprime residue classes  $a \pmod{q}$ .*

# The distribution of the middle prime factor

Does  $P^{(\alpha)}(n)$  follow Benford's law? Is it equidistributed in coprime residue classes? **Yes!**

Theorem (M., P., S.,  $P^{(\alpha)}(n)$  follows Benford's law)

*Fix an integer  $b \geq 2$ , and fix a positive integer  $D$ . The set of  $n$  for which  $P^{(\alpha)}(n)$  begins with  $D$  in base  $b$  has asymptotic density  $\log(1 + D^{-1})/\log b$ .*

Theorem (M., P., S.,  $P^{(\alpha)}(n)$  is equidistributed in residue classes)

*The number of  $n \leq x$  for which  $P^{(\frac{1}{2})}(n) \equiv a \pmod{q}$  is  $(1 + o(1))x/\phi(q)$  as  $x \rightarrow \infty$ , uniformly in moduli  $q \leq (\log x)^{\frac{1}{2}-\epsilon}$  and in coprime residue classes  $a \pmod{q}$ .*

The uniformity is essentially optimal: we can show it fails for  $q > (\log x)^{\frac{1}{2}+\epsilon}$ .

# Proof Methods

We found two different methods of proving each of these results.

# Proof Methods

We found two different methods of proving each of these results.

**Method 1:** General result about  $P^{(\alpha)}(n)$  in sets of primes with logarithmic weights.

# Proof Methods

We found two different methods of proving each of these results.

**Method 1:** General result about  $P^{(\alpha)}(n)$  in sets of primes with logarithmic weights.

## Theorem

*Let  $\mathcal{P}$  be any subset of the primes such that whenever  $X$ ,  $Y$ , and  $Y/X \rightarrow \infty$ ,*

$$\sum_{\substack{p \in (X, Y] \\ p \in \mathcal{P}}} \frac{1}{p} \sim \nu \sum_{p \in (X, Y]} \frac{1}{p}$$

*for a fixed  $\nu \in (0, 1]$ . Then the set of  $n$  with  $P^{(\alpha)}(n) \in \mathcal{P}$  has asymptotic density  $\nu$ .*



# Proof Methods

We found two different methods of proving each of these results.

**Method 1:** General result about  $P^{(\alpha)}(n)$  in sets of primes with logarithmic weights.

## Theorem

*Let  $\mathcal{P}$  be any subset of the primes such that whenever  $X$ ,  $Y$ , and  $Y/X \rightarrow \infty$ ,*

$$\sum_{\substack{p \in (X, Y] \\ p \in \mathcal{P}}} \frac{1}{p} \sim \nu \sum_{p \in (X, Y]} \frac{1}{p}$$

*for a fixed  $\nu \in (0, 1]$ . Then the set of  $n$  with  $P^{(\alpha)}(n) \in \mathcal{P}$  has asymptotic density  $\nu$ .*

Primes starting with  $D$  in base  $b$  and those  $p \equiv a \pmod{q}$  for fixed  $q$  have this property.

# Proof Idea

For most integers  $n$ ,  $P^{(\frac{1}{2})}(n)$  isn't too close to its neighbors in the factorization of  $n$ .

# Proof Idea

For most integers  $n$ ,  $P^{(\frac{1}{2})}(n)$  isn't too close to its neighbors in the factorization of  $n$ .

Suppose  $p = P^{(\frac{1}{2})}(n)$ ,  $n = ApB$  with  $P^+(A) \leq p \leq P^-(B)$ .

# Proof Idea

For most integers  $n$ ,  $P^{(\frac{1}{2})}(n)$  isn't too close to its neighbors in the factorization of  $n$ .

Suppose  $p = P^{(\frac{1}{2})}(n)$ ,  $n = ApB$  with  $P^+(A) \leq p \leq P^-(B)$ . “Typically”  $\frac{P^-(B)}{P^+(A)} > \log_3 x$ .

# Proof Idea

For most integers  $n$ ,  $P^{(\frac{1}{2})}(n)$  isn't too close to its neighbors in the factorization of  $n$ .

Suppose  $p = P^{(\frac{1}{2})}(n)$ ,  $n = ApB$  with  $P^+(A) \leq p \leq P^-(B)$ . “Typically”  $\frac{P^-(B)}{P^+(A)} > \log_3 x$ .

This gives us wiggle room to average over all of the primes between  $P^+(A)$  and  $P^-(B)$ .

# Proof Idea

For most integers  $n$ ,  $P^{(\frac{1}{2})}(n)$  isn't too close to its neighbors in the factorization of  $n$ .

Suppose  $p = P^{(\frac{1}{2})}(n)$ ,  $n = ApB$  with  $P^+(A) \leq p \leq P^-(B)$ . “Typically”  $\frac{P^-(B)}{P^+(A)} > \log_3 x$ .

This gives us wiggle room to average over all of the primes between  $P^+(A)$  and  $P^-(B)$ .

For fixed  $A$  and  $B$ , we can multiply  $AB$  by any prime  $p \in [P^+(A), P^-(B)]$  and that prime will be the middle prime factor.

# The number of integers whose middle prime factor is $p$

**Method 2:** Count the number  $M_p^{(\frac{1}{2})}(x)$  of integers up to  $x$  with middle prime factor  $p$ .

# The number of integers whose middle prime factor is $p$

**Method 2:** Count the number  $M_p^{(\frac{1}{2})}(x)$  of integers up to  $x$  with middle prime factor  $p$ . It will be useful to define  $\beta := \frac{\log_2 p}{\log_2 x}$ .



# The number of integers whose middle prime factor is $p$

**Method 2:** Count the number  $M_p^{(\frac{1}{2})}(x)$  of integers up to  $x$  with middle prime factor  $p$ . It will be useful to define  $\beta := \frac{\log_2 p}{\log_2 x}$ . So  $p = e^{(\log x)^\beta}$ .

# The number of integers whose middle prime factor is $p$

**Method 2:** Count the number  $M_p^{(\frac{1}{2})}(x)$  of integers up to  $x$  with middle prime factor  $p$ . It will be useful to define  $\beta := \frac{\log_2 p}{\log_2 x}$ . So  $p = e^{(\log x)^\beta}$ .

## Theorem

*Suppose  $p \rightarrow \infty$  is prime and either  $\beta < \frac{1}{5} - \epsilon$  or  $\frac{1}{5} + \epsilon < \beta < 1 - \epsilon$ . Then we have*

$$M_p^{(\frac{1}{2})}(x) \sim \begin{cases} \frac{C_\beta x}{p(\log x)^{1-2\sqrt{\beta(1-\beta)}} \sqrt{\log_2 x}} & \frac{1}{5} + \epsilon < \beta < 1 - \epsilon, \\ \frac{C x}{p(\log x)^{\frac{1}{2}-\frac{3}{2}\beta}} & 0 < \beta < \frac{1}{5} - \epsilon, \end{cases}$$

# The constants

# The constants

The constants in the previous theorem are given by

$$C_\beta := \frac{\exp\left(\frac{\gamma(1-2\beta)}{\sqrt{\beta(1-\beta)}}\right)}{\Gamma\left(1 + \sqrt{\frac{\beta}{1-\beta}}\right)} \frac{\sqrt{\beta} + \sqrt{1-\beta}}{2\sqrt{\pi}\beta^{1/4}(1-\beta)^{3/4}} \prod_{q \text{ prime}} \left(1 - \frac{1}{q}\right)^{\sqrt{\frac{1-\beta}{\beta}}} \left(1 - \frac{\sqrt{\frac{1-\beta}{\beta}}}{q}\right)^{-1}$$
$$C := \frac{3e^{\frac{3\gamma}{2}}}{4\sqrt{\pi}} \prod_{q>2 \text{ prime}} \left(1 + \frac{1}{q(q-2)}\right) = 1.523555 \dots$$

# The constants

The constants in the previous theorem are given by

$$C_\beta := \frac{\exp\left(\frac{\gamma(1-2\beta)}{\sqrt{\beta(1-\beta)}}\right)}{\Gamma\left(1 + \sqrt{\frac{\beta}{1-\beta}}\right)} \frac{\sqrt{\beta} + \sqrt{1-\beta}}{2\sqrt{\pi}\beta^{1/4}(1-\beta)^{3/4}} \prod_{q \text{ prime}} \left(1 - \frac{1}{q}\right)^{\sqrt{\frac{1-\beta}{\beta}}} \left(1 - \frac{\sqrt{\frac{1-\beta}{\beta}}}{q}\right)^{-1}$$
$$C := \frac{3e^{\frac{3\gamma}{2}}}{4\sqrt{\pi}} \prod_{q>2 \text{ prime}} \left(1 + \frac{1}{q(q-2)}\right) = 1.523555\dots$$

... Note that when  $\beta = \frac{1}{2}$  we have  $C_{\frac{1}{2}} = \sqrt{\frac{2}{\pi}}$ .

# Applications

- $P^{(\frac{1}{2})}(n)$  follows Benford's law and is equidistributed in coprime residue classes.

# Applications

- $P^{(\frac{1}{2})}(n)$  follows Benford's law and is equidistributed in coprime residue classes.

## Theorem (Improved convergence to normal distribution)

*For all  $t$  we have*

$$\frac{1}{x} \# \left\{ n \leq x : \frac{\log_2 P^{(\frac{1}{2})}(n) - \frac{1}{2} \log_2 x}{\sqrt{\log_2 x}} < t \right\} = \Phi(2t) + O\left(\frac{(\log_3)^{3/2}}{\sqrt{\log_2 x}}\right).$$

# Average size of the middle prime factor

Recall that  $\frac{1}{x} \sum_{2 \leq n \leq x} P^+(n) \sim \lambda \log x$ ,  $\frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim e^{-\gamma} \log_2 x$ .



# Average size of the middle prime factor

Recall that  $\frac{1}{x} \sum_{2 \leq n \leq x} P^+(n) \sim \lambda \log x$ ,  $\frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim e^{-\gamma} \log_2 x$ .

## Theorem

Let  $\varphi = \frac{1+\sqrt{5}}{2}$  the golden ratio, and  $\varphi' = \frac{1}{\varphi} = \varphi - 1 = \frac{\sqrt{5}-1}{2} = 0.6180\dots$

$$\frac{1}{x} \sum_{2 \leq n \leq x} \log P^{(\frac{1}{2})}(n) \sim A(\log x)^{\varphi'}$$

# Average size of the middle prime factor

Recall that  $\frac{1}{x} \sum_{2 \leq n \leq x} P^+(n) \sim \lambda \log x$ ,  $\frac{1}{x} \sum_{2 \leq n \leq x} P^-(n) \sim e^{-\gamma} \log_2 x$ .

## Theorem

Let  $\varphi = \frac{1+\sqrt{5}}{2}$  the golden ratio, and  $\varphi' = \frac{1}{\varphi} = \varphi - 1 = \frac{\sqrt{5}-1}{2} = 0.6180\dots$

$$\frac{1}{x} \sum_{2 \leq n \leq x} \log P^{(\frac{1}{2})}(n) \sim A(\log x)^{\varphi'}$$

where

$$A := \frac{e^{-\gamma}}{\varphi!} \frac{\varphi + 1}{\sqrt{5}} \prod_p \left(1 - \frac{1}{p}\right)^{\varphi'} \left(1 - \frac{\varphi'}{p}\right)^{-1} = 1.313314\dots$$

# Generalizations to other $\alpha$

These results generalize to  $P^{(\alpha)}(n)$  for any  $\alpha \in (0,1)$ .

# Generalizations to other $\alpha$

These results generalize to  $P^{(\alpha)}(n)$  for any  $\alpha \in (0,1)$ . Consider  $M_p^{(\alpha)}(x)$ , the count of  $n \leq x$  with  $P^{(\alpha)}(n) = p$ .

# Generalizations to other $\alpha$

These results generalize to  $P^{(\alpha)}(n)$  for any  $\alpha \in (0,1)$ . Consider  $M_p^{(\alpha)}(x)$ , the count of  $n \leq x$  with  $P^{(\alpha)}(n) = p$ . Find a phase transition at  $\beta_\alpha := \left(1 + 2^{\frac{1}{1-\alpha}}(\alpha^{-1} - 1)\right)^{-1}$

# Generalizations to other $\alpha$

These results generalize to  $P^{(\alpha)}(n)$  for any  $\alpha \in (0, 1)$ . Consider  $M_p^{(\alpha)}(x)$ , the count of  $n \leq x$  with  $P^{(\alpha)}(n) = p$ . Find a phase transition at  $\beta_\alpha := \left(1 + 2^{\frac{1}{1-\alpha}}(\alpha^{-1} - 1)\right)^{-1}$

## Theorem

Fix  $\alpha \in (0, 1)$  and suppose  $\beta_\alpha + \epsilon < \beta < 1 - \epsilon$ . Then,

$$M_p^{(\alpha)}(x) \sim \frac{C_{\beta, \alpha} x}{p(\log x)^{1 - \left(\frac{\beta}{\alpha}\right)^\alpha \left(\frac{1-\beta}{1-\alpha}\right)^{1-\alpha}} \sqrt{\log_2 x}}$$

$$C_{\alpha,\beta}$$

Set  $\chi := \frac{(1-\alpha)\beta}{(1-\beta)\alpha},$

Set  $\chi := \frac{(1-\alpha)\beta}{(1-\beta)\alpha}$ , then

$$C_{\beta,\alpha} := \frac{e^{\gamma(\chi^{\alpha-1}-\chi^\alpha)}}{\Gamma(1+\chi^\alpha)} \frac{\chi^{\frac{3}{2}\alpha-1} \rho_{\chi,\alpha}}{\sqrt{2\pi\alpha(1-\beta)}} \prod_q \left(1 - \frac{1}{q}\right)^{\chi^{\alpha-1}} \left(1 - \frac{\chi^{\alpha-1}}{q}\right)^{-1}$$



Set  $\chi := \frac{(1-\alpha)\beta}{(1-\beta)\alpha}$ , then

$$C_{\beta,\alpha} := \frac{e^{\gamma(\chi^{\alpha-1}-\chi^\alpha)}}{\Gamma(1+\chi^\alpha)} \frac{\chi^{\frac{3}{2}\alpha-1} \rho_{\chi,\alpha}}{\sqrt{2\pi\alpha(1-\beta)}} \prod_q \left(1 - \frac{1}{q}\right)^{\chi^{\alpha-1}} \left(1 - \frac{\chi^{\alpha-1}}{q}\right)^{-1}$$

$$\rho_{\chi,\alpha} := \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \chi^{\{n(1-\alpha)\}}$$

Set  $\chi := \frac{(1-\alpha)\beta}{(1-\beta)\alpha}$ , then

$$C_{\beta,\alpha} := \frac{e^{\gamma(\chi^{\alpha-1}-\chi^\alpha)}}{\Gamma(1+\chi^\alpha)} \frac{\chi^{\frac{3}{2}\alpha-1} \rho_{\chi,\alpha}}{\sqrt{2\pi\alpha(1-\beta)}} \prod_q \left(1 - \frac{1}{q}\right)^{\chi^{\alpha-1}} \left(1 - \frac{\chi^{\alpha-1}}{q}\right)^{-1}$$

$$\rho_{\chi,\alpha} := \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \chi^{\{n(1-\alpha)\}} = \begin{cases} \sum_{i=0}^{b-1} \chi^{i/b} & \alpha = \frac{a}{b} \text{ rational} \\ \int_0^1 \chi^t dt = \frac{\chi-1}{\log \chi} & \alpha \text{ irrational} \end{cases}.$$

Set  $\chi := \frac{(1-\alpha)\beta}{(1-\beta)\alpha}$ , then

$$C_{\beta,\alpha} := \frac{e^{\gamma(\chi^{\alpha-1}-\chi^\alpha)}}{\Gamma(1+\chi^\alpha)} \frac{\chi^{\frac{3}{2}\alpha-1} \rho_{\chi,\alpha}}{\sqrt{2\pi\alpha(1-\beta)}} \prod_q \left(1 - \frac{1}{q}\right)^{\chi^{\alpha-1}} \left(1 - \frac{\chi^{\alpha-1}}{q}\right)^{-1}$$

$$\rho_{\chi,\alpha} := \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \chi^{\{n(1-\alpha)\}} = \begin{cases} \sum_{i=0}^{b-1} \chi^{i/b} & \alpha = \frac{a}{b} \text{ rational} \\ \int_0^1 \chi^t dt = \frac{\chi-1}{\log \chi} & \alpha \text{ irrational} \end{cases}.$$

As a function of  $\alpha$  (for a fixed  $\beta$ )  $C_{\beta,\alpha}$  is continuous at every irrational, but discontinuous at every rational value of  $\alpha$ , besides  $\alpha = \beta$ .

# New improvements by Rotgé, (2025)

Focuses on the middle prime factor,  $P^{\frac{1}{2}}(n)$ .

# New improvements by Rotgé, (2025)

Focuses on the middle prime factor,  $P^{\frac{1}{2}}(n)$ .

- Works out the details of the “phase transition” at  $\beta = \frac{1}{5}$ , showing that there is a Gaussian transition between the two regimes.

# New improvements by Rotgé, (2025)

Focuses on the middle prime factor,  $P^{\frac{1}{2}}(n)$ .

- Works out the details of the “phase transition” at  $\beta = \frac{1}{5}$ , showing that there is a Gaussian transition between the two regimes.
- Refines the computation of

$$\frac{1}{x} \sum_{n < x} \log P^{(\frac{1}{2})}(n) = A(\log x)^{\varphi'} \left\{ 1 + \sum_{1 \leq j \leq J} \frac{Q_j(\theta_x)}{(\log_2 x)^j} + O\left(\frac{1}{(\log_2 x)^{J+1}}\right) \right\}.$$

For any fixed  $J$  where the  $Q_j$  are polynomials,  $\theta_x = \{\log_2 x / \sqrt{5}\}$ .

# New improvements by Rotgé, (2025)

Focuses on the middle prime factor,  $P^{\frac{1}{2}}(n)$ .

- Works out the details of the “phase transition” at  $\beta = \frac{1}{5}$ , showing that there is a Gaussian transition between the two regimes.
- Refines the computation of

$$\frac{1}{x} \sum_{n < x} \log P^{(\frac{1}{2})}(n) = A(\log x)^{\varphi'} \left\{ 1 + \sum_{1 \leq j \leq J} \frac{Q_j(\theta_x)}{(\log_2 x)^j} + O\left(\frac{1}{(\log_2 x)^{J+1}}\right) \right\}.$$

For any fixed  $J$  where the  $Q_j$  are polynomials,  $\theta_x = \{\log_2 x / \sqrt{5}\}$ .

- Obtains an improved (and optimal) error term for the convergence the typical values,

$$\frac{1}{x} \# \left\{ n \leq x : \frac{\log_2 P^{(\frac{1}{2})}(n) - \frac{1}{2} \log_2 x}{\sqrt{\log_2 x}} < t \right\} = \Phi(2t) + O\left(\frac{1}{\sqrt{\log_2 x}}\right).$$

# Open Questions



# Open Questions

The median of the middle prime factor is still pretty open.

# Open Questions

The median of the middle prime factor is still pretty open.

After [Rotgé](#)'s improvements we now know this median is  $\exp(O(\sqrt{\log x}))$ .

# Open Questions

The median of the middle prime factor is still pretty open.

After [Rotgé](#)'s improvements we now know this median is  $\exp(O(\sqrt{\log x}))$ .

We also don't know the behavior of  $M_p^{(\frac{1}{2})}(x)$  as when  $\beta = \frac{\log_2 p}{\log_2 x} \rightarrow 1$ .

# Open Questions

The median of the middle prime factor is still pretty open.

After [Rotgé](#)'s improvements we now know this median is  $\exp(O(\sqrt{\log x}))$ .

We also don't know the behavior of  $M_p^{(\frac{1}{2})}(x)$  as when  $\beta = \frac{\log_2 p}{\log_2 x} \rightarrow 1$ .

# THANK YOU!