

# Counting Primitive Sets

and other statistics of the divisor graph of  $\{1, 2, \dots, n\}$

Nathan McNew  
Towson University

INTEGERS 2018  
Augusta GA  
October 5th, 2018

# Primitive Sets

A set of integers is **primitive** if no integer in the set divides another.

# Primitive Sets

A set of integers is **primitive** if no integer in the set divides another.

## Examples:

- Primes

# Primitive Sets

A set of integers is **primitive** if no integer in the set divides another.

## Examples:

- Primes
- Integers with exactly  $k$  prime factors counted with multiplicity.

# Primitive Sets

A set of integers is **primitive** if no integer in the set divides another.

## Examples:

- Primes
- Integers with exactly  $k$  prime factors counted with multiplicity.
- $\left\{ \left\lfloor \frac{n}{2} \right\rfloor + 1, \left\lfloor \frac{n}{2} \right\rfloor + 2, \dots, n \right\}$

# Primitive Sets

A set of integers is **primitive** if no integer in the set divides another.

## Examples:

- Primes
- Integers with exactly  $k$  prime factors counted with multiplicity.
- $\left\{ \left\lfloor \frac{n}{2} \right\rfloor + 1, \left\lfloor \frac{n}{2} \right\rfloor + 2, \dots, n \right\}$

A **divisor graph** is a graph with vertices labeled by integers and edges connecting each pair of integers where one divides another.

# Primitive Sets

A set of integers is **primitive** if no integer in the set divides another.

## Examples:

- Primes
- Integers with exactly  $k$  prime factors counted with multiplicity.
- $\left\{ \left\lfloor \frac{n}{2} \right\rfloor + 1, \left\lfloor \frac{n}{2} \right\rfloor + 2, \dots, n \right\}$

A **divisor graph** is a graph with vertices labeled by integers and edges connecting each pair of integers where one divides another.

**Independent** subsets of a divisor graph are primitive sets of integers.

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.



# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5



# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	13

# Finite primitive sets

Note:  $\{\}$  and  $\{1\}$  are both primitive sets.

What do other primitive subsets of the first few integers look like?

$n$	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	13

How many primitive subsets of  $\{1, 2 \dots n\}$  are there?

# Counting primitive sets

Let  $Q(n)$  count the primitive sets with largest element at most  $n$ .


# Counting primitive sets

Let  $Q(n)$  count the primitive sets with largest element at most  $n$ .

A051026	Number of primitive subsequences of $\{1, 2, \dots, n\}$ .	5
	1, 2, 3, 5, 7, 13, 17, 33, 45, 73, 103, 205, 253, 505, 733, 1133, 1529, 3057, 3897, 7793, 10241, 16513, 24593, 49185, 59265, 109297, 163369, 262489, 355729, 711457, 879937, 1759873, 2360641, 3908545, 5858113, 10534337, 12701537, 25403073, 38090337, 63299265, 81044097, 162088193, 205482593, 410965185, 570487233, 855676353 ( <a href="#">list</a> ; <a href="#">graph</a> ; <a href="#">refs</a> ; <a href="#">listen</a> ; <a href="#">history</a> ; <a href="#">text</a> ; <a href="#">internal format</a> )	
OFFSET	0, 2	
COMMENTS	$a(n)$ counts all subsequences of $\{1, \dots, n\}$ in which no term divides any other. If $n$ is a prime $a(n) = 2 \cdot a(n-1) - 1$ because for each subsequence $s$ counted by $a(n-1)$ two different subsequences are counted by $a(n)$ : $s$ and $s, n$ . There is only one exception: $1, n$ is not a primitive subsequence because 1 divides $n$ . For all $n > 1$ : $a(n) < 2 \cdot a(n-1)$ . - <a href="#">Alois P. Heinz</a> , Mar 07 2011	

# Bounds


Cameron and Erdős considered the function  $Q(n)$  in 1990.





# Bounds

Cameron and Erdős considered the function  $Q(n)$  in 1990.




(Almost) trivially:

$$(\sqrt{2})^n < Q(n) < 2^n$$

# Bounds

Cameron and Erdős considered the function  $Q(n)$  in 1990.



(Almost) trivially:

$$(\sqrt{2})^n < Q(n) < 2^n$$

Every subset of  $\left(\frac{n}{2}, n\right]$  is primitive. There are  $2^{\lceil \frac{n}{2} \rceil} > \sqrt{2}^n$  such subsets.

# Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

# Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

**Conjecture:**  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists.

# Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

**Conjecture:**  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists.

**Theorem** (Angelo, 2017)

*The limit  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists.*

# Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

**Conjecture:**  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists.

**Theorem** (Angelo, 2017)

*The limit  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists. Equivalently,  $Q(n) = c^{n+o(1)}$ .*

# Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

**Conjecture:**  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists.

**Theorem** (Angelo, 2017)

*The limit  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists. Equivalently,  $Q(n) = c^{n+o(1)}$ .*

Proof: Study sets without integers having  $k$ -smooth integer ratio.

# Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

**Conjecture:**  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists.

**Theorem** (Angelo, 2017)

*The limit  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists. Equivalently,  $Q(n) = c^{n+o(1)}$ .*

Proof: Study sets without integers having  $k$ -smooth integer ratio.  
( $n$  is  $k$ -smooth if its largest prime divisor,  $P^+(n) \leq k$ .)



# Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

**Conjecture:**  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists.

**Theorem** (Angelo, 2017)

*The limit  $\lim_{n \rightarrow \infty} Q(n)^{1/n}$  exists. Equivalently,  $Q(n) = c^{n+o(1)}$ .*

Proof: Study sets without integers having  $k$ -smooth integer ratio. ( $n$  is  $k$ -smooth if its largest prime divisor,  $P^+(n) \leq k$ .) Let  $k \rightarrow \infty$ .

Proof is not effective: Gives no insight on the value of this constant.

# Constructing Primitive Sets

# Constructing Primitive Sets

Every subset of  $(\frac{n}{2}, n]$  is primitive, so  $Q(n) \geq 2^{n/2}$ .

# Constructing Primitive Sets

Every subset of  $(\frac{n}{2}, n]$  is primitive, so  $Q(n) \geq 2^{n/2}$ .

Try to make this bigger by considering integers in  $(\frac{n}{3}, \frac{n}{2}]$ .

# Constructing Primitive Sets

Every subset of  $(\frac{n}{2}, n]$  is primitive, so  $Q(n) \geq 2^{n/2}$ .

Try to make this bigger by considering integers in  $(\frac{n}{3}, \frac{n}{2}]$ .

- For  $k \in (\frac{n}{3}, \frac{n}{2}]$  we can include  $k, 2k$  or neither.

# Constructing Primitive Sets

Every subset of  $(\frac{n}{2}, n]$  is primitive, so  $Q(n) \geq 2^{n/2}$ .

Try to make this bigger by considering integers in  $(\frac{n}{3}, \frac{n}{2}]$ .

- For  $k \in (\frac{n}{3}, \frac{n}{2}]$  we can include  $k, 2k$  or neither. (3 possibilities)
- Equivalent to counting primitive subsets of  $\{1, 2\}$

# Constructing Primitive Sets

Every subset of  $(\frac{n}{2}, n]$  is primitive, so  $Q(n) \geq 2^{n/2}$ .

Try to make this bigger by considering integers in  $(\frac{n}{3}, \frac{n}{2}]$ .

- For  $k \in (\frac{n}{3}, \frac{n}{2}]$  we can include  $k, 2k$  or neither. (3 possibilities)
- Equivalent to counting primitive subsets of  $\{1, 2\}$
- Also equivalent to counting independent subsets of the graph:

# Constructing Primitive Sets

Every subset of  $(\frac{n}{2}, n]$  is primitive, so  $Q(n) \geq 2^{n/2}$ .

Try to make this bigger by considering integers in  $(\frac{n}{3}, \frac{n}{2}]$ .

- For  $k \in (\frac{n}{3}, \frac{n}{2}]$  we can include  $k, 2k$  or neither. (3 possibilities)
- Equivalent to counting primitive subsets of  $\{1, 2\}$
- Also equivalent to counting independent subsets of the graph:





# Constructing Primitive Sets

Every subset of  $(\frac{n}{2}, n]$  is primitive, so  $Q(n) \geq 2^{n/2}$ .

Try to make this bigger by considering integers in  $(\frac{n}{3}, \frac{n}{2}]$ .

- For  $k \in (\frac{n}{3}, \frac{n}{2}]$  we can include  $k, 2k$  or neither. (3 possibilities)
- Equivalent to counting primitive subsets of  $\{1, 2\}$
- Also equivalent to counting independent subsets of the graph:



An **independent** set of vertices in a graph has no adjacent vertices.

# Constructing Primitive Sets

We can use this to improve the lower bound  $Q(n) > 2^{n/2}$ .

# Constructing Primitive Sets

We can use this to improve the lower bound  $Q(n) > 2^{n/2}$ .

For each  $k \in (\frac{n}{3}, \frac{n}{2}]$  we can multiply by  $\frac{3}{2}$ , the ratio of allowed subsets of  $\{k, 2k\}$ , to those of just  $\{2k\}$ . ( $2k$  was an element of  $(\frac{n}{2}, n]$ , so those subsets were already taken into account.)

# Constructing Primitive Sets

We can use this to improve the lower bound  $Q(n) > 2^{n/2}$ .

For each  $k \in (\frac{n}{3}, \frac{n}{2}]$  we can multiply by  $\frac{3}{2}$ , the ratio of allowed subsets of  $\{k, 2k\}$ , to those of just  $\{2k\}$ . ( $2k$  was an element of  $(\frac{n}{2}, n]$ , so those subsets were already taken into account.)

Thus

$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} = 2^{n/3} 3^{n/6} \approx 1.5131^n.$$

# Constructing Primitive Sets

Pressing on!

# Constructing Primitive Sets

Pressing on! Working backward, considering progressively smaller  $k$ .

# Constructing Primitive Sets

Pressing on! Working backward, considering progressively smaller  $k$ .

Lets consider  $k \in (\frac{n}{4}, \frac{n}{3}]$ .

# Constructing Primitive Sets

Pressing on! Working backward, considering progressively smaller  $k$ .

Lets consider  $k \in (\frac{n}{4}, \frac{n}{3}]$ . Now We need to worry about  $k, 2k, 3k$ .



# Constructing Primitive Sets

Pressing on! Working backward, considering progressively smaller  $k$ .

Lets consider  $k \in (\frac{n}{4}, \frac{n}{3}]$ . Now We need to worry about  $k, 2k, 3k$ .

Actually two scenarios depending on the parity of  $k$ .


# Constructing Primitive Sets

Pressing on! Working backward, considering progressively smaller  $k$ .

Lets consider  $k \in (\frac{n}{4}, \frac{n}{3}]$ . Now We need to worry about  $k, 2k, 3k$ .

Actually two scenarios depending on the parity of  $k$ .

For  $k$  odd, 5 primitive subsets of  $\{k, 2k, 3k\}$  replace 4 of just  $\{2k, 3k\}$ .



# Constructing Primitive Sets

If  $k = 2\ell$  is even then  $\frac{3k}{2} = 3\ell$  is also an integer.

# Constructing Primitive Sets

If  $k = 2\ell$  is even then  $\frac{3k}{2} = 3\ell$  is also an integer.


We have to consider all of  $2\ell, 3\ell, 4\ell, 6\ell$ .

# Constructing Primitive Sets

If  $k = 2\ell$  is even then  $\frac{3k}{2} = 3\ell$  is also an integer.

We have to consider all of  $2\ell, 3\ell, 4\ell, 6\ell$ .

Like counting primitive subsets of  $\{2, 3, 4, 6\}$ .



# Constructing Primitive Sets


If  $k = 2\ell$  is even then  $\frac{3k}{2} = 3\ell$  is also an integer.

We have to consider all of  $2\ell, 3\ell, 4\ell, 6\ell$ .

Like counting primitive subsets of  $\{2, 3, 4, 6\}$ .

Now we get 8 subsets of  $\{2, 3, 4, 6\}$ :

$\{\}, \{2\}, \{3\}, \{4\}, \{6\}, \{2, 3\}, \{3, 4\}, \{4, 6\}$



# Constructing Primitive Sets

If  $k = 2\ell$  is even then  $\frac{3k}{2} = 3\ell$  is also an integer.


We have to consider all of  $2\ell, 3\ell, 4\ell, 6\ell$ .

Like counting primitive subsets of  $\{2, 3, 4, 6\}$ .

Now we get 8 subsets of  $\{2, 3, 4, 6\}$ :

$\{\}, \{2\}, \{3\}, \{4\}, \{6\}, \{2, 3\}, \{3, 4\}, \{4, 6\}$

Replacing the 6 primitive subsets of only  $\{3, 4, 6\}$ .



# Constructing Primitive Sets

If  $k = 2\ell$  is even then  $\frac{3k}{2} = 3\ell$  is also an integer.


We have to consider all of  $2\ell, 3\ell, 4\ell, 6\ell$ .

Like counting primitive subsets of  $\{2, 3, 4, 6\}$ .

Now we get 8 subsets of  $\{2, 3, 4, 6\}$ :

$\{\}, \{2\}, \{3\}, \{4\}, \{6\}, \{2, 3\}, \{3, 4\}, \{4, 6\}$

Replacing the 6 primitive subsets of only  $\{3, 4, 6\}$ .



$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} \left(\frac{5}{4}\right)^{n/24} \left(\frac{8}{6}\right)^{n/24}$$



# Constructing Primitive Sets

If  $k = 2\ell$  is even then  $\frac{3k}{2} = 3\ell$  is also an integer.


We have to consider all of  $2\ell, 3\ell, 4\ell, 6\ell$ .

Like counting primitive subsets of  $\{2, 3, 4, 6\}$ .

Now we get 8 subsets of  $\{2, 3, 4, 6\}$ :

$\{\}, \{2\}, \{3\}, \{4\}, \{6\}, \{2, 3\}, \{3, 4\}, \{4, 6\}$

Replacing the 6 primitive subsets of only  $\{3, 4, 6\}$ .



$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} \left(\frac{5}{4}\right)^{n/24} \left(\frac{8}{6}\right)^{n/24} = 2^{5n/24} 3^{n/8} 5^{n/24}$$

# Constructing Primitive Sets

If  $k = 2\ell$  is even then  $\frac{3k}{2} = 3\ell$  is also an integer.


We have to consider all of  $2\ell, 3\ell, 4\ell, 6\ell$ .

Like counting primitive subsets of  $\{2, 3, 4, 6\}$ .

Now we get 8 subsets of  $\{2, 3, 4, 6\}$ :

$\{\}, \{2\}, \{3\}, \{4\}, \{6\}, \{2, 3\}, \{3, 4\}, \{4, 6\}$

Replacing the 6 primitive subsets of only  $\{3, 4, 6\}$ .



$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} \left(\frac{5}{4}\right)^{n/24} \left(\frac{8}{6}\right)^{n/24} = 2^{5n/24} 3^{n/8} 5^{n/24} \approx 1.5456^n$$

# One more wrinkle


# One more wrinkle

Consider  $k \in \left(\frac{n}{5}, \frac{n}{4}\right]$ , the next interval.

# One more wrinkle

Consider  $k \in \left(\frac{n}{5}, \frac{n}{4}\right]$ , the next interval.


If  $k = 2\ell$  is even we must consider  $2\ell, 4\ell, 6\ell, 8\ell$ , as well as  $3\ell$ .




# One more wrinkle

Consider  $k \in (\frac{n}{5}, \frac{n}{4}]$ , the next interval.

If  $k = 2\ell$  is even we must consider  $2\ell, 4\ell, 6\ell, 8\ell$ , as well as  $3\ell$ .




However, if  $k \leq \frac{2n}{9}$  then  $9\ell \leq n$  must also be considered.




# One more wrinkle

Consider  $k \in (\frac{n}{5}, \frac{n}{4}]$ , the next interval.

If  $k = 2\ell$  is even we must consider  $2\ell, 4\ell, 6\ell, 8\ell$ , as well as  $3\ell$ .



However, if  $k \leq \frac{2n}{9}$  then  $9\ell \leq n$  must also be considered.



So now we consider  $2\ell, 3\ell, 4\ell, 6\ell, 8\ell$  and  $9\ell$ .

# Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$



# Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

# Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

- $1 \leq r(k, n) \leq 2$

# Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

- $1 \leq r(k, n) \leq 2$
- $r(k, n)$  only depends on the structure of the connected component of  $k$  in the divisibility graph of  $[k, n]$ .

# Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

- $1 \leq r(k, n) \leq 2$
- $r(k, n)$  only depends on the structure of the connected component of  $k$  in the divisibility graph of  $[k, n]$ .
- $Q(n) = \prod_{i=1}^n r(i, n)$

# Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

- $1 \leq r(k, n) \leq 2$
- $r(k, n)$  only depends on the structure of the connected component of  $k$  in the divisibility graph of  $[k, n]$ .
- $Q(n) = \prod_{i=1}^n r(i, n)$  (Telescoping product)

# Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

- $1 \leq r(k, n) \leq 2$
- $r(k, n)$  only depends on the structure of the connected component of  $k$  in the divisibility graph of  $[k, n]$ .
- $Q(n) = \prod_{i=1}^n r(i, n)$  (Telescoping product)
- Goal: group together equal terms in this product.

# Key Observation

Fix  $k, n$ .

# Key Observation

Fix  $k, n$ . Let  $i = \lfloor \frac{n}{k} \rfloor$ .



# Key Observation

Fix  $k, n$ . Let  $i = \lfloor \frac{n}{k} \rfloor$ . The multiples of  $k$  up to  $n$  are  $k, 2k, \dots ik$ .

# Key Observation

Fix  $k, n$ . Let  $i = \lfloor \frac{n}{k} \rfloor$ . The multiples of  $k$  up to  $n$  are  $k, 2k, \dots ik$ .

If one integer from  $[k, n]$  divides another the ratio must be  $i$ -smooth.

# Key Observation

Fix  $k, n$ . Let  $i = \lfloor \frac{n}{k} \rfloor$ . The multiples of  $k$  up to  $n$  are  $k, 2k, \dots ik$ .

If one integer from  $[k, n]$  divides another the ratio must be  $i$ -smooth.

Let  $d = \max\{e|k : P^+(e) \leq i\}$ , the largest  $i$ -smooth divisor of  $k$ .

# Key Observation

Fix  $k, n$ . Let  $i = \lfloor \frac{n}{k} \rfloor$ . The multiples of  $k$  up to  $n$  are  $k, 2k, \dots ik$ .

If one integer from  $[k, n]$  divides another the ratio must be  $i$ -smooth.

Let  $d = \max\{e|k : P^+(e) \leq i\}$ , the largest  $i$ -smooth divisor of  $k$ .

Let  $\ell = \frac{k}{d}$  the “rough” part of  $k$ .

# Key Observation

Fix  $k, n$ . Let  $i = \lfloor \frac{n}{k} \rfloor$ . The multiples of  $k$  up to  $n$  are  $k, 2k, \dots ik$ .

If one integer from  $[k, n]$  divides another the ratio must be  $i$ -smooth.

Let  $d = \max\{e|k : P^+(e) \leq i\}$ , the largest  $i$ -smooth divisor of  $k$ .

Let  $\ell = \frac{k}{d}$  the “rough” part of  $k$ .

Every integer in the connected component of  $k$  is divisible by  $\ell$ .

# Key Observation

Fix  $k, n$ . Let  $i = \lfloor \frac{n}{k} \rfloor$ . The multiples of  $k$  up to  $n$  are  $k, 2k, \dots ik$ .

If one integer from  $[k, n]$  divides another the ratio must be  $i$ -smooth.

Let  $d = \max\{e|k : P^+(e) \leq i\}$ , the largest  $i$ -smooth divisor of  $k$ .

Let  $\ell = \frac{k}{d}$  the “rough” part of  $k$ .

Every integer in the connected component of  $k$  is divisible by  $\ell$ .

Let  $t = \lfloor \frac{n}{\ell} \rfloor$ , so  $t\ell$  is the largest integer up to  $n$  divisible by  $\ell$ .

# Key Observation

Fix  $k, n$ . Let  $i = \lfloor \frac{n}{k} \rfloor$ . The multiples of  $k$  up to  $n$  are  $k, 2k, \dots ik$ .

If one integer from  $[k, n]$  divides another the ratio must be  $i$ -smooth.

Let  $d = \max\{e|k : P^+(e) \leq i\}$ , the largest  $i$ -smooth divisor of  $k$ .

Let  $\ell = \frac{k}{d}$  the “rough” part of  $k$ .

Every integer in the connected component of  $k$  is divisible by  $\ell$ .

Let  $t = \lfloor \frac{n}{\ell} \rfloor$ , so  $t\ell$  is the largest integer up to  $n$  divisible by  $\ell$ .

Then  $r(k, n) = r(d\ell, t\ell)$

# Key Observation

Fix  $k, n$ . Let  $i = \lfloor \frac{n}{k} \rfloor$ . The multiples of  $k$  up to  $n$  are  $k, 2k, \dots ik$ .

If one integer from  $[k, n]$  divides another the ratio must be  $i$ -smooth.

Let  $d = \max\{e|k : P^+(e) \leq i\}$ , the largest  $i$ -smooth divisor of  $k$ .

Let  $\ell = \frac{k}{d}$  the “rough” part of  $k$ .

Every integer in the connected component of  $k$  is divisible by  $\ell$ .

Let  $t = \lfloor \frac{n}{\ell} \rfloor$ , so  $t\ell$  is the largest integer up to  $n$  divisible by  $\ell$ .

Then  $r(k, n) = r(d\ell, t\ell) = r(d, t)$ .



# Approximating $Q(n)$

## Theorem

*For any  $\epsilon > 0$ , the number of primitive subsets of  $[1, n]$  is*

$$Q(n) = c^{n(1+O(\exp(-\sqrt{(\frac{1}{6}-\epsilon)\log n \log \log n})))}$$

# Approximating $Q(n)$

## Theorem

*For any  $\epsilon > 0$ , the number of primitive subsets of  $[1, n]$  is*

$$Q(n) = c^{n(1+O(\exp(-\sqrt{(\frac{1}{6}-\epsilon)\log n \log \log n})))}$$

*or*

$$\log Q(n) = n \log(c) + O\left(\frac{n}{e^{\sqrt{(\frac{1}{6}-\epsilon)\log n \log \log n}}}\right).$$

# Approximating $Q(n)$

## Theorem

*For any  $\epsilon > 0$ , the number of primitive subsets of  $[1, n]$  is*

$$Q(n) = c^{n(1+O(\exp(-\sqrt{(\frac{1}{6}-\epsilon)\log n \log \log n})))}$$

*or*

$$\log Q(n) = n \log(c) + O\left(\frac{n}{e^{\sqrt{(\frac{1}{6}-\epsilon)\log n \log \log n}}}\right).$$

*The constant  $c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ P^+(d) < i}} \prod_{t \in [id, (i+1)d)} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}$*

*is effectively computable and  $1.5729 < c < 1.5745$ .*

# Numerical Bound

Get a lower bound for  $c$  by computing terms in this product.

# Numerical Bound

Get a lower bound for  $c$  by computing terms in this product. Using Sage to compute  $r(d,t)$  for all  $i,d,t$  with  $di^5 \leq 10^8$  gives  $1.5729 < c$ .

# Numerical Bound

Get a lower bound for  $c$  by computing terms in this product. Using Sage to compute  $r(d, t)$  for all  $i, d, t$  with  $di^5 \leq 10^8$  gives  $1.5729 < c$ .

**#24418** closed defect (fixed)



Opened 5 months ago

Closed 4 months ago

## Doctest: bug numerical\_approx( $2^{450232897/4888643760}$ )

Reported by:	<a href="#">vdelecroix</a>	Owned by:	
Priority:	<a href="#">major</a>	Milestone:	<a href="#">sage-8.2</a>
Component:	<a href="#">symbolics</a>	Keywords:	<a href="#">bug</a>
Cc:	<a href="#">rws</a>	Merged in:	
Authors:	Ralf Stephan	Reviewers:	Jeroen Demeyer
Report Upstream:	<a href="#">N/A</a>	Work issues:	
Branch:	<a href="#">821f7d9 (Commits)</a>	Commit:	821f7d9f3568316bc0b8b1f5619bce...
Dependencies:		Stopgaps:	

Description (last modified by [vdelecroix](#))  $\Delta$

```
sage: numerical_approx(2^(450232897/4888643760))
```

```
-----  
RuntimeError                                Traceback (most recent call last)
```

```
<ipython-input-2-3c4e30ac02c1> in <module>()  
----> 1 numerical_approx(Integer(2)**(Integer(450232897)/Integer(4888643760)))
```

```
/opt/sage/local/lib/python2.7/site-packages/sage/misc/functional.pyc in numerical_ap  
1406         return numerical_approx_generic(x, prec)
```

# Numerical Bound

Get a lower bound for  $c$  by computing terms in this product. Using Sage to compute  $r(d,t)$  for all  $i,d,t$  with  $di^5 \leq 10^8$  gives  $1.5729 < c$ .

This technique doesn't give very good upper bounds since any unestimated terms need to be replaced by 2 (or 1.5).

# Numerical Bound

Get a lower bound for  $c$  by computing terms in this product. Using Sage to compute  $r(d,t)$  for all  $i,d,t$  with  $di^5 \leq 10^8$  gives  $1.5729 < c$ .

This technique doesn't give very good upper bounds since any unestimated terms need to be replaced by 2 (or 1.5). Gives  $c < 1.58$ .



# Numerical Bound

Get a lower bound for  $c$  by computing terms in this product. Using Sage to compute  $r(d,t)$  for all  $i,d,t$  with  $di^5 \leq 10^8$  gives  $1.5729 < c$ .

This technique doesn't give very good upper bounds since any unestimated terms need to be replaced by 2 (or 1.5). Gives  $c < 1.58$ .

The bound  $c < 1.5745$  was recently obtained by [Liu](#), [Pach](#) and [Palincza](#) (2018) who also prove that  $c$  is effectively computable.

# (Conjectural) Upper Bound

# (Conjectural) Upper Bound

Do the same thing including integers from the beginning of the interval rather than the end.

# (Conjectural) Upper Bound

Do the same thing including integers from the beginning of the interval rather than the end.

Let

$$g(k) = \frac{\# \text{Primitive subsets of } [1, k]}{\# \text{Primitive subsets of } [1, k-1]} - 1$$

# (Conjectural) Upper Bound

Do the same thing including integers from the beginning of the interval rather than the end.

Let

$$\begin{aligned} g(k) &= \frac{\# \text{Primitive subsets of } [1, k]}{\# \text{Primitive subsets of } [1, k-1]} - 1 \\ &= \frac{\# \text{Primitive subsets of } [1, k] \text{ including } k}{\# \text{Primitive subsets of } [1, k-1]} \end{aligned}$$

# (Conjectural) Upper Bound

Do the same thing including integers from the beginning of the interval rather than the end.

Let

$$\begin{aligned} g(k) &= \frac{\# \text{Primitive subsets of } [1, k]}{\# \text{Primitive subsets of } [1, k-1]} - 1 \\ &= \frac{\# \text{Primitive subsets of } [1, k] \text{ including } k}{\# \text{Primitive subsets of } [1, k-1]} \end{aligned}$$

As before,

$$Q(n) = \prod_{k=1}^n (g(k) + 1)$$

# (Conjectural) Upper Bound

# (Conjectural) Upper Bound

## Conjecture

*The function  $g(n)$  is submultiplicative.*



# (Conjectural) Upper Bound

## Conjecture

*The function  $g(n)$  is submultiplicative. If  $(n,m)=1$  then*

$$g(nm) \leq g(n)g(m).$$

# (Conjectural) Upper Bound

## Conjecture

*The function  $g(n)$  is submultiplicative. If  $(n,m)=1$  then*

$$g(nm) \leq g(n)g(m).$$

*Furthermore, for any prime  $p$ ,  $g(p^i) > g(p^{i+1})$ .*

# (Conjectural) Upper Bound

## Conjecture

*The function  $g(n)$  is submultiplicative. If  $(n,m)=1$  then*

$$g(nm) \leq g(n)g(m).$$

*Furthermore, for any prime  $p$ ,  $g(p^i) > g(p^{i+1})$ .*

The conjecture holds for all values of  $n < 899$ .

# (Conjectural) Upper Bound

## Conjecture

*The function  $g(n)$  is submultiplicative. If  $(n,m)=1$  then*

$$g(nm) \leq g(n)g(m).$$

*Furthermore, for any prime  $p$ ,  $g(p^i) > g(p^{i+1})$ .*

The conjecture holds for all values of  $n < 899$ .

Assuming the conjecture, and using values of  $g(n)$ ,  $n < 899$  gives

$$1.5729 < c < 1.5735.$$

# A general theorem

# A general theorem

## Theorem (M.)

Fix  $\epsilon > 0$ ,  $A \geq 0$ . Suppose  $|f(k, n)| \leq A$  and  $f(k, n)$  depends only on the connected component of  $k$  in the divisor graph of  $[k, n]$ . Then

$$\sum_{a=1}^n f(a, n) = nC_f + O_A \left( n \exp \left( -\sqrt{\left(\frac{1}{6} - \epsilon\right) \log n \log \log n} \right) \right)$$

where

$$C_f = \sum_{i=1}^{\infty} \sum_{\substack{d \\ P^+(d) \leq i}} \sum_{t \in [id, (i+1)d)} \left( \frac{f(d, t)}{t(t+1)} \prod_{p \leq i} \frac{p-1}{p} \right).$$

# Other Applications: Maximum Primitive Subsets

# Other Applications: Maximum Primitive Subsets

Recall that the largest primitive subset of this interval has size  $\left\lceil \frac{n}{2} \right\rceil$ .



# Other Applications: Maximum Primitive Subsets

Recall that the largest primitive subset of this interval has size  $\left\lceil \frac{n}{2} \right\rceil$ .

Let  $M(n)$  count the primitive subsets of  $\{1, \dots, n\}$  of size  $\left\lceil \frac{n}{2} \right\rceil$ .

# Other Applications: Maximum Primitive Subsets

Recall that the largest primitive subset of this interval has size  $\left\lceil \frac{n}{2} \right\rceil$ .

Let  $M(n)$  count the primitive subsets of  $\{1, \dots, n\}$  of size  $\left\lceil \frac{n}{2} \right\rceil$ .

Theorem ([Vijay](#), 2018)

$$1.141^n < M(n) < 1.187^n$$

# Other Applications: Maximum Primitive Subsets

Recall that the largest primitive subset of this interval has size  $\left\lceil \frac{n}{2} \right\rceil$ .

Let  $M(n)$  count the primitive subsets of  $\{1, \dots, n\}$  of size  $\left\lceil \frac{n}{2} \right\rceil$ .

Theorem (Vijay, 2018)

$$1.141^n < M(n) < 1.187^n$$

Theorem (Liu, Pach, Palincza, 2018)

$$M(n) = \alpha^{n+o(n)} \text{ as } n \rightarrow \infty \text{ and } 1.14817 < \alpha < 1.14823.$$

Theorem (M., 2018)

$$M(n) = \alpha^{n \left( 1 + O \left( \exp \left( -\sqrt{\left( \frac{1}{6} - \epsilon \right) \log n \log \log n} \right) \right) \right)} \text{ and } 1.14819 < \alpha.$$

# Other Applications: Maximal Primitive Sets

A primitive subset of  $[1, n]$  is maximal if it is not contained in another primitive subset. Let  $m(n)$  count maximal primitive subsets of  $[1, n]$ .

## Theorem

$$m(n) = \beta^n \left(1 + O\left(\exp\left(-\sqrt{\left(\frac{1}{6} - \epsilon\right) \log n \log \log n}\right)\right)\right) \text{ and } 1.2125 < \beta < 1.2409.$$

# Other Applications: Maximal Primitive Sets

A primitive subset of  $[1, n]$  is maximal if it is not contained in another primitive subset. Let  $m(n)$  count maximal primitive subsets of  $[1, n]$ .

## Theorem

$$m(n) = \beta^n \left(1 + O\left(\exp\left(-\sqrt{\left(\frac{1}{6} - \epsilon\right) \log n \log \log n}\right)\right)\right) \text{ and } 1.2125 < \beta < 1.2409.$$

## Corollary

Let  $V(n)$  denote the median size of primitive subsets of  $[1, n]$ . Then

$$0.1681n < V(n) < 0.3918n$$

# Other Applications: Maximal Primitive Sets

A primitive subset of  $[1, n]$  is maximal if it is not contained in another primitive subset. Let  $m(n)$  count maximal primitive subsets of  $[1, n]$ .

## Theorem

$$m(n) = \beta^n \left(1 + O\left(\exp\left(-\sqrt{\left(\frac{1}{6} - \epsilon\right) \log n \log \log n}\right)\right)\right) \text{ and } 1.2125 < \beta < 1.2409.$$

## Corollary

Let  $V(n)$  denote the median size of primitive subsets of  $[1, n]$ . Then

$$0.1681n < V(n) < 0.3918n$$

## Question

Is  $V(n) \sim vn$  for some  $v$ ? If so, is  $v$  computable?

# Path covers of the divisor graph

# Path covers of the divisor graph

Pomerance, Erdős, Saias and others study the the length of the longest path in the divisor graph of  $[1, n]$ , showing it is  $\asymp \frac{n}{\log n}$ .




# Path covers of the divisor graph

Pomerance, Erdős, Saias and others study the the length of the longest path in the divisor graph of  $[1, n]$ , showing it is  $\asymp \frac{n}{\log n}$ .

Let  $C(n)$  be the least number of disjoint paths that contain all the vertices of this graph.

**Example:**  $C(7) = 2$




# Path covers of the divisor graph

Pomerance, Erdős, Saias and others study the the length of the longest path in the divisor graph of  $[1, n]$ , showing it is  $\asymp \frac{n}{\log n}$ .

Let  $C(n)$  be the least number of disjoint paths that contain all the vertices of this graph.

**Example:**  $C(7) = 2$

The divisor graph of  $[1, 7]$  can be covered by  $\{7, 1, 5\}$  and  $\{3, 6, 2, 4\}$  but it is not possible to use a single path.



# Path covers of the divisor graph

Bounds on the path cover number  $C(n)$  of this divisor graph have improved over time.

# Path covers of the divisor graph

Bounds on the path cover number  $C(n)$  of this divisor graph have improved over time.

**Saias** (2003):  $\frac{n}{6} \leq C(n) \leq \frac{n}{4}$  for sufficiently large  $n$ .

# Path covers of the divisor graph

Bounds on the path cover number  $C(n)$  of this divisor graph have improved over time.

**Saias** (2003):  $\frac{n}{6} \leq C(n) \leq \frac{n}{4}$  for sufficiently large  $n$ .

**Mazet** (2006):  $C(n) \sim \nu n$  for some  $0.1706 \leq \nu \leq 0.2289$ .

# Path covers of the divisor graph

Bounds on the path cover number  $C(n)$  of this divisor graph have improved over time.

**Saias** (2003):  $\frac{n}{6} \leq C(n) \leq \frac{n}{4}$  for sufficiently large  $n$ .

**Mazet** (2006):  $C(n) \sim \nu n$  for some  $0.1706 \leq \nu \leq 0.2289$ .

**Chadozeau** (2008):  $C(n) = \nu n \left( 1 + O \left( \frac{1}{\log \log n \log \log \log n} \right) \right)$ .

# Path covers of the divisor graph

Bounds on the path cover number  $C(n)$  of this divisor graph have improved over time.

**Saias** (2003):  $\frac{n}{6} \leq C(n) \leq \frac{n}{4}$  for sufficiently large  $n$ .

**Mazet** (2006):  $C(n) \sim \nu n$  for some  $0.1706 \leq \nu \leq 0.2289$ .

**Chadozeau** (2008):  $C(n) = \nu n \left( 1 + O \left( \frac{1}{\log \log n \log \log \log n} \right) \right)$ .

## Theorem

$$C(n) = \nu n \left( 1 + O \left( \exp \left( -\sqrt{\left( \frac{1}{6} - \epsilon \right) \log n \log \log n} \right) \right) \right).$$

*The constant  $\nu$  is effectively computable and  $0.17644 < \nu$ .*

THANK YOU!



# Proof of the error term (outline)

# Proof of the error term (outline)

Recall  $c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ P^+(d) < i}} \prod_{t \in [id, (i+1)d)} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$

# Proof of the error term (outline)

Recall  $c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ P^+(d) < i}} \prod_{t \in [id, (i+1)d)} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$

We have

$$Q(n) = \prod_{k=1}^n r(k, n)$$

# Proof of the error term (outline)

$$\text{Recall } c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ P^+(d) < i}} \prod_{t \in [id, (i+1)d)} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$$

We have

$$\begin{aligned} Q(n) &= \prod_{k=1}^n r(k, n) \\ &= c^n \left( \prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\ &\quad \times \exp \left( O \left( \frac{n}{N} + \sum_{i=1}^N \#\{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right) \end{aligned}$$

# Proof Outline

# Proof Outline

$$c^n \left( \prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\ \times \exp \left( O \left( \frac{n}{N} + \sum_{i=1}^N \#\{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right)$$

# Proof Outline

$$\begin{aligned}
 & c^n \left( \prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\
 & \quad \times \exp \left( O \left( \frac{n}{N} + \sum_{i=1}^N \# \{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right) \\
 & = c^n \exp \left( O \left( \sum_{i=1}^N \sum_{d < m} d + \frac{n}{N} + N \# \{k \leq n : d|k, d > M, P^+(d) < N\} \right) \right)
 \end{aligned}$$

# Proof Outline

$$\begin{aligned}
 & c^n \left( \prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\
 & \quad \times \exp \left( O \left( \frac{n}{N} + \sum_{i=1}^N \# \{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right) \\
 & = c^n \exp \left( O \left( \sum_{i=1}^N \sum_{d < m} d + \frac{n}{N} + N \# \{k \leq n : d|k, d > M, P^+(d) < N\} \right) \right)
 \end{aligned}$$



# Proof Outline

$$\begin{aligned}
 & c^n \left( \prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\
 & \quad \times \exp \left( O \left( \frac{n}{N} + \sum_{i=1}^N \# \{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right) \\
 & = c^n \exp \left( O \left( \sum_{i=1}^N \sum_{d < m} d + \frac{n}{N} + N \# \{k \leq n : d|k, d > M, P^+(d) < N\} \right) \right) \\
 & = c^n \exp \left( O \left( NM^2 + \frac{n}{N} + Nn \exp \left( -\frac{\log M}{2 \log N} \right) \right) \right)
 \end{aligned}$$

# Proof Outline

$$\begin{aligned}
 & c^n \left( \prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\
 & \quad \times \exp \left( O \left( \frac{n}{N} + \sum_{i=1}^N \# \{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right) \\
 & = c^n \exp \left( O \left( \sum_{i=1}^N \sum_{d < m} d + \frac{n}{N} + N \# \{k \leq n : d|k, d > M, P^+(d) < N\} \right) \right) \\
 & = c^n \exp \left( O \left( NM^2 + \frac{n}{N} + Nn \exp \left( -\frac{\log M}{2 \log N} \right) \right) \right)
 \end{aligned}$$

Take  $M = n^{1/2-\epsilon}$ ,  $N = \exp((\log n)^{1/2-\epsilon})$ .