

Counting primitive sets and divisor-permutations using divisor graphs

Nathan McNew
Towson University

Loyola Math Department Colloquium
March 22nd, 2023

Primitive Sets

Definition

A set $S \subset \mathbb{N}$ is **primitive** if no element of the set divides another: if $m, n \in S$ are distinct then $m \nmid n$.

Primitive Sets

Definition

A set $S \subset \mathbb{N}$ is **primitive** if no element of the set divides another: if $m, n \in S$ are distinct then $m \nmid n$.

Examples:

- Prime numbers $\mathcal{P} = \{2, 3, 5, \dots\}$.

Primitive Sets

Definition

A set $S \subset \mathbb{N}$ is **primitive** if no element of the set divides another: if $m, n \in S$ are distinct then $m \nmid n$.

Examples:

- Prime numbers $\mathcal{P} = \{2, 3, 5, \dots\}$.
- $\{m, m+1, m+2, \dots, 2m-1\}$.

Primitive Sets

Definition

A set $S \subset \mathbb{N}$ is **primitive** if no element of the set divides another: if $m, n \in S$ are distinct then $m \nmid n$.

Examples:

- Prime numbers $\mathcal{P} = \{2, 3, 5, \dots\}$.
- $\{m, m+1, m+2, \dots, 2m-1\}$.
- Integers with exactly k prime factors,

Primitive Sets

Definition

A set $S \subset \mathbb{N}$ is **primitive** if no element of the set divides another: if $m, n \in S$ are distinct then $m \nmid n$.

Examples:

- Prime numbers $\mathcal{P} = \{2, 3, 5, \dots\}$.
- $\{m, m+1, m+2, \dots, 2m-1\}$.
- Integers with exactly k prime factors,
 $\mathcal{P}_2 = \{4, 6, 9, 10, 14, 15, 21, 22, \dots\}$.

Primitive Sets

Definition

A set $S \subset \mathbb{N}$ is **primitive** if no element of the set divides another: if $m, n \in S$ are distinct then $m \nmid n$.

Examples:

- Prime numbers $\mathcal{P} = \{2, 3, 5, \dots\}$.
- $\{m, m+1, m+2, \dots, 2m-1\}$.
- Integers with exactly k prime factors,
 $\mathcal{P}_2 = \{4, 6, 9, 10, 14, 15, 21, 22, \dots\}$.
- $\{1\}$.

Primitive Sets

Definition

A set $S \subset \mathbb{N}$ is **primitive** if no element of the set divides another: if $m, n \in S$ are distinct then $m \nmid n$.

Examples:

- Prime numbers $\mathcal{P} = \{2, 3, 5, \dots\}$.
- $\{m, m+1, m+2, \dots, 2m-1\}$.
- Integers with exactly k prime factors,
 $\mathcal{P}_2 = \{4, 6, 9, 10, 14, 15, 21, 22, \dots\}$.
- $\{1\}$.
- Primitive abundant numbers.

Primitive abundant numbers

Recall $\sigma(n) = \sum_{d|n} d$ is the sum of divisors function.

Primitive abundant numbers

Recall $\sigma(n) = \sum_{d|n} d$ is the sum of divisors function.

Ancient Greeks called n *perfect* if $\sigma(n) = 2n$, since the divisors less than n sum to n .

Primitive abundant numbers

Recall $\sigma(n) = \sum_{d|n} d$ is the sum of divisors function.

Ancient Greeks called n *perfect* if $\sigma(n) = 2n$, since the divisors less than n sum to n . For example $1+2+3 = 6$ or $1+2+4+7+14 = 28$.

Primitive abundant numbers

Recall $\sigma(n) = \sum_{d|n} d$ is the sum of divisors function.

Ancient Greeks called n *perfect* if $\sigma(n) = 2n$, since the divisors less than n sum to n . For example $1+2+3 = 6$ or $1+2+4+7+14 = 28$.

If $\sigma(n) > 2n$ then n is called *abundant*, if $\sigma(n) < 2n$ it is *deficient*.

Primitive abundant numbers

Recall $\sigma(n) = \sum_{d|n} d$ is the sum of divisors function.

Ancient Greeks called n *perfect* if $\sigma(n) = 2n$, since the divisors less than n sum to n . For example $1+2+3 = 6$ or $1+2+4+7+14 = 28$.

If $\sigma(n) > 2n$ then n is called *abundant*, if $\sigma(n) < 2n$ it is *deficient*.
 $\sigma(20) = 1+2+4+5+10+20 = 42$

Primitive abundant numbers

Recall $\sigma(n) = \sum_{d|n} d$ is the sum of divisors function.

Ancient Greeks called n *perfect* if $\sigma(n) = 2n$, since the divisors less than n sum to n . For example $1+2+3 = 6$ or $1+2+4+7+14 = 28$.

If $\sigma(n) > 2n$ then n is called *abundant*, if $\sigma(n) < 2n$ it is *deficient*.
 $\sigma(20) = 1+2+4+5+10+20 = 42$ $\sigma(21) = 1+3+7+21 = 32$.

Primitive abundant numbers

Recall $\sigma(n) = \sum_{d|n} d$ is the sum of divisors function.

Ancient Greeks called n *perfect* if $\sigma(n) = 2n$, since the divisors less than n sum to n . For example $1+2+3 = 6$ or $1+2+4+7+14 = 28$.

If $\sigma(n) > 2n$ then n is called *abundant*, if $\sigma(n) < 2n$ it is *deficient*.
 $\sigma(20) = 1+2+4+5+10+20 = 42$ $\sigma(21) = 1+3+7+21 = 32$.

Any multiple of an abundant number is abundant (σ is multiplicative)

Primitive abundant numbers

Recall $\sigma(n) = \sum_{d|n} d$ is the sum of divisors function.

Ancient Greeks called n *perfect* if $\sigma(n) = 2n$, since the divisors less than n sum to n . For example $1+2+3 = 6$ or $1+2+4+7+14 = 28$.

If $\sigma(n) > 2n$ then n is called *abundant*, if $\sigma(n) < 2n$ it is *deficient*.
 $\sigma(20) = 1+2+4+5+10+20 = 42$ $\sigma(21) = 1+3+7+21 = 32$.

Any multiple of an abundant number is abundant (σ is multiplicative)

A *primitive abundant number* is an abundant number, all of whose divisors are deficient (20, 70, 88, 104, 272, 304, 368, 464, 550...)

Primitive abundant numbers

Erdős showed that if S is a set of integers whose reciprocal sum $\sum_{s \in S} \frac{1}{s}$ converges, then the set of multiples $\{ns \mid n \in \mathbb{N}, s \in S\}$ has a density.

Primitive abundant numbers

Erdős showed that if S is a set of integers whose reciprocal sum $\sum_{s \in S} \frac{1}{s}$ converges, then the set of multiples $\{ns \mid n \in \mathbb{N}, s \in S\}$ has a density.

He also showed that the reciprocal sum of primitive abundant numbers converges.

Primitive abundant numbers

[Erdős](#) showed that if S is a set of integers whose reciprocal sum $\sum_{s \in S} \frac{1}{s}$ converges, then the set of multiples $\{ns \mid n \in \mathbb{N}, s \in S\}$ has a density.

He also showed that the reciprocal sum of primitive abundant numbers converges. So the abundant numbers have a density. We now know, due to [Kobayashi](#) that this density is about 0.2476...

Primitive abundant numbers

[Erdős](#) showed that if S is a set of integers whose reciprocal sum $\sum_{s \in S} \frac{1}{s}$ converges, then the set of multiples $\{ns \mid n \in \mathbb{N}, s \in S\}$ has a density.

He also showed that the reciprocal sum of primitive abundant numbers converges. So the abundant numbers have a density. We now know, due to [Kobayashi](#) that this density is about 0.2476...

What about primitive sets where the reciprocal sum diverges? For example, the reciprocal sum of prime numbers $\sum_{p \in \mathcal{P}} \frac{1}{p}$ diverges.

Large primitive sets

In 1935 Erdős defined primitive sets as a generalization of primitive abundant numbers, and showed:

Large primitive sets

In 1935 Erdős defined primitive sets as a generalization of primitive abundant numbers, and showed:

Theorem (Erdős)

If $S \neq \{1\}$ is a primitive set then
$$\sum_{n \in S} \frac{1}{n \log n} < \infty.$$

Large primitive sets

In 1935 Erdős defined primitive sets as a generalization of primitive abundant numbers, and showed:

Theorem (Erdős)

If $S \neq \{1\}$ is a primitive set then
$$\sum_{n \in S} \frac{1}{n \log n} < \infty.$$

Considering primes: even though $\sum_{p \in \mathcal{P}} \frac{1}{p}$ diverges, $\sum_{p \in \mathcal{P}} \frac{1}{p \log p}$ converges!

Large primitive sets

In 1935 Erdős defined primitive sets as a generalization of primitive abundant numbers, and showed:

Theorem (Erdős)

If $S \neq \{1\}$ is a primitive set then
$$\sum_{n \in S} \frac{1}{n \log n} < \infty.$$

Considering primes: even though $\sum_{p \in \mathcal{P}} \frac{1}{p}$ diverges, $\sum_{p \in \mathcal{P}} \frac{1}{p \log p}$ converges!

Conjecture (Erdős, 1988)

If $S \neq \{1\}$ is a primitive set then
$$\sum_{n \in S} \frac{1}{n \log n} \leq \sum_{p \in \mathcal{P}} \frac{1}{p \log p} = 1.63 \dots$$

Large primitive sets

In 1935 Erdős defined primitive sets as a generalization of primitive abundant numbers, and showed:

Theorem (Erdős)

If $S \neq \{1\}$ is a primitive set then
$$\sum_{n \in S} \frac{1}{n \log n} < \infty.$$

Considering primes: even though $\sum_{p \in \mathcal{P}} \frac{1}{p}$ diverges, $\sum_{p \in \mathcal{P}} \frac{1}{p \log p}$ converges!

~~Conjecture (Erdős, 1988)~~ Theorem (Lichtman, 2022)

If $S \neq \{1\}$ is a primitive set then
$$\sum_{n \in S} \frac{1}{n \log n} \leq \sum_{p \in \mathcal{P}} \frac{1}{p \log p} = 1.63 \dots$$

The divisor graph

The divisor graph

The **divisor graph** of $\{1, 2, \dots, n\}$, denoted $\mathcal{D}_{[1,n]}$, is the graph on vertices v_1, v_2, \dots, v_n and an edge between v_i and v_j if $i \mid j$ (or $j \mid i$).

The divisor graph


The **divisor graph** of $\{1, 2, \dots, n\}$, denoted $\mathcal{D}_{[1,n]}$, is the graph on vertices v_1, v_2, \dots, v_n and an edge between v_i and v_j if $i \mid j$ (or $j \mid i$).

Example: $n = 9$

The divisor graph

The **divisor graph** of $\{1, 2, \dots, n\}$, denoted $\mathcal{D}_{[1,n]}$, is the graph on vertices v_1, v_2, \dots, v_n and an edge between v_i and v_j if $i \mid j$ (or $j \mid i$).

Example: $n = 9$




The divisor graph

The **divisor graph** of $\{1, 2, \dots, n\}$, denoted $\mathcal{D}_{[1,n]}$, is the graph on vertices v_1, v_2, \dots, v_n and an edge between v_i and v_j if $i \mid j$ (or $j \mid i$).

Example: $n = 9$

In general $\mathcal{D}_{[k,n]}$ is the subgraph of $\mathcal{D}_{[1,n]}$ with only vertices in $[k, n]$.




The divisor graph

The **divisor graph** of $\{1, 2, \dots, n\}$, denoted $\mathcal{D}_{[1,n]}$, is the graph on vertices v_1, v_2, \dots, v_n and an edge between v_i and v_j if $i \mid j$ (or $j \mid i$).

Example: $n = 9$

In general $\mathcal{D}_{[k,n]}$ is the subgraph of $\mathcal{D}_{[1,n]}$ with only vertices in $[k, n]$.




Independent subsets of a divisor graph are primitive sets of integers.

The divisor graph

The **divisor graph** of $\{1, 2, \dots, n\}$, denoted $\mathcal{D}_{[1,n]}$, is the graph on vertices v_1, v_2, \dots, v_n and an edge between v_i and v_j if $i \mid j$ (or $j \mid i$).

Example: $n = 9$

In general $\mathcal{D}_{[k,n]}$ is the subgraph of $\mathcal{D}_{[1,n]}$ with only vertices in $[k, n]$.




Independent subsets of a divisor graph are primitive sets of integers.

The divisor graph

The **divisor graph** of $\{1, 2, \dots, n\}$, denoted $\mathcal{D}_{[1,n]}$, is the graph on vertices v_1, v_2, \dots, v_n and an edge between v_i and v_j if $i \mid j$ (or $j \mid i$).

Example: $n = 9$

In general $\mathcal{D}_{[k,n]}$ is the subgraph of $\mathcal{D}_{[1,n]}$ with only vertices in $[k, n]$.




Independent subsets of a divisor graph are primitive sets of integers.

The divisor graph

The **divisor graph** of $\{1, 2, \dots, n\}$, denoted $\mathcal{D}_{[1,n]}$, is the graph on vertices v_1, v_2, \dots, v_n and an edge between v_i and v_j if $i \mid j$ (or $j \mid i$).

Example: $n = 9$

In general $\mathcal{D}_{[k,n]}$ is the subgraph of $\mathcal{D}_{[1,n]}$ with only vertices in $[k, n]$.



Independent subsets of a divisor graph are primitive sets of integers.

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
-----	---	-------

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	13

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	13
6		

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	13
6		17

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	13
6		17

How many primitive subsets of $\{1, 2 \dots n\}$ are there?

Counting primitive sets

Let $Q(n)$ count the primitive sets with largest element at most n .


Counting primitive sets

Let $Q(n)$ count the primitive sets with largest element at most n .

A051026	Number of primitive subsequences of {1, 2, ..., n}.	5
	1, 2, 3, 5, 7, 13, 17, 33, 45, 73, 103, 205, 253, 505, 733, 1133, 1529, 3057, 3897, 7793, 10241, 16513, 24593, 49185, 59265, 109297, 163369, 262489, 355729, 711457, 879937, 1759873, 2360641, 3908545, 5858113, 10534337, 12701537, 25403073, 38090337, 63299265, 81044097, 162088193, 205482593, 410965185, 570487233, 855676353 (list ; graph ; refs ; listen ; history ; text ; internal format)	
OFFSET	0, 2	
COMMENTS	$a(n)$ counts all subsequences of $\{1, \dots, n\}$ in which no term divides any other. If n is a prime $a(n) = 2^*a(n-1)-1$ because for each subsequence s counted by $a(n-1)$ two different subsequences are counted by $a(n)$: s and s, n . There is only one exception: $1, n$ is not a primitive subsequence because 1 divides n . For all $n > 1$: $a(n) < 2^*a(n-1)$. - Alois P. Heinz , Mar 07 2011	


Bounds

Cameron and Erdős considered the function $Q(n)$ in 1990.



Bounds

Cameron and Erdős considered the function $Q(n)$ in 1990.




(Almost) trivially:

$$(\sqrt{2})^n \leq Q(n) \leq 2^n$$

Bounds

Cameron and Erdős considered the function $Q(n)$ in 1990.



(Almost) trivially:

$$(\sqrt{2})^n \leq Q(n) \leq 2^n$$

Every subset of $\left(\frac{n}{2}, n\right]$ is primitive. There are $2^{\lceil \frac{n}{2} \rceil} \geq \sqrt{2}^n$ such subsets.

Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

Conjecture: $\lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists.

Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

Conjecture: $\lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists.

Theorem (Angelo, 2017)

$c = \lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists. Equivalently, $Q(n) = c^{n+o(n)}$.



Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

Conjecture: $\lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists.

Theorem (Angelo, 2017)

$c = \lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists. Equivalently, $Q(n) = c^{n+o(n)}$.



$o(n)$ means an error term $E(n)$ where $\lim_{n \rightarrow \infty} \frac{E(n)}{n} = 0$.

Bounds

Cameron and Erdős (1990) improve these bounds to:

$$1.5596^n < Q(n) < 1.60^n$$

Conjecture: $\lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists.

Theorem (Angelo, 2017)

$c = \lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists. Equivalently, $Q(n) = c^{n+o(n)}$.



$o(n)$ means an error term $E(n)$ where $\lim_{n \rightarrow \infty} \frac{E(n)}{n} = 0$.

His proof gives no insight on the value of this constant.


Idea: count primitive sets working backward from n

Idea: count primitive sets working backward from n

Every subset of $(\frac{n}{2}, n]$ is primitive, so $Q(n) \geq 2^{n/2}$.


Idea: count primitive sets working backward from n

Every subset of $(\frac{n}{2}, n]$ is primitive, so $Q(n) \geq 2^{n/2}$. As a graph, $\mathcal{D}_{(\frac{n}{2}, n]}$:



Idea: count primitive sets working backward from n


Every subset of $(\frac{n}{2}, n]$ is primitive, so $Q(n) \geq 2^{n/2}$. As a graph, $\mathcal{D}_{(\frac{n}{2}, n]}$:



Each integer can either be included or not included (2 possibilities).

Idea: count primitive sets working backward from n

Every subset of $(\frac{n}{2}, n]$ is primitive, so $Q(n) \geq 2^{n/2}$. As a graph, $\mathcal{D}_{(\frac{n}{2}, n]}$:




Each integer can either be included or not included (2 possibilities).

Try to make this bigger by considering integers in $(\frac{n}{3}, \frac{n}{2}]$.


Idea: count primitive sets working backward from n

Every subset of $(\frac{n}{2}, n]$ is primitive, so $Q(n) \geq 2^{n/2}$. As a graph, $\mathcal{D}_{(\frac{n}{2}, n]}$:



Each integer can either be included or not included (2 possibilities).

Try to make this bigger by considering integers in $(\frac{n}{3}, \frac{n}{2}]$. $\mathcal{D}_{(\frac{n}{3}, \frac{n}{2}]}$:



Constructing Primitive Sets

For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we can include $k, 2k$ or neither.

Constructing Primitive Sets

For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we can include $k, 2k$ or neither. (3 possibilities)

Constructing Primitive Sets

For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we can include $k, 2k$ or neither. (3 possibilities)

Each new connected component of the graph is equivalent to counting primitive subsets of $\{1, 2\}$

Constructing Primitive Sets

For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we can include $k, 2k$ or neither. (3 possibilities)

Each new connected component of the graph is equivalent to counting primitive subsets of $\{1, 2\}$ or independent subsets of:

Constructing Primitive Sets

For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we can include $k, 2k$ or neither. (3 possibilities)

Each new connected component of the graph is equivalent to counting primitive subsets of $\{1, 2\}$ or independent subsets of:



Constructing Primitive Sets

For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we can include $k, 2k$ or neither. (3 possibilities)

Each new connected component of the graph is equivalent to counting primitive subsets of $\{1, 2\}$ or independent subsets of:



Use this to improve the lower bound for $Q(n)$.

Constructing Primitive Sets

For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we can include $k, 2k$ or neither. (3 possibilities)

Each new connected component of the graph is equivalent to counting primitive subsets of $\{1, 2\}$ or independent subsets of:



Use this to improve the lower bound for $Q(n)$. For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we multiply by $\frac{3}{2}$, the ratio of primitive subsets of $\{k, 2k\}$, to those of just $\{2k\}$. ($2k$ was in $(\frac{n}{2}, n]$, so those were already counted.)

Constructing Primitive Sets

For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we can include $k, 2k$ or neither. (3 possibilities)

Each new connected component of the graph is equivalent to counting primitive subsets of $\{1, 2\}$ or independent subsets of:



Use this to improve the lower bound for $Q(n)$. For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we multiply by $\frac{3}{2}$, the ratio of primitive subsets of $\{k, 2k\}$, to those of just $\{2k\}$. ($2k$ was in $(\frac{n}{2}, n]$, so those were already counted.)

$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6}$$

Constructing Primitive Sets

For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we can include $k, 2k$ or neither. (3 possibilities)

Each new connected component of the graph is equivalent to counting primitive subsets of $\{1, 2\}$ or independent subsets of:



Use this to improve the lower bound for $Q(n)$. For each $k \in (\frac{n}{3}, \frac{n}{2}]$ we multiply by $\frac{3}{2}$, the ratio of primitive subsets of $\{k, 2k\}$, to those of just $\{2k\}$. ($2k$ was in $(\frac{n}{2}, n]$, so those were already counted.)

$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} = 2^{n/3} 3^{n/6} \approx 1.5131^n.$$

Constructing primitive sets: pressing on!

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k .

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$.

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:


Odd k :

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:

Odd k :




5 primitive subsets of $\{k, 2k, 3k\}$ replace 4 of just $\{2k, 3k\}$.

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:

Odd k :



5 primitive subsets of $\{k, 2k, 3k\}$ replace 4 of just $\{2k, 3k\}$.


Even $k = 2\ell$:

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:

Odd k :



5 primitive subsets of $\{k, 2k, 3k\}$ replace 4 of just $\{2k, 3k\}$.


Even $k = 2\ell$: Now $\frac{3k}{2} = 3\ell$ is also an integer.

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:

Odd k :



5 primitive subsets of $\{k, 2k, 3k\}$ replace 4 of just $\{2k, 3k\}$.

Even $k = 2\ell$: Now $\frac{3k}{2} = 3\ell$ is also an integer.


Consider subsets of $k=2\ell, 3\ell, 2k=4\ell, 3k=6\ell$.

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:


Odd k :



5 primitive subsets of $\{k, 2k, 3k\}$ replace 4 of just $\{2k, 3k\}$.

Even $k = 2\ell$: Now $\frac{3k}{2} = 3\ell$ is also an integer.

Consider subsets of $k=2\ell$, 3ℓ , $2k=4\ell$, $3k=6\ell$.




Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:

Odd k :




5 primitive subsets of $\{k, 2k, 3k\}$ replace 4 of just $\{2k, 3k\}$.

Even $k = 2\ell$: Now $\frac{3k}{2} = 3\ell$ is also an integer.

Consider subsets of $k=2\ell$, 3ℓ , $2k=4\ell$, $3k=6\ell$.

Find 8 primitive subsets:

$\{\}, \{2\ell\}, \{3\ell\}, \{4\ell\}, \{6\ell\},$
 $\{2\ell, 3\ell\}, \{3\ell, 4\ell\}, \{4\ell, 6\ell\}$




Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:

Odd k :




5 primitive subsets of $\{k, 2k, 3k\}$ replace 4 of just $\{2k, 3k\}$.

Even $k = 2\ell$: Now $\frac{3k}{2} = 3\ell$ is also an integer.

Consider subsets of $k=2\ell$, 3ℓ , $2k=4\ell$, $3k=6\ell$.

Find 8 primitive subsets:

$\{\}, \{2\ell\}, \{3\ell\}, \{4\ell\}, \{6\ell\},$
 $\{2\ell, 3\ell\}, \{3\ell, 4\ell\}, \{4\ell, 6\ell\}$




Replacing the 6 subsets of only $\{3\ell, 4\ell, 6\ell\}$.

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:

Odd k :




5 primitive subsets of $\{k, 2k, 3k\}$ replace 4 of just $\{2k, 3k\}$.

Even $k = 2\ell$: Now $\frac{3k}{2} = 3\ell$ is also an integer.

Consider subsets of $k=2\ell$, 3ℓ , $2k=4\ell$, $3k=6\ell$.

Find 8 primitive subsets:

$\{\}, \{2\ell\}, \{3\ell\}, \{4\ell\}, \{6\ell\},$
 $\{2\ell, 3\ell\}, \{3\ell, 4\ell\}, \{4\ell, 6\ell\}$



Replacing the 6 subsets of only $\{3\ell, 4\ell, 6\ell\}$.


$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} \left(\frac{5}{4}\right)^{n/24} \left(\frac{8}{6}\right)^{n/24}$$

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:

Odd k :




5 primitive subsets of $\{k, 2k, 3k\}$ replace 4 of just $\{2k, 3k\}$.

Even $k = 2\ell$: Now $\frac{3k}{2} = 3\ell$ is also an integer.

Consider subsets of $k=2\ell$, 3ℓ , $2k=4\ell$, $3k=6\ell$.

Find 8 primitive subsets:

$\{\}, \{2\ell\}, \{3\ell\}, \{4\ell\}, \{6\ell\},$
 $\{2\ell, 3\ell\}, \{3\ell, 4\ell\}, \{4\ell, 6\ell\}$



Replacing the 6 subsets of only $\{3\ell, 4\ell, 6\ell\}$.


$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} \left(\frac{5}{4}\right)^{n/24} \left(\frac{8}{6}\right)^{n/24} = 2^{5n/24} 3^{n/8} 5^{n/24}$$

Constructing primitive sets: pressing on!

Work backward, considering progressively smaller k . Take $k \in (\frac{n}{4}, \frac{n}{3}]$.

Now we need to worry about $k, 2k, 3k$. Actually two scenarios:

Odd k :




5 primitive subsets of $\{k, 2k, 3k\}$ replace 4 of just $\{2k, 3k\}$.

Even $k = 2\ell$: Now $\frac{3k}{2} = 3\ell$ is also an integer.

Consider subsets of $k=2\ell$, 3ℓ , $2k=4\ell$, $3k=6\ell$.

Find 8 primitive subsets:

$\{\}, \{2\ell\}, \{3\ell\}, \{4\ell\}, \{6\ell\},$
 $\{2\ell, 3\ell\}, \{3\ell, 4\ell\}, \{4\ell, 6\ell\}$



Replacing the 6 subsets of only $\{3\ell, 4\ell, 6\ell\}$.

$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} \left(\frac{5}{4}\right)^{n/24} \left(\frac{8}{6}\right)^{n/24} = 2^{5n/24} 3^{n/8} 5^{n/24} \approx 1.5456^n$$

One more wrinkle


One more wrinkle

Consider $k \in \left(\frac{n}{5}, \frac{n}{4}\right]$, the next interval.

One more wrinkle

Consider $k \in (\frac{n}{5}, \frac{n}{4}]$, the next interval.


If $k = 2\ell$ is even we must consider $2\ell, 4\ell, 6\ell, 8\ell$, as well as 3ℓ .




One more wrinkle

Consider $k \in (\frac{n}{5}, \frac{n}{4}]$, the next interval.

If $k = 2\ell$ is even we must consider $2\ell, 4\ell, 6\ell, 8\ell$, as well as 3ℓ .




However, if $k \leq \frac{2n}{9}$ then $9\ell \leq n$ must also be considered.




One more wrinkle

Consider $k \in (\frac{n}{5}, \frac{n}{4}]$, the next interval.

If $k = 2\ell$ is even we must consider $2\ell, 4\ell, 6\ell, 8\ell$, as well as 3ℓ .



However, if $k \leq \frac{2n}{9}$ then $9\ell \leq n$ must also be considered.




So now we consider $2\ell, 3\ell, 4\ell, 6\ell, 8\ell$ and 9ℓ .


One more wrinkle

Consider $k \in (\frac{n}{5}, \frac{n}{4}]$, the next interval.

If $k = 2\ell$ is even we must consider $2\ell, 4\ell, 6\ell, 8\ell$, as well as 3ℓ .



However, if $k \leq \frac{2n}{9}$ then $9\ell \leq n$ must also be considered.



So now we consider $2\ell, 3\ell, 4\ell, 6\ell, 8\ell$ and 9ℓ .

So $k \in (\frac{n}{5}, \frac{2n}{9}]$ behave differently than those $k \in (\frac{2n}{9}, \frac{n}{4}]$

Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

- $1 \leq r(k, n) \leq 2$

Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

- $1 \leq r(k, n) \leq 2$
- $r(k, n)$ only depends on the structure of the connected component of k in the divisibility graph of $[k, n]$.

Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

- $1 \leq r(k, n) \leq 2$
- $r(k, n)$ only depends on the structure of the connected component of k in the divisibility graph of $[k, n]$.
- $Q(n) = \prod_{i=1}^n r(i, n)$

Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

- $1 \leq r(k, n) \leq 2$
- $r(k, n)$ only depends on the structure of the connected component of k in the divisibility graph of $[k, n]$.
- $Q(n) = \prod_{i=1}^n r(i, n)$ (Telescoping product)

Computing $Q(n)$

$$\text{Let } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}.$$

Observations:

- $1 \leq r(k, n) \leq 2$
- $r(k, n)$ only depends on the structure of the connected component of k in the divisibility graph of $[k, n]$.
- $Q(n) = \prod_{i=1}^n r(i, n)$ (Telescoping product)
- Goal: group together equal terms in this product.

Key Observations about $r(k, n)$

Fix k, n .

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$.

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$. Multiples of k in $[k, n]$ are $k, 2k, \dots ik$.

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$. Multiples of k in $[k, n]$ are $k, 2k, \dots, ik$.

If one integer from $[k, n]$ divides another the ratio is an integer $\leq i$.

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$. Multiples of k in $[k, n]$ are $k, 2k, \dots, ik$.

If one integer from $[k, n]$ divides another the ratio is an integer $\leq i$.

Connected components of $\mathcal{D}_{[k, n]}$ are divisible by the same primes $> i$.

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$. Multiples of k in $[k, n]$ are $k, 2k, \dots ik$.

If one integer from $[k, n]$ divides another the ratio is an integer $\leq i$.

Connected components of $\mathcal{D}_{[k, n]}$ are divisible by the same primes $> i$.

Let ℓ be the the product of all of the primes dividing k greater than i .

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$. Multiples of k in $[k, n]$ are $k, 2k, \dots, ik$.

If one integer from $[k, n]$ divides another the ratio is an integer $\leq i$.

Connected components of $\mathcal{D}_{[k, n]}$ are divisible by the same primes $> i$.

Let ℓ be the the product of all of the primes dividing k greater than i .

Every integer in the connected component of k is divisible by ℓ .

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$. Multiples of k in $[k, n]$ are $k, 2k, \dots, ik$.

If one integer from $[k, n]$ divides another the ratio is an integer $\leq i$.

Connected components of $\mathcal{D}_{[k, n]}$ are divisible by the same primes $> i$.

Let ℓ be the the product of all of the primes dividing k greater than i .

Every integer in the connected component of k is divisible by ℓ .

Let $d = \frac{k}{\ell}$ be the i -**smooth** part of k .

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$. Multiples of k in $[k, n]$ are $k, 2k, \dots, ik$.

If one integer from $[k, n]$ divides another the ratio is an integer $\leq i$.

Connected components of $\mathcal{D}_{[k, n]}$ are divisible by the same primes $> i$.

Let ℓ be the product of all of the primes dividing k greater than i .

Every integer in the connected component of k is divisible by ℓ .

Let $d = \frac{k}{\ell}$ be the i -**smooth** part of k .

Let $t = \lfloor \frac{n}{\ell} \rfloor$. So that $t\ell$ is the largest integer $\leq n$ divisible by ℓ .

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$. Multiples of k in $[k, n]$ are $k, 2k, \dots, ik$.

If one integer from $[k, n]$ divides another the ratio is an integer $\leq i$.

Connected components of $\mathcal{D}_{[k, n]}$ are divisible by the same primes $> i$.

Let ℓ be the product of all of the primes dividing k greater than i .

Every integer in the connected component of k is divisible by ℓ .

Let $d = \frac{k}{\ell}$ be the i -**smooth** part of k .

Let $t = \lfloor \frac{n}{\ell} \rfloor$. So that $t\ell$ is the largest integer $\leq n$ divisible by ℓ .

Then $r(k, n) = r(d\ell, t\ell)$

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$. Multiples of k in $[k, n]$ are $k, 2k, \dots ik$.

If one integer from $[k, n]$ divides another the ratio is an integer $\leq i$.

Connected components of $\mathcal{D}_{[k, n]}$ are divisible by the same primes $> i$.

Let ℓ be the the product of all of the primes dividing k greater than i .

Every integer in the connected component of k is divisible by ℓ .

Let $d = \frac{k}{\ell}$ be the i -**smooth** part of k .

Let $t = \lfloor \frac{n}{\ell} \rfloor$. So that $t\ell$ is the largest integer $\leq n$ divisible by ℓ .

Then $r(k, n) = r(d\ell, t\ell) = r(d, t)$.

Key Observations about $r(k, n)$

Fix k, n . Let $i = \lfloor \frac{n}{k} \rfloor$. Multiples of k in $[k, n]$ are $k, 2k, \dots, ik$.

If one integer from $[k, n]$ divides another the ratio is an integer $\leq i$.

Connected components of $\mathcal{D}_{[k, n]}$ are divisible by the same primes $> i$.

Let ℓ be the the product of all of the primes dividing k greater than i .

Every integer in the connected component of k is divisible by ℓ .

Let $d = \frac{k}{\ell}$ be the i -**smooth** part of k .

Let $t = \lfloor \frac{n}{\ell} \rfloor$. So that $t\ell$ is the largest integer $\leq n$ divisible by ℓ .

Then $r(k, n) = r(d\ell, t\ell) = r(d, t)$.

As $n \rightarrow \infty$, the number of $k \leq n$ sharing the same “ d ” and “ t ” is

$$\frac{n + o(n)}{t(t+1)} \prod_{p \leq i} \left(1 - \frac{1}{p}\right).$$

Approximating $Q(n)$

$r(k, n) = r(d, t)$, $i = \lfloor \frac{n}{k} \rfloor$, d is the i -smooth part of k .

Approximating $Q(n)$

$r(k, n) = r(d, t)$, $i = \lfloor \frac{n}{k} \rfloor$, d is the i -smooth part of k .

$$Q(n) = \prod_{k=1}^n r(k, n)$$

Approximating $Q(n)$

$r(k, n) = r(d, t)$, $i = \lfloor \frac{n}{k} \rfloor$, d is the i -smooth part of k .

$$Q(n) = \prod_{k=1}^n r(k, n) = \prod_{i=1}^n$$

Approximating $Q(n)$

$r(k, n) = r(d, t)$, $i = \lfloor \frac{n}{k} \rfloor$, d is the i -smooth part of k .

$$Q(n) = \prod_{k=1}^n r(k, n) = \prod_{i=1}^n \prod_{\substack{d \\ d \text{ is } i\text{-smooth}}}^n$$

Approximating $Q(n)$

$r(k, n) = r(d, t)$, $i = \lfloor \frac{n}{k} \rfloor$, d is the i -smooth part of k .

$$Q(n) = \prod_{k=1}^n r(k, n) = \prod_{i=1}^n \prod_{\substack{d \\ d \text{ is } i\text{-smooth}}} \prod_{t=id}^{(i+1)d-1}$$

Approximating $Q(n)$

$r(k, n) = r(d, t)$, $i = \lfloor \frac{n}{k} \rfloor$, d is the i -smooth part of k .

$$Q(n) = \prod_{k=1}^n r(k, n) = \prod_{i=1}^n \prod_{\substack{d \\ d \text{ is } i\text{-smooth}}} \prod_{t=id}^{(i+1)d-1} r(d, t)^{\frac{n+o(n)}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$$

Approximating $Q(n)$

$r(k, n) = r(d, t)$, $i = \lfloor \frac{n}{k} \rfloor$, d is the i -smooth part of k .

$$Q(n) = \prod_{k=1}^n r(k, n) = \prod_{i=1}^n \prod_{\substack{d \\ d \text{ is } i\text{-smooth}}} \prod_{t=id}^{(i+1)d-1} r(d, t)^{\frac{n+o(n)}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$$

Set $c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ d \text{ is } i\text{-smooth}}} \prod_{t=id}^{(i+1)d-1} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$

Approximating $Q(n)$

$r(k, n) = r(d, t)$, $i = \lfloor \frac{n}{k} \rfloor$, d is the i -smooth part of k .

$$Q(n) = \prod_{k=1}^n r(k, n) = \prod_{i=1}^n \prod_{\substack{d \\ d \text{ is } i\text{-smooth}}} \prod_{t=id}^{(i+1)d-1} r(d, t)^{\frac{n+o(n)}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$$

$$\text{Set } c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ d \text{ is } i\text{-smooth}}} \prod_{t=id}^{(i+1)d-1} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$$

We find that $Q(n) = c^{n+o(n)}$.

Approximating $Q(n)$

Theorem

As $n \rightarrow \infty$ the number of primitive subsets of $[1, n]$ is

$$Q(n) = c^{n+o\left(n \exp\left(-\sqrt{\frac{1}{2} \log n \log \log n}\right)\right)}$$

Approximating $Q(n)$

Theorem

As $n \rightarrow \infty$ the number of primitive subsets of $[1, n]$ is

$$Q(n) = c^{n+o\left(n \exp\left(-\sqrt{\frac{1}{2} \log n \log \log n}\right)\right)}$$

or

$$\log Q(n) = n \log c + o\left(\frac{n}{e^{\sqrt{\frac{1}{2} \log n \log \log n}}}\right).$$

Approximating $Q(n)$

Theorem

As $n \rightarrow \infty$ the number of primitive subsets of $[1, n]$ is

$$Q(n) = c^{n+o\left(n \exp\left(-\sqrt{\frac{1}{2} \log n \log \log n}\right)\right)}$$

or

$$\log Q(n) = n \log c + o\left(\frac{n}{e^{\sqrt{\frac{1}{2} \log n \log \log n}}}\right).$$

The constant $c = \prod_{i=1}^{\infty} \prod_{\substack{d \text{ is } i\text{-smooth}}} \prod_{t \in [id, (i+1)d)} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}$
is bounded between $1.5729 < c < 1.5745$.

A general theorem

A general theorem

Theorem (M.)

Suppose $f(k, n)$ depends only on the connected component of k in $\mathcal{D}_{[k, n]}$ and $|f(k, n)| \leq A$ for some fixed A . Then

$$\sum_{a=1}^n f(a, n) = nC_f + o\left(A n \exp\left(-\sqrt{\frac{1}{2} \log n \log \log n}\right)\right)$$

where

$$C_f = \sum_{i=1}^{\infty} \sum_{\substack{d \\ d \text{ is } i\text{-smooth}}} \sum_{t \in [id, (i+1)d)} \left(\frac{f(d, t)}{t(t+1)} \prod_{p \leq i} \frac{p-1}{p} \right).$$

An application: Path covers of the divisor graph

An application: Path covers of the divisor graph

Pomerance, Erdős, Saias and others study the the length of the longest path in the divisor graph of $[1, n]$, showing it is $\asymp \frac{n}{\log n}$.

An application: Path covers of the divisor graph

Pomerance, Erdős, Saias and others study the the length of the longest path in the divisor graph of $[1, n]$, showing it is $\asymp \frac{n}{\log n}$.


Let $C(n)$ be the least number of disjoint paths that contain all the vertices of this graph.

An application: Path covers of the divisor graph

Pomerance, Erdős, Saias and others study the the length of the longest path in the divisor graph of $[1, n]$, showing it is $\asymp \frac{n}{\log n}$.

Let $C(n)$ be the least number of disjoint paths that contain all the vertices of this graph.

Example: $C(9) = 2$




An application: Path covers of the divisor graph

Pomerance, Erdős, Saias and others study the the length of the longest path in the divisor graph of $[1, n]$, showing it is $\asymp \frac{n}{\log n}$.

Let $C(n)$ be the least number of disjoint paths that contain all the vertices of this graph.

Example: $C(9) = 2$

The divisor graph $\mathcal{D}_{[1,9]}$ can be covered by $\{7, 1, 5\}$ and $\{9, 3, 6, 2, 4, 8\}$ but it is not possible to use a single path.



An application: Path covers of the divisor graph

Bounds on the path cover number $C(n)$ of $\mathcal{D}_{[1,n]}$ have improved.

An application: Path covers of the divisor graph

Bounds on the path cover number $C(n)$ of $\mathcal{D}_{[1,n]}$ have improved.



Theorem (Saia (2003))

$$\frac{n}{6} \leq C(n) \leq \frac{n}{4} \text{ for sufficiently large } n.$$

An application: Path covers of the divisor graph

Bounds on the path cover number $C(n)$ of $\mathcal{D}_{[1,n]}$ have improved.



Theorem ([Saia](#)s (2003))

$$\frac{n}{6} \leq C(n) \leq \frac{n}{4} \text{ for sufficiently large } n.$$

Theorem ([Mazet](#) (2006))

$$C(n) = \nu n + o(n) \text{ for} \\ 0.1706 \leq \nu \leq 0.2289.$$

An application: Path covers of the divisor graph

Bounds on the path cover number $C(n)$ of $\mathcal{D}_{[1,n]}$ have improved.



Theorem ([Saia](#)s (2003))

$$\frac{n}{6} \leq C(n) \leq \frac{n}{4} \text{ for sufficiently large } n.$$

Theorem ([Mazet](#) (2006))

$$C(n) = \nu n + o(n) \text{ for } 0.1706 \leq \nu \leq 0.2289.$$

Theorem ([Chadozeau](#) (2008))

$$C(n) = \nu n \left(1 + O\left(\frac{1}{\log \log n \log \log \log n}\right)\right).$$



A Telescoping Sum

Theorem ([Mazet](#), [Chadozeau](#))

$$C(n) = \nu n + O\left(\frac{n}{\log \log n \log \log \log n}\right), \quad 0.1706 \leq \nu \leq 0.2289.$$

A Telescoping Sum

Theorem (Mazet, Chadozeau)

$$C(n) = \nu n + O\left(\frac{n}{\log \log n \log \log \log n}\right), \quad 0.1706 \leq \nu \leq 0.2289.$$

Set $f(k, n) = \#\{\text{paths to cover } \mathcal{D}_{[k, n]}\} - \#\{\text{paths to cover } \mathcal{D}_{[k+1, n]}\}.$

A Telescoping Sum

Theorem (Mazet, Chadozeau)

$$C(n) = \nu n + O\left(\frac{n}{\log \log n \log \log \log n}\right), \quad 0.1706 \leq \nu \leq 0.2289.$$

Set $f(k, n) = \#\{\text{paths to cover } \mathcal{D}_{[k, n]}\} - \#\{\text{paths to cover } \mathcal{D}_{[k+1, n]}\}$.
 $f(k, n) \leq 1$: Can always cover the vertex k using a new path $\{k\}$.

A Telescoping Sum

Theorem (Mazet, Chadozeau)

$$C(n) = \nu n + O\left(\frac{n}{\log \log n \log \log \log n}\right), \quad 0.1706 \leq \nu \leq 0.2289.$$

Set $f(k, n) = \#\{\text{paths to cover } \mathcal{D}_{[k,n]}\} - \#\{\text{paths to cover } \mathcal{D}_{[k+1,n]}\}.$

$f(k, n) \leq 1$: Can always cover the vertex k using a new path $\{k\}.$

$f(k, n) \geq -1$: Can decrease by one if two paths are connected by $k.$

A Telescoping Sum

Theorem (Mazet, Chadozeau)

$$C(n) = \nu n + O\left(\frac{n}{\log \log n \log \log \log n}\right), \quad 0.1706 \leq \nu \leq 0.2289.$$

Set $f(k, n) = \#\{\text{paths to cover } \mathcal{D}_{[k,n]}\} - \#\{\text{paths to cover } \mathcal{D}_{[k+1,n]}\}.$

$f(k, n) \leq 1$: Can always cover the vertex k using a new path $\{k\}.$

$f(k, n) \geq -1$: Can decrease by one if two paths are connected by $k.$

$f(k, n)$ depends only on the connected component of k in $\mathcal{D}_{[k,n]}.$

A Telescoping Sum

Theorem (Mazet, Chadozeau)

$$C(n) = \nu n + O\left(\frac{n}{\log \log n \log \log \log n}\right), \quad 0.1706 \leq \nu \leq 0.2289.$$

Set $f(k, n) = \#\{\text{paths to cover } \mathcal{D}_{[k,n]}\} - \#\{\text{paths to cover } \mathcal{D}_{[k+1,n]}\}.$

$f(k, n) \leq 1$: Can always cover the vertex k using a new path $\{k\}.$

$f(k, n) \geq -1$: Can decrease by one if two paths are connected by $k.$

$f(k, n)$ depends only on the connected component of k in $\mathcal{D}_{[k,n]}.$

$$C(n) = \sum_{k=1}^n f(k, n)$$

A Telescoping Sum

Theorem (Mazet, Chadozeau)

$$C(n) = \nu n + O\left(\frac{n}{\log \log n \log \log \log n}\right), \quad 0.1706 \leq \nu \leq 0.2289.$$

Set $f(k, n) = \#\{\text{paths to cover } \mathcal{D}_{[k,n]}\} - \#\{\text{paths to cover } \mathcal{D}_{[k+1,n]}\}$.

$f(k, n) \leq 1$: Can always cover the vertex k using a new path $\{k\}$.

$f(k, n) \geq -1$: Can decrease by one if two paths are connected by k .

$f(k, n)$ depends only on the connected component of k in $\mathcal{D}_{[k,n]}$.

$$C(n) = \sum_{k=1}^n f(k, n)$$

Theorem

$$C(n) = \nu n + o\left(\frac{n}{\exp\left(\sqrt{\frac{1}{2} \log n \log \log n}\right)}\right) \text{ and } 0.1909 < \nu < 0.2179.$$

Divisor-permutations

Recently, [Pomerance](#) asked how many permutations σ have the property that either $\sigma(i)|i$ or $i|\sigma(i)$ for every index i .

Divisor-permutations

Recently, [Pomerance](#) asked how many permutations σ have the property that either $\sigma(i)|i$ or $i|\sigma(i)$ for every index i .

Call these **divisor-permutations** and denote their count by $D(n)$.

Divisor-permutations

Recently, [Pomerance](#) asked how many permutations σ have the property that either $\sigma(i)|i$ or $i|\sigma(i)$ for every index i .

Call these **divisor-permutations** and denote their count by $D(n)$.

Example: $n = 4$, 3412

Divisor-permutations

Recently, [Pomerance](#) asked how many permutations σ have the property that either $\sigma(i)|i$ or $i|\sigma(i)$ for every index i .

Call these **divisor-permutations** and denote their count by $D(n)$.

Example: $n = 4$, 3412 1234 1432 2431 2134 3214 4231 4132

Divisor-permutations

Recently, [Pomerance](#) asked how many permutations σ have the property that either $\sigma(i)|i$ or $i|\sigma(i)$ for every index i .

Call these **divisor-permutations** and denote their count by $D(n)$.

Example: $n = 4$, 3412 1234 1432 2431 2134 3214 4231 4132
8 total $D(4) = 8$.

Divisor-permutations

Recently, [Pomerance](#) asked how many permutations σ have the property that either $\sigma(i)|i$ or $i|\sigma(i)$ for every index i .

Call these **divisor-permutations** and denote their count by $D(n)$.

Example: $n = 4$, 3412 1234 1432 2431 2134 3214 4231 4132
8 total $D(4) = 8$.

In a subsequent paper he proves

$$1.93^n \leq D(n) \leq 13.6^n$$

for sufficiently large n .

The connection to the divisor graph

$$\left\{ \text{Permutations } \sigma \text{ of } n \text{ where } \right. \\ \left. \forall i \text{ either } \sigma(i) \mid i \text{ or } i \mid \sigma(i) \right\} \iff \left\{ \text{Directed, vertex-disjoint cycle} \right. \\ \left. \text{covers of } \mathcal{D}_{[1,n]} \text{ with self-loops} \right\}$$

The connection to the divisor graph

$$\left\{ \text{Permutations } \sigma \text{ of } n \text{ where } \right. \\ \left. \forall i \text{ either } \sigma(i) \mid i \text{ or } i \mid \sigma(i) \right\} \iff \left\{ \text{Directed, vertex-disjoint cycle} \right. \\ \left. \text{covers of } \mathcal{D}_{[1,n]} \text{ with self-loops} \right\}$$

Example: $\sigma = 916852743$

The connection to the divisor graph


$$\left\{ \text{Permutations } \sigma \text{ of } n \text{ where } \right. \\ \left. \forall i \text{ either } \sigma(i) \mid i \text{ or } i \mid \sigma(i) \right\} \iff \left\{ \text{Directed, vertex-disjoint cycle} \right. \\ \left. \text{covers of } \mathcal{D}_{[1,n]} \text{ with self-loops} \right\}$$

Example: $\sigma = 916852743 = (1\ 9\ 3\ 6\ 2)(4\ 8)(5)(7)$

The connection to the divisor graph

$$\left\{ \text{Permutations } \sigma \text{ of } n \text{ where } \right. \\ \left. \forall i \text{ either } \sigma(i) \mid i \text{ or } i \mid \sigma(i) \right\} \iff \left\{ \text{Directed, vertex-disjoint cycle} \right. \\ \left. \text{covers of } \mathcal{D}_{[1,n]} \text{ with self-loops} \right\}$$

Example: $\sigma = 916852743 = (1\ 9\ 3\ 6\ 2)(4\ 8)(5)(7)$



$$d(k, n)$$

$$d(k, n) = \frac{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k,n]}}{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k+1,n]}}, \quad D(n) = \prod_{k=1}^n d(k, n).$$

$$d(k, n)$$

$$d(k, n) = \frac{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k,n]}}{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k+1,n]}}, \quad D(n) = \prod_{k=1}^n d(k, n).$$

Seems like the right setup to apply the general theorem....

$$d(k, n)$$

$$d(k, n) = \frac{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k,n]}}{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k+1,n]}}, \quad D(n) = \prod_{k=1}^n d(k, n).$$

Seems like the right setup to apply the general theorem....

...Except it isn't clear that $d(k, n)$ is bounded.

$$d(k, n)$$

$$d(k, n) = \frac{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k, n]}}{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k+1, n]}}, \quad D(n) = \prod_{k=1}^n d(k, n).$$

Seems like the right setup to apply the general theorem....

...Except it isn't clear that $d(k, n)$ is bounded.

In fact it is unbounded, $d(k, n) > \pi\left(\frac{n}{k}\right) - \pi\left(\frac{n}{2k}\right)$.

$$d(k, n)$$

We can adapt the proof of the general theorem to handle a function $f(k, n)$ that doesn't grow too quickly with $\frac{n}{k}$.

$$d(k, n)$$

We can adapt the proof of the general theorem to handle a function $f(k, n)$ that doesn't grow too quickly with $\frac{n}{k}$.

Doesn't seem trivial to find any sub-exponential bound for $d(k, n)$.

$$d(k, n)$$

We can adapt the proof of the general theorem to handle a function $f(k, n)$ that doesn't grow too quickly with $\frac{n}{k}$.

Doesn't seem trivial to find any sub-exponential bound for $d(k, n)$.

For any graph G let $D(G)$ denote the count of directed vertex-disjoint cycle covers of G (allowing loops at each vertex.)

$d(k, n)$

We can adapt the proof of the general theorem to handle a function $f(k, n)$ that doesn't grow too quickly with $\frac{n}{k}$.

Doesn't seem trivial to find any sub-exponential bound for $d(k, n)$.

For any graph G let $D(G)$ denote the count of directed vertex-disjoint cycle covers of G (allowing loops at each vertex.) $D(n) = D(\mathcal{D}_{[1,n]})$.

$$\text{Let } R(G, v) = \frac{D(G)}{D(G \setminus \{v\})}.$$

$$d(k, n)$$

We can adapt the proof of the general theorem to handle a function $f(k, n)$ that doesn't grow too quickly with $\frac{n}{k}$.

Doesn't seem trivial to find any sub-exponential bound for $d(k, n)$.

For any graph G let $D(G)$ denote the count of directed vertex-disjoint cycle covers of G (allowing loops at each vertex.) $D(n) = D(\mathcal{D}_{[1,n]})$.

$$\text{Let } R(G, v) = \frac{D(G)}{D(G \setminus \{v\})}.$$

Theorem

For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) \leq 1 + \frac{d^2 - d}{2}.$$

Proof outline

Theorem

For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) = \frac{D(G)}{D(G \setminus \{v\})} \leq 1 + \frac{d^2 - d}{2}.$$

Proof outline

Theorem

For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) = \frac{D(G)}{D(G \setminus \{v\})} \leq 1 + \frac{d^2 - d}{2}.$$

Let $W = \{w_1, w_2, \dots, w_{d-1}\}$ be the neighbors of v in G .

Proof outline

Theorem

For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) = \frac{D(G)}{D(G \setminus \{v\})} \leq 1 + \frac{d^2 - d}{2}.$$

Let $W = \{w_1, w_2, \dots, w_{d-1}\}$ be the neighbors of v in G .

Write $D(G) = C_v + \sum_{i=1}^{d-1} C_{vw_i} + 2 \sum_{1 \leq i < j < d} C_{w_i v w_j}$.

- $C_v = D(G \setminus \{v\})$ counts cycle covers where v is fixed (a 1-cycle).
- C_{vw_i} counts cycle covers where v is in a 2-cycle with w_i .
- $C_{w_i v w_j}$ counts cycle covers containing the edges $w_i \rightarrow v \rightarrow w_j$.

Proof outline

Theorem


For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) = \frac{D(G)}{D(G \setminus \{v\})} \leq 1 + \frac{d^2 - d}{2}.$$

Let $W = \{w_1, w_2, \dots, w_{d-1}\}$ be the neighbors of v in G .

Write $D(G) = C_v + \sum_{i=1}^{d-1} C_{vw_i} + 2 \sum_{1 \leq i < j < d} C_{w_i v w_j}$.

- $C_v = D(G \setminus \{v\})$ counts cycle covers where v is fixed (a 1-cycle).
- C_{vw_i} counts cycle covers where v is in a 2-cycle with w_i .
- $C_{w_i v w_j}$ counts cycle covers containing the edges $w_i \rightarrow v \rightarrow w_j$.



Proof outline

Theorem

For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) = \frac{D(G)}{D(G \setminus \{v\})} \leq 1 + \frac{d^2 - d}{2}.$$

Let $W = \{w_1, w_2, \dots, w_{d-1}\}$ be the neighbors of v in G .

Write $D(G) = C_v + \sum_{i=1}^{d-1} C_{vw_i} + 2 \sum_{1 \leq i < j < d} C_{w_i v w_j}$.

- $C_v = D(G \setminus \{v\})$ counts cycle covers where v is fixed (a 1-cycle).
- C_{vw_i} counts cycle covers where v is in a 2-cycle with w_i .
- $C_{w_i v w_j}$ counts cycle covers containing the edges $w_i \rightarrow v \rightarrow w_j$.

Proof outline

Theorem

For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) = \frac{D(G)}{D(G \setminus \{v\})} \leq 1 + \frac{d^2 - d}{2}.$$

Let $W = \{w_1, w_2, \dots, w_{d-1}\}$ be the neighbors of v in G .

Write $D(G) = C_v + \sum_{i=1}^{d-1} C_{vw_i} + 2 \sum_{1 \leq i < j < d} C_{w_i v w_j}$.

- $C_v = D(G \setminus \{v\})$ counts cycle covers where v is fixed (a 1-cycle).
- C_{vw_i} counts cycle covers where v is in a 2-cycle with w_i .
- $C_{w_i v w_j}$ counts cycle covers containing the edges $w_i \rightarrow v \rightarrow w_j$.

$$R(G, v) = \frac{D(G)}{C_v}$$

Proof outline

Theorem

For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) = \frac{D(G)}{D(G \setminus \{v\})} \leq 1 + \frac{d^2 - d}{2}.$$

Let $W = \{w_1, w_2, \dots, w_{d-1}\}$ be the neighbors of v in G .

Write $D(G) = C_v + \sum_{i=1}^{d-1} C_{vw_i} + 2 \sum_{1 \leq i < j < d} C_{w_i v w_j}$.

- $C_v = D(G \setminus \{v\})$ counts cycle covers where v is fixed (a 1-cycle).
- C_{vw_i} counts cycle covers where v is in a 2-cycle with w_i .
- $C_{w_i v w_j}$ counts cycle covers containing the edges $w_i \rightarrow v \rightarrow w_j$.

$$R(G, v) = \frac{D(G)}{C_v} \leq 1 + (d-1) + 2 \sum_{1 \leq i < j < d} \frac{C_{w_i v w_j}}{C_v}$$

Proof outline

$$R(G, v) = \frac{D(G)}{C_v} \leq 1 + (d-1) + 2 \sum_{1 \leq i < j < d} \frac{C_{w_i v w_j}}{C_v}$$

Proof outline

$$R(G, v) = \frac{D(G)}{C_v} \leq 1 + (d-1) + 2 \sum_{1 \leq i < j < d} \frac{C_{w_i v w_j}}{C_v}$$

Claim: $(C_{w_i v w_j})^2 \leq \frac{1}{4} (C_v)^2$.

Proof outline

$$R(G, v) = \frac{D(G)}{C_v} \leq 1 + (d-1) + 2 \sum_{1 \leq i < j < d} \frac{C_{w_i v w_j}}{C_v}$$

Claim: $(C_{w_i v w_j})^2 \leq \frac{1}{4} (C_v)^2$.

Write $C_v = X + Y_i + Y_j + Z$

- X counts the cases where v , w_i , and w_j are all part of 1-cycles
- Y_i counts those where only v and w_i are fixed (not w_j).
- Z counts those where v is fixed but neither w_i nor w_j is fixed.

Then write

$$(C_v)^2 = (X + Y_i + Y_j + Z)^2 \geq (X + Z)^2 + (Y_i + Y_j)^2$$

Proof outline

$$R(G, v) = \frac{D(G)}{C_v} \leq 1 + (d-1) + 2 \sum_{1 \leq i < j < d} \frac{C_{w_i v w_j}}{C_v}$$

Claim: $(C_{w_i v w_j})^2 \leq \frac{1}{4} (C_v)^2$.

Write $C_v = X + Y_i + Y_j + Z$

- X counts the cases where v , w_i , and w_j are all part of 1-cycles
- Y_i counts those where only v and w_i are fixed (not w_j).
- Z counts those where v is fixed but neither w_i nor w_j is fixed.

Then write

$$(C_v)^2 = (X + Y_i + Y_j + Z)^2 \geq (X + Z)^2 + (Y_i + Y_j)^2 \geq 4(XZ + Y_i Y_j)$$

by the AM-GM inequality.

Proof outline

$$R(G, v) = \frac{D(G)}{C_v} \leq 1 + (d-1) + 2 \sum_{1 \leq i < j < d} \frac{C_{w_i v w_j}}{C_v}$$

Claim: $(C_{w_i v w_j})^2 \leq \frac{1}{4} (C_v)^2$.

Write $C_v = X + Y_i + Y_j + Z$

- X counts the cases where v , w_i , and w_j are all part of 1-cycles
- Y_i counts those where only v and w_i are fixed (not w_j).
- Z counts those where v is fixed but neither w_i nor w_j is fixed.

Then write

$$(C_v)^2 = (X + Y_i + Y_j + Z)^2 \geq (X + Z)^2 + (Y_i + Y_j)^2 \geq 4(XZ + Y_i Y_j)$$

by the AM-GM inequality. It remains to show $XZ + Y_i Y_j \geq (C_{w_i v w_j})^2$.

$$XZ + Y_i Y_j \geq (C_{w_i v w_j})^2$$

Find an injection from objects counted by $(C_{w_i v w_j})^2$ to objects counted by $XZ + Y_i Y_j$.

$$XZ + Y_i Y_j \geq (C_{w_i v w_j})^2$$


Find an injection from objects counted by $(C_{w_i v w_j})^2$ to objects counted by $XZ + Y_i Y_j$.

Each of XZ , $Y_i Y_j$ and $(C_{w_i v w_j})^2$ count pairs of directed vertex-disjoint cycle covers. In each pair, color the first one blue, and the second red.

$$XZ + Y_i Y_j \geq (C_{w_i v w_j})^2$$

Find an injection from objects counted by $(C_{w_i v w_j})^2$ to objects counted by $XZ + Y_i Y_j$.


Each of XZ , $Y_i Y_j$ and $(C_{w_i v w_j})^2$ count pairs of directed vertex-disjoint cycle covers. In each pair, color the first one blue, and the second red.



$$XZ + Y_i Y_j \geq (C_{w_i v w_j})^2$$

Find an injection from objects counted by $(C_{w_i v w_j})^2$ to objects counted by $XZ + Y_i Y_j$.


Each of XZ , $Y_i Y_j$ and $(C_{w_i v w_j})^2$ count pairs of directed vertex-disjoint cycle covers. In each pair, color the first one blue, and the second red.



$$XZ + Y_i Y_j \geq (C_{w_i v w_j})^2$$

Find an injection from objects counted by $(C_{w_i v w_j})^2$ to objects counted by $XZ + Y_i Y_j$.

Each of XZ , $Y_i Y_j$ and $(C_{w_i v w_j})^2$ count pairs of directed vertex-disjoint cycle covers. In each pair, color the first one blue, and the second red.




$$XZ + Y_i Y_j \geq (C_{w_i v w_j})^2$$


Find an injection from objects counted by $(C_{w_i v w_j})^2$ to objects counted by $XZ + Y_i Y_j$.

Each of XZ , $Y_i Y_j$ and $(C_{w_i v w_j})^2$ count pairs of directed vertex-disjoint cycle covers. In each pair, color the first one blue, and the second red.


Draw both on the same graph. Get a colored, directed multigraph, every vertex has one inward pointing edge of each color.



Colorings!




Colorings!




Take a colored multigraph obtained from $(C_{w_i v w_j})^2$. (It has both blue and red edges $w_i \rightarrow v \rightarrow w_j$.)

Colorings!




Take a colored multigraph obtained from $(C_{w_i v w_j})^2$. (It has both blue and red edges $w_i \rightarrow v \rightarrow w_j$.) Remove all four of these edges.

Colorings!



Take a colored multigraph obtained from $(C_{w_i v w_j})^2$. (It has both blue and red edges $w_i \rightarrow v \rightarrow w_j$.) Remove all four of these edges.


Colorings!



Take a colored multigraph obtained from $(C_{w_i v w_j})^2$. (It has both blue and red edges $w_i \rightarrow v \rightarrow w_j$.) Remove all four of these edges.

Add two loops to v one of each color.


Colorings!



Take a colored multigraph obtained from $(C_{w_i v w_j})^2$. (It has both blue and red edges $w_i \rightarrow v \rightarrow w_j$.) Remove all four of these edges.

Add two loops to v one of each color.


Colorings!




Take a colored multigraph obtained from $(C_{w_i v w_j})^2$. (It has both blue and red edges $w_i \rightarrow v \rightarrow w_j$.) Remove all four of these edges.

Add two loops to v one of each color.


Now every vertex (except w_i, w_j) has in- and out-edges of each color. w_i has in-edges of each color, and w_j has out-edges of each color.



The colored multigraph consists of alternating-color cycles, plus two alternating-color paths $w_j \rightarrow w_i$.




The colored multigraph consists of alternating-color cycles, plus two alternating-color paths $w_j \rightarrow w_i$. Call the path ending in a blue edge P .




The colored multigraph consists of alternating-color cycles, plus two alternating-color paths $w_j \rightarrow w_i$. Call the path ending in a blue edge P .

Now add a (initially uncolored) loop to each of the vertices w_i and w_j .



The colored multigraph consists of alternating-color cycles, plus two alternating-color paths $w_j \rightarrow w_i$. Call the path ending in a blue edge P .


Now add a (initially uncolored) loop to each of the vertices w_i and w_j .



The colored multigraph consists of alternating-color cycles, plus two alternating-color paths $w_j \rightarrow w_i$. Call the path ending in a blue edge P .

Now add a (initially uncolored) loop to each of the vertices w_i and w_j .

Color the new loop on w_i blue.




The colored multigraph consists of alternating-color cycles, plus two alternating-color paths $w_j \rightarrow w_i$. Call the path ending in a blue edge P .

Now add a (initially uncolored) loop to each of the vertices w_i and w_j .

Color the new loop on w_i blue. Recolor and reverse every edge along P . Every vertex along P (except w_j) has a consistent coloring.

Both edges adjacent to w_j have the same color and opposite orientations. Color the new loop at w_j the opposite color.




The colored multigraph consists of alternating-color cycles, plus two alternating-color paths $w_j \rightarrow w_i$. Call the path ending in a blue edge P .

Now add a (initially uncolored) loop to each of the vertices w_i and w_j .

Color the new loop on w_i blue. Recolor and reverse every edge along P . Every vertex along P (except w_j) has a consistent coloring.

Both edges adjacent to w_j have the same color and opposite orientations. Color the new loop at w_j the opposite color.




The colored multigraph consists of alternating-color cycles, plus two alternating-color paths $w_j \rightarrow w_i$. Call the path ending in a blue edge P .

Now add a (initially uncolored) loop to each of the vertices w_i and w_j .

Color the new loop on w_i blue. Recolor and reverse every edge along P . Every vertex along P (except w_j) has a consistent coloring.


Both edges adjacent to w_j have the same color and opposite orientations. Color the new loop at w_j the opposite color.

Wrapping up



The final result is a coloring in which v is fixed by both colors, every vertex has exactly one in-edge and out-edge of each color.


Wrapping up



The final result is a coloring in which v is fixed by both colors, every vertex has exactly one in-edge and out-edge of each color.

If both the self-loops on w_i and w_j have the same color (blue) the result is counted by XZ , otherwise it is counted by $Y_i Y_j$.

Wrapping up




The final result is a coloring in which v is fixed by both colors, every vertex has exactly one in-edge and out-edge of each color.

If both the self-loops on w_i and w_j have the same color (blue) the result is counted by XZ , otherwise it is counted by $Y_i Y_j$.

We have shown an injection from objects counted by $(C_{w_i v w_j})^2$ to objects counted by $XZ + Y_i Y_j$.

Wrapping up




The final result is a coloring in which v is fixed by both colors, every vertex has exactly one in-edge and out-edge of each color.

If both the self-loops on w_i and w_j have the same color (blue) the result is counted by XZ , otherwise it is counted by $Y_i Y_j$.

We have shown an injection from objects counted by $(C_{w_i v w_j})^2$ to objects counted by $XZ + Y_i Y_j$. So $(C_{w_i v w_j})^2 \leq XZ + Y_i Y_j$

Wrapping up



The final result is a coloring in which v is fixed by both colors, every vertex has exactly one in-edge and out-edge of each color.

If both the self-loops on w_i and w_j have the same color (blue) the result is counted by XZ , otherwise it is counted by $Y_i Y_j$.

We have shown an injection from objects counted by $(C_{w_i v w_j})^2$ to objects counted by $XZ + Y_i Y_j$. So $(C_{w_i v w_j})^2 \leq XZ + Y_i Y_j \leq \frac{1}{4}(C_v)^2$

Wrapping up

Theorem

For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) \leq 1 + \frac{d^2 - d}{2}.$$

Wrapping up

Theorem

For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) \leq 1 + \frac{d^2 - d}{2}.$$

Using this, we find that

$$d(k, n) = \frac{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k, n]}}{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k+1, n]}} \leq 1 + \frac{\lfloor \frac{n}{k} \rfloor^2 - \lfloor \frac{n}{k} \rfloor}{2}.$$

Wrapping up

Theorem

For any graph G and vertex v of degree d , (counting a self loop)

$$R(G, v) \leq 1 + \frac{d^2 - d}{2}.$$

Using this, we find that

$$d(k, n) = \frac{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k, n]}}{\# \text{ vertex-disjoint cycle covers of } \mathcal{D}_{[k+1, n]}} \leq 1 + \frac{\lfloor \frac{n}{k} \rfloor^2 - \lfloor \frac{n}{k} \rfloor}{2}.$$

$$D(n) = \prod_{k=1}^n d(k, n).$$

Counting divisor-permutations

$$D(n) = \#\{\sigma \in S_n : \forall i \text{ either } \sigma(i)|i \text{ or } i|\sigma(i)\}.$$

Counting divisor-permutations

$$D(n) = \#\{\sigma \in S_n : \forall i \text{ either } \sigma(i)|i \text{ or } i|\sigma(i)\}.$$



Theorem ([Pomerance](#) (2022))

$$1.93^n \leq D(n) \leq 13.6^n \text{ (} n \text{ sufficiently large.)}$$

Counting divisor-permutations

$$D(n) = \#\{\sigma \in S_n : \forall i \text{ either } \sigma(i)|i \text{ or } i|\sigma(i)\}.$$



Theorem (Pomerance (2022))

$$1.93^n \leq D(n) \leq 13.6^n \text{ (} n \text{ sufficiently large.)}$$

Theorem

$$\text{As } n \rightarrow \infty, \quad D(n) = c_d^{n+o\left(n \exp\left(-\sqrt{\frac{1}{2} \log n \log \log n}\right)\right)}.$$

Counting divisor-permutations

$$D(n) = \#\{\sigma \in S_n : \forall i \text{ either } \sigma(i)|i \text{ or } i|\sigma(i)\}.$$



Theorem (Pomerance (2022))

$$1.93^n \leq D(n) \leq 13.6^n \text{ (} n \text{ sufficiently large.)}$$

Theorem

$$\text{As } n \rightarrow \infty, D(n) = c_d^{n+o\left(n \exp\left(-\sqrt{\frac{1}{2} \log n \log \log n}\right)\right)}.$$

Using $d(k, n) \leq 1 + \frac{\lfloor \frac{n}{k} \rfloor^2 - \lfloor \frac{n}{k} \rfloor}{2}$ we improve $c_d < 13.6$ to $c_d < 3.32$.

Counting divisor-permutations

$$D(n) = \#\{\sigma \in S_n : \forall i \text{ either } \sigma(i)|i \text{ or } i|\sigma(i)\}.$$



Theorem (Pomerance (2022))

$$1.93^n \leq D(n) \leq 13.6^n \text{ (} n \text{ sufficiently large.)}$$

Theorem

$$\text{As } n \rightarrow \infty, D(n) = c_d^{n+o\left(n \exp\left(-\sqrt{\frac{1}{2} \log n \log \log n}\right)\right)}.$$

Using $d(k, n) \leq 1 + \frac{\lfloor \frac{n}{k} \rfloor^2 - \lfloor \frac{n}{k} \rfloor}{2}$ we improve $c_d < 13.6$ to $c_d < 3.32$.

Using numerical computation we improve this to $2.069 < c_d < 2.694$.

Open questions

What is the best constant C , $R(G, v) \leq (C + o(1))d^2$ as $d \rightarrow \infty$?

Open questions

What is the best constant C , $R(G, v) \leq (C + o(1))d^2$ as $d \rightarrow \infty$?

We have $C \leq \frac{1}{2}$.

Open questions

What is the best constant C , $R(G, \nu) \leq (C + o(1))d^2$ as $d \rightarrow \infty$?

We have $C \leq \frac{1}{2}$.

On the other hand, looking at complete bipartite graphs $K_{\frac{n}{2}, n}$ we find

$$R(K_{\frac{d}{2}, d}, \nu) \geq \left(\frac{1}{4} + o(1) \right) d^2.$$

Open questions

What is the length $l(n)$ of the longest path in $\mathcal{D}_{[1,n]}$?

Open questions

What is the length $l(n)$ of the longest path in $\mathcal{D}_{[1,n]}$?

Saia: $3 \frac{x}{\log x} \leq l(x) \leq 7 \frac{x}{\log x}$, for sufficiently large x .

Open questions

What is the length $l(n)$ of the longest path in $\mathcal{D}_{[1,n]}$?

Saia: $3 \frac{x}{\log x} \leq l(x) \leq 7 \frac{x}{\log x}$, for sufficiently large x .

What is the length of the longest cycle?

Open questions

What is the length $l(n)$ of the longest path in $\mathcal{D}_{[1,n]}$?

Saia: $3 \frac{x}{\log x} \leq l(x) \leq 7 \frac{x}{\log x}$, for sufficiently large x .

What is the length of the longest cycle? Its at most $l(x)$...

Open questions

What is the length $l(n)$ of the longest path in $\mathcal{D}_{[1,n]}$?

Saia: $3 \frac{x}{\log x} \leq l(x) \leq 7 \frac{x}{\log x}$, for sufficiently large x .

What is the length of the longest cycle? Its at most $l(x) \dots$

Let L be a path of length $l(n)$ in $\mathcal{D}_{[1,n]}$. What is the length of the longest path in $\mathcal{D}_{[1,n]} \setminus L$?

Open questions

What is the length $l(n)$ of the longest path in $\mathcal{D}_{[1,n]}$?

Saia: $3 \frac{x}{\log x} \leq l(x) \leq 7 \frac{x}{\log x}$, for sufficiently large x .

What is the length of the longest cycle? Its at most $l(x)$...

Let L be a path of length $l(n)$ in $\mathcal{D}_{[1,n]}$. What is the length of the longest path in $\mathcal{D}_{[1,n]} \setminus L$? It is $\gg \frac{\sqrt{n}}{\log n}$.

THANK YOU!