

Counting primitive subsets of $\{1, 2, \dots, n\}$

Nathan McNew
Towson University

CANT 2018
CUNY Graduate Center
May 25th, 2018

Primitive Sets

A **Primitive Set** is a set of integers in which no two integers divide one another.

Primitive Sets

A **Primitive Set** is a set of integers in which no two integers divide one another.

Examples:

- Primes

Primitive Sets

A **Primitive Set** is a set of integers in which no two integers divide one another.

Examples:

- Primes
- Semiprimes, $\{pq \mid p, q \text{ prime}\}$

Primitive Sets

A **Primitive Set** is a set of integers in which no two integers divide one another.

Examples:

- Primes
- Semiprimes, $\{pq \mid p, q \text{ prime}\}$
- Integers with exactly k prime factors counted with multiplicity.

Primitive Sets

A **Primitive Set** is a set of integers in which no two integers divide one another.

Examples:

- Primes
- Semiprimes, $\{pq \mid p, q \text{ prime}\}$
- Integers with exactly k prime factors counted with multiplicity.
- $\{\lfloor \frac{n}{2} \rfloor + 1, \lfloor \frac{n}{2} \rfloor + 2, \dots, n\}$

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets.

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	13

Finite primitive sets

Note: $\{\}$ and $\{1\}$ are both primitive sets. What do other primitive subsets of the first few integers look like?

n	primitive subsets of $\{1, 2 \dots n\}$	count
1	$\{\}, \{1\}$	2
2	$\{\}, \{1\}, \{2\}$	3
3	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}$	5
4	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}$	7
5	$\{\}, \{1\}, \{2\}, \{3\}, \{2, 3\}, \{4\}, \{3, 4\}, \{5\}, \{2, 5\}, \{3, 5\}, \{2, 3, 5\}, \{4, 5\}, \{3, 4, 5\}$	13

How many primitive subsets of $\{1, 2 \dots n\}$ are there?

Counting primitive sets

Let $Q(n)$ count the primitive sets with largest element at most n .


Counting primitive sets

Let $Q(n)$ count the primitive sets with largest element at most n .

A051026	Number of primitive subsequences of $\{1, 2, \dots, n\}$.	5
	1, 2, 3, 5, 7, 13, 17, 33, 45, 73, 103, 205, 253, 505, 733, 1133, 1529, 3057, 3897, 7793, 10241, 16513, 24593, 49185, 59265, 109297, 163369, 262489, 355729, 711457, 879937, 1759873, 2360641, 3908545, 5858113, 10534337, 12701537, 25403073, 38090337, 63299265, 81044097, 162088193, 205482593, 410965185, 570487233, 855676353 (list ; graph ; refs ; listen ; history ; text ; internal format)	
OFFSET	0, 2	
COMMENTS	$a(n)$ counts all subsequences of $\{1, \dots, n\}$ in which no term divides any other. If n is a prime $a(n) = 2 \cdot a(n-1) - 1$ because for each subsequence s counted by $a(n-1)$ two different subsequences are counted by $a(n)$: s and s, n . There is only one exception: $1, n$ is not a primitive subsequence because 1 divides n . For all $n > 1$: $a(n) < 2 \cdot a(n-1)$. - Alois P. Heinz , Mar 07 2011	


Bounds

Cameron and Erdős considered the function $Q(n)$ in 1990.



Bounds

Cameron and Erdős considered the function $Q(n)$ in 1990.




(Almost) trivially:

$$(\sqrt{2})^n < Q(n) < 2^n$$

Bounds

Cameron and Erdős considered the function $Q(n)$ in 1990.



(Almost) trivially:

$$(\sqrt{2})^n < Q(n) < 2^n$$

Every subset of integers contained in $\{\lfloor \frac{n}{2} \rfloor + 1, \lfloor \frac{n}{2} \rfloor + 2, \dots, n\}$ is primitive, and there are $2^{\lceil \frac{n}{2} \rceil} > \sqrt{2}^n$ such subsets.

Bounds

Cameron and Erdős improve these bounds to:

$$1.55967^n < Q(n) < 1.59^n$$

Bounds

Cameron and Erdős improve these bounds to:

$$1.55967^n < Q(n) < 1.59^n$$

Bounds

Cameron and Erdős improve these bounds to:

$$1.55967^n < Q(n) < 1.59^n$$

Conjecture: $\lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists.

New Developments

Angelo (2017) proves Cameron and Erdős's conjecture.

New Developments

Angelo (2017) proves Cameron and Erdős's conjecture.

Theorem

The limit $\lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists.

New Developments

Angelo (2017) proves Cameron and Erdős's conjecture.

Theorem

The limit $\lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists. Equivalently, $Q(n) = c^{n+o(1)}$.

New Developments

Angelo (2017) proves Cameron and Erdős's conjecture.

Theorem

The limit $\lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists. Equivalently, $Q(n) = c^{n+o(1)}$.

Proof is by considering sets in which the ratio between any two terms is not a k -smooth integer.

New Developments

Angelo (2017) proves Cameron and Erdős's conjecture.

Theorem

The limit $\lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists. Equivalently, $Q(n) = c^{n+o(1)}$.

Proof is by considering sets in which the ratio between any two terms is not a k -smooth integer. Then let $k \rightarrow \infty$.

New Developments

Angelo (2017) proves Cameron and Erdős's conjecture.

Theorem

The limit $\lim_{n \rightarrow \infty} Q(n)^{1/n}$ exists. Equivalently, $Q(n) = c^{n+o(1)}$.

Proof is by considering sets in which the ratio between any two terms is not a k -smooth integer. Then let $k \rightarrow \infty$.

Say n is k -smooth if the largest prime divisor $P^+(n) \leq k$.

Proof gives no insight to the value of the limit nor improved bounds.

Constructing Primitive Sets

Constructing Primitive Sets

Recall, any subset of the integers in $(\frac{n}{2}, n]$ is primitive. Since each element of this set can either be included or not, $Q(n) \geq 2^{n/2}$.

Constructing Primitive Sets

Recall, any subset of the integers in $(\frac{n}{2}, n]$ is primitive. Since each element of this set can either be included or not, $Q(n) \geq 2^{n/2}$.

Make this bigger by considering integers in $(\frac{n}{3}, \frac{n}{2}]$.

Constructing Primitive Sets

Recall, any subset of the integers in $(\frac{n}{2}, n]$ is primitive. Since each element of this set can either be included or not, $Q(n) \geq 2^{n/2}$.

Make this bigger by considering integers in $(\frac{n}{3}, \frac{n}{2}]$. For each such k in this range we can either include $k, 2k$ or neither.

Constructing Primitive Sets

Recall, any subset of the integers in $(\frac{n}{2}, n]$ is primitive. Since each element of this set can either be included or not, $Q(n) \geq 2^{n/2}$.

Make this bigger by considering integers in $(\frac{n}{3}, \frac{n}{2}]$. For each such k in this range we can either include $k, 2k$ or neither. 3 possibilities.

Constructing Primitive Sets

Recall, any subset of the integers in $(\frac{n}{2}, n]$ is primitive. Since each element of this set can either be included or not, $Q(n) \geq 2^{n/2}$.

Make this bigger by considering integers in $(\frac{n}{3}, \frac{n}{2}]$. For each such k in this range we can either include $k, 2k$ or neither. 3 possibilities.


This is the same as counting primitive subsets of $\{1, 2\}$

Constructing Primitive Sets

Recall, any subset of the integers in $(\frac{n}{2}, n]$ is primitive. Since each element of this set can either be included or not, $Q(n) \geq 2^{n/2}$.

Make this bigger by considering integers in $(\frac{n}{3}, \frac{n}{2}]$. For each such k in this range we can either include $k, 2k$ or neither. 3 possibilities.

This is the same as counting primitive subsets of $\{1, 2\}$ or independent subsets of the graph:



Constructing Primitive Sets

For each such k we can then multiply by 3, but must divide by 2, the number of possible sets already obtained before k was considered.

Constructing Primitive Sets

For each such k we can then multiply by 3, but must divide by 2, the number of possible sets already obtained before k was considered.

Thus $Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} = 2^{n/3} 3^{n/6} \approx 1.5131^n$.

Constructing Primitive Sets

For each such k we can then multiply by 3, but must divide by 2, the number of possible sets already obtained before k was considered.

Thus $Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} = 2^{n/3} 3^{n/6} \approx 1.5131^n$.

Likewise consider $k \in \left(\frac{n}{4}, \frac{n}{3}\right]$.

Constructing Primitive Sets

For each such k we can then multiply by 3, but must divide by 2, the number of possible sets already obtained before k was considered.

Thus $Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} = 2^{n/3} 3^{n/6} \approx 1.5131^n$.

Likewise consider $k \in \left(\frac{n}{4}, \frac{n}{3}\right]$.

Now we have to worry about $k, 2k, 3k$.

Constructing Primitive Sets

For each such k we can then multiply by 3, but must divide by 2, the number of possible sets already obtained before k was considered.

Thus $Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} = 2^{n/3} 3^{n/6} \approx 1.5131^n$.

Likewise consider $k \in \left(\frac{n}{4}, \frac{n}{3}\right]$.

Now we have to worry about $k, 2k, 3k$. Actually 2 possibilities.

Constructing Primitive Sets


For each such k we can then multiply by 3, but must divide by 2, the number of possible sets already obtained before k was considered.

Thus $Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} = 2^{n/3} 3^{n/6} \approx 1.5131^n$.

Likewise consider $k \in \left(\frac{n}{4}, \frac{n}{3}\right]$.

Now we have to worry about $k, 2k, 3k$. Actually 2 possibilities.

If k odd this is equivalent to counting primitive subsets of $\{1, 2, 3\}$, which is 5, replacing the 4 possibilities before.



Constructing Primitive Sets

Constructing Primitive Sets

If $k = 2\ell$ is even however then $\frac{3k}{2} = 3\ell$ is also an integer that is greater than k . $3k = 6\ell$ can only be included if 3ℓ is not.

Constructing Primitive Sets


If $k = 2\ell$ is even however then $\frac{3k}{2} = 3\ell$ is also an integer that is greater than k . $3k = 6\ell$ can only be included if 3ℓ is not.

Must consider $2\ell, 3\ell, 4\ell, 6\ell$

Constructing Primitive Sets

If $k = 2\ell$ is even however then $\frac{3k}{2} = 3\ell$ is also an integer that is greater than k . $3k = 6\ell$ can only be included if 3ℓ is not.

Must consider $2\ell, 3\ell, 4\ell, 6\ell$ like counting primitive subsets of $\{2, 3, 4, 6\}$.




8 possibilities $\{\}, \{2\}, \{3\}, \{4\}, \{6\}, \{2, 3\}, \{3, 4\}, \{4, 6\}$ replace 6 before.

Constructing Primitive Sets

If $k = 2\ell$ is even however then $\frac{3k}{2} = 3\ell$ is also an integer that is greater than k . $3k = 6\ell$ can only be included if 3ℓ is not.

Must consider $2\ell, 3\ell, 4\ell, 6\ell$ like counting primitive subsets of $\{2, 3, 4, 6\}$.




8 possibilities $\{\}, \{2\}, \{3\}, \{4\}, \{6\}, \{2, 3\}, \{3, 4\}, \{4, 6\}$ replace 6 before.

$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} \left(\frac{5}{4}\right)^{n/24} \left(\frac{8}{6}\right)^{n/24}$$

Constructing Primitive Sets

If $k = 2\ell$ is even however then $\frac{3k}{2} = 3\ell$ is also an integer that is greater than k . $3k = 6\ell$ can only be included if 3ℓ is not.

Must consider $2\ell, 3\ell, 4\ell, 6\ell$ like counting primitive subsets of $\{2, 3, 4, 6\}$.




8 possibilities $\{\}, \{2\}, \{3\}, \{4\}, \{6\}, \{2, 3\}, \{3, 4\}, \{4, 6\}$ replace 6 before.

$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} \left(\frac{5}{4}\right)^{n/24} \left(\frac{8}{6}\right)^{n/24} = 2^{5n/24} 3^{n/8} 5^{n/24}$$

Constructing Primitive Sets

If $k = 2\ell$ is even however then $\frac{3k}{2} = 3\ell$ is also an integer that is greater than k . $3k = 6\ell$ can only be included if 3ℓ is not.

Must consider $2\ell, 3\ell, 4\ell, 6\ell$ like counting primitive subsets of $\{2, 3, 4, 6\}$.



8 possibilities $\{\}, \{2\}, \{3\}, \{4\}, \{6\}, \{2, 3\}, \{3, 4\}, \{4, 6\}$ replace 6 before.

$$Q(n) \geq 2^{n/2} \left(\frac{3}{2}\right)^{n/6} \left(\frac{5}{4}\right)^{n/24} \left(\frac{8}{6}\right)^{n/24} = 2^{5n/24} 3^{n/8} 5^{n/24} \approx 1.5456^n$$


One more wrinkle

One more wrinkle


Consider $k \in \left(\frac{n}{5}, \frac{n}{4}\right]$, the next interval.

One more wrinkle

Consider $k \in (\frac{n}{5}, \frac{n}{4}]$, the next interval. If $k = 2\ell$ is even we must consider $2\ell, 4\ell, 6\ell, 8\ell$, as well as 3ℓ .




However, if $k \leq \frac{2n}{9}$ then $9\ell \leq n$ must also be considered.




One more wrinkle

Consider $k \in (\frac{n}{5}, \frac{n}{4}]$, the next interval. If $k = 2\ell$ is even we must consider $2\ell, 4\ell, 6\ell, 8\ell$, as well as 3ℓ .



However, if $k \leq \frac{2n}{9}$ then $9\ell \leq n$ must also be considered.



So we consider $2\ell, 3\ell, 4\ell, 6\ell, 8\ell$ and 9ℓ .

Key observation

Let $r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}$ be the contribution of k to our potential primitive set.

Key observation

Let $r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}$ be the contribution of k to our potential primitive set. This ratio depends only on the connected component of k in the divisibility graph of $[k, n]$.

Key observation

Let $r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}$ be the contribution of k to our potential primitive set. This ratio depends only on the connected component of k in the divisibility graph of $[k, n]$.

Let $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$.

Key observation

Let $r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}$ be the contribution of k to our potential primitive set. This ratio depends only on the connected component of k in the divisibility graph of $[k, n]$.

Let $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$. (All included in connected component of k .)

Key observation

Let $r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}$ be the contribution of k to our potential primitive set. This ratio depends only on the connected component of k in the divisibility graph of $[k, n]$.

Let $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$. (All included in connected component of k .)

None of the ratios in this component can involve a prime greater than or equal to i (except possibly the edge from k to ik if i is prime.)

Key observation

Let $r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}$ be the contribution of k to our potential primitive set. This ratio depends only on the connected component of k in the divisibility graph of $[k, n]$.

Let $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$. (All included in connected component of k .)

None of the ratios in this component can involve a prime greater than or equal to i (except possibly the edge from k to ik if i is prime.)

Let $d = \max\{e | i : P^+(e) < i\}$, the largest $(i-1)$ -smooth divisor of k .

Key observation

Let $r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}$ be the contribution of k to our potential primitive set. This ratio depends only on the connected component of k in the divisibility graph of $[k, n]$.

Let $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$. (All included in connected component of k .)

None of the ratios in this component can involve a prime greater than or equal to i (except possibly the edge from k to ik if i is prime.)

Let $d = \max\{e | i : P^+(e) < i\}$, the largest $(i-1)$ -smooth divisor of k .

Let $\ell = \frac{k}{d}$ the “rough” part of k .

Key observation

Let $r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}$ be the contribution of k to our potential primitive set. This ratio depends only on the connected component of k in the divisibility graph of $[k, n]$.

Let $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$. (All included in connected component of k .)

None of the ratios in this component can involve a prime greater than or equal to i (except possibly the edge from k to ik if i is prime.)

Let $d = \max\{e | i : P^+(e) < i\}$, the largest $(i-1)$ -smooth divisor of k .

Let $\ell = \frac{k}{d}$ the “rough” part of k .

Every integer in the connected component of k is divisible by ℓ .

Computing $Q(n)$

Computing $Q(n)$

$$\text{Since } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}, \quad Q(n) = \prod_{k=1}^n r(k, n).$$

Computing $Q(n)$

$$\text{Since } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}, \quad Q(n) = \prod_{k=1}^n r(k, n).$$

Goal: Group together those values of $r(k, n)$ that are equal.

Computing $Q(n)$

$$\text{Since } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}, \quad Q(n) = \prod_{k=1}^n r(k, n).$$

Goal: Group together those values of $r(k, n)$ that are equal.

Recall $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$.

Computing $Q(n)$

$$\text{Since } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}, \quad Q(n) = \prod_{k=1}^n r(k, n).$$

Goal: Group together those values of $r(k, n)$ that are equal.

Recall $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$. d is the largest $(i-1)$ -smooth divisor of k ,

Computing $Q(n)$

$$\text{Since } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}, \quad Q(n) = \prod_{k=1}^n r(k, n).$$

Goal: Group together those values of $r(k, n)$ that are equal.

Recall $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$. d is the largest $(i-1)$ -smooth divisor of k , $\ell = \frac{k}{d}$ is the “rough” part of k .

Computing $Q(n)$

$$\text{Since } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}, \quad Q(n) = \prod_{k=1}^n r(k, n).$$

Goal: Group together those values of $r(k, n)$ that are equal.

Recall $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$. d is the largest $(i-1)$ -smooth divisor of k , $\ell = \frac{k}{d}$ is the “rough” part of k .

Let $t = \lfloor \frac{n}{\ell} \rfloor$, so $t\ell$ is the largest integer up to n divisible by ℓ .

Computing $Q(n)$

$$\text{Since } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}, \quad Q(n) = \prod_{k=1}^n r(k, n).$$

Goal: Group together those values of $r(k, n)$ that are equal.

Recall $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$. d is the largest $(i-1)$ -smooth divisor of k , $\ell = \frac{k}{d}$ is the “rough” part of k .

Let $t = \lfloor \frac{n}{\ell} \rfloor$, so $t\ell$ is the largest integer up to n divisible by ℓ .

Note: $ik \leq t\ell < (i+1)k$ and $k = d\ell$, so $id \leq t < (i+1)d$.

Computing $Q(n)$

$$\text{Since } r(k, n) = \frac{\# \text{Primitive subsets of } [k, n]}{\# \text{Primitive subsets of } [k+1, n]}, \quad Q(n) = \prod_{k=1}^n r(k, n).$$

Goal: Group together those values of $r(k, n)$ that are equal.

Recall $i = \lfloor \frac{n}{k} \rfloor$, so the multiples of k up to n are $k, 2k, \dots, ik$. d is the largest $(i-1)$ -smooth divisor of k , $\ell = \frac{k}{d}$ is the “rough” part of k .

Let $t = \lfloor \frac{n}{\ell} \rfloor$, so $t\ell$ is the largest integer up to n divisible by ℓ .

Note: $ik \leq t\ell < (i+1)k$ and $k = d\ell$, so $id \leq t < (i+1)d$.

Then $r(i, n) = r(d\ell, t\ell) = r(d, t)$.

Computing $Q(n)$

$$\text{Let } r(i, n) = \frac{\# \text{Primitive subsets of } [i, n]}{\# \text{Primitive subsets of } [i+1, n]}.$$

Computing $Q(n)$

$$\text{Let } r(i, n) = \frac{\#\text{Primitive subsets of } [i, n]}{\#\text{Primitive subsets of } [i+1, n]}.$$

$$\text{So that } Q(n) = \prod_{i=1}^n r(i, n).$$

Computing $Q(n)$

$$\text{Let } r(i, n) = \frac{\# \text{Primitive subsets of } [i, n]}{\# \text{Primitive subsets of } [i+1, n]}.$$

$$\text{So that } Q(n) = \prod_{i=1}^n r(i, n).$$

Goal: group together those values of $r(i, n)$ that are equal.

Let $k = \lfloor \frac{n}{i} \rfloor$, so the multiples of i up to n are $i, 2i, \dots, ki$. Then

Computing $Q(n)$

$$\text{Let } c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ P^+(d) < i}} \prod_{t \in [id, (i+1)d)} r(d, t)^{\frac{1}{t(t+1)} \prod_{p < i} \frac{p-1}{p}}.$$

Theorem (M.)

For any $\epsilon > 0$, the number of primitive subsets of $[1, n]$ is

$$Q(n) = c^{n + O\left(\frac{n}{\exp((\log n)^{1/2 - \epsilon})}\right)}$$

Computing $Q(n)$

$$\text{Let } c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ P^+(d) < i}} \prod_{t \in [id, (i+1)d)} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$$

Theorem (M.)

For any $\epsilon > 0$, the number of primitive subsets of $[1, n]$ is

$$Q(n) = c^{n + O\left(\frac{n}{\exp((\log n)^{1/2 - \epsilon})}\right)}$$

or

$$\log Q(n) = n \log(c) \left(1 + O\left(e^{-(\log n)^{1/2 - \epsilon}}\right)\right).$$

Proof Outline

Proof Outline

$$\text{Recall } c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ P^+(d) < i}} \prod_{t \in [id, (i+1)d)} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$$

Proof Outline

$$\text{Recall } c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ P^+(d) < i}} \prod_{t \in [id, (i+1)d)} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$$

We have

$$Q(n) = \prod_{k=1}^n r(k, n)$$

Proof Outline

$$\text{Recall } c = \prod_{i=1}^{\infty} \prod_{\substack{d \\ P^+(d) < i}} \prod_{t \in [id, (i+1)d)} r(d, t)^{\frac{1}{t(t+1)}} \prod_{p < i} \frac{p-1}{p}.$$

We have

$$\begin{aligned} Q(n) &= \prod_{k=1}^n r(k, n) \\ &= c^n \left(\prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\ &\quad \times \exp \left(O \left(\frac{n}{N} + \sum_{i=1}^N \#\{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right) \end{aligned}$$

Proof Outline

Proof Outline

$$c^n \left(\prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\ \times \exp \left(O \left(\frac{n}{N} + \sum_{i=1}^N \#\{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right)$$

Proof Outline

$$\begin{aligned}
 & c^n \left(\prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\
 & \times \exp \left(O \left(\frac{n}{N} + \sum_{i=1}^N \# \{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right) \\
 & = c^n \exp \left(O \left(\sum_{i=1}^N \sum_{d < m} d + \frac{n}{N} + N \# \{k \leq n : d|k, d > M, P^+(d) < N\} \right) \right)
 \end{aligned}$$

Proof Outline

$$\begin{aligned}
 & c^n \left(\prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\
 & \times \exp \left(O \left(\frac{n}{N} + \sum_{i=1}^N \# \{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right) \\
 & = c^n \exp \left(O \left(\sum_{i=1}^N \sum_{d < m} d + \frac{n}{N} + N \# \{k \leq n : d|k, d > M, P^+(d) < N\} \right) \right)
 \end{aligned}$$

Proof Outline

$$\begin{aligned}
 & c^n \left(\prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\
 & \quad \times \exp \left(O \left(\frac{n}{N} + \sum_{i=1}^N \# \{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right) \\
 & = c^n \exp \left(O \left(\sum_{i=1}^N \sum_{d < m} d + \frac{n}{N} + N \# \{k \leq n : d|k, d > M, P^+(d) < N\} \right) \right) \\
 & = c^n \exp \left(O \left(NM^2 + \frac{n}{N} + Nn \exp \left(-\frac{\log M}{2 \log N} \right) \right) \right)
 \end{aligned}$$

Proof Outline

$$\begin{aligned}
 & c^n \left(\prod_{i=1}^N \prod_{\substack{d \\ P^+(d) < i \\ d < M}} \prod_{t \in [id, (i+1)d)} r(d, t)^{O(1)} \right) \\
 & \quad \times \exp \left(O \left(\frac{n}{N} + \sum_{i=1}^N \# \{k \leq n : d|k, d > M, P^+(d) < i\} \right) \right) \\
 & = c^n \exp \left(O \left(\sum_{i=1}^N \sum_{d < m} d + \frac{n}{N} + N \# \{k \leq n : d|k, d > M, P^+(d) < N\} \right) \right) \\
 & = c^n \exp \left(O \left(NM^2 + \frac{n}{N} + Nn \exp \left(-\frac{\log M}{2 \log N} \right) \right) \right)
 \end{aligned}$$

Take $M = n^{1/2-\epsilon}$, $N = \exp((\log n)^{1/2-\epsilon})$.

Lower Bound

Lower Bound

We can get a lower bound for c by computing terms in this product.

Lower Bound

We can get a lower bound for c by computing terms in this product.

All computations were done in Sage.

Lower Bound

We can get a lower bound for c by computing terms in this product.
All computations were done in Sage.

#24418 closed defect (fixed) 8.2 β 4 Opened 5 months ago Closed 4 months ago

Doctest: bug numerical_approx($2^{450232897/4888643760}$)

Reported by:	vdelecroix	Owned by:	
Priority:	major	Milestone:	sage-8.2
Component:	symbolics	Keywords:	bug
Cc:	rws	Merged in:	
Authors:	Ralf Stephan	Reviewers:	Jeroen Demeyer
Report Upstream:	N/A	Work issues:	
Branch:	821f7d9 (Commits)	Commit:	821f7d9f3568316bc0b8b1f5619bce...
Dependencies:		Stopgaps:	

Description (last modified by [vdelecroix](#)) Δ

```
sage: numerical_approx(2^(450232897/4888643760))
-----
RuntimeError                                Traceback (most recent call last)
<ipython-input-2-3c4e30ac02c1> in <module>()
----> 1 numerical_approx(Integer(2)**(Integer(450232897)/Integer(4888643760)))

/opt/sage/local/lib/python2.7/site-packages/sage/misc/functional.pyc in numerical_ap
1406         return numerical_approx_generic(x, prec)
1407     else:
```

Lower Bound

We can get a lower bound for c by computing terms in this product.
All computations were done in Sage.

Lower Bound

We can get a lower bound for c by computing terms in this product.
All computations were done in Sage.

Computing $r(d, t)$ with $dk \leq t \leq d(k+1)$ for all pairs (d, k) with $d \times k \leq 720$ gives

$$1.572071 < c.$$

Lower Bound

We can get a lower bound for c by computing terms in this product.
All computations were done in Sage.

Computing $r(d, t)$ with $dk \leq t \leq d(k+1)$ for all pairs (d, k) with $d \times k \leq 720$ gives

$$1.572071 < c.$$

This technique doesn't give very good upper bounds since any unestimated terms need to be replaced by 2 (or 1.5).

Lower Bound

We can get a lower bound for c by computing terms in this product.
All computations were done in Sage.

Computing $r(d, t)$ with $dk \leq t \leq d(k+1)$ for all pairs (d, k) with $d \times k \leq 720$ gives

$$1.572071 < c.$$

This technique doesn't give very good upper bounds since any unestimated terms need to be replaced by 2 (or 1.5). Gives $c < 1.60$.

(Conjectural) Upper Bound

(Conjectural) Upper Bound

Do the same thing including integers from the beginning of the interval rather than the end.

(Conjectural) Upper Bound

Do the same thing including integers from the beginning of the interval rather than the end.

Let

$$g(k) = \frac{\# \text{Primitive subsets of } [1, k]}{\# \text{Primitive subsets of } [1, k-1]} - 1$$

(Conjectural) Upper Bound

Do the same thing including integers from the beginning of the interval rather than the end.

Let

$$\begin{aligned} g(k) &= \frac{\# \text{Primitive subsets of } [1, k]}{\# \text{Primitive subsets of } [1, k-1]} - 1 \\ &= \frac{\# \text{Primitive subsets of } [1, k] \text{ including } k}{\# \text{Primitive subsets of } [1, k-1]} \end{aligned}$$

(Conjectural) Upper Bound

Do the same thing including integers from the beginning of the interval rather than the end.

Let

$$\begin{aligned} g(k) &= \frac{\# \text{Primitive subsets of } [1, k]}{\# \text{Primitive subsets of } [1, k-1]} - 1 \\ &= \frac{\# \text{Primitive subsets of } [1, k] \text{ including } k}{\# \text{Primitive subsets of } [1, k-1]} \end{aligned}$$

As before,

$$Q(n) = \prod_{k=1}^n (g(k) + 1)$$

(Conjectural) Upper Bound

(Conjectural) Upper Bound

Conjecture

The function $g(n)$ is submultiplicative.

(Conjectural) Upper Bound

Conjecture

The function $g(n)$ is submultiplicative. If $(n,m)=1$ then

$$g(nm) \leq g(n)g(m).$$

(Conjectural) Upper Bound

Conjecture

The function $g(n)$ is submultiplicative. If $(n,m)=1$ then

$$g(nm) \leq g(n)g(m).$$

Furthermore, for any prime p , $g(p^i) > g(p^{i+1})$.

(Conjectural) Upper Bound

Conjecture

The function $g(n)$ is submultiplicative. If $(n,m)=1$ then

$$g(nm) \leq g(n)g(m).$$

Furthermore, for any prime p , $g(p^i) > g(p^{i+1})$.

The conjecture holds for all values of $n < 899$.

(Conjectural) Upper Bound

Conjecture

The function $g(n)$ is submultiplicative. If $(n,m)=1$ then

$$g(nm) \leq g(n)g(m).$$

Furthermore, for any prime p , $g(p^i) > g(p^{i+1})$.

The conjecture holds for all values of $n < 899$.

For example, $g(2) = \frac{1}{2}$, and $g(n) \leq \frac{1}{2}$ for all even n .

(Conjectural) Upper Bound

Conjecture

The function $g(n)$ is submultiplicative. If $(n,m)=1$ then

$$g(nm) \leq g(n)g(m).$$

Furthermore, for any prime p , $g(p^i) > g(p^{i+1})$.

The conjecture holds for all values of $n < 899$.

For example, $g(2) = \frac{1}{2}$, and $g(n) \leq \frac{1}{2}$ for all even n .

Note: $g(p) = \frac{Q(p-1)-1}{Q(p-1)}$.

(Conjectural) Upper Bound

(Conjectural) Upper Bound

Assuming the conjecture, using the computed values of $g(n)$, $n < 899$, and taking into account all 31-smooth numbers gives

(Conjectural) Upper Bound

Assuming the conjecture, using the computed values of $g(n)$, $n < 899$, and taking into account all 31-smooth numbers gives

$$c < 1.573487.$$

(Conjectural) Upper Bound

Assuming the conjecture, using the computed values of $g(n)$, $n < 899$, and taking into account all 31-smooth numbers gives

$$c < 1.573487.$$

This gives (conjecturally)

$$1.572071 < c < 1.573487.$$

Extensions

Extensions

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$.

Extensions

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$.
Recall that the largest primitive subset of this interval has size $\left\lceil \frac{n}{2} \right\rceil$.

Extensions

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$. Recall that the largest primitive subset of this interval has size $\left\lceil \frac{n}{2} \right\rceil$. Let $M(n)$ count the primitive subsets of $\{1, \dots, n\}$ of size $\left\lceil \frac{n}{2} \right\rceil$.

Extensions

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$. Recall that the largest primitive subset of this interval has size $\left\lceil \frac{n}{2} \right\rceil$. Let $M(n)$ count the primitive subsets of $\{1, \dots, n\}$ of size $\left\lceil \frac{n}{2} \right\rceil$.

Theorem ([Vijay](#), 2018)

$$1.303^n < M(2n) < 1.408^n$$

Extensions

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$. Recall that the largest primitive subset of this interval has size $\left\lceil \frac{n}{2} \right\rceil$. Let $M(n)$ count the primitive subsets of $\{1, \dots, n\}$ of size $\left\lceil \frac{n}{2} \right\rceil$.

Theorem (Vijay, 2018)

$$1.303^n < M(2n) < 1.408^n$$

Theorem (M., 2018)

For $\epsilon > 0$, there exists c' defined analogously to c such that

$$M(n) = c'^{n+O\left(\frac{n}{\exp((\log n)^{1/2-\epsilon})}\right)}$$

Extensions

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$. Recall that the largest primitive subset of this interval has size $\left\lceil \frac{n}{2} \right\rceil$. Let $M(n)$ count the primitive subsets of $\{1, \dots, n\}$ of size $\left\lceil \frac{n}{2} \right\rceil$.

Theorem (Vijay, 2018)

$$1.303^n < M(2n) < 1.408^n$$

Theorem (M., 2018)

For $\epsilon > 0$, there exists c' defined analogously to c such that

$$M(n) = c'^{n+O\left(\frac{n}{\exp((\log n)^{1/2-\epsilon})}\right)}$$

and $1.148070 < c'$ or $1.308066 < c'^2$

Extensions: Maximal Primitive Sets

Extensions: Maximal Primitive Sets

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$.

Extensions: Maximal Primitive Sets

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$.
Recall that the largest primitive subset of this interval has size $\left\lceil \frac{n}{2} \right\rceil$.

Extensions: Maximal Primitive Sets

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$. Recall that the largest primitive subset of this interval has size $\left\lceil \frac{n}{2} \right\rceil$. Let $M(n)$ count the primitive subsets of $\{1, \dots, n\}$ of size $\left\lceil \frac{n}{2} \right\rceil$.

Extensions: Maximal Primitive Sets

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$. Recall that the largest primitive subset of this interval has size $\left\lceil \frac{n}{2} \right\rceil$. Let $M(n)$ count the primitive subsets of $\{1, \dots, n\}$ of size $\left\lceil \frac{n}{2} \right\rceil$.

Theorem ([Vijay](#), 2018)

$$1.303^n < M(2n) < 1.408^n$$

Extensions: Maximal Primitive Sets

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$. Recall that the largest primitive subset of this interval has size $\left\lceil \frac{n}{2} \right\rceil$. Let $M(n)$ count the primitive subsets of $\{1, \dots, n\}$ of size $\left\lceil \frac{n}{2} \right\rceil$.

Theorem (Vijay, 2018)

$$1.303^n < M(2n) < 1.408^n$$

Theorem (M., 2018)

There exists c' defined analogously to c such that for $\epsilon > 0$,

$$M(n) = c'^{n+O\left(\frac{n}{\exp((\log n)^{1/2-\epsilon})}\right)}$$

Extensions: Maximal Primitive Sets

Use the same idea to count maximal primitive subsets of $\{1, \dots, n\}$. Recall that the largest primitive subset of this interval has size $\left\lceil \frac{n}{2} \right\rceil$. Let $M(n)$ count the primitive subsets of $\{1, \dots, n\}$ of size $\left\lceil \frac{n}{2} \right\rceil$.

Theorem (Vijay, 2018)

$$1.303^n < M(2n) < 1.408^n$$

Theorem (M., 2018)

There exists c' defined analogously to c such that for $\epsilon > 0$,

$$M(n) = c'^{n+O\left(\frac{n}{\exp((\log n)^{1/2-\epsilon})}\right)}$$

and $1.148070 < c'$ or $1.308066 < c'^2$.

Extensions: Geometric Progression Free Sets

Let $G(n)$ denote the number of subsets of $1, \dots, n$ that avoid 3-term geometric progressions with integral ratio.

Extensions: Geometric Progression Free Sets

Let $G(n)$ denote the number of subsets of $1, \dots, n$ that avoid 3-term geometric progressions with integral ratio.

Since the squarefree numbers avoid geometric progressions we get trivially that $G(n) > 2^{6n/\pi^2} \approx 1.5424^n$.

Theorem (M., 2018)

For $\epsilon > 0$, there exists g such that

$$G(n) = g^{n + O\left(\frac{n}{\exp((\log n)^{1/2 - \epsilon})}\right)}$$

Extensions: Geometric Progression Free Sets

Let $G(n)$ denote the number of subsets of $1, \dots, n$ that avoid 3-term geometric progressions with integral ratio.

Since the squarefree numbers avoid geometric progressions we get trivially that $G(n) > 2^{6n/\pi^2} \approx 1.5424^n$.

Theorem (M., 2018)

For $\epsilon > 0$, there exists g such that

$$G(n) = g^{n + O\left(\frac{n}{\exp((\log n)^{1/2 - \epsilon})}\right)}$$

and $1.8192 < g$.

Thank you!