

Bayesian and Frequentist Multilevel Modeling

Andy Grogan-Kaylor

2020-03-09

Contents

1	Introduction	1
2	Conceptual Appropriateness	1
3	Prior Information	2
4	Smaller Samples	2
5	Full Distribution of Parameters	2
6	Distributional Models	2
7	Non-Linear Terms	2
8	Maximal Models	2
	References	3

1 Introduction

$$y_{ij} = \beta_0 + \beta_1 x_{1i} + u_{0j} + e_{ij}$$

All multilevel models account for group structure, in estimating the association of x and y , by including a random intercept (u_0), and possibly one or more random intercept terms ($u_1, u_2, etc....$).

Bayesian models may offer some advantages over frequentist models, but may be substantially slower to converge.

2 Conceptual Appropriateness

Following (Kruschke 2014) all Bayesian models have a *conceptual appropriateness*.

In frequentist reasoning we are estimating the probability of observing data at least as extreme as our data, while assuming a null hypothesis (H_0). Quite often, H_0 , e.g. $\beta = 0$, or $\bar{x}_A - \bar{x}_B = 0$, is not a substantively interesting or substantively meaningful hypothesis.

In Bayesian analysis, we are not rejecting a null hypothesis. Instead, we are *directly estimating the value of a parameter* such as β and are indeed estimating a *full probability distribution* for this parameter.

Such an approach means that we have the ability to accept the null hypothesis H_0 (Kruschke and Liddell 2018). This ability to accept H_0 might possibly lead to theory simplification (Morey, Homer, and

Proulx 2018), as well as to a lower likelihood of the publication bias that results from frequentist methods predicated upon the rejection of H_0 (Kruschke and Liddell 2018).

3 Prior Information

Bayesian models allow one to incorporate prior information about a parameter of interest.

Prior information may come from the prior research literature, e.g. from systematic reviews or meta-analyses, or expert opinion or clinical wisdom.

4 Smaller Samples

Bayesian multilevel models may be better with small samples, especially samples with small numbers of Level 2 units (Hox et al. 2012). It is not clear to what degree this improvement in performance is dependent upon the use of informative priors.

5 Full Distribution of Parameters

Bayesian models of all kinds provide full distributions of the parameters (e.g. β 's and random effects (u 's))—both singly and jointly—rather than only point estimates.

Information about the full distribution of a parameter, such as the estimate of the probability distribution of values of a risk factor, a protective factor, or the effect of an intervention, may be substantively meaningful.

6 Distributional Models

Bayesian estimators allow one to directly model σ_{u_0} , the variance of the Level 2 units, as a function of covariates (Burkner 2018).

7 Non-Linear Terms

Bayesian estimators allow for the incorporation of non-linear terms (Burkner 2018).¹

8 Maximal Models

Bayesian estimators allow for the estimation of so called *maximal models* (Barr et al. 2013; Frank 2018), which allow for the inclusion




Figure 1: Distribution of a Single Parameter

Joint Distribution of Random Effects




Figure 2: Joint Distribution of Parameters




Figure 3: Non-Linear Terms

¹ Such non-linear terms offer ways of non-parametrically fitting curvature. Do they represent over fitting? Do they

of a large number of random slopes, e.g. u_1, u_2, u_3, \dots , etc. even when some of those estimated slopes are close to 0.

In contrast, (Matuschek et al. 2017) argue that such a *maximal* approach may lead to a loss of statistical power and further argue that one should adhere to “a random effect structure that is supported by the data.”

References

- Barr, Dale J., Roger Levy, Christoph Scheepers, and Harry J. Tily. 2013. “Random Effects Structure for Confirmatory Hypothesis Testing: Keep It Maximal.” *Journal of Memory and Language* 68 (3): 255–78. <https://doi.org/10.1016/j.jml.2012.11.001>.
- Burkner, Paul-Christian. 2018. “Advanced Bayesian Multilevel Modeling with the R Package Brms.” *The R Journal* 10 (1): 395–411.
- Frank, Michael. 2018. “Mixed Effects Models: Is It Time to Go Bayesian by Default?” <http://babieslearninglanguage.blogspot.com/2018/02/mixed-effects-models-is-it-time-to-go.html>.
- Hox, Joop, Joop J. C.M. Hox, Rens van de Schoot, and Suzette Matthijsse. 2012. “How Few Countries Will Do? Comparative Survey Analysis from a Bayesian Perspective.” *Survey Research Methods*. <https://doi.org/10.18148/srm/2012.v6i2.5033>.
- Kruschke, John K. 2014. *Doing Bayesian Data Analysis: A Tutorial with R, Jags, and Stan, Second Edition*. Elsevier Science. <https://doi.org/10.1016/B978-0-12-405888-0.09999-2>.
- Kruschke, John K, and Torrin M Liddell. 2018. “The Bayesian New Statistics: Hypothesis Testing, Estimation, Meta-Analysis, and Power Analysis from a Bayesian Perspective.” *Psychonomic Bulletin & Review* 25 (1): 178–206. <https://doi.org/10.3758/s13423-016-1221-4>.
- Matuschek, Hannes, Reinhold Kliegl, Shravan Vasishth, Harald Baayen, and Douglas Bates. 2017. “Balancing Type I Error and Power in Linear Mixed Models.” *Journal of Memory and Language*. <https://doi.org/10.1016/j.jml.2017.01.001>.
- Morey, Richard D., Saskia Homer, and Travis Proulx. 2018. “Beyond Statistics: Accepting the Null Hypothesis in Mature Sciences.” *Advances in Methods and Practices in Psychological Science*. <https://doi.org/10.1177/2515245918776023>.