# A Quick Introduction to R

## Andy Grogan-Kaylor

#### 2024-12-16

#### Table of contents

1	Why Use R?	1
2	Get R	2
3	Get Data	2
4	Process and Clean Data	2
5		3
6	Analyze Data: Descriptive Statistics	6

# 1 Why Use R?

R has a reputation for being difficult to learn, and a lot of that reputation is deserved. However, it is possible to teach R in an accessible way, and a little bit of R can take you a long way.

R is open source, and therefore free, statistical software that is particularly good at obtaining, analyzing and visualizing data.

R Commands are stored in a *script* or *code* file that usually ends in .R, e.g. myscript.R. The command file is distinct from your actual data, stored in an .RData file, e.g. mydata.RData.

A great deal of data analysis and visualization involves the same core set of steps.

Given the fact that we often want to apply the same core set of tasks to new questions and new data, there are ways to overcome the steep learning curve and learn a replicable set of commands that can be applied to problem after problem. The same 5 to 10 lines of R code can often be tweaked over and over again for multiple projects.



Figure 1: A common data workflow

#### 2 Get R

R is available at https://www.r-project.org/. R is a lot easier to run if you run it from RStudio, http://www.rstudio.com.

#### 3 Get Data

Data often comes from other types of data files like SPSS, Stata, or Excel. Especially in beginning R programming, getting the data into R can be the most complicated part of your program.

```
load("the/path/to/mydata.Rdata") # data in R format

library(haven) # library for importing data

mydata <- read_sav("the/path/to/mySPSSfile.sav") # SPSS

mydata <- read_dta("the/path/to/myStatafile.dta") # Stata

library(readxl) # library for importing Excel files

mydata <- read_excel("the/path/to/mySpreadsheet.xls")

save(mydata, file = "mydata.RData") # save in R format</pre>
```

#### 4 Process and Clean Data

The \$ sign is a kind of "connector". mydata\$x means: "The variable x in the dataset called mydata".

```
mydata$x[mydata$x == -9] <- NA # missing to NA
```

R makes a strong distinction between *continuous numeric* variables that measure scales like mental health or neighborhood safety, and *categorical factor variables* that measure non-ordered categories like religious identity or gender identity.


Many statistical and graphical procedures are designed to recognize and work with different variable types. You often *don't* need to use all of the options. e.g. mydata\$w <-factor(mydata\$z) will often work just fine. Changing variables from factor to numeric, and vice versa can sometimes be the simple solution that solves a lot of problems when you are trying to graph your variables.

#### 5 Visualize Data

#### 5.1 Histogram

```
hist(mydata$x, # what I'm graphing
  main = "your title goes here", # title
  xlab = "income", # label for x axis
  col = "blue") # color
```

# your title goes here




# **?** Tip

You often don't need to use all of the options. e.g. hist(mydata\$x) will work just fine.

### 5.2 Barplot

```
barplot(table(mydata$z), # what I'm graphing
    names.arg = c("Group A", "Group B"), # names
    main = "your title goes here", # title
    xlab = "group", # label for x axis
    col = "gold") # color
```

## your title goes here




## **?** Tip

You often *don't* need to use all of the options. e.g. barplot(table(mydata\$z)) will work just fine.

#### 5.3 Scatterplot

```
plot(mydata$x, mydata$y, # plot x and y
    main = "your title goes here", # title
    xlab = "income", # label for x axis
    ylab = "mental health", # label for y axis
    pch = 19, # Plot CHaracter, 19 is filled dots
    las = 2, # LAbel Style, 2 is "perpendicular"
    col = "darkgreen") # color
```

# your title goes here



## **?** Tip

You often don't need to use all of the options. e.g. plot(mydata\$x, mydata\$y) will work just fine.

### **?** Tip

When scatterplots have fewer dots than you think they should have, often due to "over-printing", adding some random noise, or "jittering" the dots in the scatterplot may help: plot(jitter(mydata\$y, factor = 5000) ~ mydata\$x). Experiment with different sizes of factor.

# 6 Analyze Data: Descriptive Statistics

```
summary(mydata$x) # for continuous or factor variables
```

```
Min. 1st Qu. Median Mean 3rd Qu. Max. 81.66 94.50 100.28 100.68 106.70 124.84
```

table(mydata\$z) # especially suitable for factor variables

1 2

### 78 22

For another approach to summarizing your data, try:

```
library(skimr)
skim(mydata)
skim(mydata$x)
```