

Multilevel Models, Equations, Syntax & English

Andy Grogan-Kaylor

2024-06-13

model	equation	Stata	English
Intercept Only	$y = \beta_0 + e_{ij}$	<code>mixed y</code>	We estimated the mean of [outcome]
Intercept Independent Variable(s)	$y = \beta_0 + \beta_1 x + e_{ij}$ $y = \beta_0 + \beta_1 x + \beta_2 z + e_{ij}$	<code>mixed y x</code> <code>mixed y x z</code>	We estimated the relationship of [independent variable(s)] with [outcome]
Intercept Random variation of the intercept	$y = \beta_0 + e_{ij} + u_{0j}$	<code>mixed y groupid:</code>	We estimated the mean of [outcome]. We allowed the intercept of the model to vary by [groupid].
Unconditional intraclass correlation coefficient (ICC)	$\frac{var(u_{0j})}{var(u_{0j}) + var(e_{ij})}$	<code>mixed y groupid: estat icc</code>	XX% of the variation in [outcome] was explained by clustering of participants in [groupid]

model	equation	Stata	English
Intercept Independent variable(s) Random variation of the intercept	$y = \beta_0 + \beta_1 x + e_{ij} + u_{0j}$ $y = \beta_0 + \beta_1 x + \beta_2 z + e_{ij} + u_{0j}$	<pre>mixed y x groupid: mixed y x z groupid:</pre>	<p>We estimated the relationship of [independent variable(s)] with [outcome]. We allowed the intercept of the model to vary by group.</p>
Intercept Independent variable Random intercept Random slope	$y = \beta_0 + \beta_1 x + e_{ij} + u_{0j} + u_{1j} x$	<pre>mixed y x groupid: x</pre>	<p>We estimated the relationship of [independent variable] with [outcome]. We allowed the intercept of the model to vary by group. We also allowed the relationship of [independent variable] with [outcome] to vary by group.</p>
We can estimate multilevel models with more than 1 random slope.	$y = \beta_0 + \beta_1 x + \beta_2 z + e_{ij} + u_{0j} + u_{1j} x + u_{2j} z$	<pre>mixed y x z groupid: x z</pre>	