

Why Multilevel Models Are Good Models For Longitudinal Data

Multilevel Models Offer An Incredibly Flexible Treatment of Time and Time Varying Processes and Covariates

Andy Grogan-Kaylor

2025-06-12

Table of contents

| | | |
|---|--|---|
| 1 | Visually | 1 |
| 2 | Data Structures | 1 |
| 3 | Equation | 2 |
| 4 | Advantages of the Multilevel Model for Longitudinal Data | 3 |
| | References | 3 |

1 Visually

2 Data Structures

Multilevel models for longitudinal data prefer data in long format.

Table 1: Data in WIDE format

| id | x1 | x2 | x3 | y1 | y2 | y3 |
|----|----|----|----|----|----|----|
| 1 | | | | | | |
| 2 | | | | | | |
| 3 | | | | | | |




Figure 1: A Multilevel Model For Longitudinal Data

Table 2: Data in LONG format

| id | t | x | y |
|----|---|---|---|
| 1 | 1 | | |
| 1 | 2 | | |
| 1 | 3 | | |
| 2 | 1 | | |
| 2 | 2 | | |
| 2 | 3 | | |
| 3 | 1 | | |
| 3 | 2 | | |
| 3 | 3 | | |

3 Equation

$$y_{it} = \beta_0 + \beta_1 t_{it} + \beta_2 x_{it} + u_{0i} + e_{it} \quad (1)$$

💡 Person-Observations

Every row is a *person-observation* (person i observed at time t). Every person has *multiple rows*.

4 Advantages of the Multilevel Model for Longitudinal Data

1. Using the algebra in Equation 1, these models can easily accommodate both time varying and time invariant coefficients (Hox, 2010; Hox et al., 2018; Rabe-Hesketh & Skrondal, 2022; Raudenbush & Bryk, 2002; Singer & Willett, 2003).
2. There is no multicollinearity issue with multiple β coefficients for multiple waves of data. By inspection of Equation 1, we see that there is only a single β coefficient for each variable, \therefore no multicollinearity problem.
3. *Unbalanced data is less of a problem*, the data structure and estimation are robust to these possibilities (Raudenbush, 1995; Raudenbush & Bryk, 2002; Singer & Willett, 2003).
4. *Missing data is less of a problem* (assuming MCAR). When a person observation is missing, that person simply has fewer rows of data (Hox, 2010; Luke, 2004; Rabe-Hesketh & Skrondal, 2022; Raudenbush, 1995; Raudenbush & Bryk, 2002). But all rows of data are “matched” to the same person by i .
5. We have an *explicit function of time* $\beta_1 t$, and could treat time more flexibly, by creating a polynomial function of time e.g. by adding $\beta_2 t^2$, etc. (Raudenbush & Bryk, 2002; Singer & Willett, 2003). (We could even substitute $\beta \ln(t)$.)
6. Again, by inspection of Equation 1, we see that *multiple or many time-points are not a problem*. We would use the same algebra for 2 time points or for 10,000 time points. (Helpful when we start to think about intensive longitudinal data e.g. George Holden’s *recording study*).
7. We are *measuring exactly the time at which events take place* for each individual (Luke, 2004; Singer & Willett, 2003). Not simply saying *Wave 1, Wave 2, Wave 3*, etc...
8. Unequally spaced time points are not a problem (Raudenbush, 1995). Every individual could have a *completely different set of time points* and even a *completely different number of time points* (Hox, 2010; Hox et al., 2018; Luke, 2004; Singer & Willett, 2003).

Caution

We do need to think carefully about what is the appropriate variable for time. Is it the variable we used to reshape the data—often **wave**—or some other more appropriate metric, like **age** or **time in study** (Singer & Willett, 2003)?

References

- Hox, J. (2010). *Multilevel analysis: Techniques and applications* (2nd ed.). Routledge.
- Hox, J., Moerbeek, M., & van de Schoot, R. (2018). *Multilevel analysis: Techniques and applications* (Third edition.). Routledge, Taylor & Francis Group.
- Luke, D. (2004). *Multilevel modeling*. SAGE Publications, Inc. <https://doi.org/10.4135/9781412985147>
- Rabe-Hesketh, S., & Skrondal, A. (2022). Multilevel and longitudinal modeling using stata. In *Stata Press* (4th ed.). Stata Press.
- Raudenbush, S. W. (1995). Hierarchical linear models to study the effects of social context on development. In J. M. Gottman (Ed.), *The analysis of change* (pp. 165–199). Lawrence Erlbaum Associates.
- Raudenbush, S. W., & Bryk, A. S. (2002). *Hierarchical linear models: Applications and data analysis methods*. Sage Publications.

Singer, J. D., & Willett, J. B. (2003). *Applied longitudinal data analysis : Modeling change and event occurrence*. Oxford University Press.