

Multilevel Multilingual

Multilevel Models in Stata, R and Julia

Andrew Grogan-Kaylor

2024-06-27

Table of contents

1	Multilevel Multilingual	4
1.1	Introduction	4
1.2	The Data	6
1.3	An Introduction To Equations and Syntax	7
1.3.1	Stata	7
1.3.2	R	7
1.3.3	Julia	7
2	Statistical Workflows	8
2.1	Statistical Software Is Best Run Using a Script	8
2.2	Scripts	9
2.3	Script Flow	9
2.4	Storing Statistical Data	9
2.5	It Is Possible To Use Multiple Statistical Packages	9
2.6	Good Statistical Workflows Require Safe Workspaces	10
2.7	Good Statistical Workflows Require Patience And Can Be Psychologically Demanding	10
3	Descriptive Statistics	11
3.1	Descriptive Statistics	11
3.1.1	Stata	11
3.1.2	R	12
3.1.3	Julia	13
3.2	Interpretation	14
4	Unconditional Models	15
4.1	Two Level Model	15
4.1.1	The Equation	15
4.1.2	Run Models	15
4.1.3	Interpretation	18
4.2	Three Level Model	19
4.2.1	The Equation	19
4.2.2	Run Models	19
4.2.3	Interpretation	23

5	Cross Sectional Multilevel Models	24
5.1	The Equation	24
5.2	Correlated and Uncorrelated Random Effects	24
5.3	Run Models	25
5.3.1	Stata	25
5.3.2	R	27
5.3.3	Julia	29
5.4	Interpretation	30
6	Longitudinal Multilevel Models	31
6.1	The Data	31
6.2	The Equation	31
6.3	Growth Trajectories	31
6.4	Run Models	32
6.4.1	Stata	32
6.4.2	R	36
6.4.3	Julia	40
6.5	Interpretation	42
	References	44

List of Tables

1.1	Software for Multilevel Modeling	4
1.2	Sample of Simulated Multilevel Data	6
1.2	Table continues below	6
1.3	Sample of Simulated Multilevel Data	6
5.1	Correlated and Uncorrelated Random Effects	25
6.1	Slope and Intercept for Each Group	32

1 Multilevel Multilingual

“This curious world which we inhabit is more wonderful than it is convenient...”
(Thoreau, 1975)

“Mathematics is my secret. My secret weakness. I feel like a stubborn, helpless fool in the middle of a problem. Trapped and crazed. Also, thrilled.” (Schanen, 2021)

1.1 Introduction

Below, I describe the use of [Stata](#) (StataCorp, 2023), [R](#) (Bates et al., 2015; R Core Team, 2023), and [Julia](#) (Bates, 2024; Bezanson et al., 2017) to estimate multilevel models.

All of these software packages can estimate multilevel models and can visualize relationships in the data. However, there are substantial differences between the different packages: Stata is proprietary *for cost* software, which is very well documented and very intuitive. While it costs money to purchase Stata, the price is often very reasonable for academic and educational use. R is free open source software which is less intuitive, but there are many excellent resources for learning R. There is often a cost associated with purchasing books and other materials for learning R, which sometimes feels like it offsets the fact that R is free. Julia is newer open source software, and ostensibly much faster than either Stata or R, which may be an important advantage when running multilevel models with very large data sets. At this point in time, both Stata and R feel much more *stable* than Julia which is still evolving software.

While any of these software packages can be used for learning and estimating multilevel models, I will offer my own opinion—based upon 15 years of teaching multilevel models at the doctoral level—that Stata offers the quickest pathway for learning the basic and advanced uses of multilevel models. I also believe the intuitive nature of Stata syntax contributes to accurate and replicable work in this area.

Table 1.1: Software for Multilevel Modeling

Software	Cost	Ease of Use
Stata	some cost	learning curve, but very intuitive for both multilevel modeling and graphing.

Software	Cost	Ease of Use
R	free	learning curve: intuitive for multilevel modeling; but steeper learning curve for graphing (<code>ggplot</code>).
Julia	free	steep learning curve in general: steep learning curve for multilevel modeling; and very steep learning curve for graphing. Graphics libraries are very much under development and in flux.

💡 Results Will Vary Somewhat

Estimating multilevel models is a complex endeavor. The software details of how this is accomplished are beyond the purview of this book. Suffice it to say that across different software packages there will be differences in estimation routines, resulting in some numerical differences in the results provided by different software packages. Substantively speaking, however, results should agree across software.

💡 Multi-Line Commands

Sometimes I have written commands out over multiple lines. I have done this for especially long commands, but have also sometimes done this simply for the sake of clarity. The different software packages have different approaches to multi-line commands.

1. By default, *Stata* ends a command at the end of a line. If you are going to write a multi-line command you should use the `///` line continuation characters.
2. *R* is the software that most naturally can be written using multiple lines, as *R* commands are usually clearly encased in parentheses `()` or continued with `+` signs.
3. Like *Stata*, *Julia* expects commands to end at the end of a line. If you are going to write a multi-line command, all commands except for the last line should end in a character that clearly indicates continuation, like a `+` sign. An alternative is to encase the entire *Julia* command in an outer set of parentheses `()`.

💡 Running Statistical Packages in Quarto

I used Quarto (Allaire et al., 2024) (<https://quarto.org/>) to create this Appendix. Quarto is a programming and publishing environment that can run multiple programming languages, including *Stata*, *R* and *Julia*, and that can write to multiple output formats including HTML, PDF, and MS Word. To run *Stata*, I used the `Statamarkdown` library (Hemken, 2023) in *R* to connect *Stata* to Quarto. Quarto has a built in connection to *R*,

and runs R without issue. To run Julia, I used the `JuliaCall` library (Li, 2019) in R to connect Quarto to Julia.

Of course, each of these programs can be run by itself, if you have them installed on your computer.

1.2 The Data

Datasets

The examples use the `simulated_multilevel_data.dta` and `simulated_multilevel_longitudinal_data.dta` files.

Here is a [direct link](#) to download the cross-sectional data.

Here is a [direct link](#) to download the longitudinal data.

Table 1.2: Sample of Simulated Multilevel Data

Table 1.2: Table continues below

country	HDI	family	id	identity	intervention	physical_punishment
1	69	1	1.1	1	0	3
1	69	2	1.2	1	1	2
1	69	3	1.3	0	1	3
1	69	4	1.4	1	0	0
1	69	5	1.5	1	0	4
1	69	6	1.6	0	1	5

Table 1.3: Sample of Simulated Multilevel Data

warmth	outcome
3	57.47
1	50.1
2	52.92
5	60.17
4	55.05
3	49.81

1.3 An Introduction To Equations and Syntax

To explain statistical syntax for each software, I consider the general case of a multilevel model with dependent variable y , independent variables \mathbf{x} and \mathbf{z} , clustering variable `group`, and a random slope for \mathbf{x} . i is the index for the person, while j is the index for the group.

$$y = \beta_0 + \beta_1 x_{ij} + \beta_2 z_{ij} + u_{0j} + u_{1j} \times x_{ij} + e_{ij} \quad (1.1)$$

1.3.1 Stata

In Stata `mixed`, the syntax for a multilevel model of the form described in Equation 1.1 is:

```
mixed y x z || group: x
```

1.3.2 R

In R `lme4`, the syntax for a multilevel model of the form described in Equation 1.1 is:

```
library(lme4)

lmer(y ~ x + z + (1 + x || group), data = ...)
```

1.3.3 Julia

In Julia `MixedModels`, the syntax for a multilevel model of the form described in Equation 1.1 is:

```
using MixedModels

fit(MixedModel, @formula(y ~ x + z + (1 + x | group)), data)
```


2 Statistical Workflows

2.1 Statistical Software Is Best Run Using a Script

Many statistical workflows—whatever the statistical package being used—follow the same conceptual pattern.




Figure 2.1: A Common Statistical Workflow

Increasingly, we want to think about workflows that are

- **documentable, transparent, and auditable:** We have a record of what we did if we want to double check our work, clarify a result, or develop a new project with a similar process. We, or others, can find the inevitable errors in our work, **and correct them**.
- **replicable:** Others can replicate our findings with the same or new data.

- **scalable:** We are developing a process that can be as easily used with *thousands* or *millions* of rows of data as it can with *ten* rows of data. We are developing a process that can be easily repeated if we are *constantly getting new or updated data*, e.g. getting new data every week, or every month.

2.2 Scripts

For most statistical workflows, we will often want to write a script or code. Data analysis scripts can be stored in a Quarto document (Allaire et al., 2024) as they are in this Appendix, or every statistical package has its own unique format for storing scripts as a text file: in Stata, scripts are stored in `.do` files; in R, scripts are stored in `.R` files, and in Julia, scripts are stored in `.jl` files.

2.3 Script Flow

A good practice when writing a script, is to have a script that begins with the raw data, moves through any necessary re-coding or cleaning of the data, generates descriptive statistics, generates the appropriate multivariate results, and then generates any necessary visualizations.

2.4 Storing Statistical Data

It is usually best to store quantitative data in a statistical format such as R (`.Rdata`), or Stata (`.dta`), or even a text format such as `.csv`. Spreadsheets are likely to be a bad tool for storing quantitative data.

2.5 It Is Possible To Use Multiple Statistical Packages

While this Appendix focuses on the use of each individual statistical package on its own, it is certainly possible to use multiple statistical packages as part of the same workflow. For example, one might employ Stata to carry out data management tasks, and then possibly use R to run a multilevel model with a more complicated multilevel structure, such as a cross-classified model, or Julia to more quickly run a model with a large data.

2.6 Good Statistical Workflows Require Safe Workspaces

It is also *very important* to be aware that good complex workflows are *highly iterative* and *highly collaborative*. Good complex workflows require a *safe workspace* in which team members feel free to admit their own errors, and help with others' mistakes in a non-judgmental fashion. Such a *safe environment* is necessary to build an environment where the *overall error rate* is low.

2.7 Good Statistical Workflows Require Patience And Can Be Psychologically Demanding

Developing a good documented and auditable workflow that is implemented in code requires a lot of patience, and often, **many iterations**. Working through these many iterations can be psychologically demanding. It is important to remember that careful attention to getting the details right early in the research process, while sometimes tiring and frustrating, will pay large dividends later on when the research is reviewed, presented, published and read.

3 Descriptive Statistics

3.1 Descriptive Statistics

3.1.1 Stata

```
use simulated_multilevel_data.dta // use data
```

We use `summarize` for *continuous* variables, and `tabulate` for *categorical* variables.

```
summarize outcome warmth physical_punishment HDI
```

```
tabulate identity
```

```
tabulate intervention
```

Variable	Obs	Mean	Std. dev.	Min	Max
-----+-----					
outcome	3,000	52.43327	6.530996	29.60798	74.83553
warmth	3,000	3.521667	1.888399	0	7
physical_p~t	3,000	2.478667	1.360942	0	5
HDI	3,000	64.76667	17.24562	33	87

hypothetica			
l identity			
group			
variable	Freq.	Percent	Cum.
-----+-----			
0	1,507	50.23	50.23
1	1,493	49.77	100.00
-----+-----			
Total	3,000	100.00	

recieved interventio	n	Freq.	Percent	Cum.
0	1,547	51.57	51.57	
1	1,453	48.43	100.00	
Total	3,000	100.00		

3.1.2 R

```
library(haven) # read data in Stata format

df <- read_dta("simulated_multilevel_data.dta")
```

R's descriptive statistics functions rely heavily on whether a variable is a *numeric* variable, or a *factor* variable. Below, I convert two variables to factors (`factor`) before using `summary`¹ to generate descriptive statistics.

```
df$country <- factor(df$country)

df$identity <- factor(df$identity)

df$intervention <- factor(df$intervention)

summary(df)
```

```

country      HDI      family      id      identity
1      : 100  Min.   :33.00  Min.   : 1.00  Length:3000  0:1507
2      : 100  1st Qu.:53.00  1st Qu.: 25.75  Class :character 1:1493
3      : 100  Median :70.00  Median : 50.50  Mode  :character
4      : 100  Mean   :64.77  Mean   : 50.50
5      : 100  3rd Qu.:81.00  3rd Qu.: 75.25
6      : 100  Max.   :87.00  Max.   :100.00
(Other):2400
intervention physical_punishment  warmth      outcome
0:1547      Min.   :0.000      Min.   :0.000  Min.   :29.61
1:1453      1st Qu.:2.000      1st Qu.:2.000  1st Qu.:48.02
```

¹`skimr` is an excellent new alternative library for generating descriptive statistics in R.

Median	:2.000	Median	:4.000	Median	:52.45
Mean	:2.479	Mean	:3.522	Mean	:52.43
3rd Qu.	:3.000	3rd Qu.	:5.000	3rd Qu.	:56.86
Max.	:5.000	Max.	:7.000	Max.	:74.84

3.1.3 Julia

```
using Tables, MixedModels, MixedModelsExtras, StatFiles, DataFrames, CategoricalArrays, DataAPI

df = DataFrame(load("simulated_multilevel_data.dta"))
```

Similarly to R, Julia relies on the idea of *variable type*. I use `transform` to convert the appropriate variables to *categorical* variables.

```
@transform!(df, :country = categorical(:country))

@transform!(df, :identity = categorical(:identity))

@transform!(df, :intervention = categorical(:intervention))
```

```
describe(df) # descriptive statistics
```

9×7 DataFrame

Row	variable Symbol	mean Union...	min Any	median Union...	max Any	nmissing Int64	eltype Union
1	country		1.0		30.0	0	Union{
2	HDI	64.7667	33.0	70.0	87.0	0	Union{
3	family	50.5	1.0	50.5	100.0	0	Union{
4	id		1.1		9.99	0	Union{
5	identity		0.0		1.0	0	Union{
6	intervention		0.0		1.0	0	Union{
7	physical_punishment	2.47867	0.0	2.0	5.0	0	Union{
8	warmth	3.52167	0.0	4.0	7.0	0	Union{
9	outcome	52.4333	29.608	52.449	74.8355	0	Union{

1 column omitted

3.2 Interpretation

Examining descriptive statistics is an important first step in any analysis. It is important to examine your descriptive statistics before skipping ahead to more sophisticated analyses, such as multilevel models.

In examining the descriptive statistics for this data, we get a sense of the data.

- `outcome` has a mean of approximately 52 and ranges from approximately 30 to 75.
- `warmth` and `physical punishment` are both variables that represent the number of times that parents use each of these forms of discipline in a week. The average of the former is about 3.5, while the average of the latter is about 2.5.
- `HDI`, the Human Development Index has an average of about 65, and a wide range.
- `identity` is a categorical variable for a hypothetical identity group, and has values of 0 and 1.
- `intervention` is also a categorical variable, and has values of 0 and 1.

4 Unconditional Models

4.1 Two Level Model

An *unconditional* multilevel model is a model with no independent variables. One should always run an unconditional model as the first step of a multilevel model in order to get a sense of the way that variation is apportioned in the model across the different levels.

4.1.1 The Equation

$$\text{outcome}_{ij} = \beta_0 + u_{0j} + e_{ij} \quad (4.1)$$

The Intraclass Correlation Coefficient (ICC) is given by:

$$\text{ICC} = \frac{\text{var}(u_{0j})}{\text{var}(u_{0j}) + \text{var}(e_{ij})} \quad (4.2)$$

In a two level multilevel model, the ICC provides a measure of the proportion of variation attributable to Level 2.

4.1.2 Run Models

4.1.2.1 Stata

```
use simulated_multilevel_data.dta // use data

mixed outcome || country: // unconditional model

estat icc // ICC
```


Performing EM optimization ...

Performing gradient-based optimization:

Iteration 0: Log likelihood = -9802.8371

Iteration 1: Log likelihood = -9802.8371

Computing standard errors ...

Mixed-effects ML regression

Group variable: country

Number of obs = 3,000

Number of groups = 30

Obs per group:

min = 100

avg = 100.0

max = 100

Wald chi2(0) = .

Prob > chi2 = .

Log likelihood = -9802.8371

outcome	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
-----+-----						
_cons	52.43327	.3451217	151.93	0.000	51.75685	53.1097

Random-effects parameters		Estimate	Std. err.	[95% conf. interval]	
country: Identity					
	var(_cons)	3.178658	.9226737	1.799552	5.614658
	var(Residual)	39.46106	1.024013	37.50421	41.52

LR test vs. linear model: chibar2(01) = 166.31 Prob >= chibar2 = 0.0000

Intraclass correlation

Level	ICC	Std. err.	[95% conf. interval]	
country	.0745469	.0201254	.0434963	.1248696

4.1.2.2 R

```
library(haven)

df <- read_dta("simulated_multilevel_data.dta")
```

```
library(lme4) # estimate multilevel models

fit0 <- lmer(outcome ~ (1 | country),
             data = df) # unconditional model

summary(fit0)
```

Linear mixed model fit by REML ['lmerMod']
Formula: outcome ~ (1 | country)
Data: df

REML criterion at convergence: 19605.9

Scaled residuals:

Min	1Q	Median	3Q	Max
-3.3844	-0.6655	-0.0086	0.6725	3.6626

Random effects:

Groups	Name	Variance	Std.Dev.
country	(Intercept)	3.302	1.817
Residual		39.461	6.282

Number of obs: 3000, groups: country, 30

Fixed effects:

	Estimate	Std. Error	t value
(Intercept)	52.433	0.351	149.4

```
library(performance)
```

```
performance::icc(fit0) # ICC
```

Intraclass Correlation Coefficient

Adjusted ICC: 0.077
Unadjusted ICC: 0.077

4.1.2.3 Julia

```
using Tables, MixedModels, MixedModelsExtras,  
StatFiles, DataFrames, CategoricalArrays, DataFramesMeta
```

```
df = DataFrame(load("simulated_multilevel_data.dta"))
```

```
@transform!(df, :country = categorical(:country))
```

```
m0 = fit(MixedModel,  
         @formula(outcome ~ (1 | country)), df) # unconditional model
```

Linear mixed model fit by maximum likelihood

outcome ~ 1 + (1 | country)

	logLik	-2 logLik	AIC	AICc	BIC
	-9802.8371	19605.6742	19611.6742	19611.6822	19629.6933

Variance components:

	Column	Variance	Std.Dev.
country	(Intercept)	3.17863	1.78287
Residual		39.46106	6.28180

Number of obs: 3000; levels of grouping factors: 30

Fixed-effects parameters:

	Coef.	Std. Error	z	Pr(> z)
(Intercept)	52.4333	0.345121	151.93	<1e-99

```
icc(m0) # ICC
```

```
0.07454637475695493
```

4.1.3 Interpretation

In each case, the software finds that nearly 8% of the variation in the outcome is explainable by the clustering of the observations in each country.

4.2 Three Level Model

4.2.1 The Equation

$$\text{outcome}_{ij} = \beta_0 + u_{0j} + v_{0i} + e_{ij} \quad (4.3)$$

As discussed in the main text, in a three level model, there are two intraclass correlation coefficients (StataCorp, 2023). The formulas for the Intraclass Correlation Coefficient (ICC) are given by (StataCorp, 2023):

$$\text{ICC} = \frac{\text{var}(u_{0j})}{\text{var}(u_{0j}) + \text{var}(v_{0i}) + \text{var}(e_{ij})} \quad (4.4)$$

Following StataCorp (2023), Equation 4.4 is the correlation of responses for person-timepoints from the same country but different persons.

$$\text{ICC} = \frac{\text{var}(u_{0j}) + \text{var}(v_{0i})}{\text{var}(u_{0j}) + \text{var}(v_{0i}) + \text{var}(e_{ij})} \quad (4.5)$$

Again, closely following StataCorp (2023), Equation 4.5 is the correlation of responses for person-timepoints from the same country and same person.

4.2.2 Run Models

4.2.2.1 Stata

```
use simulated_multilevel_longitudinal_data.dta // use data

mixed outcome || country: || id: // unconditional model

estat icc // ICC
```

Performing EM optimization ...

Performing gradient-based optimization:
Iteration 0: Log likelihood = -29058.266
Iteration 1: Log likelihood = -29058.259
Iteration 2: Log likelihood = -29058.259

Computing standard errors ...

Mixed-effects ML regression

Number of obs = 9,000

Grouping information

Group variable		No. of groups	Observations per group		
			Minimum	Average	Maximum
country		30	300	300.0	300
id		3,000	3	3.0	3

Log likelihood = -29058.259

Wald chi2(0) = .
Prob > chi2 = .

outcome	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
_cons	53.37768	.3387943	157.55	0.000	52.71366	54.04171

Random-effects parameters		Estimate	Std. err.	[95% conf. interval]	
country: Identity					
	var(_cons)	3.232092	.8891367	1.885043	5.54174
id: Identity					
	var(_cons)	11.72403	.5747501	10.64996	12.90641
	var(Residual)	28.23424	.5154843	27.24178	29.26287

LR test vs. linear model: chi2(2) = 1314.88 Prob > chi2 = 0.0000

Note: LR test is conservative and provided only for reference.

Intraclass correlation

Level	ICC	Std. err.	[95% conf. interval]
-------	-----	-----------	----------------------

country	.0748336	.0190847	.0450028	.1219141
id country	.3462837	.0171461	.3134867	.3806097

4.2.2.2 R

In R, the ICC for a three level model is easiest to estimate “by hand”.

```
library(haven)

dfL <- read_dta("simulated_multilevel_longitudinal_data.dta")
```

```
library(lme4) # estimate multilevel models

fit0L <- lmer(outcome ~ (1 | country/id),
              data = dfL) # unconditional model

summary(fit0L)
```

```
Linear mixed model fit by REML ['lmerMod']
Formula: outcome ~ (1 | country/id)
Data: dfL
```

REML criterion at convergence: 58116.8

Scaled residuals:

Min	1Q	Median	3Q	Max
-3.7858	-0.6059	-0.0062	0.6017	3.4348

Random effects:

Groups	Name	Variance	Std.Dev.
id:country	(Intercept)	11.724	3.424
country	(Intercept)	3.351	1.830
Residual		28.234	5.314

Number of obs: 9000, groups: id:country, 3000; country, 30

Fixed effects:

	Estimate	Std. Error	t value
(Intercept)	53.3777	0.3446	154.9

```
3.351 / (11.724 + 3.351 + 28.234)
```

```
[1] 0.07737422
```

```
(3.351 + 11.724) / (11.724 + 3.351 + 28.234)
```

```
[1] 0.3480801
```

4.2.2.3 Julia

In Julia, the ICC for a three level model is also easiest to estimate “by hand”.

```
using Tables, MixedModels, StatFiles, DataFrames, CategoricalArrays, DataFramesMeta  
dfL = DataFrame(load("simulated_multilevel_longitudinal_data.dta"))
```

```
@transform!(dfL, :country = categorical(:country))
```

```
mOL = fit(MixedModel, @formula(outcome ~  
                               (1 | country) +  
                               (1 | id)), dfL)
```

Linear mixed model fit by maximum likelihood

outcome ~ 1 + (1 | country) + (1 | id)

	logLik	-2 logLik	AIC	AICc	BIC
	-29058.2592	58116.5184	58124.5184	58124.5229	58152.9384

Variance components:

	Column	Variance	Std.Dev.
id	(Intercept)	11.72401	3.42403
country	(Intercept)	3.23190	1.79775
Residual		28.23426	5.31359

Number of obs: 9000; levels of grouping factors: 3000, 30

Fixed-effects parameters:

	Coef.	Std. Error	z	Pr(> z)
(Intercept)	53.3777	0.338785	157.56	<1e-99

$$3.23190 / (11.72401 + 3.23190 + 28.23426)$$

0.07482952718176382

$$(3.23190 + 11.72401) / (11.72401 + 3.23190 + 28.23426)$$

0.34628041519632824

4.2.3 Interpretation

Each software suggests that almost 8% of the variation in the outcome is within time points for different individuals within the same country, while almost 35% of the variation in the outcome is within time points for the same individual within the same country.

5 Cross Sectional Multilevel Models

5.1 The Equation

Recall the general model of Equation 1.1, and the syntax outlined in Section 1.3. Below in Equation 5.1, we consider a more substantive example.

$$\begin{aligned} \text{outcome}_{ij} = & \beta_0 + \beta_1 \text{warmth}_{ij} + \\ & \beta_2 \text{physical punishment}_{ij} + \\ & \beta_3 \text{identity}_{ij} + \beta_4 \text{intervention}_{ij} + \beta_5 \text{HDI}_j + \\ & u_{0j} + u_{1j} \times \text{warmth}_{ij} + e_{ij} \end{aligned} \tag{5.1}$$

5.2 Correlated and Uncorrelated Random Effects

Consider the covariance matrix of random effects (e.g. u_{0j} and u_{1j}). In Equation 5.2 the covariances of the random effects are constrained to be zero.

$$\begin{bmatrix} \text{var}(u_{0j}) & 0 \\ 0 & \text{var}(u_{1j}) \end{bmatrix} \tag{5.2}$$

As discussed in the Chapter on multilevel models with cross-sectional data, however, one can consider a multilevel model in which the random effects are correlated, as is the case in Equation 5.3.

$$\begin{bmatrix} \text{var}(u_{0j}) & \text{cov}(u_{0j}, u_{1j}) \\ \text{cov}(u_{0j}, u_{1j}) & \text{var}(u_{1j}) \end{bmatrix} \tag{5.3}$$

Procedures for estimating models with uncorrelated and correlated random effects are detailed below (Bates et al., 2015; Bates, 2024; StataCorp, 2023).

Table 5.1: Correlated and Uncorrelated Random Effects

Software	Uncorrelated Random Effects	Correlated Random Effects
Stata	default	add option: <code>, cov(uns)</code>
R	separate random effects from grouping variable with <code> </code>	separate random effects from grouping variable with <code> </code>
Julia	separate terms for each random effect e.g. <code>(1 group) + (0 + x group)</code>	separate random effects from grouping variable with <code> </code> .

All models in the examples below are run with *uncorrelated* random effects, but could just as easily be run with *correlated* random effects.

5.3 Run Models

Continuous and Categorical Variables

Statistically—as noted in the main text—it is important to be clear on whether independent variables in one’s model are continuous or categorical. *Continuous* variables can be entered straightforwardly into statistical syntax. *Categorical* variables, on the other hand usually require specific attention in statistical software. In Stata, categorical variables are indicated in a statistical model by prefixing them with an `i.`. In R, categorical variables are distinguished by making them into factors e.g. `x <- factor(x)`. In Julia, categorical variables are created by using the `@transform` syntax detailed below.

5.3.1 Stata

5.3.1.1 Get The Data

```
use simulated_multilevel_data.dta
```

5.3.1.2 Run The Model

```
mixed outcome warmth physical_punishment i.identity i.intervention HDI || ///
country: warmth
```

Performing EM optimization ...

Performing gradient-based optimization:

Iteration 0: Log likelihood = -9626.6279

Iteration 1: Log likelihood = -9626.607

Iteration 2: Log likelihood = -9626.607

Computing standard errors ...

Mixed-effects ML regression

Group variable: country

Number of obs = 3,000

Number of groups = 30

Obs per group:

min = 100

avg = 100.0

max = 100

Wald chi2(5) = 334.14

Prob > chi2 = 0.0000

Log likelihood = -9626.607

	outcome	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
	warmth	.8345368	.0637213	13.10	0.000	.7096453	.9594282
	physical_punishment	-.9916657	.0797906	-12.43	0.000	-1.148052	-.8352791
	1.identity	-.3004767	.2170295	-1.38	0.166	-.7258466	.1248933
	1.intervention	.6396427	.2174519	2.94	0.003	.2134448	1.065841
	HDI	-.003228	.0199257	-0.16	0.871	-.0422817	.0358256
	_cons	51.99991	1.371257	37.92	0.000	49.3123	54.68753

Random-effects parameters		Estimate	Std. err.	[95% conf. interval]	
country: Independent					
	var(warmth)	.0227504	.0257784	.0024689	.2096436
	var(_cons)	2.963975	.9737647	1.556777	5.643163

var(Residual)	34.97499	.9097109	33.23668	36.80422
---------------	----------	----------	----------	----------

LR test vs. linear model: $\chi^2(2) = 205.74$ Prob > $\chi^2 = 0.0000$

Note: LR test is conservative and provided only for reference.

5.3.2 R

5.3.2.1 Get The Data

```
library(haven)

df <- read_dta("simulated_multilevel_data.dta")
```

5.3.2.2 Change Some Variables To Categorical

```
df$identity <- factor(df$identity)

df$intervention <- factor(df$intervention)
```

5.3.2.3 Run The Model

Caution

`lme4` does not directly provide p values in results, because of some disagreement over exactly how these p values should be calculated. Therefore, in this Appendix, I also call `library(lmerTest)` to provide p values for `lme4` results.

Tip

R prefers to use scientific notation when possible. I find that the use of scientific notation can be confusing in reading results. I turn off scientific notation by setting a penalty for its use: `options(scipen = 999)`.

```
library(lme4)

library(lmerTest)
```

```
options(scipen = 999)

fit1 <- lmer(outcome ~ warmth + physical_punishment +
             identity + intervention + HDI +
             (1 + warmth || country),
             data = df)

summary(fit1)
```

Linear mixed model fit by REML. t-tests use Satterthwaite's method [lmerModLmerTest]
 Formula: outcome ~ warmth + physical_punishment + identity + intervention + HDI + (1 + warmth || country)
 Data: df

REML criterion at convergence: 19268.8

Scaled residuals:

Min	1Q	Median	3Q	Max
-3.9774	-0.6563	0.0186	0.6645	3.6730

Random effects:

Groups	Name	Variance	Std.Dev.
country	(Intercept)	3.19120	1.786
country.1	warmth	0.02464	0.157
Residual		35.01779	5.918

Number of obs: 3000, groups: country, 30

Fixed effects:

	Estimate	Std. Error	df	t value
(Intercept)	52.011324	1.414976	30.293141	36.758
warmth	0.834562	0.064250	41.896457	12.989
physical_punishment	-0.991893	0.079845	2968.012381	-12.423
identity1	-0.300354	0.217179	2970.108153	-1.383
intervention1	0.639060	0.217603	2971.186718	2.937
HDI	-0.003394	0.020598	27.592814	-0.165

	Pr(> t)
(Intercept)	< 0.0000000000000002 ***
warmth	0.000000000000000277 ***
physical_punishment	< 0.0000000000000002 ***
identity1	0.16678

```
intervention1          0.00334 **
HDI                    0.87030
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Correlation of Fixed Effects:

```
(Intr) warmth physc_ idntt1 intrv1
warmth      -0.124
physcl_pnsh -0.149 -0.003
identity1    -0.072 -0.012 -0.003
interventn1 -0.082  0.034  0.022 -0.018
HDI          -0.943 -0.006  0.009 -0.001  0.000
```

5.3.3 Julia

5.3.3.1 Get The Data

```
using Tables, MixedModels, StatFiles, DataFrames, CategoricalArrays, DataFramesMeta

df = DataFrame(load("simulated_multilevel_data.dta"))
```

5.3.3.2 Change Some Variables To Categorical

```
@transform!(df, :country = categorical(:country))

@transform!(df, :identity = categorical(:identity))

@transform!(df, :intervention = categorical(:intervention))
```

5.3.3.3 Run The Model

```
m1 = fit(MixedModel, @formula(outcome ~ warmth + physical_punishment +
                             identity + intervention + HDI +
                             (1 | country) +
                             (0 + warmth | country)), df)
```

Linear mixed model fit by maximum likelihood

outcome ~ 1 + warmth + physical_punishment + identity + intervention + HDI + (1 | country)

	logLik	-2 logLik	AIC	AICc	BIC
	-9626.6070	19253.2140	19271.2140	19271.2742	19325.2713

Variance components:

	Column	Variance	Std.Dev.	Corr.
country	(Intercept)	2.963849	1.721583	
	warmth	0.022756	0.150852	.
Residual		34.974984	5.913965	

Number of obs: 3000; levels of grouping factors: 30

Fixed-effects parameters:

	Coef.	Std. Error	z	Pr(> z)
(Intercept)	51.9999	1.37124	37.92	<1e-99
warmth	0.834537	0.0637228	13.10	<1e-38
physical_punishment	-0.991665	0.0797906	-12.43	<1e-34
identity: 1.0	-0.300475	0.217029	-1.38	0.1662
intervention: 1.0	0.639641	0.217452	2.94	0.0033
HDI	-0.0032286	0.0199255	-0.16	0.8713

5.4 Interpretation

Models suggest that parental warmth is associated with increases in the beneficial outcome, while physical punishment is associated with decreases in the beneficial outcome. The intervention is associated with increases in the outcome. There is insufficient evidence that either identity group or the Human Development Index are associated with the outcome.

6 Longitudinal Multilevel Models

6.1 The Data

The data employed in these examples are a longitudinal extension of the data described in Section 1.2.

6.2 The Equation

$$\text{outcome}_{itj} = \beta_0 + \beta_1 \text{parental warmth}_{itj} + \beta_2 \text{physical punishment}_{itj} + \beta_3 \text{time}_{itj} + \quad (6.1)$$

$$\beta_4 \text{identity}_{itj} + \beta_5 \text{intervention}_{itj} + \beta_6 \text{HDI}_j +$$

$$u_{0j} + u_{1j} \times \text{parental warmth}_{itj} +$$

$$v_{0ij} + v_{1ij} \times \text{time}_{itj} + e_{itj}$$

6.3 Growth Trajectories

Remember, following the discussion in the main text, that in longitudinal multilevel models, the variable for *time* assumes an important role as we are often thinking of a *growth trajectory* over time.

As discussed in the main text, think about a model where *identity* is a (1/0) variable for membership in one of two groups:

$$\text{outcome} = \beta_0 + \beta_t \text{time} + \beta_{\text{identity}} \text{identity} + \beta_{\text{interaction}} \text{identity} \times \text{time} + u_{0i} + e_{it}$$

Then, each identity group has its own intercept and time trajectory:

Table 6.1: Slope and Intercept for Each Group

Group	Intercept	Slope (Time Trajectory)
0	β_0	β_t
1	$\beta_0 + \beta_{\text{identity}}$	$\beta_t + \beta_{\text{interaction}}$

💡 Main Effects and Interactions

Thus, again following the main text, in longitudinal multilevel models, *main effects* modify the *intercept* of the time trajectory, while *interactions with time*, modify the *slope* of the time trajectory. Below, we run models with *main effects only*, then models with *main effects, and interactions with time*.

6.4 Run Models

⚠ Warning

Remember that we are estimating a model in which time points are nested inside families, who are in turn nested inside countries. For each software package, it is accordingly important to specify the way in which different levels of the data are nested. Pay careful attention to the syntax examples below with regard to `id` and `country`

6.4.1 Stata

6.4.1.1 Get The Data

```
use simulated_multilevel_longitudinal_data.dta
```

6.4.1.2 Run The Models

6.4.1.2.1 Main Effects Only

```
mixed outcome t warmth physical_punishment i.identity i.intervention HDI || ///
country: warmth || id: t
```

Performing EM optimization ...

Performing gradient-based optimization:

Iteration 0: Log likelihood = -28523.49

Iteration 1: Log likelihood = -28499.987

Iteration 2: Log likelihood = -28499.739

Iteration 3: Log likelihood = -28499.604

Iteration 4: Log likelihood = -28499.603

Computing standard errors ...

Mixed-effects ML regression

Number of obs = 9,000

Grouping information

Group variable		No. of groups	Observations per group		
			Minimum	Average	Maximum
country		30	300	300.0	300
id		3,000	3	3.0	3

Log likelihood = -28499.603

Wald chi2(6) = 1096.15

Prob > chi2 = 0.0000

outcome	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
t	.943864	.0658716	14.33	0.000	.814758	1.07297
warmth	.9134959	.0423732	21.56	0.000	.830446	.9965459
physical_punishment	-1.007897	.0497622	-20.25	0.000	-1.105429	-.9103647
1.identity	-.1276926	.1515835	-0.84	0.400	-.4247908	.1694057
1.intervention	.8589966	.1519095	5.65	0.000	.5612596	1.156734
HDI	-.0005657	.0196437	-0.03	0.977	-.0390666	.0379352
_cons	50.46724	1.338318	37.71	0.000	47.84418	53.09029

Random-effects parameters	Estimate	Std. err.	[95% conf. interval]	
country: Independent				
var(warmth)	.0107586	.0127845	.0010478	.1104703
var(_cons)	3.167085	.9146761	1.798154	5.578181

```

-----+-----
id: Independent |
      var(t) | 3.58e-09 7.06e-07 3.5e-177 3.7e+159
      var(_cons) | 8.387275 .4724188 7.510631 9.366242
-----+-----
      var(Residual) | 26.02733 .4753701 25.11211 26.97592
-----+-----
LR test vs. linear model: chi2(4) = 1247.03          Prob > chi2 = 0.0000

```

Note: LR test is conservative and provided only for reference.

6.4.1.2.2 Interactions With Time

```
mixed outcome c.t##(c.warmth c.physical_punishment i.identity i.intervention c.HDI) || count:
```

Performing EM optimization ...

Performing gradient-based optimization:

```

Iteration 0: Log likelihood = -28522.21
Iteration 1: Log likelihood = -28498.677
Iteration 2: Log likelihood = -28498.468
Iteration 3: Log likelihood = -28498.31
Iteration 4: Log likelihood = -28498.309

```

Computing standard errors ...

Mixed-effects ML regression Number of obs = 9,000

Grouping information

```

-----+-----
Group variable | No. of      Observations per group
               | groups      Minimum   Average   Maximum
-----+-----
      country |      30      300      300.0      300
        id |    3,000       3       3.0       3
-----+-----

```

```

Log likelihood = -28498.309
Wald chi2(11) = 1100.25
Prob > chi2    = 0.0000

```

outcome	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
t	.7582075	.326177	2.32	0.020	.1189123	1.397503
warmth	.8170757	.082662	9.88	0.000	.6550611	.9790903
physical_punishment	-1.009031	.1112932	-9.07	0.000	-1.227162	-.7909007
1.identity	-.2387167	.3039964	-0.79	0.432	-.8345387	.3571053
1.intervention	.6607606	.3044503	2.17	0.030	.064049	1.257472
HDI	.0013614	.0210842	0.06	0.949	-.0399628	.0426856
c.t#c.warmth	.0483637	.0356074	1.36	0.174	-.0214255	.1181529
c.t#						
c.physical_punishment	.0005421	.0494354	0.01	0.991	-.0963496	.0974338
identity#c.t						
1	.0554389	.1317444	0.42	0.674	-.2027754	.3136532
intervention#c.t						
1	.0992811	.131925	0.75	0.452	-.1592872	.3578493
c.t#c.HDI	-.0009551	.0038216	-0.25	0.803	-.0084453	.0065352
_cons	50.83632	1.483548	34.27	0.000	47.92862	53.74402

Random-effects parameters	Estimate	Std. err.	[95% conf. interval]	
country: Independent				
var(warmth)	.0106014	.0127458	.0010046	.1118779
var(_cons)	3.170088	.9153355	1.80009	5.582753
id: Independent				
var(t)	9.47e-10	2.07e-07	1.5e-195	6.0e+176
var(_cons)	8.39189	.4724106	7.515234	9.370809
var(Residual)	26.01583	.4751602	25.10101	26.964

LR test vs. linear model: $\chi^2(4) = 1247.84$ Prob > $\chi^2 = 0.0000$

Note: LR test is conservative and provided only for reference.

6.4.2 R

6.4.2.1 Get The Data

```
library(haven)

dfL <- read_dta("simulated_multilevel_longitudinal_data.dta")
```

6.4.2.2 Change Some Variables To Categorical

```
dfL$identity <- factor(dfL$identity)

dfL$intervention <- factor(dfL$intervention)
```

6.4.2.3 Run The Models

Caution

`lme4` does not directly provide p values in results, because of some disagreement over exactly how these p values should be calculated. Therefore, in this Appendix, I also call library `lmerTest` to provide p values for `lme4` results.

Tip

R prefers to use scientific notation when possible. I find that the use of scientific notation can be confusing in reading results. I turn off scientific notation by setting a penalty for its use: `options(scipen = 999)`.

6.4.2.3.1 Main Effects Only

```
library(lme4)

library(lmerTest)

options(scipen = 999)

fit2A <- lmer(outcome ~ t + warmth + physical_punishment +
```

```

        identity + intervention + HDI +
        (1 | country/id),
        data = dfL)

summary(fit2A)

```

Linear mixed model fit by REML. t-tests use Satterthwaite's method [lmerModLmerTest]

Formula:

```
outcome ~ t + warmth + physical_punishment + identity + intervention +
        HDI + (1 | country/id)
```

Data: dfL

REML criterion at convergence: 57022.7

Scaled residuals:

Min	1Q	Median	3Q	Max
-3.6850	-0.6094	-0.0035	0.6133	3.6792

Random effects:

Groups	Name	Variance	Std.Dev.
id:country	(Intercept)	8.438	2.905
country	(Intercept)	3.675	1.917
Residual		26.036	5.103

Number of obs: 9000, groups: id:country, 3000; country, 30

Fixed effects:

	Estimate	Std. Error	df	t value
(Intercept)	50.3842343	1.4139114	29.8246912	35.635
t	0.9433806	0.0658755	5998.3764548	14.321
warmth	0.9140307	0.0379336	4745.3497493	24.096
physical_punishment	-1.0087537	0.0497972	6483.6771808	-20.257
identity1	-0.1319548	0.1517350	2968.7828107	-0.870
intervention1	0.8591494	0.1520510	2971.8111995	5.650
HDI	0.0007909	0.0207656	28.0001855	0.038

Pr(>|t|)

(Intercept)	< 0.0000000000000002	***
t	< 0.0000000000000002	***
warmth	< 0.0000000000000002	***
physical_punishment	< 0.0000000000000002	***
identity1	0.385	
intervention1	0.0000000175	***

```

HDI                                0.970
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Correlation of Fixed Effects:
              (Intr) t      warmth physc_ idntt1 intrv1
t              -0.092
warmth         -0.091 -0.002
physcl_pnsh    -0.092 -0.007 -0.012
identity1      -0.051  0.000 -0.013 -0.003
interventn1    -0.058  0.000  0.039  0.019 -0.018
HDI            -0.951  0.000 -0.004  0.005  0.000  0.002

```

6.4.2.3.2 Interactions With Time

```

fit2B <- lmer(outcome ~ t *(warmth + physical_punishment +
                        identity + intervention + HDI) +
              (1 | country/id),
              data = dfL)

summary(fit2B)

```

Linear mixed model fit by REML. t-tests use Satterthwaite's method [
lmerModLmerTest]

Formula:

```
outcome ~ t * (warmth + physical_punishment + identity + intervention +
              HDI) + (1 | country/id)
```

Data: dfL

REML criterion at convergence: 57042.8

Scaled residuals:

	Min	1Q	Median	3Q	Max
	-3.7118	-0.6092	-0.0024	0.6150	3.6779

Random effects:

Groups	Name	Variance	Std.Dev.
id:country	(Intercept)	8.436	2.905
country	(Intercept)	3.675	1.917
Residual		26.046	5.104

Number of obs: 9000, groups: id:country, 3000; country, 30

Fixed effects:

	Estimate	Std. Error	df	t value
(Intercept)	50.7590272	1.5518360	43.2608620	32.709
t	0.7552909	0.3263028	6176.7440549	2.315
warmth	0.8170912	0.0805355	8274.9995422	10.146
physical_punishment	-1.0097729	0.1113557	8084.6084915	-9.068
identity1	-0.2446453	0.3041604	8695.8966197	-0.804
intervention1	0.6604671	0.3046286	8697.0843430	2.168
HDI	0.0026692	0.0221295	36.1037733	0.121
t:warmth	0.0486211	0.0356217	6404.8723333	1.365
t:physical_punishment	0.0004964	0.0494590	6753.0158441	0.010
t:identity1	0.0563140	0.1318043	5993.4518022	0.427
t:intervention1	0.0995037	0.1319917	5994.1433001	0.754
t:HDI	-0.0009379	0.0038233	5993.9090880	-0.245

Pr(>|t|)

(Intercept)	<0.0000000000000002 ***
t	0.0207 *
warmth	<0.0000000000000002 ***
physical_punishment	<0.0000000000000002 ***
identity1	0.4212
intervention1	0.0302 *
HDI	0.9047
t:warmth	0.1723
t:physical_punishment	0.9920
t:identity1	0.6692
t:intervention1	0.4510
t:HDI	0.8062

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Correlation of Fixed Effects:

	(Intr) t	warmth	physc_	idntt1	intrv1	HDI	t:wrmt	t:phy_
t	-0.421							
warmth	-0.178	0.331						
physcl_pnsh	-0.190	0.360	-0.005					
identity1	-0.093	0.166	-0.013	-0.002				
interventn1	-0.107	0.192	0.039	0.019	-0.017			
HDI	-0.925	0.264	-0.007	0.012	-0.001	0.003		
t:warmth	0.158	-0.377	-0.882	0.001	0.011	-0.035	0.006	
t:physcl_pn	0.170	-0.402	0.004	-0.894	-0.001	-0.017	-0.010	-0.003
t:identity1	0.081	-0.192	0.011	0.000	-0.867	0.014	0.001	-0.013
t:intrvntn1	0.093	-0.222	-0.035	-0.017	0.014	-0.867	-0.003	0.041


```

t:HDI          0.322 -0.765  0.015 -0.027  0.002 -0.007 -0.346 -0.016  0.029
               t:dnt1 t:ntr1
t
warmth
physcl_pnsh
identity1
intervntn1
HDI
t:warmth
t:physcl_pn
t:identity1
t:intrvntn1 -0.016
t:HDI       -0.002  0.008

```

6.4.3 Julia

6.4.3.1 Get The Data

```

using Tables, MixedModels, StatFiles, DataFrames, CategoricalArrays, DataFramesMeta

dfL = DataFrame(load("simulated_multilevel_longitudinal_data.dta"))

```

6.4.3.2 Change Some Variables To Categorical

```

@transform!(dfL, :country = categorical(:country))

@transform!(dfL, :identity = categorical(:identity))

@transform!(dfL, :intervention = categorical(:intervention))

```

6.4.3.3 Run The Models

6.4.3.3.1 Main Effects Only

```

m2A = fit(MixedModel, @formula(outcome ~ t + warmth +
                                physical_punishment +
                                identity + intervention +
                                HDI +

```

```
(1 | country) +
(0 + warmth | country) +
(1 | id)), dfL)
```

Linear mixed model fit by maximum likelihood

```
outcome ~ 1 + t + warmth + physical_punishment + identity + intervention + HDI + (1 | country)
logLik    -2 logLik      AIC      AICc      BIC
-28499.6031 56999.2063 57021.2063 57021.2356 57099.3610
```

Variance components:

	Column	Variance	Std.Dev.	Corr.
id	(Intercept)	8.387214	2.896069	
country	(Intercept)	3.167143	1.779647	
	warmth	0.010762	0.103739	.
Residual		26.027363	5.101702	

Number of obs: 9000; levels of grouping factors: 3000, 30

Fixed-effects parameters:

	Coef.	Std. Error	z	Pr(> z)
(Intercept)	50.4673	1.33833	37.71	<1e-99
t	0.943864	0.0658717	14.33	<1e-45
warmth	0.913496	0.0423744	21.56	<1e-99
physical_punishment	-1.0079	0.0497622	-20.25	<1e-90
identity: 1.0	-0.127692	0.151583	-0.84	0.3996
intervention: 1.0	0.858997	0.151909	5.65	<1e-07
HDI	-0.000566026	0.0196439	-0.03	0.9770

6.4.3.3.2 Interactions With Time

```
m2B = fit(MixedModel, @formula(outcome ~ t * (warmth +
physical_punishment +
identity + intervention +
HDI) +
(1 | country) +
(0 + warmth | country) +
(1 | id)), dfL)
```

Linear mixed model fit by maximum likelihood

```
outcome ~ 1 + t + warmth + physical_punishment + identity + intervention + HDI + t & warmth
      logLik   -2 logLik       AIC       AICc       BIC
-28498.3091  56996.6182  57028.6182  57028.6788  57142.2979
```

Variance components:

	Column	Variance	Std.Dev.	Corr.
id	(Intercept)	8.391746	2.896851	
country	(Intercept)	3.170032	1.780458	
	warmth	0.010609	0.102999	.
Residual		26.015906	5.100579	

Number of obs: 9000; levels of grouping factors: 3000, 30

Fixed-effects parameters:

	Coef.	Std. Error	z	Pr(> z)
(Intercept)	50.8364	1.48355	34.27	<1e-99
t	0.758209	0.326177	2.32	0.0201
warmth	0.817076	0.0826636	9.88	<1e-22
physical_punishment	-1.00903	0.111293	-9.07	<1e-18
identity: 1.0	-0.238714	0.303996	-0.79	0.4323
intervention: 1.0	0.660761	0.30445	2.17	0.0300
HDI	0.00136065	0.0210842	0.06	0.9485
t & warmth	0.0483635	0.0356074	1.36	0.1744
t & physical_punishment	0.000542203	0.0494355	0.01	0.9912
t & identity: 1.0	0.0554385	0.131745	0.42	0.6739
t & intervention: 1.0	0.0992809	0.131925	0.75	0.4517
t & HDI	-0.000955067	0.00382162	-0.25	0.8027

6.5 Interpretation

The *main effects only model* suggests that time is associated with increases in the outcome. In the main effects model, main effects other than time, indicate whether a particular variable is associated with higher or lower intercepts of the time trajectory, at the beginning of the study time. Warmth is again associated with increases in the outcome, while physical punishment is associated with decreases in the outcome. Identity is again not associated with the outcome, while the intervention is associated with higher levels of the outcome. The Human Development Index is again not associated with the outcome.

The second model adds interactions with time to the first model. Results are largely similar to the prior model. However, here we not only examine whether main effects other than time

are associated with higher or lower time trajectories, but also whether particular variables are associated with differences in the slope of the time trajectory. In this case, we find insufficient evidence that any independent variable is associated with changes in the slope of the time trajectory.

Which Interactions To Test?

In this example—for the sake of illustration—I test the interaction of *every* independent variable with time. In many cases, it may make sense to test only one or two interactions of time with particular variables of key interest. Also, after finding, as I did in this model, that none of the interactions of other independent variables with time are significant, I might report the model with interactions, or might report only the results of the model with only main effects.

It may be illustrative to imagine how we would interpret the results had a particular interaction term been statistically significant. Let us consider one of the interaction terms with the largest coefficient, **intervention#time**. The interaction of the intervention with time is positive. Had this coefficient been statistically significant, it would have indicated that the intervention was associated with more rapid increases in the outcome over time *in addition to* the fact that the intervention is associated with higher initial levels of the outcome.

References

- Allaire, J. J., Teague, C., Scheidegger, C., Xie, Y., & Dervieux, C. (2024). *Quarto* (Version 1.4). <https://doi.org/10.5281/zenodo.5960048>
- Bates, D. (2024). *MixedModels.jl Documentation*. <https://juliastats.org/MixedModels.jl/stable/>
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. B. (2017). Julia: A fresh approach to numerical computing. *SIAM Review*, 59(1), 65–98. <https://doi.org/10.1137/141000671>
- Hemken, D. (2023). *Statamarkdown: 'Stata' markdown*. <https://CRAN.R-project.org/package=Statamarkdown>
- Li, C. (2019). JuliaCall: An R package for seamless integration between R and Julia. *The Journal of Open Source Software*, 4(35), 1284. <https://doi.org/10.21105/joss.01284>
- R Core Team. (2023). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Schanen, J. (2021). *Math person (Strogatz Prize entry)*. National Museum of Mathematics.
- StataCorp. (2023). *Stata 18 mixed effects reference manual*. Stata Press.
- Thoreau, H. D. (1975). The commercial spirit of modern times [1837]. In J. J. Moldenhauer, E. Moser, & A. C. Kern (Eds.), *Early essays and miscellanies*. Princeton University Press.