

Multilevel Workshop

Andrew Grogan-Kaylor

2025-11-10

Table of contents

1	Introduction	6
1.1	Background	6
1.2	Simulated Multilevel Data	6
2	The Importance of Accounting for Clustered Data	8
3	Grouped and Individual Data	9
4	Analyses	10
4.1	OLS	10
4.2	MLM	10
4.3	Compare OLS and MLM	11
5	Two Level Cross Sectional; And Three Level Longitudinal Models	12
5.1	Cross Sectional Model	12
5.1.1	Get Data	12
5.1.2	The Equation	12
5.1.3	Descriptive Statistics	12
5.1.4	Spaghetti Plot	13
5.1.5	Unconditional Model	13
5.1.6	Conditional Model	15
5.2	Longitudinal Model	16
5.2.1	Get Data	16
5.2.2	The Equation	16
5.2.3	Descriptive Statistics	16
5.2.4	Alternate Plot	17
5.2.5	Unconditional Model	17
5.2.6	Conditional Model	18
5.3	Nice Table of Results	19
5.4	QUESTIONS???	20
6	Cross-Classified Models	21
6.1	Introduction	21
6.2	Get Data	21
6.3	Cross Classified Model	21
6.3.1	Standard (Less Computationally Efficient) Syntax	21

6.3.2	Cross Classified With Computationally Efficient Syntax	22
6.4	Three Level Model	23
6.5	Nice Table of Results of Three Level and Cross Classified Model	25
6.6	Verification of Syntax Equivalence for Cross Classified Model	25
6.7	QUESTIONS???	26
References		27
License and Citation		28
	License	28
	Citation	28
	BibTeX Citation	28

List of Figures

5.1	Spaghetti Plot of Outcome by Warmth by Country	13
5.2	Alternate Plot of Outcome by Time by Individual; First 10 Observations	17

List of Tables

1.1	Variables in Simulated Multilevel Data	7
1.2	Sample Data From Simulated Multilevel Data	7

1 Introduction

“Listening to the world. Well, I did that, and I still do it. I still do it.” (Mary Oliver in Oliver and Tippett 2015)

This site contains materials for a workshop on multilevel modeling.

1.1 Background

Multilevel models are useful when you have data that are nested or clustered inside social units such as schools, neighborhoods, states, or countries.

Multilevel models are also useful when you have longitudinal data where repeated measures are collected for study participants.

The Importance of Multilevel Models

Multilevel models may improve one’s statistical inferences in two important substantive ways.

- Multilevel models adjust standard errors for clustering, and thus calculate appropriate p values. Failure to use a model that accounts for the clustering in the data may lead to improperly calculated standard errors and p values, possibly leading to false attributions of statistical significance (false positives). (See Chapter 2).
- Multilevel models adjust regression coefficients (β ’s) for the presence of clustering. Failure to use a model that accounts for the clustering in the data may lead to improperly calculated regression coefficients (β ’s) which may have the wrong magnitude, the wrong statistical significance, and even the wrong sign.

1.2 Simulated Multilevel Data

The data used in these workshop materials are *simulated* data on parents, children and families. The data are simulated to come from 30 hypothetical countries around the world. These are the same data used and discussed in my book *Multilevel Thinking: Discovering Variation, Universals, and Particulars in Cross-Cultural Research*.

There are two versions of the data: a *cross-sectional* data set from a single point in time; a *longitudinal* version of the data spanning several time points.

i The Data Can Be Downloaded Here:

- [Cross Sectional Data](#)
- [Longitudinal Data](#)

Table 1.1: Variables in Simulated Multilevel Data

pos	variable	label
1	country	country id
2	HDI	Human Development Index
3	family	family id
4	id	unique country family id
5	identity	hypothetical identity group variable
6	intervention	recieved intervention
7	physical_punishment	physical punishment in past week
8	warmth	parental warmth in past week
9	outcome	beneficial outcome

Table 1.2: Sample Data From Simulated Multilevel Data

country	HDI	family	id	identity	intervention	physical_punishment	warmth	outcome
5	83	52	5.52	0	1	0	3	55.99
4	37	89	4.89	0	1	4	4	48.05
10	59	90	10.90	0	0	4	5	60.69
24	82	90	24.90	1	0	3	5	51.63
19	72	86	19.86	1	1	4	2	38.08

2 The Importance of Accounting for Clustered Data

3 Grouped and Individual Data


Bland and Altman (1994) suggested the following procedure for simulating some data:

“The data were generated from random numbers, and there is no relation between X and Y at all. Firstly, values of X and Y were generated for each ‘subject,’ then a further random number was added to make the individual observation.” (Bland and Altman 1994)

So... we follow their procedure.

Simulating The Data

The graph below illustrates the process of simulating the data.



4 Analyses

4.1 OLS

An OLS analysis indicates that there is a statistically significant association of x and y .

```

                                OLS
-----
x_individual          1.046 **
Intercept             4.488
Number of observations    25
-----
** p<.01, * p<.05
```

4.2 MLM

In contrast, an MLM analysis (correctly) finds that there is no statistically significant association of x and y .

```

                                MLM
-----
x_individual          0.039
Intercept             97.005 **
var(_cons)            74.523
var(e)                0.594
Number of observations    25
-----
** p<.01, * p<.05
```

4.3 Compare OLS and MLM

	OLS	MLM
x_individual	1.046 **	0.039
Intercept	4.488	97.005 **
var(_cons)		74.523
var(e)		0.594
Number of observations	25	25

** p<.01, * p<.05

5 Two Level Cross Sectional; And Three Level Longitudinal Models

5.1 Cross Sectional Model

5.1.1 Get Data

```
use "simulated_multilevel_data.dta", clear
```

5.1.2 The Equation

$$\text{outcome}_{ij} = \beta_0 + \beta_1 \text{parental warmth} + \beta_2 \text{physical punishment} + \beta_3 \text{time} +$$

$$\beta_4 \text{identity}_2 + \beta_5 \text{intervention} + \beta_6 \text{HDI} +$$

$$u_{0j} + u_{1j} \times \text{parental warmth} + e_{ij}$$

5.1.3 Descriptive Statistics

```
summarize // descriptive statistics
```

Variable	Obs	Mean	Std. dev.	Min	Max
country	3,000	15.5	8.656884	1	30
HDI	3,000	64.76667	17.24562	33	87
family	3,000	50.5	28.87088	1	100
id	0				
identity	3,000	.4976667	.5000779	0	1

intervention		3,000	.4843333	.4998378	0	1
physical_p~t		3,000	2.478667	1.360942	0	5
warmth		3,000	3.521667	1.888399	0	7
outcome		3,000	52.43327	6.530996	29.60798	74.83553

5.1.4 Spaghetti Plot

```
spagplot outcome warmth, id(country) scheme(stcolor)
graph export spagplot1.png, width(1000) replace
```




Figure 5.1: Spaghetti Plot of Outcome by Warmth by Country

5.1.5 Unconditional Model

5.1.5.1 Model

```
mixed outcome || country: // unconditional model
```

Performing EM optimization ...

Performing gradient-based optimization:

Iteration 0: Log likelihood = -9802.8371

Iteration 1: Log likelihood = -9802.8371

Computing standard errors ...

Mixed-effects ML regression
Group variable: country

Number of obs = 3,000
Number of groups = 30
Obs per group:
min = 100
avg = 100.0
max = 100
Wald chi2(0) = .
Prob > chi2 = .

Log likelihood = -9802.8371

outcome	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
-----+-----						
_cons	52.43327	.3451217	151.93	0.000	51.75685	53.1097
-----+-----						

Random-effects parameters		Estimate	Std. err.	[95% conf. interval]	
-----+-----					
country: Identity					
var(_cons)		3.178658	.9226736	1.799552	5.614658
-----+-----					
var(Residual)		39.46106	1.024013	37.50421	41.52
-----+-----					

LR test vs. linear model: chibar2(01) = 166.31 Prob >= chibar2 = 0.0000

5.1.5.2 ICC

```
estat icc
```

Intraclass correlation

Level	ICC	Std. err.	[95% conf. interval]	
-----+-----				
country	.0745469	.0201254	.0434963	.1248696
-----+-----				

5.1.6 Conditional Model

```
mixed outcome warmth physical_punishment identity i.intervention HDI || country: warmth // ml
est store crosssectional // store estimates
```

Performing EM optimization ...

Performing gradient-based optimization:

Iteration 0: Log likelihood = -9626.6279

Iteration 1: Log likelihood = -9626.607

Iteration 2: Log likelihood = -9626.607

Computing standard errors ...

Mixed-effects ML regression

Group variable: country

Number of obs = 3,000

Number of groups = 30

Obs per group:

min = 100

avg = 100.0

max = 100

Wald chi2(5) = 334.14

Prob > chi2 = 0.0000

Log likelihood = -9626.607

outcome	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
warmth	.8345368	.0637213	13.10	0.000	.7096453	.9594282
physical_punishm~t	-.9916657	.0797906	-12.43	0.000	-1.148052	-.8352791
identity	-.3004767	.2170295	-1.38	0.166	-.7258466	.1248933
1.intervention	.6396427	.2174519	2.94	0.003	.2134448	1.065841
HDI	-.003228	.0199257	-0.16	0.871	-.0422817	.0358256
_cons	51.99991	1.371257	37.92	0.000	49.3123	54.68753

Random-effects parameters	Estimate	Std. err.	[95% conf. interval]	
country: Independent				
var(warmth)	.0227504	.0257784	.0024689	.2096436
var(_cons)	2.963975	.9737647	1.556777	5.643163

```
-----+-----
var(Residual) |    34.97499    .9097109    33.23668    36.80422
-----+-----
LR test vs. linear model: chi2(2) = 205.74          Prob > chi2 = 0.0000
```

Note: LR test is conservative and provided only for reference.

5.2 Longitudinal Model

5.2.1 Get Data

```
use "simulated_multilevel_longitudinal_data.dta", clear
```

5.2.2 The Equation

$$\text{outcome}_{ij} = \beta_0 + \beta_1 \text{parental warmth} + \beta_2 \text{physical punishment} + \beta_3 \text{time} +$$

$$\beta_4 \text{identity}_2 + \beta_5 \text{intervention} + \beta_5 \text{HDI} +$$

$$u_{0j} + u_{1j} \times \text{parental warmth} +$$

$$v_{0i} + v_{1i} \times t + e_{ij}$$

5.2.3 Descriptive Statistics

```
summarize // descriptive statistics
```

Variable	Obs	Mean	Std. dev.	Min	Max
country	9,000	15.5	8.655922	1	30
HDI	9,000	64.76667	17.2437	33	87
family	9,000	50.5	28.86767	1	100
id	0				

identity	9,000	.4976667	.5000223	0	1
intervention	9,000	.4843333	.4997823	0	1
t	9,000	2	.8165419	1	3
physical_p~t	9,000	2.485333	1.373639	0	5
warmth	9,000	3.514222	1.8839	0	7
outcome	9,000	53.37768	6.572285	29.60798	79.02199

5.2.4 Alternate Plot

```

encode id, generate(idNUMERIC) // numeric version of id

* spagplot outcome t if idNUMERIC <= 10, id(idNUMERIC) scheme(stcolor)

twoway (lfit outcome t) (scatter outcome t) if idNUMERIC <= 10, by(idNUMERIC) scheme(stcolor)

graph export spagplot2.png, width(1000) replace

```




Figure 5.2: Alternate Plot of Outcome by Time by Individual; First 10 Observations

5.2.5 Unconditional Model

5.2.5.1 Model

```

mixed outcome || country: || id: // unconditional model

```

5.2.5.2 ICC

```
estat icc
```

Intraclass correlation

Level	ICC	Std. err.	[95% conf. interval]	
country	.0748336	.0190847	.0450028	.1219141
id country	.3462837	.0171461	.3134867	.3806097

5.2.6 Conditional Model

```
mixed outcome t warmth physical_punishment i.identity i.intervention HDI || country: warmth  
est store longitudinal // store estimates
```

Performing EM optimization ...

Performing gradient-based optimization:

Iteration 0: Log likelihood = -28523.49

Iteration 1: Log likelihood = -28499.953

Iteration 2: Log likelihood = -28499.735

Iteration 3: Log likelihood = -28499.604

Iteration 4: Log likelihood = -28499.603

Computing standard errors ...

Mixed-effects ML regression

Number of obs = 9,000

Grouping information

Group variable	No. of groups	Observations per group		
		Minimum	Average	Maximum
country	30	300	300.0	300

```

            id |      3,000      3      3.0      3
-----
Log likelihood = -28499.603
Wald chi2(6) = 1096.15
Prob > chi2 = 0.0000

-----
outcome | Coefficient Std. err. z P>|z| [95% conf. interval]
-----+-----
t | .943864 .0658716 14.33 0.000 .814758 1.07297
warmth | .9134959 .0423732 21.56 0.000 .8304461 .9965458
physical_punishm~t | -1.007897 .0497622 -20.25 0.000 -1.105429 -.9103647
1.identity | -.1276926 .1515835 -0.84 0.400 -.4247908 .1694056
1.intervention | .8589966 .1519094 5.65 0.000 .5612596 1.156734
HDI | -.0005657 .0196437 -0.03 0.977 -.0390666 .0379352
_cons | 50.46724 1.338318 37.71 0.000 47.84418 53.09029
-----

Random-effects parameters | Estimate Std. err. [95% conf. interval]
-----+-----
country: Independent |
var(warmth) | .0107585 .0127845 .0010477 .1104712
var(_cons) | 3.167087 .9146767 1.798155 5.578185
-----
id: Independent |
var(t) | 5.69e-10 1.29e-07 1.1e-202 3.0e+183
var(_cons) | 8.387268 .4724189 7.510624 9.366236
-----
var(Residual) | 26.02734 .4753703 25.11211 26.97592
-----

LR test vs. linear model: chi2(4) = 1247.03 Prob > chi2 = 0.0000

```

Note: LR test is conservative and provided only for reference.

5.3 Nice Table of Results

```

etable, estimates(crosssectional longitudinal) ///
showstars showstarsnote /// show stars and note
column(estimate) // column is modelname

```

	crosssectional	longitudinal
parental warmth in past week	0.835 ** (0.064)	0.913 ** (0.042)
physical punishment in past week	-0.992 ** (0.080)	-1.008 ** (0.050)
hypothetical identity group variable	-0.300 (0.217)	
recieved intervention		
1	0.640 ** (0.217)	0.859 ** (0.152)
Human Development Index	-0.003 (0.020)	-0.001 (0.020)
time		0.944 ** (0.066)
hypothetical identity group variable		
1		-0.128 (0.152)
Intercept	52.000 ** (1.371)	50.467 ** (1.338)
var(warmth)	0.023 (0.026)	0.011 (0.013)
var(_cons)	2.964 (0.974)	3.167 (0.915)
var(e)	34.975 (0.910)	26.027 (0.475)
var(_cons)		8.387 (0.472)
var(t)		0.000 (0.000)
Number of observations	3000	9000

** p<.01, * p<.05

5.4 QUESTIONS???

6 Cross-Classified Models

6.1 Introduction

A two level multilevel model imagines that *Level 1* units are nested in *Level 2* units. A three level multilevel model imagines that *Level 1* units are nested in *Level 2* units, which are in turn nested in *Level 3* units.

A cross-classified model imagines that the nesting is not hierarchical, but rather that there are two sets of clusters or nestings which overlap, but are not hierarchical.

6.2 Get Data

```
use "simulated_multilevel_longitudinal_data.dta", clear
```

6.3 Cross Classified Model

We can treat these random effects as being *cross classified*.

This might be useful if we had data where individuals lived in different countries at different times.

However, because `id` is in fact nested inside `country`, in this case, estimating the random effects as cross classified will be more time consuming, but will give us equivalent results to a three level model.

6.3.1 Standard (Less Computationally Efficient) Syntax

The below syntax will take a very long time to run with the full sample, and thus we have commented it out.

```
* mixed outcome t warmth physical_punishment || _all: R.country || _all: R.id
* est store crossed1
```

The documentation notes that we can use a *much* more computationally efficient version of the above command, which is what we do in these notes. The user can verify that both versions of the command will produce equivalent results.

In fact, at the end of handout we verify the similarity of both sets of syntax using a random sample.

6.3.2 Cross Classified With Computationally Efficient Syntax

```
mixed outcome t warmth physical_punishment || _all: R.country || id:
est store crossed2 // store crossed effects result
```

Performing EM optimization ...

Performing gradient-based optimization:
 Iteration 0: Log likelihood = -28516.314
 Iteration 1: Log likelihood = -28516.277
 Iteration 2: Log likelihood = -28516.277

Computing standard errors ...

Mixed-effects ML regression Number of obs = 9,000

Grouping information

Group variable		No. of groups	Observations per group		
			Minimum	Average	Maximum
-----+-----					
_all		1	9,000	9,000.0	9,000
id		3,000	3	3.0	3

Log likelihood = -28516.277 Wald chi2(3) = 1168.69
Prob > chi2 = 0.0000

	outcome	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
	t	.9434605	.065866	14.32	0.000	.8143654	1.072556
	warmth	.9053924	.0380439	23.80	0.000	.8308277	.9799572
physical_punishm~t		-1.014385	.0499354	-20.31	0.000	-1.112257	-.916514
	_cons	50.8301	.4123007	123.28	0.000	50.022	51.63819

Random-effects parameters		Estimate	Std. err.	[95% conf. interval]	
_all: Identity					
	var(R.country)	3.429974	.930313	2.015668	5.836634
id: Identity					
	var(_cons)	8.608872	.4757699	7.725107	9.59374
	var(Residual)	26.02862	.4752444	25.11363	26.97695

LR test vs. linear model: $\chi^2(2) = 1260.84$ Prob > $\chi^2 = 0.0000$

Note: LR test is conservative and provided only for reference.

6.4 Three Level Model

```
mixed outcome t warmth physical_punishment || country: || id: // 3 level w/ random intercept
est store threelevel // store random intercept model
```

Performing EM optimization ...

Performing gradient-based optimization:

Iteration 0: Log likelihood = -28516.314

Iteration 1: Log likelihood = -28516.277

Iteration 2: Log likelihood = -28516.277

Computing standard errors ...

Mixed-effects ML regression

Number of obs = 9,000

Grouping information

Group variable		No. of groups	Observations per group		
			Minimum	Average	Maximum
country		30	300	300.0	300
id		3,000	3	3.0	3

Log likelihood = -28516.277

Wald chi2(3) = 1168.69

Prob > chi2 = 0.0000

outcome	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
t	.9434605	.065866	14.32	0.000	.8143654	1.072556
warmth	.9053924	.0380439	23.80	0.000	.8308277	.9799572
physical_punishm~t	-1.014385	.0499354	-20.31	0.000	-1.112257	-.916514
_cons	50.8301	.4123007	123.28	0.000	50.022	51.63819

Random-effects parameters		Estimate	Std. err.	[95% conf. interval]	
country: Identity					
	var(_cons)	3.429974	.930313	2.015668	5.836634
id: Identity					
	var(_cons)	8.608872	.4757699	7.725107	9.59374
	var(Residual)	26.02862	.4752444	25.11363	26.97695

LR test vs. linear model: chi2(2) = 1260.84

Prob > chi2 = 0.0000

Note: LR test is conservative and provided only for reference.

6.5 Nice Table of Results of Three Level and Cross Classified Model

```
etable, estimates(threelevel crossed2), ///  
showstars showstarsnote /// show stars and note  
column(estimate) // column is modelname
```

```
invalid 'showstars'  
r(198);
```

```
r(198);
```

6.6 Verification of Syntax Equivalence for Cross Classified Model

```
keep if family <= 5 // random sample of families  
  
quietly mixed outcome t warmth physical_punishment || _all: R.country || _all: R.id  
  
est store crossed1A // less efficient syntax  
  
quietly mixed outcome t warmth physical_punishment || _all: R.country || id:  
  
est store crossed2A // more efficient syntax  
  
etable, estimates(crossed1A crossed2A) ///  
showstars showstarsnote /// show stars and note  
column(estimate) // column is modelname
```

(8,550 observations deleted)

crossed1A crossed2A

```

-----
time                0.745 **    0.745 **
                   (0.281)    (0.281)
parental warmth in past week  0.871 **    0.871 **
                   (0.160)    (0.160)
physical punishment in past week -1.262 ** -1.262 **
                   (0.206)    (0.206)
Intercept           51.755 **    51.755 **
                   (1.009)    (1.009)
var(R_country)       2.245
                   (1.319)    (1.319)
var(R_id)            5.425
                   (1.843)
var(e)              23.638    23.638
                   (1.933)    (1.933)
var(_cons)           5.425
                   (1.843)
Number of observations      450    450
-----
** p<.01, * p<.05

```

6.7 QUESTIONS???

References

- Bland, J M, and D G Altman. 1994. “Statistics Notes: Correlation, Regression, and Repeated Data.” *BMJ* 308 (April): 896. <https://doi.org/10.1136/bmj.308.6933.896>.
- Oliver, Mary, and Krista Tippett. 2015. “Mary Oliver: ‘I Got Saved by the Beauty of the World.’” The On Being Project. <https://onbeing.org/programs/mary-oliver-i-got-saved-by-the-beauty-of-the-world/>.

License and Citation

License



Multilevel Workshop by [Andrew Grogan-Kaylor](#) is licensed under a [Creative Commons Attribution 4.0 International License](#)

Citation

For attribution, please cite as:

Grogan-Kaylor (2025). *Multilevel Workshop*. Retrieved from <https://agrogan1.github.io/multilevel-workshop>

BibTeX Citation

```
@book{,
  author = {Andrew Grogan-Kaylor},
  city = {Ann Arbor, MI},
  title = {Multilevel Workshop},
  url = {https://agrogan1.github.io/multilevel-workshop},
  year = {2025},
}
```