Why OLS Is (Often) A Bad Model For Longitudinal Data

And Why The Equations and Data Structure of MLM Are The Answer

Andy Grogan-Kaylor

2026-10-27

Table of contents

1 Some Beginning Ideas 2

2 An Empirical Example 2

3 Introduction 2

4 A First Longitudinal Model 3

5 What About Change Scores? 3

6 What If We Have More Than Two Time Points? 3

7 Additionally ... 3

8 Our Answer To the Problem 4
- 8.1 Data in Long Format . . . 4
- 8.2 Equation . . . 4
  - 8.2.1 Simple MLM . . . 4
  - 8.2.2 Longitudinal MLM . . . 5

9 This Has The Following Advantages 5
- 9.1 First... . . . 5
- 9.2 How To Address Missing Data? . . . 5
- 9.3 Further... . . . 6
- 9.4 Appropriate Metric For Time . . . 6
- 9.5 Visually . . . 6
- 9.6 Lastly . . . 7
- 9.7 Let's continue to explore how this model works. . . . 7

10 Appendix: Two Conceptual Diagrams 7
- 10.1 OLS or MLM for 2 Timepoints . . . 7
- 10.2 Cross-Lagged Model . . . 7

References 7

1 Some Beginning Ideas

> "Despite the incredible diversity existing among and within human cultures, there are many phenomena that occur regularly in all known societies. These commonalities, or universals, while deriving in part from human nature, may also have specific social, cultural, and systemic sources. We need to develop a working understanding of these universals so that we might advance legitimate, empirically based human science set on creating knowledge that is politically relevant to fostering real solutions to the problems that complicate human co-existence in the Age of the Anthropocene." (Antweiler 2016)

> "The language we have in that world is not large enough for the territory that we've already entered." (Whyte and Tippett 2016)

2 An Empirical Example

😃 = life satisfaction
⏰ = time
👥 = social support

$$😃 = \beta_0 + \beta_1 ⏰ + \beta_2 👥 + \beta_4 👥 \times ⏰ + u_{0i} + u_{1i} \times ⏰ + u_{2i} \times 👥 + e_{it}$$

Figure 1: Life Satisfaction as a Function of Time And Social Support

3 Introduction

We are all familiar with the idea of:

$y_i = \beta_0 + \beta_1 x + e_i$ (OLS)

> Substantive Example
>
> Let's use the substantive example in Figure 1.

Table 1: Data in WIDE format

id	x1	x2	x3	y1	y2	y3
1						
2						
3						

4 A First Longitudinal Model

We could imagine a longitudinal model where we regress y_i at time 2 on y_i at time 1….

$y_{i2} = \beta_0 + \beta_1 x + \beta_2 y_{i1} + e_i$

And we could even make this (*perhaps confusingly*) a multilevel model for individual i in social unit j:

$y_{i2j} = \beta_0 + \beta_1 x + \beta_2 y_{i1j} + u_{0j} + e_{ij}$

… and add all of the usual random slope terms…

> **Thought Question**
>
> Any problems yet?

5 What About Change Scores?

$y_{i2} - y_{i1} = \beta_0 + \beta_1 x + e_i$

> **What Happens To The Regression Coefficients in a Change Score Model?**
>
> βy_{i1}

6 What If We Have More Than Two Time Points?

$y_{i3} = \beta_0 + \beta_1 x + \beta_2 y_{i1} + \beta_3 y_{i2} + e_i$

> **Multicollinearity**
>
> What is the problem here? We have 2 terms that are likely to be collinear:
> β_2 & β_3

This issue only becomes worse the more time points we add.

As a result, we are not really modeling y_2 and y_1.

7 Additionally …

> **No Explicit Function of Time**
>
> *Additionally*, we do not have an explicit function of time. We don't know really have a clear idea of whether our outcome increases with time, or decreases with time. Or whether the effect is curvilinear e.g. t^2 or $\ln(t)$.

⚠ *Unbalanced* Data Are A Problem

Additionally, any data that is *unbalanced* i.e. study participants enter the study late, or leave the study early are going to be difficult for this kind of model to deal with.

⚠ Missing Data Are A Problem

Similarly, data that is *missing at one time point, but present at other time points*, is going to be a problem for this kind of model. (and it is going to be difficult for many of our colleagues to see how we can get around this issue.)

8 Our Answer To the Problem

💡 We Reshape The Data and Use the SAME Notation!!!

"Mathematics is the art of giving the same name to different things." (Poincare 1908)

8.1 Data in Long Format

Table 2: Data in LONG format

id	t	x	y
1	1		
1	2		
1	3		
2	1		
2	2		
2	3		
3	1		
3	2		
3	3		

8.2 Equation

So.... we take our standard multilevel notation.

8.2.1 Simple MLM

$$y_{ij} = \beta_0 + \beta_1 x + u_{0j} + e_{ij} \tag{1}$$

cross out j write in t.

8.2.2 Longitudinal MLM

$$y_{it} = \beta_0 + \beta_1 t + u_{0i} + e_{it} \tag{2}$$

> Person-Observations
>
> Every row is a *person-observation* (person i observed at time t). Every person has *multiple rows*.

9 This Has The Following Advantages

> Key Intuition
>
> The *key intuition* is that any statistical procedure–whether it is aware of the clustering of the data or not–operates on the data on a *row by row* basis.

9.1 First...

1. Using the algebra in Equation 2, these models can easily accommodate both time varying and time invariant coefficients (Hox 2010; Hox et al. 2018; Singer and Willett 2003; Raudenbush and Bryk 2002; Rabe-Hesketh and Skrondal 2022).
2. No multicollinearity issue. By inspection of Equation 2, we see that there is only a single β coefficient for each variable, $\therefore$ no multicollinearity problem.
3. *Unbalanced data is less of a problem*, the data structure and estimation are robust to these possibilities (Singer and Willett 2003; Raudenbush and Bryk 2002).
4. *Missing data is less of a problem* (assuming *MCAR*). When a person observation is missing, that person simply has fewer rows of data (Hox 2010; Luke 2004; Raudenbush and Bryk 2002; Rabe-Hesketh and Skrondal 2022). But all rows of data are "matched" to the same person by i.

9.2 How To Address Missing Data?

> Addressing Missing Data is Complicated!!!
>
> It is sometimes best to (a) do nothing; (b) do something complicated.

- Ignore it.
- Fill in the mean.
- Use previous observation.
- Use next observation.
- Linearly interpolate previous and next observation.
- Regression imputation.
- Multiple imputation.

See: https://agrogan1.github.io/posts/missing-data/index.html

9.3 Further...

5. We now have an *explicit function of time* $\beta_1 t$, and could treat time more flexibly, by creating a polynomial function of time e.g. by adding $\beta_2 t^2$, etc. (Raudenbush and Bryk 2002; Singer and Willett 2003). (We could even substitute $\beta \ln(t)$.)
6. Again, by inspection of Equation 2, we see that *multiple or many time-points are not a problem*. Same algebra for 2 time points as for 10,000 time points. (Helpful when we start to think about intensive longitudinal data *e.g.* George Holden's *recording study*).
7. We are *measuring exactly the time at which events take place* for each individual (Singer and Willett 2003; Luke 2004). Not simply saying *Wave 1*, *Wave 2*, *Wave 3*, etc...
8. Every individual could have a *completely different set of time points* and even a *completely different number of time points* (Hox 2010; Hox et al. 2018; Singer and Willett 2003; Luke 2004).

And we can even add βx back into the model.

9.4 Appropriate Metric For Time

> **Caution**
>
> We do need to think carefully about what is the appropriate variable for time. Is it the variable we used to reshape the data–often `wave`–or some other more appropriate metric, like `age` or `time in study` (Singer and Willett 2003)?

9.5 Visually




Figure 2: A Multilevel Model For Longitudinal Data

9.6 Lastly

> **Caution**
>
> Generating appropriate descriptive statistics can be a question.

9.7 Let's continue to explore how this model works.

10 Appendix: Two Conceptual Diagrams

10.1 OLS or MLM for 2 Timepoints




Figure 3: An OLS Or Multilevel Model For 2 Timepoints

10.2 Cross-Lagged Model




Figure 4: A Cross Lagged Model For 3 Timepoints

References

Antweiler, Christoph. 2016. *Our Common Denominator: Human Universals Revisited*. Berghahn.

Hox, Joop. 2010. *Multilevel Analysis: Techniques and Applications*. 2nd ed. Routledge.

Hox, Joop, Mirjam Moerbeek, and Rens van de Schoot. 2018. *Multilevel Analysis: Techniques and Applications*. Third edition. Routledge, Taylor & Francis Group,.

Luke, Douglas. 2004. *Multilevel Modeling*. SAGE Publications, Inc. https://doi.org/10.4135/9781412985147.

Poincare, Henri. 1908. *Science Et Methode*. Flammarion.

Rabe-Hesketh, Sophia, and Anders Skrondal. 2022. *Multilevel and Longitudinal Modeling Using Stata*. In *Stata Press*, 4th ed. Stata Press.

Raudenbush, Stephen W, and Anthony S Bryk. 2002. *Hierarchical Linear Models: Applications and Data Analysis Methods*. Sage Publications.

Singer, Judith D, and John B Willett. 2003. *Applied Longitudinal Data Analysis : Modeling Change and Event Occurrence*. Oxford University Press.

Whyte, David, and Krista Tippett. 2016. *David Whyte: Seeking Language Large Enough*. The On Being Project. https://onbeing.org/programs/david-whyte-seeking-language-large-enough/.