

Logistic Regression The Basics

Andy Grogan-Kaylor

2025-07-07

Table of contents

1	Logistic Regression	1
2	Get The Data	1
3	Describe The Data	2
4	The Equation	2
5	Estimate Logistic Regression (logit y x)	3
6	Odds Ratios (logit y x, or)	3
7	β Coefficients and Odds Ratios	4
8	Coefficients, Standard Errors, p values, and Confidence Intervals	4

1 Logistic Regression

Basic handout on logistic regression for a binary dependent variable.

2 Get The Data

We start by obtaining *simulated data* from StataCorp.

```
clear all

graph close _all

use http://www.stata-press.com/data/r15/margex, clear
```

(Artificial data for margins)

3 Describe The Data

The variables are as follows:

```
describe
```

Contains data from <http://www.stata-press.com/data/r15/margex.dta>

```
Observations:      3,000      Artificial data for margins
Variables:         11        27 Nov 2016 14:27
```

Variable name	Storage type	Display format	Value label	Variable label
y	float	%6.1f		
outcome	byte	%2.0f		
sex	byte	%6.0f	sexlbl	
group	byte	%2.0f		
age	float	%3.0f		
distance	float	%6.2f		
ycn	float	%6.1f		
yc	float	%6.1f		
treatment	byte	%2.0f		
agegroup	byte	%8.0g	agelab	
arm	byte	%8.0g		

Sorted by: group

4 The Equation

$$\ln \left(\frac{p(outcome)}{1-p(outcome)} \right) = \beta_0 + \beta_1 x_1$$

Here $p(outcome)$ is the probability of the outcome.

$\frac{p(outcome)}{1-p(outcome)}$ is the *odds* of the outcome.

Hence, $\ln \left(\frac{p(outcome)}{1-p(outcome)} \right)$ is the *log odds*.

Logistic regression returns a β coefficient for each independent variable x .

These β coefficients can then be *exponentiated* to obtain *odds ratios*:

$$OR = e^{\beta}$$

5 Estimate Logistic Regression (logit y x)

We then run a logistic regression model in which `outcome` is the dependent variable. `sex`, `age` and `group` are the independent variables.

```
logit outcome i.sex c.age i.group
```

```
Iteration 0: Log likelihood = -1366.0718
Iteration 1: Log likelihood = -1111.4595
Iteration 2: Log likelihood = -1069.588
Iteration 3: Log likelihood = -1068
Iteration 4: Log likelihood = -1067.9941
Iteration 5: Log likelihood = -1067.9941
```

Logistic regression

Number of obs = 3,000
LR chi2(4) = 596.16
Prob > chi2 = 0.0000
Pseudo R2 = 0.2182

Log likelihood = -1067.9941

outcome	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
sex						
female	.4991622	.1347463	3.70	0.000	.2350643	.76326
age	.0902429	.0064801	13.93	0.000	.0775421	.1029437
group						
2	-.5855242	.1350192	-4.34	0.000	-.850157	-.3208915
3	-1.360208	.2914263	-4.67	0.000	-1.931393	-.7890228
_cons	-5.553038	.3498204	-15.87	0.000	-6.238674	-4.867403

6 Odds Ratios (logit y x, or)

We re-run the model with exponentiated coefficients (e^{β} to obtain odds ratios.

```
logit outcome i.sex c.age i.group, or
```

```
Iteration 0: Log likelihood = -1366.0718
Iteration 1: Log likelihood = -1111.4595
Iteration 2: Log likelihood = -1069.588
Iteration 3: Log likelihood = -1068
Iteration 4: Log likelihood = -1067.9941
```

Iteration 5: Log likelihood = -1067.9941

Logistic regression

Number of obs = 3,000

LR chi2(4) = 596.16

Prob > chi2 = 0.0000

Pseudo R2 = 0.2182

Log likelihood = -1067.9941

outcome	Odds ratio	Std. err.	z	P> z	[95% conf. interval]	
sex						
female	1.64734	.221973	3.70	0.000	1.26499	2.145258
age	1.09444	.0070921	13.93	0.000	1.080628	1.108429
group						
2	.5568139	.0751806	-4.34	0.000	.4273478	.725502
3	.2566074	.0747822	-4.67	0.000	.1449462	.4542885
_cons	.0038757	.0013558	-15.87	0.000	.0019524	.0076933

Note: _cons estimates baseline odds.

7 β Coefficients and Odds Ratios

Substantively	β	OR
x is associated with an increase in y	> 0.0	> 1.0
no association	0.0	1.0
x is associated with a decrease in y	< 0.0	< 1.0

8 Coefficients, Standard Errors, p values, and Confidence Intervals

- z statistic: $z = \frac{\beta}{se}$.
- p value if $z_{\text{observed}} > 1.96$ then $p < .05$.
- $CI = \beta \pm 1.96 * se$

Hence for the coefficient for **sex**, the confidence interval is:

$$.4991622 \pm (1.959964 * .1347463) = (.2350643, .7632601)$$

Confidence intervals for *odds ratios* (e^β) are obtained by exponentiating the confidence interval for the β coefficients. As a result of this non-linear transformation, confidence intervals for odds ratios are not symmetric.