



Improving Nitrogen Use Efficiency in Wheat by Genome Wide and Candidate Genes Targeted Association Studies

Agathe Mini, Gaetan Touzy, Catherine Ravel, Titouan Bonnot,
Etienne Paux, Jacques Le Gouis, Stéphane Lafarge

The BreedWheat Consortium



BreedWheat: French research project on wheat

2011-2020



15 public research laboratories (INRA, Universities...)


1 technical institute (Arvalis)

10 Breeding companies

1 competitiveness cluster (Céréales Vallée)

Genetic of wheat adaptation to biotic and abiotic stresses

Analyse genetic diversity and use genomic data to better understand wheat adaptation to biotic and abiotic stresses



- Nitrogen Use Efficiency: capacity for a plant to use nitrogen inputs
 - Improve NUE in wheat to better valorize fertilizer inputs
 - For ecological issue (leaching, fertilizer production...)
 - For economical issue: nitrogen input account for 25% of operational cost
 - For wheat, NUE is around 65%
 - A target for breeding to have a higher NUE for a better yield and grain protein content

Genetics of wheat adaptation to nitrogen stresses

Candidate genes

Markers associated with trait(s) linked to yield stability
under stress (GWAS results)

Knowledge about French/European germplasm

New phenotypic traits/methods

Candidate genes
from -omics data

GWAS

GWAS

Genotypic Data

420k Axiom array






Phenotypic Data



Genomic areas of interest



Tools for breeding



Associated markers

Search for new variabilities in diversity panel

Field validation

GWAS: Materials

- Panel Composition

- 220 winter wheat elite varieties




- Genotypic data

⇒ 197K SNPs available for GWAS (polymorphic...)

⇒ With physical position on Wheat RefSeqV1


- Analysis of structuration, LD


- Mean = 2,4 cM
- K & Q matrix for GWAS analysis




GWAS: Phenotypic Data

- 26 trials / 3 years, dedicated to:
 - Nitrogen: 12 => Opt. & Stress conditions
 - Drought: 6
 - FHB: 4
 - *Septoria*: 4
- 26 trials / 1.10^6 data points
 - Plant phenology
 - Yield, yield components, GPC...
 - Biomass, nitrogen content, senescence...



- For each environment, determination of
 - Timing of stress
 - Length of stress
 - Stress intensity
 - Using
 - Environmental data
 - Plant development informations
 - Phenotypic data
- 
- A PCA plot with 'Dim 1 (16.16%)' on the x-axis and 'Dim 2 (16.16%)' on the y-axis. The x-axis ranges from -1 to 4, and the y-axis ranges from -1 to 4. A dashed horizontal line is at y=0. A dashed yellow curve starts near (0, -0.5) and goes up and to the right. A point labeled '2012_A' is located near the origin (0, 0).




10 Field trials validated and kept for the global analysis

		Intensity		
		High	Medium	Low
Duration	High	4		
	Medium	2	1	
	Low			3


Genotypes characterization





- Stress tolerance index => knowledge on genotypes behaviour regarding Nitrogen stresses
 - Calculate for each field trials
 - Classification made by type of stress: mean of index tolerance
 - High
 - Medium
 - Low

Génotype	GYRN 2012_Artycla	GYRN 2012_DECap	GYRN 2012_NRMmon	GYRN 2012_ZZhou	GYRN 2012_SICmau	GYRN 2013_Artycla	GYRN 2013_Inicie	GYRN 2013_NRMmon	GYRN 2013_Libret	GYRN 2013_SYNand	MOYENNE des résidus normalisés	Ecart type résidus normalisés	NOMBRE DE FOIS OU RN > 0
PBW030	0.13	1.30	1.48	1.06	0.75	1.78	0.41	1.20	3.26	1.57	1.29	0.82	10
PBW184	0.78	1.29	0.34	-0.06	2.59	1.94	2.48	2.19	-0.11	1.40	1.29	0.96	8
PBW011	1.00	1.94	2.31	-0.94	0.91	1.82	0.89	0.10	1.14	1.85	1.10	0.92	9
PBW130	1.44	0.59	0.68	2.83	0.95	2.50	0.63	1.12	0.16	0.09	1.10	0.87	10
PBW012	2.50	1.81	1.43	1.61	0.09	1.43	-0.96	1.83	-0.54	0.85	1.01	1.07	8
PBW060	-1.30	1.57	1.23	1.32	0.50	1.30	-0.73	2.67	0.77	1.39	0.87	1.09	8
PBW142	1.18	1.82	0.31	1.22	1.27	1.09	-0.93	1.59	0.48	0.54	0.86	0.75	9
PBW015	1.86	1.28	0.95	-1.06	0.73	1.35	0.62	1.10	0.26	1.36	0.84	0.76	9
PBW129	1.52	1.18	2.85	1.74	0.90	-0.37	-0.22	0.14	1.72	-1.07	0.84	1.14	7
PBW125	2.04	2.31	0.44	-0.12	0.10	0.74	-0.06	0.75	0.68	1.23	0.81	0.79	8
PBW195	-0.10	1.88	0.42	-0.27	2.45	0.66	1.72	0.80	-0.32	0.26	0.75	0.91	7
PBW159	0.17	0.61	0.43	-0.09	0.80	0.03	0.47	2.13	2.02	0.58	0.71	0.73	9
PBW156	0.68	0.82	0.65	-1.20	0.50	2.64	0.88	-0.05	1.57	0.62	0.71	0.94	8
PBW114	0.85	-0.24	0.02	1.18	1.17	1.75	-1.29	1.17	1.47	1.01	0.71	0.88	8
PBW008	0.78	2.07	1.48	0.40	1.26	-0.80	0.64	0.73	-0.89	1.14	0.68	0.89	8
PBW033	0.38	0.40	2.02	0.50	1.25	-1.15	0.11	1.34	0.13	1.50	0.65	0.86	9
PBW160	0.07	0.52	1.01	-0.11	1.50	0.04	1.58	0.87	1.15	-0.17	0.65	0.63	8
PBW090	-0.65	0.45	0.78	1.78	0.86	0.34	-1.01	0.86	1.53	1.36	0.63	0.85	8
PBW128	0.84	-0.37	0.37	0.09	1.53	1.68	0.37	1.46	-0.46	0.54	0.61	0.73	8
PBW021	0.10	1.45	0.78	-0.05	0.15	1.29	0.34	-0.51	1.61	0.67	0.58	0.67	8



GWAS: a pipe line of analysis

- Analysis from the raw data to GWAS results
 - Workflow analysis
 - R script
 - Common way to analyze field trials
 - Graphical and statistical analysis
 - Capacity to
 - Analyze phenotypic field data in order to:
 - Determine heterogeneity, errors, outliers...
 - Genotype x Treatment interaction
 - Heritabilities
 - ...
 - Calculate adjusted means corrected by spatial effects
 - Produce GWAS results
- 
- A histogram showing the distribution of field trial data. The x-axis is labeled 'valeur SA' and ranges from 350 to 700. The y-axis is labeled 'Effectif' and ranges from 0.000 to 0.008. The histogram bars are light blue. A red normal distribution curve is overlaid on the histogram, peaking at approximately 500 on the x-axis.

[illegible]

GWAS: Numbers

Traits Classification	Traits Number
Nitrogen	17
Biomass	5
Others	3
Height	1
Disease	11
Phenology	4
GPC	5
Yield and Components	11
Senescence	22
Lodging	10
Tolerance Index	9

In HN & LN

181 traits

10 trials

1 075 traits x treatments x trials

196 464 SNP
Usable for GWAS

211 198 800 tested associations

1st selection: 73
LD blocks
selected on
traits,
robustness...


16193

SNP associated in at least 2 trials and
for the same traits (624 LD blocks)


91151

Associated SNPs
with traits of
interest

GWAS: examples




- A robust areas detected in 10 trials



- Selection of genetic areas
- Definition of boundaries using LD decay
- Candidate genes in the area
 - Need to densify => Sequence capture on genes in the confidence interval based on RefSeqV1 prediction

Allelic effect

- Effect on traits of the minor allele in the panel




Next ?

- Densification
 - To have better associated markers (causal SNPs)
 - SequenceCapture targeted to genes in the area => SNP discovery & association studies
- Identification of candidate gene
 - *A priori* => by annotation analysis
 - Without *a priori* => transcriptomic data

Transcriptomic Data:

- NUE:

- Response of grain protein composition to N and S supply in *Triticum monococcum*



(Bonnot et al. in prep)

- **18 StageCond** analyzed x 3 rep (54 samples)
 - 2 stages (100, 200° Cd)
 - 4 stages (300, 400, 500, 600° Cd) x 4 nutritions
- Paired-End sequencing, HiSeq 2500 – Illumina
- **Expression** : 22 443 genes expressed

Transcriptomic Data:

- Analysis

- GO: stress response, amino acids transport...

- Differential expression regarding:

- Time points
 - Nutrition

- GeneNetwork:

- RuNet: a tool to make connections

- has been developed to infer regulatory networks from high throughput –omics data, phenotypic data...

- Identification of master regulators linked to traits of interest...

- In combination with proteomic/metabolomic data

- Genes, proteins, metabolites responding similarly to the nutrition?


- Genes, proteins, metabolites strongly co-expressed /co-accumulated ?

- Central genes in the response to the nutrition ?

RESEARCH ARTICLE

RuNet: A Web-Oriented Platform for Regulatory Network Inference, Application to Wheat –Omics Data

Jonathan Vincent^{1,2,3}, Pierre Martre^{2,3,4,*}, Benjamin Gouriou¹, Catherine Ravel^{2,3}, Zhanwu Dai^{2,3,4}, Jean-Marc Petit⁴, Marie Pailloux^{1,*}




Transcriptomic Data:

- Analysis



– 152 Candidate genes

- Understanding of specific stress mechanism
- SNP discovery / Association studies
- Co-localization between transcriptomic candidate genes and GWAS results



GWAS: Conclusions

- GWAS
 - A very interesting set of phenotypic data and GWAS results
 - Need to valorize all these data-set:
 - Meta-analysis, GxE, combination of positive alleles etc...
- ToDo
 - Densification to have access to better associated markers
 - Analysis of natural variabilities in a diversity panel from Core collection (450 acc.)
 - Validation in the field of the most promising regions



BreedWheat - Conclusions

- Candidate genes:
 - Gene lists, knowledge on molecular/physiology mechanisms of nitrogen stress response...
- Tools for breeding, transgenesis/genome editing etc...
 - Associated-markers
 - Genotypes classification
 - Identification of interesting candidate genes

For NUE, but also drought, FHB and *Septoria*...

P 276
Gaetan Touzy

Thanks to all people involved in these work



Gaetan Touzy, Katia Beauchêne, Jean Pierre Cohan



Celine Zimmerli, Guenter Wertz, Michael Schmolke



Franck Lacoudre, Jayne Stragliati, Celine Duque, Jérémy Derory



Agathe Mini, Guylaine Besnier-Hebert, Sébastien Praud



Stephen Sunderwith, Benoit Foucault, Patrice Senellard



Sylvie Dutriez, Alain Lacassagne



Olivier Lucas, Laure Duchalais, Christophe Michelet



Denis Beghin, Ellen Goudemand, Olivier Robert



Pascal Giraudeau, Sébastien Ducroquet



Jacques le Gouis, **Catherine Ravel**,
Titouan Bonnot, Etienne Paux,
Emmanuel Heumez
All teams involved in research projects



Sebastien Caiveau, Odile Argilier,



Thank You

