

# Genome-wide significance

In genome-wide association studies, **genome-wide significance** (abbreviated **GWS**) is a specific threshold for determining the statistical significance of a reported association between a given single-nucleotide polymorphism (SNP) and a given trait. The most commonly accepted threshold is  $p < 5 \times 10^{-8}$ , which is based on performing a Bonferroni correction for all the independent common SNPs across the human genome.<sup>[1]</sup> If a *p*-value is found to be lower than this threshold in a genome-wide association study, the null hypothesis of no true SNP-association will typically be rejected.<sup>[2]</sup> However, there has been some criticism of this threshold, with a 2012 study suggesting that a significant number of associations with *p*-values just above this threshold are genuine, replicable associations. The authors of this study concluded that their finding "...suggests a possible relaxation in the current GWS threshold."<sup>[3]</sup> More recently, it has been suggested that the conventional threshold should be modified to take into account the increasing prevalence of low-frequency genetic variants in genome-wide association studies.<sup>[4]</sup>

## References

1. Xu, ChangJiang; Tachmazidou, Ioanna; Walter, Klaudia; Ciampi, Antonio; Zeggini, Eleftheria; Greenwood, Celia M. T.; the UK10K Consortium (April 2014). "Estimating Genome-Wide Significance for Whole-Genome Sequencing Studies: Genome-Wide Significance for Rare Variants" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4489336>). *Genetic Epidemiology*. **38** (4): 281–290. doi:10.1002/gepi.21797 (<https://doi.org/10.1002%2Fgepi.21797>). PMC 4489336 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4489336>). PMID 24676807 (<https://pubmed.ncbi.nlm.nih.gov/24676807>).
2. Roeder, Kathryn; Wasserman, Larry (November 2009). "Genome-Wide Significance Levels and Weighted Hypothesis Testing" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2920568>). *Statistical Science*. **24** (4): 398–413. doi:10.1214/09-STS289 (<https://doi.org/10.1214%2F09-ST289>). ISSN 0883-4237 (<https://www.worldcat.org/issn/0883-4237>). PMC 2920568 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2920568>). PMID 20711421 (<https://pubmed.ncbi.nlm.nih.gov/20711421>).
3. Panagiotou, Orestis A; Ioannidis, John P A; for the Genome-Wide Significance Project (February 2012). "What should the genome-wide significance threshold be? Empirical replication of borderline genetic associations". *International Journal of Epidemiology*. **41** (1): 273–286. doi:10.1093/ije/dyr178 (<https://doi.org/10.1093%2Fije%2Fdyr178>). ISSN 0300-5771 (<https://www.worldcat.org/issn/0300-5771>). PMID 22253303 (<https://pubmed.ncbi.nlm.nih.gov/22253303>).
4. Fadista, João; Manning, Alisa K.; Florez, Jose C.; Groop, Leif (August 2016). "The (in)famous GWAS P-value threshold revisited and updated for low-frequency variants" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4970684>). *European Journal of Human Genetics*. **24** (8): 1202–1205. doi:10.1038/ejhg.2015.269 (<https://doi.org/10.1038%2Fejhg.2015.269>). ISSN 1476-5438 (<https://www.worldcat.org/issn/1476-5438>). PMC 4970684 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4970684>). PMID 26733288 (<https://pubmed.ncbi.nlm.nih.gov/26733288>).

Retrieved from "[https://en.wikipedia.org/w/index.php?title=Genome-wide\\_significance&oldid=950555465](https://en.wikipedia.org/w/index.php?title=Genome-wide_significance&oldid=950555465)"

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use and Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.