

# Vector Spaces

Arthur Ryman, arthur.ryman@gmail.com

June 25, 2022

## Abstract

This article contains Z Notation type declarations for vector spaces and some related objects. It has been type checked by *fUZZ*.

## Contents

<b>1</b>	<b>Real Vector Spaces</b>	<b>1</b>
1.1	Notation for Vector Addition, Zero, and Negative: $+$ <code>\addV</code> , $\mathbf{0}$ <code>\zeroV</code> , and $-$ <code>\negV</code> . . . . .	1
1.2	Real Scalar Multiplication: $*$ <code>\mulS</code> , $\times$ <code>\timesS</code> , and <i>RealScalarMultiplication</i> . . . . .	1
1.3	The Set of All Real Vector Spaces: $\text{vec}_{\mathbb{R}}$ <code>\vecR</code> . . . . .	2
1.4	Real Linear Transformations: <i>RealLinearTransformation</i> . . . . .	2
1.5	The Set of All Real Linear Transformations: $L_{\mathbb{R}}$ <code>\homVecR</code> . . . . .	3
<b>2</b>	<b>Real <math>n</math>-tuples</b>	<b>3</b>
2.1	The Set of All Finite Sequences of Real Numbers: $\mathbb{R}^{\infty}$ <code>\Rinf</code> . . . . .	3
2.2	The Component Projection Function: $\pi$ <code>\piRinf</code> . . . . .	4
2.3	The Set of All Well-Dimensioned Subsets of $\mathbb{R}^{\infty}$ : $\Delta_{\mathbb{R}}$ <code>\DeltaRinf</code> . . . . .	4
2.4	The Dimension of a Well-Dimensioned Set of Tuples: $\dim$ <code>\dimRinf</code> . . . . .	4
2.5	The Set of All Compatible Pairs of Tuples: $\mathbb{R}^{\Delta}$ <code>\RinfDelta</code> . . . . .	4
2.6	Addition of Compatible Tuples: $+$ <code>\addRinf</code> . . . . .	4
2.7	Subtraction of Compatible Tuples: $-$ <code>\subRinf</code> . . . . .	5
2.8	The Negative of a Tuple: $-$ <code>\negRinf</code> . . . . .	5
2.9	Scalar Multiplication of a Tuple: $*$ <code>\smulRinf</code> . . . . .	5
2.10	The Set of All Real $n$ -tuples: $\mathbb{R}$ <code>\Rtup</code> . . . . .	6
2.11	Addition of $n$ -tuples: <i>addRtup</i> . . . . .	6
2.12	Subtraction of $n$ -tuples: <i>subRtup</i> . . . . .	6
2.13	The Negative of an $n$ -tuple: <i>negRtup</i> . . . . .	7
2.14	The Zero Real $n$ -tuple: $\mathbf{0}$ <code>\zeroRtup</code> . . . . .	7
2.15	Scalar Multiplication of an $n$ -tuple: <i>smulRtup</i> . . . . .	7

2.16	The Real Vector Space of $n$ -tuples: <i>vecRtup</i> . . . . .	8
2.17	Linear Transformations of $n$ -tuples: $L_{\mathbb{R}}$ \iLinRtup . . . . .	8
2.18	The Identity Transformation of $n$ -tuples: $I$ \idRtup . . . . .	8
<b>3</b>	<b>The Metric Topology on Real <math>n</math>-tuples</b>	<b>8</b>
3.1	The Dot Product of Tuples: $\cdot$ \dotRinf . . . . .	8
3.2	The Norm of a Tuple: <i>norm</i> \normRinf . . . . .	9
3.3	The Open Ball at a Tuple: <i>ball</i> \ballRinf . . . . .	9
3.4	The Set of All Open Balls at an $n$ -tuple: <i>balls</i> \ballsRtup . . . . .	9
3.5	The Usual Topology on $n$ -tuples: $\tau_{\mathbb{R}}$ \tauRtup . . . . .	10
3.6	The Set of All Neighbourhoods of a Tuple: <i>neigh</i> \neighRinf . . . . .	10
3.7	The Topological Space of $n$ -tuples: $\mathbb{R}_{\tau}$ \tsRtup . . . . .	10
<b>4</b>	<b>Continuity</b>	<b>10</b>
4.1	Real-Valued Functions That Are Continuous on the Set of All $n$ -tuples: $C^0$ \CzeroRtup . . . . .	10
4.2	Real-Valued Functions That Are Continuous on a Subset of $n$ -tuples: $C^0$ \CzeroSubsetRtup . . . . .	11
4.3	Real-Valued Functions That Are Continuous at an $n$ -tuple: $C^0$ \CzeroPointRtup	11
4.4	$m$ -tuple-Valued Functions That Are Continuous on the Set of All $n$ -tuples: $C^0$ \CzeroRtupRtup . . . . .	11
4.5	$m$ -tuple-Valued Functions That Are Continuous on a Subset of $n$ -tuples: $C^0$ \CzeroSubsetRtupRtup . . . . .	12
4.6	$m$ -tuple-Valued Functions That Are Continuous at an $n$ -tuple: <i>VectorConinuous</i> , $C^0$ \CzeroPointRtupRtup . . . . .	12
<b>5</b>	<b>Differentiability</b>	<b>13</b>
5.1	The Difference Quotient: <i>DifferenceQuotient</i> and <i>diffQuot</i> . . . . .	13
5.2	The Derivative of a Continuous $m$ -tuple-Valued Function: <i>VectorDifferentiable</i>	14

## 1 Real Vector Spaces

Real vector spaces are multidimensional generalizations of real numbers. They are the objects studied in linear algebra and are foundational to differential geometry.

In the following let  $\mathbf{t}$  denote a set of elements which we'll refer to as *vectors* and let  $A$  denote an Abelian group over the vectors in which the binary operation is denoted as addition. Let  $v$  and  $w$  denote vectors and let  $x$  and  $y$  denote real numbers.

### 1.1 Notation for Vector Addition, Zero, and Negative: $+$ \addV, $\mathbf{0}$ \zeroV, and $-$ \negV

Let  $v + w$  denote vector addition, let  $\mathbf{0}$  denote the zero vector, and let  $-v$  denote the negative vector.

### 1.2 Real Scalar Multiplication: $*$ \mulS, $\times$ \timesS, and *RealScalarMultiplication*

A *real scalar multiplication* operation on the vectors is an operation *smul* that maps the pair  $(x, v)$  to another vector, typically denoted  $x * v$  or  $x \times v$ , such that multiplication by 0 maps all vectors to the group identity element, multiplication by 1 maps each vector to itself, multiplication preserves group addition, and multiplication distributes over both real and group addition.

Let *RealScalarMultiplication* denote this situation.

<i>RealScalarMultiplication</i> [t]	_____
$A : \text{abgroup } t$	
$smul : \mathbb{R} \times t \rightarrow t$	
<b>let</b> $(- + -) == A;$	
$\mathbf{0} == \text{identity\_element } A;$	
$(- * -) == smul \bullet$	
$\forall x, y : \mathbb{R}; v, w : t \bullet$	
$\mathbf{0} * v = \mathbf{0} \wedge$	
$1 * v = v \wedge$	
$(x * y) * v = x * (y * v) \wedge$	
$(x + y) * v = x * v + y * v \wedge$	
$x * (v + w) = x * v + x * w$	

- Multiplying by 0 gives the zero vector.
- Multiplying by 1 gives the same vector.
- Scalar multiplication is associative.
- Scalar addition distributes over scalar multiplication.
- Vector addition distributes over scalar multiplication.

### 1.3 The Set of All Real Vector Spaces: $\text{vec}_{\mathbb{R}}$ \vecR

A *real vector space* is a pair  $(A, smul)$  where  $A$  is an Abelian group and *smul* is a real scalar multiplication on the elements of  $A$ . The elements of  $A$  are referred to as *vectors*.

Let  $\text{vec}_{\mathbb{R}} t$  denote the set of all real vector spaces over  $t$ ,

$$\text{vec}_{\mathbb{R}} t == \{ \text{RealScalarMultiplication}[t] \bullet (A, smul) \}$$

## 1.4 Real Linear Transformations: *RealLinearTransformation*

Let  $V_1$  and  $V_2$  be real vector spaces and let  $f$  be a homomorphism of the underlying Abelian groups. The map  $f$  is said to be a *linear transformation* if  $f$  maps scalar multiples of vectors to the scalar multiple of the mapped vectors.

Let *RealLinearTransformation* denote this situation.

$RealLinearTransformation[t, u]$	_____
$f : t \rightarrow u$ $V_1 : \text{vec}_{\mathbb{R}} t$ $V_2 : \text{vec}_{\mathbb{R}} u$	
<b>let</b> $A_1 == \text{first } V_1; (- * -) == \text{second } V_1;$ $A_2 == \text{first } V_2; (- \times -) == \text{second } V_2 \bullet$ $f \in \text{hom}_{\text{grp}}(A_1, A_2) \wedge$ $(\forall x : \mathbb{R}; v : t \bullet$ $\quad f(x * v) = x \times (f v))$	

- The vector space  $V_1$  has Abelian group  $A_1$  and scalar multiplication  $(- * -)$ .
- The vector space  $V_2$  has Abelian group  $A_2$  and scalar multiplication  $(- \times -)$ .
- The map  $f$  is a homomorphism of the underlying Abelian groups.
- The map  $f$  maps scalar multiples of vectors in  $t$  to scalar multiples of the mapped vectors in  $u$ .

## 1.5 The Set of All Real Linear Transformations: $L_{\mathbb{R}} \setminus \text{homVecR}$

Let  $V_1$  and  $V_2$  be real vector spaces. Let  $L_{\mathbb{R}}(V_1, V_2)$  denote the set of all linear transformations from  $V_1$  to  $V_2$ . A linear transformation is also referred to as a *homomorphism* of vector spaces.

$$L_{\mathbb{R}}[t, u] == (\lambda V_1 : \text{vec}_{\mathbb{R}} t; V_2 : \text{vec}_{\mathbb{R}} u \bullet \{ f : t \rightarrow u \mid RealLinearTransformation[t, u] \})$$

## 2 Real $n$ -tuples

The preceding section described real vector spaces abstractly. In this section we define a family of finite-dimensional real vector spaces whose elements are finite sequences of real numbers, also referred to as *real tuples*.

## 2.1 The Set of All Finite Sequences of Real Numbers: $\mathbb{R}^\infty \setminus \text{Rinf}$

Let  $n$  be a natural number. A finite sequence of  $n$  real numbers is called a *real  $n$ -tuple*. Let  $\mathbb{R}^\infty$  denote the set of all real  $n$ -tuples for any  $n$ .

$$\mathbb{R}^\infty == \text{seq } \mathbb{R}$$

## 2.2 The Component Projection Function: $\pi \setminus \text{piRinf}$

The real numbers that comprise an  $n$ -tuple are called its *components*. Let  $v$  be a real  $n$ -tuple and let  $i$  be an integer where  $1 \leq i \leq n$ . The real number  $v(i)$  is the  $i$ -th component of  $v$ . Let  $\pi(i)$  be the projection function that maps an  $n$ -tuple  $v$  to its  $i$ -th component  $v(i)$ .

$$\begin{array}{|l} \pi : \mathbb{N}_1 \longrightarrow \mathbb{R}^\infty \rightarrow \mathbb{R} \\ \hline \forall i : \mathbb{N}_1 \bullet \\ \pi(i) = (\lambda v : \mathbb{R}^\infty \mid i \in \text{dom } v \bullet v(i)) \end{array}$$

## 2.3 The Set of All Well-Dimensioned Subsets of $\mathbb{R}^\infty$ : $\Delta_{\mathbb{R}} \setminus \text{DeltaRinf}$

A non-empty subset of  $\mathbb{R}^\infty$  is said to be *well-dimensioned* if each of its elements has the same number of components. Let  $\Delta_{\mathbb{R}}$  denote the family of all well-dimensioned subsets of  $\mathbb{R}^\infty$ .

$$\begin{array}{|l} \Delta_{\mathbb{R}} : \mathcal{F} \mathbb{R}^\infty \\ \hline \Delta_{\mathbb{R}} = \{ S : \mathbb{P}_1 \mathbb{R}^\infty \mid \forall v, w : S \bullet \#v = \#w \} \end{array}$$

## 2.4 The Dimension of a Well-Dimensioned Set of Tuples: $\text{dim} \setminus \text{dimRinf}$

Let  $S \in \Delta_{\mathbb{R}}$  be a well-dimensioned set of tuples. The number of components of each tuple in  $S$  is called its *dimension*. Let  $\text{dim}(S)$  denote the dimension of  $S$ .

$$\begin{array}{|l} \text{dim} : \Delta_{\mathbb{R}} \longrightarrow \mathbb{N} \\ \hline \forall S : \Delta_{\mathbb{R}} \bullet \\ \text{dim } S = (\mu v : S \bullet \#v) \end{array}$$

## 2.5 The Set of All Compatible Pairs of Tuples: $\mathbb{R}^\Delta \setminus \text{RinfDelta}$

The pair of real tuples  $(v, w)$  is said to be *compatible* if each member has the same number of components. Let  $\mathbb{R}^\Delta$  denote the set of all compatible pairs of real tuples. If the pair  $(v, w)$  is compatible then  $v$  and  $w$  are said to be compatible with each other.

$$\begin{array}{|l} \mathbb{R}^\Delta : \mathbb{R}^\infty \leftrightarrow \mathbb{R}^\infty \\ \hline \mathbb{R}^\Delta = \{ v, w : \mathbb{R}^\infty \mid \#v = \#w \} \end{array}$$

## 2.6 Addition of Compatible Tuples: $+$ \addRinf

Let  $v$  and  $w$  be  $n$ -tuples. Vector addition of  $v$  and  $w$  is the  $n$ -tuple  $v + w$  defined by component-wise addition.

$$\left| \begin{array}{l} \_ + \_ : \mathbb{R}^\Delta \longrightarrow \mathbb{R}^\infty \\ \hline \langle \rangle + \langle \rangle = \langle \rangle \\ \forall n : \mathbb{N}_1; v, w : \mathbb{R}^\infty \mid n = \#v = \#w \bullet \\ \quad v + w = (\lambda i : 1 \dots n \bullet v\ i + w\ i) \end{array} \right.$$

## 2.7 Subtraction of Compatible Tuples: $-$ \subRinf

Vector subtraction is defined similarly.

$$\left| \begin{array}{l} \_ - \_ : \mathbb{R}^\Delta \longrightarrow \mathbb{R}^\infty \\ \hline \langle \rangle - \langle \rangle = \langle \rangle \\ \forall n : \mathbb{N}_1; v, w : \mathbb{R}^\infty \mid n = \#v = \#w \bullet \\ \quad v - w = (\lambda i : 1 \dots n \bullet v\ i - w\ i) \end{array} \right.$$

## 2.8 The Negative of a Tuple: $-$ \negRinf

Let  $-v$  denote the negative of  $v$ .

$$\left| \begin{array}{l} \_ : \mathbb{R}^\infty \longrightarrow \mathbb{R}^\infty \\ \hline -\langle \rangle = \langle \rangle \\ \forall n : \mathbb{N}_1; v : \mathbb{R}^\infty \mid n = \#v \bullet \\ \quad -v = (\lambda i : 1 \dots n \bullet -(v\ i)) \end{array} \right.$$

## 2.9 Scalar Multiplication of a Tuple: $*$ \smulRinf

Let  $v$  be an  $n$ -tuple and let  $c$  be a real number. Scalar multiplication of  $v$  by  $c$  is the  $n$ -tuple  $c * v$  defined by component-wise multiplication.

$$\left| \begin{array}{l} \_ * \_ : \mathbb{R} \times \mathbb{R}^\infty \longrightarrow \mathbb{R}^\infty \\ \hline \forall c : \mathbb{R} \bullet \\ \quad c * \langle \rangle = \langle \rangle \\ \forall c : \mathbb{R}; n : \mathbb{N}_1; v : \mathbb{R}^\infty \mid n = \#v \bullet \\ \quad c * v = (\lambda i : 1 \dots n \bullet c * (v\ i)) \end{array} \right.$$

**Remark.** *Scalar multiplication is associative in the sense that  $(a * b) * v = a * (b * v)$*

$$\begin{aligned} \forall a, b : \mathbb{R}; v : \mathbb{R}^\infty \bullet \\ (a * b) * v = a * (b * v) \end{aligned}$$

## 2.10 The Set of All Real $n$ -tuples: $\mathbb{R} \setminus \text{Rtup}$

Let  $\mathbb{R}(n)$  denote  $\mathbb{R}^n$ , the set of all  $n$ -tuples for some given  $n$ .

$$\begin{array}{|l} \mathbb{R} : \mathbb{N} \longrightarrow \mathbb{P} \mathbb{R}^\infty \\ \hline \forall n : \mathbb{N} \bullet \\ \mathbb{R}(n) = \{ v : \mathbb{R}^\infty \mid \#v = n \} \end{array}$$

**Remark.**

$$\mathbb{R}^\infty = \bigcup \{ n : \mathbb{N} \bullet \mathbb{R}(n) \}$$

**Remark.** *The subset  $\mathbb{R}(n)$  is well-dimensioned.*

$$\begin{aligned} \forall n : \mathbb{N} \bullet \\ \mathbb{R}(n) \in \Delta_{\mathbb{R}} \end{aligned}$$

**Remark.** *The dimension of  $\mathbb{R}(n)$  is  $n$ .*

$$\begin{aligned} \forall n : \mathbb{N} \bullet \\ \dim(\mathbb{R}(n)) = n \end{aligned}$$

## 2.11 Addition of $n$ -tuples: $\text{addRtup}$

Let  $\text{addRtup}(n)$  denote the restriction of addition to  $\mathbb{R}(n)$ .

$$\begin{aligned} \text{addRtup} == \\ (\lambda n : \mathbb{N} \bullet \\ (\lambda v, w : \mathbb{R}(n) \bullet v + w)) \end{aligned}$$

**Example.** *The binary operation  $\text{addRtup}(n)$  defines an Abelian group over  $\mathbb{R}(n)$ .*

$$\begin{aligned} \forall n : \mathbb{N} \bullet \\ \text{addRtup}(n) \in \text{abgroup}(\mathbb{R}(n)) \end{aligned}$$

## 2.12 Subtraction of $n$ -tuples: $\text{subRtup}$

Let  $\text{subRtup}(n)$  denote the restriction of subtraction to  $\mathbb{R}(n)$ .

$$\begin{aligned} \text{subRtup} == \\ (\lambda n : \mathbb{N} \bullet \\ (\lambda v, w : \mathbb{R}(n) \bullet v - w)) \end{aligned}$$

### 2.13 The Negative of an $n$ -tuple: $negRtup$

Let  $negRtup(n)$  denote the restriction of the negative operation to  $\mathbb{R}(n)$ .

$$\begin{aligned} negRtup == \\ (\lambda n : \mathbb{N} \bullet \\ (\lambda v : \mathbb{R}(n) \bullet -v)) \end{aligned}$$

**Remark.** The operation  $negRtup(n)$  is the inverse operation of the Abelian group  $addRtup(n)$ .

$$\begin{aligned} \forall n : \mathbb{N} \bullet \\ negRtup(n) = inverse\_operation(addRtup(n)) \end{aligned}$$

### 2.14 The Zero Real $n$ -tuple: $\mathbf{0} \setminus zeroRtup$

Let  $\mathbf{0}(n)$  denote the  $n$ -tuple consisting of all zeroes.

$$\begin{array}{|l} \mathbf{0} : \mathbb{N} \rightarrow \mathbb{R}^\infty \\ \hline \mathbf{0}(0) = \langle \rangle \\ \forall n : \mathbb{N}_1 \bullet \\ \mathbf{0}(n) = (\lambda i : 1 \dots n \bullet 0) \end{array}$$

**Remark.** Every component of  $\mathbf{0}(n)$  is 0.

$$\begin{aligned} \forall n : \mathbb{N} \bullet \\ \forall i : 1 \dots n \bullet \\ (\pi i)(\mathbf{0} n) = 0 \end{aligned}$$

**Remark.** The tuple  $\mathbf{0}(n)$  is in  $\mathbb{R}(n)$ .

$$\begin{aligned} \forall n : \mathbb{N} \bullet \\ \mathbf{0}(n) \in \mathbb{R}(n) \end{aligned}$$

**Remark.** The tuple  $\mathbf{0}(n)$  is the identity element of the Abelian group  $addRtup(n)$ .

$$\begin{aligned} \forall n : \mathbb{N} \bullet \\ \mathbf{0}(n) = identity\_element(addRtup(n)) \end{aligned}$$

### 2.15 Scalar Multiplication of an $n$ -tuple: $smulRtup$

Let  $smulRtup(n)$  denote scalar multiplication restricted to  $\mathbb{R}(n)$ .

$$\begin{aligned} smulRtup == \\ (\lambda n : \mathbb{N} \bullet \\ (\lambda c : \mathbb{R}; v : \mathbb{R}(n) \bullet c * v)) \end{aligned}$$



## 2.16 The Real Vector Space of $n$ -tuples: $vecRtup$

Let  $vecRtup(n)$  denote the real vector space of  $n$ -tuples.

$$vecRtup == (\lambda n : \mathbb{N} \bullet (addRtup(n), smulRtup(n)))$$

**Remark.** The pair  $vecRtup(n)$  defines a vector space over  $\mathbb{R}(n)$ .

$$\forall n : \mathbb{N} \bullet \\ vecRtup(n) \in vec_{\mathbb{R}}(\mathbb{R}(n))$$

## 2.17 Linear Transformations of $n$ -tuples: $L_{\mathbb{R}} \setminus linRtup$

Define  $L_{\mathbb{R}}(n, m)$  to be the set of all linear transformations from  $\mathbb{R}^n$  to  $\mathbb{R}^m$ .

$$\begin{array}{|l} L_{\mathbb{R}} : \mathbb{N} \times \mathbb{N} \longrightarrow \mathcal{P}(\mathbb{R}^{\infty} \rightarrow \mathbb{R}^{\infty}) \\ \hline \forall n, m : \mathbb{N} \bullet \\ L_{\mathbb{R}}(n, m) = L_{\mathbb{R}}(vecRtup(n), vecRtup(m)) \end{array}$$

## 2.18 The Identity Transformation of $n$ -tuples: $I \setminus idRtup$

Let  $I(n)$  denote the identity function on  $\mathbb{R}(n)$ .

$$\begin{array}{|l} I : \mathbb{N} \longrightarrow \mathbb{R}^{\infty} \rightarrow \mathbb{R}^{\infty} \\ \hline \forall n : \mathbb{N} \bullet \\ I(n) = id(\mathbb{R}(n)) \end{array}$$

**Remark.** The function  $I(n)$  is a linear transformation.

$$\forall n : \mathbb{N} \bullet \\ I(n) \in L_{\mathbb{R}}(n, n)$$

# 3 The Metric Topology on Real $n$ -tuples

## 3.1 The Dot Product of Tuples: $\cdot \setminus dotRinf$

The *inner* or *dot* product of  $n$ -tuples  $v$  and  $w$  is the real number  $v \cdot w$  defined by the sum of the component-wise products.

$$\begin{array}{|l} \_ \cdot \_ : \mathbb{R}^{\Delta} \longrightarrow \mathbb{R} \\ \hline \langle \rangle \cdot \langle \rangle = 0 \\ \forall x, y : \mathbb{R}; v, w : \mathbb{R}^{\infty} \mid \#v = \#w \bullet \\ (\langle x \rangle \frown v) \cdot (\langle y \rangle \frown w) = x * y + v \cdot w \end{array}$$

Each  $\mathbb{R}(n)$  is a real inner product space under the operation of dot product defined above.

### 3.2 The Norm of a Tuple: `norm \normRinf`

The norm  $\|v\|$  of the  $n$ -tuple  $v$  is the positive square root of its dot product with itself.

$$\|v\| = \sqrt{v \cdot v}$$

Define  $\text{norm}(v)$  to be  $\|v\|$ .

$$\left| \begin{array}{l} \text{norm} : \mathbb{R}^\infty \longrightarrow \mathbb{R} \\ \hline \forall v : \mathbb{R}^\infty \bullet \\ \quad \text{norm}(v) = \text{sqrt}(v \cdot v) \end{array} \right|$$

The concepts of continuity, limits, and differentiability extend to functions between normed vector spaces such as  $\mathbb{R}^n$ .

### 3.3 The Open Ball at a Tuple: `ball \ballRinf`

Let  $\text{ball}(v, r)$  denote the *open ball* in  $\mathbb{R}(n)$  of radius  $r \in \mathbb{R}_+$  centred at  $v \in \mathbb{R}(n)$ .

$$\left| \begin{array}{l} \text{ball} : \mathbb{R}^\infty \times \mathbb{R}_+ \longrightarrow \mathcal{P} \mathbb{R}^\infty \\ \hline \forall v : \mathbb{R}^\infty; r : \mathbb{R}_+ \bullet \\ \quad \text{let } n == \#v \bullet \\ \quad \quad \text{ball}(v, r) = \{ w : \mathbb{R}(n) \mid \text{norm}(v - w) < r \} \end{array} \right|$$

### 3.4 The Set of All Open Balls at an $n$ -tuple: `balls \ballsRtup`

Let  $\text{balls}(n)$  denote the family of all open balls in  $\mathbb{R}(n)$ .

$$\left| \begin{array}{l} \text{balls} : \mathbb{N} \longrightarrow \mathcal{F} \mathbb{R}^\infty \\ \hline \forall n : \mathbb{N} \bullet \\ \quad \text{balls}(n) = \{ v : \mathbb{R}(n); r : \mathbb{R}_+ \bullet \text{ball}(v, r) \} \end{array} \right|$$

**Remark.** *The set of all open balls in  $\mathbb{R}(n)$  is a family of sets in  $\mathbb{R}(n)$ .*

$$\forall n : \mathbb{N} \bullet \\ \quad \text{balls}(n) \in \mathcal{F}(\mathbb{R}(n))$$

### 3.5 The Usual Topology on $n$ -tuples: $\tau_{\mathbb{R}} \setminus \text{tauRtup}$

The *usual topology* on  $\mathbb{R}(n)$  is the topology generated by the open balls in  $\mathbb{R}(n)$ . Let  $\tau_{\mathbb{R}}(n)$  denote the usual topology on  $\mathbb{R}(n)$ .

$$\begin{array}{|l} \tau_{\mathbb{R}} : \mathbb{N} \longrightarrow \mathcal{F}\mathbb{R}^{\infty} \\ \hline \forall n : \mathbb{N} \bullet \\ \tau_{\mathbb{R}}(n) = \text{topGen}[\mathbb{R}(n)](\text{balls}(n)) \end{array}$$

**Remark.** If  $n \in \mathbb{N}$  then  $\tau_{\mathbb{R}}(n)$  is a topology on  $\mathbb{R}(n)$ .

$$\forall n : \mathbb{N} \bullet \tau_{\mathbb{R}}(n) \in \text{top}[\mathbb{R}(n)]$$

### 3.6 The Set of All Neighbourhoods of a Tuple: $\text{neigh} \setminus \text{neighRinf}$

Let  $v \in \mathbb{R}(n)$ . An open set  $U$  in the usual topology  $\tau_{\mathbb{R}}(n)$  that contains  $v$  is called a *neighbourhood* of  $v$ . Let  $\text{neigh}(v)$  denote the set of all neighbourhoods of  $x$ .

$$\begin{array}{|l} \text{neigh} : \mathbb{R}^{\infty} \longrightarrow \mathcal{F}\mathbb{R}^{\infty} \\ \hline \forall n : \mathbb{N}; v : \mathbb{R}^{\infty} \mid n = \#v \bullet \\ \text{neigh}(v) = \{ U : \tau_{\mathbb{R}}(n) \mid v \in U \} \end{array}$$

**Remark.** The set of all neighbourhoods of  $v \in \mathbb{R}(n)$  is a family of sets in  $\mathbb{R}(n)$ .

$$\begin{array}{l} \forall n : \mathbb{N}; v : \mathbb{R}^{\infty} \mid n = \#v \bullet \\ \text{neigh}(v) \in \mathcal{F}(\mathbb{R}(n)) \end{array}$$

### 3.7 The Topological Space of $n$ -tuples: $\mathbb{R}_{\tau} \setminus \text{tsRtup}$

Let  $\mathbb{R}_{\tau}(n)$  denote the topological space defined by the usual topology on  $\mathbb{R}(n)$ .

$$\begin{array}{|l} \mathbb{R}_{\tau} : \mathbb{N} \longrightarrow \text{topSpaces}[\mathbb{R}^{\infty}] \\ \hline \forall n : \mathbb{N} \bullet \\ \mathbb{R}_{\tau}(n) = (\mathbb{R}(n), \tau_{\mathbb{R}}(n)) \end{array}$$

## 4 Continuity

### 4.1 Real-Valued Functions That Are Continuous on the Set of All $n$ -tuples: $\mathbb{C}^0 \setminus \text{CzeroRtup}$

A function  $f \in \mathbb{R}^n \longrightarrow \mathbb{R}$  is said to be *continuous* if it is continuous with respect to the usual topologies on  $\mathbb{R}^n$  and  $\mathbb{R}$ . Let  $\mathbb{C}^0(n)$  denote the set of these continuous functions.

$$\begin{array}{|l}
C^0 : \mathbb{N} \rightarrow \mathbb{P}(\mathbb{R}^\infty \rightarrow \mathbb{R}) \\
\hline
\forall n : \mathbb{N} \bullet \\
C^0(n) = C^0(\mathbb{R}_\tau(n), \mathbb{R}_\tau)
\end{array}$$

#### 4.2 Real-Valued Functions That Are Continuous on a Subset of $n$ -tuples: $C^0 \setminus \text{CzeroSubsetRtup}$

Let  $U$  be a subset of  $\mathbb{R}^n$ . A function  $f \in U \rightarrow \mathbb{R}$  is said to be *continuous on  $U$*  if it is continuous with respect to the topology induced on  $U$ . Let  $C^0(U)$  denote the set of these continuous functions.

$$\begin{array}{|l}
C^0 : \Delta_{\mathbb{R}} \rightarrow \mathbb{P}(\mathbb{R}^\infty \rightarrow \mathbb{R}) \\
\hline
\forall U : \Delta_{\mathbb{R}} \bullet \\
\text{let } n == \dim U \bullet \\
C^0(U) = C^0(\mathbb{R}_\tau(n) \upharpoonright_{\text{top}} U, \mathbb{R}_\tau)
\end{array}$$

#### 4.3 Real-Valued Functions That Are Continuous at an $n$ -tuple: $C^0 \setminus \text{CzeroPointRtup}$

A partial function  $f$  from  $\mathbb{R}^n$  to  $\mathbb{R}$  is said to be *continuous at  $x \in \mathbb{R}^n$*  if its domain contains a neighbourhood  $U$  of  $x$  such that its restriction to  $U$  is continuous on  $U$ . Let  $C^0(x)$  denote the set of such functions.

$$\begin{array}{|l}
C^0 : \mathbb{R}^\infty \rightarrow \mathbb{P}(\mathbb{R}^\infty \rightarrow \mathbb{R}) \\
\hline
\forall x : \mathbb{R}^\infty \bullet \\
\text{let } n == \#x \bullet \\
C^0(x) = \{ f : \mathbb{R}(n) \rightarrow \mathbb{R} \mid \exists U : \text{neigh}(x) \mid U \subseteq \text{dom } f \bullet U \triangleleft f \in C^0(U) \}
\end{array}$$

#### 4.4 $m$ -tuple-Valued Functions That Are Continuous on the Set of All $n$ -tuples: $C^0 \setminus \text{CzeroRtupRtup}$

A mapping  $f$  from  $\mathbb{R}(n)$  to  $\mathbb{R}(m)$  is said to be continuous if it is continuous with respect to the usual topologies. Let  $C^0(n, m)$  denote the set of these continuous mappings.

$$\begin{array}{|l}
C^0 : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{P}(\mathbb{R}^\infty \rightarrow \mathbb{R}^\infty) \\
\hline
\forall n, m : \mathbb{N} \bullet \\
C^0(n, m) = C^0(\mathbb{R}_\tau(n), \mathbb{R}_\tau(m))
\end{array}$$

**Example.** The function  $I(n)$  is continuous.

$$\begin{array}{|l}
\forall n : \mathbb{N} \bullet \\
I(n) \in C^0(n, n)
\end{array}$$

**Theorem 1.** *Linear functions are continuous.*

$$\forall n, m : \mathbb{N} \bullet \\ L_{\mathbb{R}}(n, m) \subseteq C^0(n, m)$$

#### 4.5 $m$ -tuple-Valued Functions That Are Continuous on a Subset of $n$ -tuples: $C^0 \setminus \text{CzeroSubsetRtupRtup}$

Let  $U$  be any subset of  $\mathbb{R}(n)$ . Let  $C^0(U, m)$  denote the set of continuous mappings from the topology induced by  $\mathbb{R}_\tau(n)$  on  $U$  to  $\mathbb{R}_\tau(m)$ .

$$\left| \begin{array}{l} C^0 : \Delta_{\mathbb{R}} \times \mathbb{N} \rightarrow \mathbb{P}(\mathbb{R}^\infty \leftrightarrow \mathbb{R}^\infty) \\ \hline \forall n, m : \mathbb{N} \bullet \\ \quad \forall U : \Delta_{\mathbb{R}} \mid \dim(U) = n \bullet \\ \quad \quad C^0(U, m) = C^0(\mathbb{R}_\tau(n) \upharpoonright_{\text{top}} U, \mathbb{R}_\tau(m)) \end{array} \right|$$

**Remark.**

$$\forall n, m : \mathbb{N} \bullet \\ C^0(\mathbb{R}(n), m) = C^0(n, m)$$

#### 4.6 $m$ -tuple-Valued Functions That Are Continuous at an $n$ -tuple: *VectorContinuous*, $C^0 \setminus \text{CzeroPointRtupRtup}$

Let  $x \in \mathbb{R}(n)$  and let  $f$  be a partial function from  $\mathbb{R}(n)$  to  $\mathbb{R}(m)$  whose domain includes some neighbourhood  $U$  of  $x$  such that  $f$  restricted to  $U$  is continuous. In this case  $f$  is said to be *continuous at  $x$* .

$$\left| \begin{array}{l} \text{VectorContinuous} \\ \hline n, m : \mathbb{N} \\ f : \mathbb{R}^\infty \leftrightarrow \mathbb{R}^\infty \\ x : \mathbb{R}^\infty \\ \hline f \in \mathbb{R}(n) \leftrightarrow \mathbb{R}(m) \\ \exists U : \text{neigh}(x) \mid \\ \quad U \subseteq \text{dom } f \bullet \\ \quad \quad U \triangleleft f \in C^0(U, m) \end{array} \right|$$

Let  $C^0(x, m)$  denote the set of all partial functions  $f$  from  $\mathbb{R}(n)$  to  $\mathbb{R}(m)$  that are continuous at  $x$ .

$$\begin{array}{|l}
C^0 : \mathbb{R}^\infty \times \mathbb{N} \rightarrow \mathbb{P}(\mathbb{R}^\infty \rightarrow \mathbb{R}^\infty) \\
\hline
\forall n, m : \mathbb{N} \bullet \forall x : \mathbb{R}(n) \bullet \\
C^0(x, m) = \\
\{ f : \mathbb{R}(n) \rightarrow \mathbb{R}(m) \mid \text{VectorContinuous} \}
\end{array}$$

**Example.** The function  $I(n)$  is continuous at every point  $x \in \mathbb{R}(n)$ .

$$\begin{array}{l}
\forall n : \mathbb{N} \bullet \forall x : \mathbb{R}(n) \bullet \\
I(n) \in C^0(x, n)
\end{array}$$

**Theorem 2.** Linear functions are continuous everywhere.

$$\begin{array}{l}
\forall n, m : \mathbb{N} \bullet \\
\forall x : \mathbb{R}(n); L : \mathbb{L}_{\mathbb{R}}(n, m) \bullet \\
L \in C^0(x, m)
\end{array}$$

## 5 Differentiability

Let  $x \in \mathbb{R}^n$  and let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be continuous at  $x$ . Then  $f$  is said to be *differentiable at  $x$*  if there exists a linear transformation  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  such that  $f(x + h) - f(x)$  is approximately linear in  $h$  for very small  $h$ .

$$f(x + h) - f(x) \approx L(h) + O(h^2) \quad \text{when} \quad \|h\| \approx 0$$

This condition can be written as a limit.

$$\lim_{h \rightarrow 0} \frac{\|f(x + h) - f(x) - L(h)\|}{\|h\|} = 0$$

### 5.1 The Difference Quotient: *DifferenceQuotient* and *diffQuot*

The limit exists when the following difference quotient function  $q : \mathbb{R}^n \rightarrow \mathbb{R}$  is continuous at 0.

$$q(h) = \begin{cases} \frac{\|f(x+h) - f(x) - L(h)\|}{\|h\|} & \text{if } h \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

Given a function  $f$  that is continuous at  $x$ , and a linear transformation  $L$ , we can define the difference quotient  $q$ . Clearly  $q$  is uniquely determined by  $f$ ,  $x$ , and  $L$ . Let *DifferenceQuotient* denote this situation.

<i>DifferenceQuotient</i> <i>VectorContinuous</i> $L : \mathbb{R}^\infty \rightarrow \mathbb{R}^\infty$ $q : \mathbb{R}^\infty \rightarrow \mathbb{R}$
$L \in \mathbb{L}_{\mathbb{R}}(n, m)$ $\text{dom } q = \{ h : \mathbb{R}(n) \mid x + h \in \text{dom } f \}$ $\forall h : \text{dom } q \mid h \neq \mathbf{0}(n) \bullet$ $q(h) = \text{norm}(f(x + h) - f(x) - L(h)) / \text{norm}(h)$ $q(\mathbf{0}(n)) = 0$

- $L$  is a linear transformation from  $\mathbb{R}(n)$  to  $\mathbb{R}(m)$ .
- The difference quotient  $q$  is defined on a subset of  $\mathbb{R}(n)$  that contains  $\mathbf{0}(n)$ .
- $q(h)$  is defined as the quotient when  $h$  is non-zero.
- $q(0)$  is defined as zero.

Let  $\text{diffQuot}(f, x, L)$  denote the difference quotient  $q$ .

$$\text{diffQuot} == \{ \text{DifferenceQuotient} \bullet (f, x, L) \mapsto q \}$$

## 5.2 The Derivative of a Continuous $m$ -tuple-Valued Function: *VectorDifferentiable*

The continuous function  $f$  is *differentiable at  $x$*  when there exists a linear transformation  $L$  such that the difference quotient  $q$  is continuous at 0. In this case  $L$  is unique and is referred to as the *derivative at  $x$* .

<i>VectorDifferentiable</i> <i>VectorContinuous</i> $L : \mathbb{R}^\infty \rightarrow \mathbb{R}^\infty$
$\text{let } q == \text{diffQuot}(f, x, L) \bullet$ $q \in C^0(\mathbf{0}(n))$

- The continuous function  $f$  is differentiable at  $x$  with derivative  $L$  if the resulting difference quotient  $q$  is continuous at  $\mathbf{0}(n)$ .

**Remark.** *If  $L$  exists then it is unique.*

Let  $C^\infty(x, m)$  denote the set of all functions  $f \in \mathbb{R}(n) \rightarrow \mathbb{R}(m)$  that are smooth at  $x \in \mathbb{R}(n)$ .