


Optimization

3.2.5 Hooke–Jeeves method (pattern search)

José Firmino Aguilar Madeira

What if, instead of the diagonals, we try the axes?


3.2.4 ended on Rosenbrock, at $(0,0)$, $f = 1$: Box tests the 4 **diagonals** and none improves.



What if, instead of the diagonals, we try the axes?

3.2.4 ended on Rosenbrock, at $(0,0)$, $f = 1$: Box tests the 4 **diagonals** and none improves.

But along the x_1 axis: $f(0.25, 0) = 0.953 < 1$.




What if, instead of the diagonals, we try the axes?

3.2.4 ended on Rosenbrock, at $(0,0)$, $f = 1$: Box tests the 4 diagonals and none improves.

But along the x_1 axis: $f(0.25, 0) = 0.953 < 1$.

Why not test $\pm \mathbf{e}_1, \dots, \pm \mathbf{e}_n$?

n	Box: 2^n	axes: $2n$
2	4	4
10	1 024	20
20	$\approx 10^6$	40



One exploration by hand

$f(x_1, x_2) = 3x_1^2 + x_2^2 - 12x_1 - 8x_2$ (minimum at $(2, 4)$, $f = -28$).

Base point $\mathbf{x}_0 = (1, 1)$, $f = -16$; steps


$\Delta = (\Delta_1, \Delta_2) = (0.5, 0.5)$.

coordinate x_1 , from $(1, 1)$

$(x_1 + \Delta_1, x_2) = (1.5, 1)$	$f^+ = -18.25$
------------------------------------	----------------

$(x_1, x_2) = (1, 1)$	$f = -16$
-----------------------	-----------

$(x_1 - \Delta_1, x_2) = (0.5, 1)$	$f^- = -12.25$
------------------------------------	----------------



One exploration by hand

$f(x_1, x_2) = 3x_1^2 + x_2^2 - 12x_1 - 8x_2$ (minimum at $(2, 4)$, $f = -28$).

Base point $\mathbf{x}_0 = (1, 1)$, $f = -16$; steps


$\Delta = (\Delta_1, \Delta_2) = (0.5, 0.5)$.

coordinate x_1 , from $(1, 1)$

$(x_1 + \Delta_1, x_2) = (1.5, 1)$	$f^+ = -18.25$
$(x_1, x_2) = (1, 1)$	$f = -16$
$(x_1 - \Delta_1, x_2) = (0.5, 1)$	$f^- = -12.25$

coordinate x_2 , from the **already improved** point $(1.5, 1)$

$(x_1, x_2 + \Delta_2) = (1.5, 1.5)$	$f^+ = -21$
$(x_1, x_2) = (1.5, 1)$	$f = -18.25$
$(x_1, x_2 - \Delta_2) = (1.5, 0.5)$	$f^- = -15$



One exploration by hand

$f(x_1, x_2) = 3x_1^2 + x_2^2 - 12x_1 - 8x_2$ (minimum at $(2, 4)$, $f = -28$).

Base point $\mathbf{x}_0 = (1, 1)$, $f = -16$; steps

$\Delta = (\Delta_1, \Delta_2) = (0.5, 0.5)$.

coordinate x_1 , from $(1, 1)$

$(x_1 + \Delta_1, x_2) = (1.5, 1) \quad f^+ = -18.25$

$(x_1, x_2) = (1, 1) \quad f = -16$

$(x_1 - \Delta_1, x_2) = (0.5, 1) \quad f^- = -12.25$

coordinate x_2 , from the **already improved** point $(1.5, 1)$


$(x_1, x_2 + \Delta_2) = (1.5, 1.5) \quad f^+ = -21$

$(x_1, x_2) = (1.5, 1) \quad f = -18.25$

$(x_1, x_2 - \Delta_2) = (1.5, 0.5) \quad f^- = -15$

In each coordinate we keep $f_{\min} = \min\{f^+, f, f^-\}$. Result
 $(1.5, 1.5) \neq \mathbf{x}_0$: **success**, $\mathbf{x}_1 = (1.5, 1.5)$, with 4 evaluations.

Coordinates explored **sequentially**, each from the **best point so far** (Deb, 2012).



Exploration around the base point $\mathbf{x}^c$, with steps $\Delta = (\Delta_1, \dots, \Delta_n)$

Set $\mathbf{x} = \mathbf{x}^c$ and $i = 1$.

Step 1 Compute $f = f(\mathbf{x})$, $f^+ = f(\mathbf{x}_i + \Delta_i)$ and $f^- = f(\mathbf{x}_i - \Delta_i)$.

Step 2 $f_{\min} = \min\{f, f^+, f^-\}$; $\mathbf{x}$ becomes the corresponding point.

Step 3 If $i < n$, set $i = i + 1$ and go to Step 1.

Step 4 If $\mathbf{x} \neq \mathbf{x}^c$: **success**; otherwise: **failure**.

$n_f = 2n$ per exploration ($f(\mathbf{x}^c)$ is already known)

- Both directions are always tested: we keep the **best** of the three points, not the first that improves.
- **Box**: **all combinations** of $\pm (2^n)$.
Hooke–Jeeves: **one coordinate at a time** ($2n$).
- The direction obtained, $\mathbf{x} - \mathbf{x}^c$, has components in $\{-\Delta_i, 0, +\Delta_i\}$: a rough indication of the direction of progress (not the best), without derivatives.

We have a direction that worked: why not extrapolate?

The exploration took us from $\mathbf{x}_0$ to $\mathbf{x}_1$ and improved. We have information the Box method did not use:


$$\mathbf{x}_1 - \mathbf{x}_0.$$

Why explore again only around $\mathbf{x}_1$? If that direction worked:

let us go further in the same direction

$$\mathbf{x}_{k+1}^P = \mathbf{x}_k + (\mathbf{x}_k - \mathbf{x}_{k-1}) = 2\mathbf{x}_k - \mathbf{x}_{k-1}.$$

In the example: $\mathbf{x}_2^P = 2(1.5, 1.5) - (1, 1) = (2, 2)$, $f = -24$.



We have a direction that worked: why not extrapolate?

The exploration took us from $\mathbf{x}_0$ to $\mathbf{x}_1$ and improved. We have information the Box method did not use:

$$\mathbf{x}_1 - \mathbf{x}_0.$$


Why explore again only around $\mathbf{x}_1$? If that direction worked:

let us go further in the same direction

$$\mathbf{x}_{k+1}^P = \mathbf{x}_k + (\mathbf{x}_k - \mathbf{x}_{k-1}) = 2\mathbf{x}_k - \mathbf{x}_{k-1}.$$

In the example: $\mathbf{x}_2^P = 2(1.5, 1.5) - (1, 1) = (2, 2)$, $f = -24$.

And we **explore again** around $\mathbf{x}_2^P$: $(2, 2) \rightarrow \mathbf{x}_2 = (2, 2.5)$,
 $f = -25.75 < f(\mathbf{x}_1) = -21$: **success**.



Exploratory + pattern: the Hooke–Jeeves method

Hooke and Jeeves (1961): the two moves we have just built, alternated.

Exploratory move

Local search, axis by axis, from the base point: perturb each coordinate by $\pm\Delta_i$ and keep the best of the three points. Cost: $2n$.

Pattern move

Extrapolation in the direction of progress: $\mathbf{x}_{k+1}^P = 2\mathbf{x}_k - \mathbf{x}_{k-1}$, followed by a new exploration around $\mathbf{x}_{k+1}^P$. Repeated while there is progress: the jumps *accumulate* the direction that works.

the pattern **accelerates**

the exploration **corrects**


The exploratory steps are aligned with the axes, but extrapolation makes it possible to follow oblique valleys.

The steps Δ_i are the *poll step*: they play the role of the step length of the other methods (and they come back in 5.1 and 6.5).

Complete example: the path curves

$\mathbf{x}_0 = (1, 1)$, $\Delta = (0.5, 0.5)$, $R = 2$, $\varepsilon = 0.2$ (a large ε , just so the example ends quickly).

1. **Step 2** (exploration at $\mathbf{x}_0 = (1, 1)$, $f = -16$): $(1.5, 1) \rightarrow -18.25$;
 $(1.5, 1.5) \rightarrow -21$. Success: $\mathbf{x}_1 = (1.5, 1.5)$.




Complete example: the path curves

$\mathbf{x}_0 = (1, 1)$, $\Delta = (0.5, 0.5)$, $R = 2$, $\varepsilon = 0.2$ (a large ε , just so the example ends quickly).

1. **Step 2** (exploration at $\mathbf{x}_0 = (1, 1)$, $f = -16$): $(1.5, 1) \rightarrow -18.25$; $(1.5, 1.5) \rightarrow -21$. Success: $\mathbf{x}_1 = (1.5, 1.5)$.
2. **Steps 4–6**: $\mathbf{x}_2^P = 2(1.5, 1.5) - (1, 1) = (2, 2)$, $f = -24$; exploration $\rightarrow \mathbf{x}_2 = (2, 2.5)$, $f = -25.75 < -21$: success.

pattern (2, 2), explor. $\rightarrow (2, 2.5)$, $f = -25.75$: accepted




Complete example: the path curves

$\mathbf{x}_0 = (1, 1)$, $\Delta = (0.5, 0.5)$, $R = 2$, $\varepsilon = 0.2$ (a large ε , just so the example ends quickly).

1. **Step 2** (exploration at $\mathbf{x}_0 = (1, 1)$, $f = -16$): $(1.5, 1) \rightarrow -18.25$; $(1.5, 1.5) \rightarrow -21$. Success: $\mathbf{x}_1 = (1.5, 1.5)$.
2. **Steps 4–6**: $\mathbf{x}_2^P = 2(1.5, 1.5) - (1, 1) = (2, 2)$, $f = -24$; exploration $\rightarrow \mathbf{x}_2 = (2, 2.5)$, $f = -25.75 < -21$: success.
3. **Steps 4–6**: $\mathbf{x}_3^P = 2(2, 2.5) - (1.5, 1.5) = (2.5, 3.5)$, $f = -27$; exploration $\rightarrow \mathbf{x}_3 = (2, 4)$, $f = -28 < -25.75$: success.

pattern $(2.5, 3.5)$, explor. $\rightarrow (2, 4)$, $f = -28$: accepted




Complete example: the path curves

$\mathbf{x}_0 = (1, 1)$, $\Delta = (0.5, 0.5)$, $R = 2$, $\varepsilon = 0.2$ (a large ε , just so the example ends quickly).

1. **Step 2** (exploration at $\mathbf{x}_0 = (1, 1)$, $f = -16$): $(1.5, 1) \rightarrow -18.25$; $(1.5, 1.5) \rightarrow -21$. Success: $\mathbf{x}_1 = (1.5, 1.5)$.
2. **Steps 4–6**: $\mathbf{x}_2^P = 2(1.5, 1.5) - (1, 1) = (2, 2)$, $f = -24$; exploration $\rightarrow \mathbf{x}_2 = (2, 2.5)$, $f = -25.75 < -21$: success.
3. **Steps 4–6**: $\mathbf{x}_3^P = 2(2, 2.5) - (1.5, 1.5) = (2.5, 3.5)$, $f = -27$; exploration $\rightarrow \mathbf{x}_3 = (2, 4)$, $f = -28 < -25.75$: success.
4. **Steps 4–6**: $\mathbf{x}_4^P = 2(2, 4) - (2, 2.5) = (2, 5.5)$, $f = -25.75$; exploration $\rightarrow (2, 5)$, $f = -27 \geq -28$: **failure**. Step 3: $\|\Delta\| = 0.707 \geq \varepsilon$, set $\Delta = (0.25, 0.25)$.

pattern $(2, 5.5)$, explor. $\rightarrow (2, 5)$, $f = -27$: rejected




Complete example: the path curves

$\mathbf{x}_0 = (1, 1)$, $\Delta = (0.5, 0.5)$, $R = 2$, $\varepsilon = 0.2$ (a large ε , just so the example ends quickly).

1. **Step 2** (exploration at $\mathbf{x}_0 = (1, 1)$, $f = -16$): $(1.5, 1) \rightarrow -18.25$; $(1.5, 1.5) \rightarrow -21$. Success: $\mathbf{x}_1 = (1.5, 1.5)$.
2. **Steps 4–6**: $\mathbf{x}_2^P = 2(1.5, 1.5) - (1, 1) = (2, 2)$, $f = -24$; exploration $\rightarrow \mathbf{x}_2 = (2, 2.5)$, $f = -25.75 < -21$: success.
3. **Steps 4–6**: $\mathbf{x}_3^P = 2(2, 2.5) - (1.5, 1.5) = (2.5, 3.5)$, $f = -27$; exploration $\rightarrow \mathbf{x}_3 = (2, 4)$, $f = -28 < -25.75$: success.
4. **Steps 4–6**: $\mathbf{x}_4^P = 2(2, 4) - (2, 2.5) = (2, 5.5)$, $f = -25.75$; exploration $\rightarrow (2, 5)$, $f = -27 \geq -28$: **failure**. Step 3: $\|\Delta\| = 0.707 \geq \varepsilon$, set $\Delta = (0.25, 0.25)$.
5. **Steps 2–3**: exploration at $(2, 4)$ fails; $\|\Delta\| = 0.354$: $\Delta = (0.125, 0.125)$; fails again; $\|\Delta\| = 0.177 < \varepsilon$: **stop**. Solution $(2, 4)$, $f = -28$, $n_f = 28$.

The pattern always points in a straight line; the exploration corrects; the path **curves**.



Hooke–Jeeves: $\mathbf{x}_0$, steps Δ , reduction factor $R > 1$, ε

Step 1 Choose $\mathbf{x}_0$, Δ_i ($i = 1, \dots, n$), $R > 1$ and ε ; set $k = 0$.

Step 2 Exploratory move with $\mathbf{x}_k$ as the base point; let $\mathbf{x}$ be the outcome. If it is a success, set $\mathbf{x}_{k+1} = \mathbf{x}$ and go to Step 4; otherwise, go to Step 3.

Step 3 If $\|\Delta\| < \varepsilon$, **stop**; otherwise, set $\Delta_i = \Delta_i/R$ ($i = 1, \dots, n$) and go to Step 2.

Step 4 Set $k = k + 1$ and perform the pattern move: $\mathbf{x}_{k+1}^P = \mathbf{x}_k + (\mathbf{x}_k - \mathbf{x}_{k-1})$.


Step 5 Exploratory move with $\mathbf{x}_{k+1}^P$ as the base point; let $\mathbf{x}_{k+1}$ be the outcome.

Step 6 If $f(\mathbf{x}_{k+1}) < f(\mathbf{x}_k)$, go to Step 4; otherwise, go to Step 3 (with $\mathbf{x}_k$ as the base point).

- Δ **only decreases**: it is not reset to its initial value after each success (on Rosenbrock, resetting Δ_0 after each successful exploration: $n_f \approx 20\,000$ to reach $f < 10^{-4}$, against 627).
- Recommended value: $R = 2$ (Deb, 2012).
- Cost: each pattern move costs $1 + 2n$ evaluations (the point $\mathbf{x}_{k+1}^P$ and the new exploration).
- In the original version (Hooke and Jeeves, 1961), when the pattern fails one first explores around $\mathbf{x}_k$ with the same Δ ; the course code has both versions.

Where it fails, and the four methods on Rosenbrock

- **The stopping test can be fooled:** if a narrow valley passes between the points $\mathbf{x} \pm \Delta_i \mathbf{e}_i$, the exploration fails along all axes although a descent direction exists. *Possible heuristic:* add a few random directions — it helps, but does not solve the problem of directional coverage.
- **Guarantee:** the coordinate exploration of HJ is the origin of the pattern search framework; formalized versions such as GPS (Torczon, 1997) make it possible to establish stationarity results under suitable assumptions (e.g. $f \in C^1$ and bounded level set $\Rightarrow \liminf \|\nabla f(\mathbf{x}_k)\| = 0$). With a finite step, stopping does not certify stationarity: it only says that none of the $2n$ directions improved with that step. HJ already uses a positive basis ($\pm \mathbf{e}_j$); modern pattern search generalizes it.
- In curved valleys it zigzags: successful pattern, failure, new exploration with a reduced step.



Rosenbrock from $(-1.5, 2)$	n_f to $f < 10^{-4}$
Nelder-Mead (edges 0.5)	244
Hooke-Jeeves ($\Delta_0 = 0.5, R = 2$)	627
localized random [†]	975 [867, 1052]
Box ($\Delta_0 = 1$)	13 445

[†] median [Q1, Q3], 30 runs. On this problem, with these parameters and this criterion. With the standard versions (no $f(\mathbf{x}_c)$; HJ of 1961): 166 and 430.

On the computer: from Hooke–Jeeves to modern pattern search

The core, in MATLAB:

```
for i = 1:n                % exploration
    xp = x; xp(i) = xp(i)+D(i); fp = f(xp); % f+
    xm = x; xm(i) = xm(i)-D(i); fm = f(xm); % f-
    if fp < fx && fp <= fm, x = xp; fx = fp; % best
    elseif fm < fx,      x = xm; fx = fm; end
end
xP = 2*xk - xkm1;          % pattern move
```

The full functions (Exploratory, HookeJeeves, with info.nfev) are in the course companion code.

In practice, the variables should be scaled and the Δ_i chosen relative to the scale of each variable (one may be in 10^{-3} m, another in 10^5 Pa).

patternsearch (MATLAB)

A modern pattern search (positive bases with $2n$ or $n + 1$ directions), which generalizes classical Hooke–Jeeves, with convergence theory (GPS) under suitable assumptions.

Hooke–Jeeves (the origin) → modern
pattern search
→ directional direct search
(poll, local component of GLODS, 5.1)
→ DMS, multiobjective (6.5)

Summary table — 3.2

Method	Information	Cost	Guarantee	Main limitation
Random (3.2.1)	values of f	1 per sample	global, in probability, as $N \rightarrow \infty$ (full support, f continuous)	$\ x - x^*\ \sim N^{-1/n}$
Grid (3.2.2)	values of f	m^n in total	$f_{\text{grid}} \rightarrow f^*$ (f continuous); $f_{\text{grid}} - f^* \leq L \frac{\sqrt{n}}{2} h$ (f L -Lipschitz)	exponential in n
Nelder–Mead (3.2.3)	values of f	2–3 per iteration (+ n if it shrinks)	no general one (McKinnon, 1998)	may stagnate
Box (3.2.4)	values of f	2^n per iteration	no general guarantee presented in this course; a small box does not certify stationarity at finite Δ	2^n ; only diagonals
Hooke–Jeeves (3.2.5)	values of f	$2n$ per exploration	formalized versions (GPS): stationarity in the limit, under suitable assumptions	axes; zigzag

random + grid = **explore**

Nelder–Mead + Box + Hooke–Jeeves = **refine locally**

NM has no general guarantee; HJ is the origin of pattern search, whose formalized versions (GPS) have stationarity in the limit under suitable assumptions (3.2.5); for Box we present no general guarantee. Only evaluations of f : the direction is inferred, and that costs evaluations; derivative-free $\neq$ robust to noise.

KEY TAKEAWAYS — 3.2.5

1. Hooke–Jeeves alternates axis-by-axis **exploration** ($\pm\Delta_i$ in each coordinate, keep the best of the three points; $2n$ evaluations) and **pattern move** ($\mathbf{x}_{k+1}^P = 2\mathbf{x}_k - \mathbf{x}_{k-1}$).
2. The pattern **accelerates** and the exploration **corrects**: the path curves and follows oblique valleys, but zigzags in curved valleys.
3. The steps Δ only decrease ($\Delta = \Delta/R$), when the exploration or the pattern fails; stop when $\|\Delta\| < \varepsilon$.
4. The coordinate exploration of HJ is the origin of pattern search; formalized versions (GPS) have **stationarity in the limit, under suitable assumptions**; with a finite step, stopping does not certify stationarity.
5. Hooke–Jeeves $\rightarrow$ modern pattern search $\rightarrow$ GLODS poll (5.1) $\rightarrow$ DMS (6.5).

Next (3.3): so far we have inferred directions only through evaluations of f , and that costs evaluations. **What if f is smooth and we have ∇f ?** — derivative-based methods.

References and resources

References Full list: *Course bibliography*, at the end of the course PDF.

- Hooke, R., Jeeves, T. A., “*Direct search*” solution of numerical and statistical problems, Journal of the ACM 8(2):212–229, 1961.
- Torczon, V., *On the convergence of pattern search algorithms*, SIAM Journal on Optimization 7(1):1–25, 1997.
- Conn, A. R., Scheinberg, K., Vicente, L. N., *Introduction to Derivative-Free Optimization*, SIAM (MPS-SIAM Series on Optimization 8), 2009 — Ch. 7.
- Deb, K., *Optimization for Engineering Design: Algorithms and Examples*, 2nd ed., PHI Learning, 2012 — Sec. 3.3.3 (the algorithm used here).

Resources

- Companion code: Exploratory, HookeJeeves (`info.nfev`). MATLAB: `patternsearch` (Global Optimization Toolbox).