

## 6. Hypothesis testing and confidence intervals

We have the following goals:

- Understand how to construct confidence intervals for parameters in a linear model.
- Understand how to test statistical hypotheses about a linear model.
- In particular, to ask and answer the question: “Are the data consistent with a hypothesis that a covariate, or a collection of covariates, are unimportant?” (What is the fundamental scientific importance of the slightly contorted logical reasoning in this question?)
- Learn to use R to carry out these tasks.
- See how the linear model includes and extends basic tests for means of one and two samples.

# Confidence intervals

- An interval  $[u, v]$  constructed using the data  $\mathbf{y}$  is said to **cover** a parameter  $\theta$  if  $u \leq \theta \leq v$ .
- $[u, v]$  is a 95% **confidence interval** (CI) for  $\theta$  if the same construction, applied to a large number of draws from the model, would cover  $\theta$  95% of the time.
- A **parameter** is a name for any unknown constant in a model. In linear models, each component  $\beta_1, \dots, \beta_p$  of the **coefficient vector**  $\beta$  is a parameter. So is the variance  $\sigma^2$  of the measurement error.
- A confidence interval is the usual way to represent the amount of uncertainty in an estimated parameter.
- The parameter is not random. According to the model, it has a fixed but unknown value. The observed interval  $[u, v]$  is also not random. An interval  $[U, V]$  constructed using a vector of random variables  $\mathbf{Y}$  defined in a probability model is random.
- If the model is appropriate, then it is reasonable to treat the data  $\mathbf{y}$  like a realization from the probability model.

# A confidence interval for the coefficient of a linear model

- Consider estimating  $\beta_1$  in the linear model  $\mathbf{Y} = \mathbb{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$ .
- Recall that  $E[\hat{\beta}_1] = \beta_1$  and  $SD(\hat{\beta}_1) = \sigma \sqrt{[(\mathbb{X}^T \mathbb{X})^{-1}]_{11}}$ .

**Question 6.1.** Supposing we can make a normal approximation, show that  $P[\hat{\beta}_1 - 1.96 SD(b_1) \leq \beta_1 \leq \hat{\beta}_1 + 1.96 SD(b_1)] = 0.95$

- Therefore, an approximate 95% CI for  $\beta_1$  is

$$[b_1 - 1.96 SE(b_1), b_1 + 1.96 SE(b_1)]$$

where  $\mathbf{y} = \mathbb{X}\mathbf{b} + \mathbf{e}$  with  $SE(b_1) = s \sqrt{[(\mathbb{X}^T \mathbb{X})^{-1}]_{11}}$ .

# A CI for association between unemployment and mortality

```
c1 <- summary(lm(L_detrended~U_detrended))$coefficients ; c1

##              Estimate Std. Error  t value    Pr(>|t|)
## (Intercept) 0.2899928 0.09343146 3.103802 0.002812739
## U_detrended 0.1313673 0.06321939 2.077959 0.041606370

beta_U <- c1["U_detrended","Estimate"]
SE_U <- c1["U_detrended","Std. Error"]
z <- qnorm(1-0.05/2) # for a 95% CI using a normal approximation
cat("CI = [", beta_U - z * SE_U, ",", beta_U + z * SE_U, "]")

## CI = [ 0.0074596 , 0.2552751 ]
```

**Interpretation.** We appear to have found evidence that each percentage point of unemployment above trend is associated with about 0.13 years of additional life expectancy. The 95% CI doesn't include zero.

**Question 6.2.** Do you believe this discovery? How could you criticize it?

# Association is not causation

“Whatever phenomenon varies in any manner whenever another phenomenon varies in some particular manner, is either a cause or an effect of that phenomenon, or is connected with it through some fact of causation.” (*John Stuart Mill, A System of Logic, Vol. 1. 1843. p. 470.*)

- Put differently: If  $A$  and  $B$  are associated statistically, we can infer that either  $A$  causes  $B$ , or  $B$  causes  $A$ , or both have some common cause  $C$ .
- A useful mantra: **Association is not causation.**
- Writing a linear model where  $A$  depends on  $B$  can show association but we need extra work to argue for causation. We need to rule out  $B$  causing  $A$  and the possibility of any common cause  $C$ .

**Question 6.3.** Discuss the extent to which the association between detrended unemployment and life expectancy can and cannot be interpreted causally.

# A review of progress so far in this course

**Producing and understanding this confidence interval for a linear model brought together all the things we've done so far in this course.**

- We needed to get the data into a computer and run statistical software.
- To understand what the computer was doing for us, and help us to command it correctly, we needed to know about:
  - ① matrices
  - ② writing a linear model and fitting it by least squares
  - ③ probability models
  - ④ expectation and variance
  - ⑤ the normal distribution

**You could run computer code by learning to follow line-by-line instructions without understanding what the instructions do. But then you wouldn't be in control of your own data analysis.**

# Hypothesis tests

- We try to see patterns in our data. We hope to discover phenomena that will advance science, or help the environment, or reduce sickness and poverty, or make us rich, ...
- How can we tell whether our new theory is like seeing animals or faces in the clouds?
- From Wikipedia: “**Pareidolia** is a psychological phenomenon in which the mind responds to a stimulus ... by perceiving a familiar pattern where none exists (e.g. in random data)”.
- The research community has set a standard: The evidence presented to support a new theory should be unlikely under a **null hypothesis** that the new theory is false. To quantify *unlikely* we need a probability model.

# Hypothesis tests and the scientific method

- From a different perspective, a standard view of scientific progress holds that scientific theories cannot be proved correct, they can only be falsified (<https://en.wikipedia.org/wiki/Falsifiability>).
- Accordingly, scientists look for evidence to refute the **null hypothesis** that data can be explained by current scientific understanding.
- If the null hypothesis is inadequate to explain data, the scientist may propose an **alternative hypothesis** which better explains these data.
- The alternative hypothesis will subsequently be challenged with new data.



# The scientific method in statistical language

- 1 **Ask a question**
- 2 **Obtain relevant data.**
- 3 **Write a null and alternative hypothesis to represent your question in a probability model.** This may involve writing a linear model so that  $\beta_1 = 0$  corresponds to the null hypothesis of “no effect” and  $\beta_1 \neq 0$  is a discovered “effect.”
- 4 **Calculate a test statistic.** The test statistic is a quantity computed using the data that summarizes the evidence against the null hypothesis. For our linear model example, the least squares coefficient  $b_1$  is a natural statistics to test the hypothesis  $\beta_1 = 0$ .
- 5 **Calculate the p-value**, which is the probability that the model generates a test statistic at least as extreme as that observed. For our linear model example, the p-value is  $P[|\hat{\beta}_1| > |b_1|]$ . We can find this probability, when  $\beta_1 = 0$ , using a normal approximation.
- 6 **Conclusions.** A small p-value (often,  $< 0.05$ ) is evidence for **rejecting** the null hypothesis. The data analysis may suggest new questions: **Return to Step 1.**

# Using confidence intervals to construct a hypothesis test

- It is often convenient to use the confidence interval as a test statistic.
- If the confidence interval doesn't cover the null hypothesis, then we have evidence to reject that null hypothesis.
- If we do this test using a 95% confidence interval, we have a 5% chance that we reject the null hypothesis if it is true. This follows from the definition of a confidence interval: whatever the true unknown value of the parameter, the confidence interval covers it with probability 0.95.

## Some notation for hypothesis tests

- The null hypothesis is  $H_0$  and the alternative is  $H_a$ .
- We write  $t$  for the test statistic calculated using the data  $\mathbf{y}$ . We write  $T$  for the random variable constructed by calculating the test statistic using a random vector  $\mathbf{Y}$  drawn from the probability model under  $H_0$ .
- The p-value is  $\text{pval} = P[|T| \geq |t|]$ . Here, we are assuming “extreme” means “large in magnitude.” Occasionally, it may make more sense to use  $\text{pval} = P[T \geq t]$ .
- We reject  $H_0$  at **significance level**  $\alpha$  if  $\text{pval} < \alpha$ . Common choices of  $\alpha$  are  $\alpha = 0.05$ ,  $\alpha = 0.01$ ,  $\alpha = 0.001$ .

**Question 6.4.** When we report the results of a hypothesis test, we can either (i) give the p-value, or (ii) say whether  $H_0$  is rejected at a particular significance level. What are the advantages and disadvantages of each?

## A hypothesis test for unemployment and mortality

**Question 6.5.** Write a formal hypothesis test of the null hypothesis that there is no association between unemployment and mortality. Compute a p-value using a normal approximation. What do you think is an appropriate significance level  $\alpha$  for deciding whether to reject the null hypothesis?

# Normal approximations versus Student's t distribution

- Notice that `'summary(lm(...))'` gives 't value' and `'Pr(>|t|)'`.
- The 't value' is the estimated coefficient divided by its standard error. This measures how many standard error units the estimated coefficient is from zero.
- `'Pr(>|t|)'` is similar, but slightly larger, than the p-value coming from the normal approximation.
- R is using Student's t distribution, which makes allowance for chance variation from using  $s$  as an approximation to  $\sigma$  when we compute the standard error.
- R uses a t random variable to model the distribution of the statistic  $t$ . Giving the full name (Student's t distribution) may add clarity.
- With sophisticated statistical methods, it is often hard to see if they work well just by reading about them. Fortunately, it is often relatively easy to do a **simulation study** to see what is going on.

# Simulating from Student's t distribution

- Suppose  $X, X_1, \dots, X_d$  are  $d + 1$  independent identically distributed (iid) normal random variables with mean zero and standard deviation  $\sigma$ .
- We write  $X, X_1, \dots, X_d \sim \text{iid } N[0, \sigma]$ .
- Student's t distribution on  $d$  degrees of freedom is defined to be the distribution of  $T = X/\hat{\sigma}$  where  $\hat{\sigma} = \sqrt{\frac{1}{d} \sum_{i=1}^d X_i^2}$ .
- A normal approximation would say  $T$  is approximately  $N[0, 1]$  since  $\hat{\sigma}$  is an estimate of  $\sigma$ .
- With a computer, we can simulate  $T$  many times, plot a histogram, and compare it to the probability density function of the normal distribution and Student's t distribution.
- The goals in doing this:
  - ① Some practice working with Student's t distribution.
  - ② Finding how the t distribution compares to the normal distribution as  $d$  varies.
  - ③ Practice the skill of designing a simulation experiment.

- Let's start by simulating a matrix  $X$  of iid normal random variables.

```
N <- 50000 ; sigma <- 1 ; d <- 10 ; set.seed(23)
X <- matrix(rnorm(N*(d+1),mean=0,sd=sigma),nrow=N)
```

- Now, we write a function that computes  $T$  given  $X_1, \dots, X_d, X$


```
T_evaluator <- function(x) x[d+1] / sqrt(sum(x[1:d]^2)/d)
```

- Then, use `apply()` to evaluate  $T$  on each row of 'X'.

```
Tsim <- apply(X,1,T_evaluator)
```

- A histogram of these simulations can be compared with the normal density and the t density

```
hist(Tsim,freq=F,main="",
     breaks=30,ylim=c(0,0.4))
x <- seq(length=200,
     min(Tsim),max(Tsim))
lines(x,dnorm(x),
     col="blue",
     lty="dashed")
lines(x,dt(x,df=d),
     col="red")
```



# Comparing the normal and t distributions

- Even with as few as  $d = 10$  degrees of freedom to estimate  $\sigma$ , the Student's t density looks similar to the normal density.
- Student's t has fatter tails. This is important for the probability of rare extreme outcomes.
- Here, the largest and smallest of the  $N = 5 \times 10^4$  simulations are

```
range(Tsim)
## [1] -6.438830  6.480262
```

- Let's check the chance of an outcome more than 5 (or 6) standard deviations from the mean for the normal distribution and the t on 10 degrees of freedom.

```
2*(1-pnorm(5))
## [1] 5.733031e-07
2*(1-pnorm(6))
## [1] 1.973175e-09
```

```
2*(1-pt(5,df=d))
## [1] 0.0005373336
2*(1-pt(6,df=d))
## [1] 0.0001321089
```