

Olympiad Test Paper — Mathematics (Class 7) — Paper 3

Chapter 12: Another Peek Beyond the Point

Paper 3 — (progressively harder)

Subject	Mathematics (NCERT Class 7)
Chapter	12 — Another Peek Beyond the Point
Total Questions	60 (Multiple Choice, single correct)
Maximum Marks	240
Suggested Time	90 minutes

Marking scheme: +4 for each correct answer · -1 for each wrong answer · 0 if unattempted.

Instructions: Each question has four options (A), (B), (C), (D); exactly one is correct. The two commonest traps are misplacing the decimal point in a product (counting too few or too many decimal places) and, in division by a decimal, scaling only one of the two numbers instead of both by the same power of ten. Do not write on this paper — mark answers on your answer sheet.

1. Compute $1.2 \times 0.15 \times 5$.

- | | |
|---------|----------|
| (A) 9 | (B) 0.09 |
| (C) 0.9 | (D) 0.18 |

2. Compute $7.2 \div 0.08$.

- | | |
|---------|---------|
| (A) 90 | (B) 9 |
| (C) 900 | (D) 0.9 |

3. Compute $0.46 \times 1000 \div 100$.

- | | |
|----------|---------|
| (A) 4.6 | (B) 46 |
| (C) 0.46 | (D) 460 |

4. Estimate first, then give the exact value of 4.9×3.1 .

- | | |
|-----------|-----------|
| (A) 1.519 | (B) 151.9 |
| (C) 15.9 | (D) 15.19 |

5. Write 0.375 as a fraction in lowest terms.

- | | |
|-------------|----------|
| (A) 375/100 | (B) 3/80 |
| (C) 37/100 | (D) 3/8 |

6. By what single power of ten must 0.0042 be multiplied to give 42?

- | | |
|----------|------------|
| (A) 1000 | (B) 10000 |
| (C) 100 | (D) 100000 |

7. Compute $2.5 \times 0.4 \times 0.3$.

- | | |
|----------|----------|
| (A) 3 | (B) 0.03 |
| (C) 0.12 | (D) 0.3 |

8. Compute $0.144 \div 0.012$.

- | | |
|----------|---------|
| (A) 1.2 | (B) 120 |
| (C) 0.12 | (D) 12 |

9. Compute $56.7 \div 1000$.

- | | |
|------------|-------------|
| (A) 0.567 | (B) 0.00567 |
| (C) 0.0567 | (D) 56700 |

10. Estimate first, then give the exact value of $0.612 \div 0.3$.

- | | |
|----------|-----------|
| (A) 20.4 | (B) 0.204 |
| (C) 2.04 | (D) 2.4 |

11. Express $7/8$ as a decimal.

- | | |
|------------|-----------|
| (A) 0.78 | (B) 0.875 |
| (C) 0.0875 | (D) 1.142 |

12. In the number 34.706, what is the place value of the digit 6?

- | | |
|-------------------------|---------------------------|
| (A) 6 hundredths (0.06) | (B) 6 thousandths (0.006) |
| (C) 6 tenths (0.6) | (D) 6 ones (6) |

13. Compute 1.25×0.16 .

- | | |
|-----------|----------|
| (A) 2 | (B) 0.02 |
| (C) 0.002 | (D) 0.2 |

14. Compute $9 \div 0.036$.

- | | |
|----------|---------|
| (A) 25 | (B) 250 |
| (C) 2500 | (D) 2.5 |

15. Compute 0.0035×10000 .

- | | |
|---------|----------|
| (A) 35 | (B) 3.5 |
| (C) 350 | (D) 0.35 |

16. Estimate first, then give the exact value of 19.8×0.51 .

- | | |
|------------|------------|
| (A) 100.98 | (B) 1.0098 |
| (C) 10.98 | (D) 10.098 |

17. Which fraction equals 0.04?

- (A) $\frac{1}{25}$ (B) $\frac{1}{4}$
(C) $\frac{4}{10}$ (D) $\frac{1}{250}$

18. A product was computed as 8.4. Which of the following correctly produces 8.4?

- (A) 2.4×3.5 (B) 24×3.5
(C) 0.24×3.5 (D) 2.4×35

19. Compute $(0.3)^3$.

- (A) 0.27 (B) 0.027
(C) 0.0027 (D) 0.9

20. Compute $4.8 \div 0.015$.

- (A) 32 (B) 3200
(C) 320 (D) 3.2

21. Compute $0.05 \times 1000 \div 10$.

- (A) 50 (B) 5
(C) 0.5 (D) 500

22. Estimate first, then give the exact value of $35.04 \div 0.6$.

- (A) 5.84 (B) 58.4
(C) 584 (D) 5840

23. Which decimal equals $\frac{9}{200}$?

- (A) 0.45 (B) 0.045
(C) 0.0045 (D) 4.5

24. In 5.28, by how much does the value change if the digit 2 is replaced by 7?

- (A) increases by 0.5 (B) increases by 5
(C) increases by 0.05 (D) increases by 0.7

25. Sugar costs ₹48.50 per kg. A shopper buys 1.5 kg and pays with a ₹100 note. The change received is:

- (A) ₹72.75 (B) ₹2.725
(C) ₹51.50 (D) ₹27.25

26. Pens cost ₹12.50 each. With ₹100, what is the greatest number of whole pens that can be bought, and how much is left?

- (A) 8 pens, ₹12.50 left (B) 7 pens, ₹12.50 left
(C) 80 pens, ₹0 left (D) 8 pens, ₹0 left

27. A plank is 2.4 m long. Three pieces of lengths 0.8 m, 0.65 m and 0.45 m are cut off. The remaining length is:

- (A) 0.5 m (B) 1.9 m
(C) 0.05 m (D) 0.55 m

28. A bag of rice weighs 25.5 kg. It is repacked equally into packets of 0.85 kg each. The number of packets is:

- | | |
|--------|------------|
| (A) 3 | (B) 300 |
| (C) 30 | (D) 21.675 |

29. Compute $(2.5 + 1.5) \times 0.6$.

- | | |
|----------|---------|
| (A) 24 | (B) 2.4 |
| (C) 0.24 | (D) 4.6 |

30. Compute $0.6 \div 0.4 + 0.5$.

- | | |
|----------|--------|
| (A) 1.1 | (B) 2 |
| (C) 0.65 | (D) 11 |

31. Compute $0.25 \times 0.25 \times 16$.

- | | |
|--------|----------|
| (A) 1 | (B) 0.1 |
| (C) 10 | (D) 0.01 |

32. Compute $1.25 \div 0.005$.

- | | |
|---------|----------|
| (A) 25 | (B) 2500 |
| (C) 2.5 | (D) 250 |

33. Compute $12.5 \times 100 \div 1000$.

- | | |
|----------|-----------|
| (A) 12.5 | (B) 1.25 |
| (C) 125 | (D) 0.125 |

34. Estimate first, then give the exact value of $6.93 \div 2.1$.

- | | |
|----------|----------|
| (A) 3.3 | (B) 33 |
| (C) 0.33 | (D) 3.03 |

35. Which of these is equal to 0.6?

- | | |
|-------------------|----------------------|
| (A) $\frac{3}{5}$ | (B) $\frac{6}{1000}$ |
| (C) $\frac{1}{6}$ | (D) $\frac{3}{50}$ |

36. A number written as 4.072 is multiplied by 100. In the result, which digit is in the tens place?

- | | |
|-------|-------|
| (A) 4 | (B) 7 |
| (C) 2 | (D) 0 |

37. Ink costs ₹2.40 per millilitre. A bottle holds 0.75 litre (= 750 mL). The bottle's contents are worth:

- | | |
|-----------|------------|
| (A) ₹180 | (B) ₹18000 |
| (C) ₹1.80 | (D) ₹1800 |

38. A roll of tape is 5.4 m long. After using 1.85 m, the rest is cut into 0.5 m strips. How many full 0.5 m strips can be cut?

(A) 10

(B) 8

(C) 3

(D) 7

39. A length of wire weighs 0.045 kg per metre. A coil weighs 1.8 kg. The coil's length is:

(A) 4 m

(B) 40 m

(C) 400 m

(D) 0.081 m

40. Compute $3.6 \div (0.4 \times 0.5)$.

(A) 1.8

(B) 18

(C) 180

(D) 0.18

41. Compute $2.4 \times 0.5 + 1.2 \times 0.5$.

(A) 1.8

(B) 18

(C) 0.18

(D) 2.16

42. Compute $(1.5 \div 0.3) \times 0.04$.

(A) 2

(B) 0.02

(C) 0.2

(D) 0.5

43. Compute 0.025×0.4 .

(A) 0.1

(B) 0.001

(C) 0.01

(D) 0.0001

44. Compute $0.084 \div 0.0021$.

(A) 4

(B) 40

(C) 400

(D) 0.4

45. Compute $0.7 \times 1000 \div 100 \times 10$.

(A) 7

(B) 700

(C) 70

(D) 0.7

46. Estimate first, then give the exact value of $0.396 \div 0.012$.

(A) 3.3

(B) 330

(C) 33

(D) 0.33

47. If 2.5 metres of cloth cost ₹93.75, what is the cost of 4 metres at the same rate?

(A) ₹15

(B) ₹1500

(C) ₹37.50

(D) ₹150

48. A tank holds 7.2 litres. A jug holds 0.45 litre. How many full jugs can be filled from the tank, and how much is left?

(A) 16 jugs, 0.45 litre left

(B) 1.6 jugs

(C) 160 jugs

(D) 16 jugs, 0 litres left

49. A nail is 0.035 m long. Laid end to end, how many nails make a line 1.4 m long?

- (A) 4 (B) 40
(C) 400 (D) 0.049

50. A medicine dose is 0.015 kg. A jar holds 0.6 kg of the medicine. The number of doses in the jar is:

- (A) 4 (B) 400
(C) 40 (D) 0.009

51. Compute $(0.6 + 0.4) \div 0.25$.

- (A) 0.4 (B) 40
(C) 4 (D) 0.04

52. Compute $4.2 \times 0.5 \div 0.07$.

- (A) 3 (B) 300
(C) 30 (D) 0.3

53. Tickets cost ₹37.50 each. A group buys tickets for ₹300 total. After the show they share ₹45 of snacks equally among the ticket-holders. Each person's snack share is:

- (A) ₹0.5625 (B) ₹56.25
(C) ₹5.65 (D) ₹5.625

54. A ribbon 1.5 m long is cut into 6 equal pieces. Each piece is then halved. The length of one final small piece is:

- (A) 0.25 m (B) 1.25 m
(C) 0.125 m (D) 0.0125 m

55. A sack of flour weighs 4.8 kg. One-third is used, and the rest is split into 0.4 kg bags. How many full 0.4 kg bags result?

- (A) 8 (B) 12
(C) 4 (D) 32

56. Compute $(0.9 \times 0.8) \div (0.6 \times 0.2)$.

- (A) 0.6 (B) 60
(C) 6 (D) 0.06

57. A pipe leaks 0.025 litre per minute. The water lost in 1.2 hours is:

- (A) 1.8 litres (B) 0.18 litre
(C) 18 litres (D) 0.03 litre

58. A block of metal weighs 0.96 kg. It is melted and cast into beads each weighing 0.012 kg. The number of beads is:

- (A) 8 (B) 800
(C) 80 (D) 0.0115

59. Compute $(2.5 \times 0.4) \div (0.5 \times 0.2)$.

(A) 10

(C) 100

(B) 1

(D) 0.1

60. Compute $0.4 \times 0.5 \times 0.6 \times 100$.

(A) 12

(C) 120

(B) 1.2

(D) 0.12

Solutions — Mathematics (Class 7), Chapter 12:

Another Peek Beyond the Point — Paper 3

Paper 3 — (progressively harder)

Marking: +4 correct, -1 wrong, 0 unattempted. Maximum marks **240**.

Answer Key

Q	Ans	Q	Ans	Q	Ans	Q	Ans	Q	Ans	Q	Ans
1	C	11	B	21	B	31	A	41	A	51	C
2	A	12	B	22	B	32	D	42	C	52	C
3	A	13	D	23	B	33	B	43	C	53	D
4	D	14	B	24	A	34	A	44	B	54	C
5	D	15	A	25	D	35	A	45	C	55	A
6	B	16	D	26	D	36	D	46	C	56	C
7	D	17	A	27	A	37	D	47	D	57	A
8	D	18	A	28	C	38	D	48	D	58	C
9	C	19	B	29	B	39	B	49	B	59	A
10	C	20	C	30	B	40	B	50	C	60	A

Answer-key distribution: A = 15, B = 15, C = 15, D = 15.

Worked Solutions

1. (C) Multiply step by step. 1.2×0.15 : digits $12 \times 15 = 180$; one decimal place plus two decimal places makes three, so $0.180 = 0.18$. Then $0.18 \times 5 = 0.90 = 0.9$. Reading the digits 9 with no decimal places, or counting only two places to get 0.09, both misplace the point; 0.18 is only the first product before multiplying by 5.

2. (A) Scale BOTH numbers by 100 so the divisor becomes a whole number: $7.2 \times 100 = 720$ and $0.08 \times 100 = 8$. Then $720 \div 8 = 90$. If you scale only the divisor (8) but leave 7.2, you get 0.9; scaling only 7.2 to 720 against 0.08 mishandled gives 9 or 900. Scaling both by the same power of ten is the rule.

3. (A) 0.46×1000 shifts the point three places right: 460. Then $\div 100$ shifts it two places left: $4.60 = 4.6$. Net effect is shifting one place right, which also gives 4.6. Stopping at 460 forgets the division; leaving 0.46 cancels the shifts wrongly; 46 shifts only one place for the multiply.

- 4. (D)** Estimate: about $5 \times 3 = 15$, so the answer is near 15. Exactly: $49 \times 31 = 1519$; one decimal place in each factor makes two, so 15.19. The estimate rules out 1.519 (near 1.5) and 151.9 (near 150). 15.9 comes from a careless digit multiplication, not 49×31 .
- 5. (D)** $0.375 = 375/1000$. Divide top and bottom by 125: $375 \div 125 = 3$ and $1000 \div 125 = 8$, giving $3/8$. The value $375/100$ misreads three decimal places as two; $3/80$ over-divides the denominator; $37/100$ drops a digit.
- 6. (B)** Moving from 0.0042 to 42 shifts the decimal point four places to the right ($0.0042 \rightarrow 0.042 \rightarrow 0.42 \rightarrow 4.2 \rightarrow 42$). Four places right means $\times 10000$. Three places ($\times 1000$) gives 4.2; two places ($\times 100$) gives 0.42; five places ($\times 100000$) gives 420.
- 7. (D)** Take 2.5×0.4 first: digits $25 \times 4 = 100$, one decimal place plus one decimal place is two places, so $1.00 = 1$. Then $1 \times 0.3 = 0.3$. Reading 0.12 stops after a wrong pairing; 3 and 0.03 misplace the point around the value 1×0.3 .
- 8. (D)** Scale both numbers by 1000 to clear the divisor: $0.144 \times 1000 = 144$ and $0.012 \times 1000 = 12$. Then $144 \div 12 = 12$. Scaling only one number, or stopping the point shift early, produces 1.2, 120, or 0.12.
- 9. (C)** Dividing by 1000 shifts the decimal point three places to the left: $56.7 \rightarrow 5.67 \rightarrow 0.567 \rightarrow 0.0567$. Only two places gives 0.567; four places gives 0.00567; multiplying instead of dividing gives 56700.
- 10. (C)** Estimate: $0.6 \div 0.3 = 2$, so the answer is near 2. Scale both by 10: $6.12 \div 3 = 2.04$. The estimate near 2 rejects 20.4 and 0.204. 2.4 comes from dividing 0.6 by 0.3 carelessly while ignoring the 12 thousandths.
- 11. (B)** $7 \div 8 = 0.875$ (8 into 70 is 8 remainder 6; 60 gives 7 remainder 4; 40 gives 5). The value 0.78 misreads the long division; 0.0875 shifts the point one place too far left; 1.142 is $8 \div 7$, the reciprocal direction.
- 12. (B)** After the point the places are tenths, hundredths, thousandths. Here 7 is tenths, 0 is hundredths, 6 is thousandths, so 6 means 0.006. Calling it 0.06 or 0.6 counts too few places past the point; 6 ones ignores the decimal point entirely.
- 13. (D)** Digits: $125 \times 16 = 2000$. Factors have two and two decimal places, four in all, so place the point four places: $0.2000 = 0.2$. Reading 2 ignores the point; counting too few or too many places gives 0.02 or 0.002. (Trailing zeros in 2000 are dropped after placing the point.)
- 14. (B)** Scale both by 1000: $9 \times 1000 = 9000$ and $0.036 \times 1000 = 36$. Then $9000 \div 36 = 250$. Scaling only the divisor, or shifting the point one place wrong, gives 25, 2500, or 2.5.

15. (A) Multiplying by 10000 shifts the point four places right: $0.0035 \rightarrow 0.035 \rightarrow 0.35 \rightarrow 3.5 \rightarrow 35$. Three places gives 3.5; five would give 350; two gives 0.35.

16. (D) Estimate: $20 \times 0.5 = 10$, so the answer is near 10. Exactly: $198 \times 51 = 10098$; one plus two decimal places make three, so 10.098. The estimate near 10 rejects 100.98 and 1.0098. 10.98 comes from a slip in the digit product.

17. (A) $0.04 = 4/100 = 1/25$ (divide top and bottom by 4). The value $1/4 = 0.25$, far too big; $4/10 = 0.4$, ten times too big; $1/250 = 0.004$, ten times too small.

18. (A) 2.4×3.5 : digits $24 \times 35 = 840$; one plus one decimal place makes two, so $8.40 = 8.4$. $24 \times 3.5 = 84$ (ten times bigger), $0.24 \times 3.5 = 0.84$ (ten times smaller), $2.4 \times 35 = 84$. Only the first lands the point to give 8.4.

19. (B) $(0.3)^3 = 0.3 \times 0.3 \times 0.3$. $0.3 \times 0.3 = 0.09$ (digits 9, two decimal places). Then $0.09 \times 0.3 = 0.027$ (digits 27, three decimal places). 0.27 counts only two places; 0.0027 counts four; 0.9 is 0.3×3 , not the cube.

20. (C) Scale both by 1000: $4.8 \times 1000 = 4800$ and $0.015 \times 1000 = 15$. Then $4800 \div 15 = 320$. Scaling only one number, or stopping the point shift early, gives 32, 3200, or 3.2.

21. (B) 0.05×1000 shifts the point three places right: 50. Then $\div 10$ shifts one place left: $5.0 = 5$. Net shift is two places right, also 5. Stopping at 50 forgets the divide; 0.5 over-divides; 500 forgets the divide and over-multiplies.

22. (B) Estimate: $36 \div 0.6 = 60$, so the answer is near 60. Scale both by 10: $350.4 \div 6 = 58.4$. The estimate near 60 rejects 5.84, 584 and 5840, which all misplace the point. So 58.4 is correct.

23. (B) $9/200$: multiply top and bottom by 5 to get $45/1000 = 0.045$. The value $0.45 = 9/20$, ten times too big; $0.0045 = 9/2000$, ten times too small; $4.5 = 9/2$.

24. (A) The digit 2 sits in the tenths place, so it stands for 0.2; replacing it with 7 makes that place 0.7. The change is $0.7 - 0.2 = 0.5$. Treating tenths as ones gives 5; as hundredths gives 0.05; 0.7 ignores the value already there.

25. (D) Cost = $48.50 \times 1.5 = 72.75$ (digits $485 \times 15 = 7275$, two decimal places). Change = $100 - 72.75 = 27.25$. Giving ₹72.75 reports the cost, not the change; ₹2.725 misplaces the point; ₹51.50 is $100 - 48.50$, ignoring the 1.5 kg.

26. (D) $100 \div 12.50$: scale both by 10 to get $1000 \div 125 = 8$ exactly. So 8 pens cost $8 \times 12.50 = 100$, leaving ₹0. Buying 7 leaves ₹12.50 (one more pen affordable); 80 pens misplaces the point in the division.

27. (A) Total cut = $0.8 + 0.65 + 0.45 = 1.9$ m. Remaining = $2.4 - 1.9 = 0.5$ m. Reporting 1.9 gives the cut length, not what remains; 0.05 misplaces the point; 0.55 comes from a

subtraction slip.

28. (C) $25.5 \div 0.85$: scale both by 100 to get $2550 \div 85 = 30$. Scaling only one number, or shifting the point wrong, gives 3 or 300. The value 21.675 is 25.5×0.85 (multiplying instead of dividing).

29. (B) Do the bracket first: $2.5 + 1.5 = 4.0$. Then $4.0 \times 0.6 = 2.4$. Skipping the bracket or misplacing the point gives 24 or 0.24; 4.6 mistakenly multiplies only the 1.5 by 0.6 and adds.

30. (B) Division first: $0.6 \div 0.4$, scale both by 10 to get $6 \div 4 = 1.5$. Then $1.5 + 0.5 = 2$. Adding before dividing gives 1.1; misplacing the quotient gives 0.65 or 11.

31. (A) $0.25 \times 0.25 = 0.0625$ (digits 625, four decimal places). Then $0.0625 \times 16 = 1.0 = 1$. Misplacing the point along the way gives 0.1, 10, or 0.01.

32. (D) Scale both by 1000: $1.25 \times 1000 = 1250$ and $0.005 \times 1000 = 5$. Then $1250 \div 5 = 250$. Scaling only one number, or stopping the point shift early, gives 25, 2500, or 2.5.

33. (B) 12.5×100 shifts the point two places right: 1250. Then $\div 1000$ shifts three places left: $1.250 = 1.25$. Net shift is one place left, also 1.25. Cancelling wrongly gives 12.5; stopping at 125 forgets a place; over-shifting gives 0.125.

34. (A) Estimate: $7 \div 2 = 3.5$, so the answer is near 3.5. Scale both by 10: $69.3 \div 21 = 3.3$. The estimate near 3.5 rejects 33 and 0.33. 3.03 comes from a long-division slip; 3.3 is correct.

35. (A) $0.6 = 6/10 = 3/5$. The value $6/1000 = 0.006$; $1/6 \approx 0.167$; $3/50 = 0.06$. Only $3/5$ reduces from $6/10$.

36. (D) $4.072 \times 100 = 407.2$ (point shifts two places right). In 407.2 the digits are 4 hundreds, 0 tens, 7 ones, 2 tenths. The tens digit is 0. Picking 4 reads hundreds; 7 reads ones; 2 reads tenths.

37. (D) $750 \text{ mL} \times ₹2.40 = 750 \times 2.4 = 1800$ (digits $750 \times 24 = 18000$, one decimal place gives 1800.0). So ₹1800. Misplacing the point gives ₹180, ₹18000, or ₹1.80.

38. (D) Remaining tape = $5.4 - 1.85 = 3.55 \text{ m}$. $3.55 \div 0.5 = 7.1$, so 7 full strips (the 0.1 leftover is not a full strip). Forgetting the subtraction gives 10 ($5.4 \div 0.5 = 10.8$); rounding up wrongly gives 8; using only 1.85 gives 3.

39. (B) Length = total weight $\div$ weight per metre = $1.8 \div 0.045$. Scale both by 1000: $1800 \div 45 = 40 \text{ m}$. Scaling only one number gives 4 or 400; 0.081 is 1.8×0.045 (multiplying instead of dividing).

- 40. (B)** Bracket first: $0.4 \times 0.5 = 0.20$. Then $3.6 \div 0.2$: scale both by 10 to get $36 \div 2 = 18$. Misplacing the point gives 1.8, 180, or 0.18.
- 41. (A)** $2.4 \times 0.5 = 1.2$ and $1.2 \times 0.5 = 0.6$. Sum = $1.2 + 0.6 = 1.8$. (Or factor: $(2.4 + 1.2) \times 0.5 = 3.6 \times 0.5 = 1.8$.) Misplacing the point gives 18 or 0.18; 2.16 multiplies the two products instead of adding.
- 42. (C)** Bracket first: $1.5 \div 0.3$, scale both by 10 to get $15 \div 3 = 5$. Then $5 \times 0.04 = 0.20 = 0.2$. Misplacing the point gives 2 or 0.02; 0.5 mishandles the division.
- 43. (C)** Digits: $25 \times 4 = 100$. Factors have three and one decimal place, four in all, so place the point four places: $0.0100 = 0.01$. Counting three places gives 0.1; five places gives 0.001; reading too far gives 0.0001.
- 44. (B)** Scale both by 10000: $0.084 \times 10000 = 840$ and $0.0021 \times 10000 = 21$. Then $840 \div 21 = 40$. Scaling only one number, or shifting the point wrong, gives 4, 400, or 0.4.
- 45. (C)** Work left to right. $0.7 \times 1000 = 700$. $700 \div 100 = 7$. $7 \times 10 = 70$. Net shift: $+3 - 2 + 1 = +2$ places right, so $0.7 \rightarrow 70$. Stopping early gives 7 or 700; cancelling all shifts wrongly gives 0.7.
- 46. (C)** Estimate: $0.4 \div 0.012 \approx 0.4 \div 0.01 = 40$, so the answer is in the tens. Scale both by 1000: $396 \div 12 = 33$. The estimate near 40 rejects 3.3 and 0.33. So 33 is correct.
- 47. (D)** Rate per metre = $93.75 \div 2.5$: scale both by 10 to get $937.5 \div 25 = 37.5$. Cost of 4 m = $37.5 \times 4 = 150$. So ₹150. ₹37.50 is just the per-metre rate; ₹15 and ₹1500 misplace the point.
- 48. (D)** $7.2 \div 0.45$: scale both by 100 to get $720 \div 45 = 16$ exactly. So 16 full jugs and nothing left. Misplacing the point gives 1.6 or 160; 16 with 0.45 left contradicts the exact division.
- 49. (B)** $1.4 \div 0.035$: scale both by 1000 to get $1400 \div 35 = 40$. Scaling only one number gives 4 or 400; 0.049 is 1.4×0.035 (multiplying instead of dividing).
- 50. (C)** $0.6 \div 0.015$: scale both by 1000 to get $600 \div 15 = 40$. Scaling only one number gives 4 or 400; 0.009 is 0.6×0.015 (multiplying instead of dividing).
- 51. (C)** Bracket first: $0.6 + 0.4 = 1.0$. Then $1 \div 0.25$: scale both by 100 to get $100 \div 25 = 4$. Misplacing the point gives 0.4, 40, or 0.04.
- 52. (C)** Left to right: $4.2 \times 0.5 = 2.1$. Then $2.1 \div 0.07$: scale both by 100 to get $210 \div 7 = 30$. Misplacing the point gives 3, 300, or 0.3.
- 53. (D)** Number of people = $300 \div 37.50$: scale both by 10 to get $3000 \div 375 = 8$. Snack share = $45 \div 8 = 5.625$. So ₹5.625. Misplacing the point gives ₹0.5625 or ₹56.25; ₹5.65 is

a division slip.

54. (C) Each of 6 pieces = $1.5 \div 6 = 0.25$ m. Halving: $0.25 \div 2 = 0.125$ m. Stopping after the first cut gives 0.25; misplacing the point gives 1.25 or 0.0125.

55. (A) One-third used means two-thirds remain: $(2/3) \times 4.8 = 3.2$ kg. Then $3.2 \div 0.4 = 8$ bags. Forgetting to subtract the used third gives 12 ($4.8 \div 0.4$); using only the used third (1.6 kg) gives 4; misplacing the point gives 32.

56. (C) Numerator: $0.9 \times 0.8 = 0.72$. Denominator: $0.6 \times 0.2 = 0.12$. Then $0.72 \div 0.12$: scale both by 100 to get $72 \div 12 = 6$. Misplacing the point gives 0.6, 60, or 0.06.

57. (A) 1.2 hours = $1.2 \times 60 = 72$ minutes. Loss = $72 \times 0.025 = 1.8$ litres (digits $72 \times 25 = 1800$, three decimal places give 1.800). Misplacing the point gives 0.18 or 18; 0.03 forgets to convert hours to minutes.

58. (C) $0.96 \div 0.012$: scale both by 1000 to get $960 \div 12 = 80$. Scaling only one number gives 8 or 800; 0.0115 is 0.96×0.012 (multiplying instead of dividing).

59. (A) Numerator: $2.5 \times 0.4 = 1.0$. Denominator: $0.5 \times 0.2 = 0.10$. Then $1 \div 0.1 = 10$. Misplacing the point gives 1, 100, or 0.1.

60. (A) $0.4 \times 0.5 = 0.20$; $0.20 \times 0.6 = 0.120 = 0.12$; $0.12 \times 100 = 12$. Forgetting the $\times 100$, or shifting the point wrong, gives 0.12, 1.2, or 120.