

Math 415 - Lecture 4

Linear Combinations and Matrix operations

Monday August 31 2015

Textbook: Chapter 1.3, 1.4

Textbook: Chapter 1.3, 1.4

Suggested Practice Exercise: Chapter 1.4 Exercise 1, 2, 10, 12, 13,
21, 30, 34, 45,

[Textbook](#): Chapter 1.3, 1.4

[Suggested Practice Exercise](#): Chapter 1.4 Exercise 1, 2, 10, 12, 13,
21, 30, 34, 45,

[Khan Academy Video](#): Matrix multiplication (part I)

A system such as

$$2x - y = 1$$

$$x + y = 5$$

can be written in vector form as

A system such as

$$2x - y = 1$$

$$x + y = 5$$

can be written in vector form as

$$x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 5 \end{bmatrix}$$

A system such as

$$2x - y = 1$$

$$x + y = 5$$

can be written in vector form as

$$x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 5 \end{bmatrix}$$

The right-hand side is a **linear combination** of the vectors $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$.

Example

We can think of the linear system

$$2x - y = 1$$


$$x + y = 5$$

in two different geometric ways. Recall **unique solution**:

$$x = 2, y = 3.$$


Row picture

- Each equation defines a line in \mathbb{R}^2 .




Row picture

- Each equation defines a line in \mathbb{R}^2 .
- Which points lie on the intersection of these lines?



Row picture


- Each equation defines a line in \mathbb{R}^2 .
- Which points lie on the intersection of these lines?
- $(2, 3)$ is the (only) intersection of the two lines $2x - y = 1$ and $x + y = 5$.



Column picture

- The system can be written as

$$x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 5 \end{bmatrix}$$




Column picture

- The system can be written as

$$x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 5 \end{bmatrix}$$

- Which linear combinations of $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ produce $\begin{bmatrix} 1 \\ 5 \end{bmatrix}$?




Column picture

- The system can be written as

$$x \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 5 \end{bmatrix}$$

- Which linear combinations of $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ produce $\begin{bmatrix} 1 \\ 5 \end{bmatrix}$?
- $(2, 3)$ are the coefficients of the (only) such linear combination.



Application: Industrial Espionage.

Suppose a candy factory produces stuff

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,
- *Nicknacks* and (of course)

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,
- *Nicknacks* and (of course)
- *Widgets*.

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,
- *Nicknacks* and (of course)
- *Widgets*.

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,
- *Nicknacks* and (of course)
- *Widgets*.

To produce these you need raw materials:

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,
- *Nicknacks* and (of course)
- *Widgets*.

To produce these you need raw materials:

- Sugar,

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,
- *Nicknacks* and (of course)
- *Widgets*.

To produce these you need raw materials:

- Sugar,
- Spice and

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,
- *Nicknacks* and (of course)
- *Widgets*.

To produce these you need raw materials:

- Sugar,
- Spice and
- Everything Nice (mystery ingredient!),

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,
- *Nicknacks* and (of course)
- *Widgets*.

To produce these you need raw materials:

- Sugar,
- Spice and
- Everything Nice (mystery ingredient!),

in different quantities.

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,
- *Nicknacks* and (of course)
- *Widgets*.

To produce these you need raw materials:

- Sugar,
- Spice and
- Everything Nice (mystery ingredient!),

in different quantities. We encode how much we need in *production vectors*.

Application: Industrial Espionage.

Suppose a candy factory produces stuff , in particular:

- *Doohickeys*,
- *Nicknacks* and (of course)
- *Widgets*.

To produce these you need raw materials:

- Sugar,
- Spice and
- Everything Nice (mystery ingredient!),

in different quantities. We encode how much we need in *production vectors*. So if the Doohickeys production vector

$v_D = \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix}$ then we need 10 units of Sugar, 5 units of Spice, 2 units of Everything Nice to produce one Doohickey.

Application: Industrial Espionage, Continued.

Similarly we have production vectors v_N, v_W for Nicknacks and Widgets.

Application: Industrial Espionage, Continued.

Similarly we have production vectors v_N, v_W for Nicknacks and Widgets. We assume

$$v_D = \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix}, \quad v_N = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad v_W = \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}$$

Application: Industrial Espionage, Continued.

Similarly we have production vectors v_N, v_W for Nicknacks and Widgets. We assume

$$v_D = \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix}, \quad v_N = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad v_W = \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}$$

Now let us think how much raw material the factory uses.

Application: Industrial Espionage, Continued.

Similarly we have production vectors v_N, v_W for Nicknacks and Widgets. We assume

$$v_D = \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix}, \quad v_N = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad v_W = \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}$$

Now let us think how much raw material the factory uses. The total consumption of raw material will be encoded in the

consumption vector $B = \begin{bmatrix} B_1 \\ B_2 \\ B_3 \end{bmatrix}$ of the factory.

Application: Industrial Espionage, Continued.

Similarly we have production vectors v_N, v_W for Nicknacks and Widgets. We assume

$$v_D = \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix}, \quad v_N = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad v_W = \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}$$

Now let us think how much raw material the factory uses. The total consumption of raw material will be encoded in the

consumption vector $B = \begin{bmatrix} B_1 \\ B_2 \\ B_3 \end{bmatrix}$ of the factory. So B_1 is the amount of sugar consumed in the production, B_2 the amount of spice, etc.

Application: Industrial Espionage, Continued.

Similarly we have production vectors v_N, v_W for Nicknacks and Widgets. We assume

$$v_D = \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix}, \quad v_N = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \quad v_W = \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}$$

Now let us think how much raw material the factory uses. The total consumption of raw material will be encoded in the

consumption vector $B = \begin{bmatrix} B_1 \\ B_2 \\ B_3 \end{bmatrix}$ of the factory. So B_1 is the amount

of sugar consumed in the production, B_2 the amount of spice, etc. If the factory produces c_D Doohickeys, c_N Nicknacks and c_W widgets then the consumption vector will be

$$B = c_D v_D + c_N v_N + c_W v_W$$

Application: Industrial Espionage, Continued.

More explicitly

$$B = c_D \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix} + c_N \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + c_W \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix} .$$

Application: Industrial Espionage, Continued.

More explicitly

$$B = c_D \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix} + c_N \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + c_W \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}.$$

So the consumption vector B is a **linear combination** of the production vectors.

Application: Industrial Espionage, Continued.

More explicitly

$$B = c_D \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix} + c_N \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + c_W \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}.$$

So the consumption vector B is a **linear combination** of the production vectors. Now suppose that somebody (say a competitor of our factory) is interested in determining how many Doohickeys are produced. It is a trade secret, so you can not go and ask the factory.

Application: Industrial Espionage, Continued.

More explicitly

$$B = c_D \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix} + c_N \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + c_W \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}.$$

So the consumption vector B is a **linear combination** of the production vectors. Now suppose that somebody (say a competitor of our factory) is interested in determining how many Doohickeys are produced. It is a trade secret, so you can not go and ask the factory. If many Doohickeys are produced the competitor might switch to the production of Whatchamacallits.

Application: Industrial Espionage, Continued.

More explicitly

$$B = c_D \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix} + c_N \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + c_W \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}.$$

So the consumption vector B is a **linear combination** of the production vectors. Now suppose that somebody (say a competitor of our factory) is interested in determining how many Doohickeys are produced. It is a trade secret, so you can not go and ask the factory. If many Doohickeys are produced the competitor might switch to the production of Whatchamacallits.

So the competitor sends a spy to the entrance of the factory and she writes down how many truck loads of raw materials enters. This determines the production vector B . Then the problem is to find the coefficients c_D, c_N, c_W , i.e., the number of Doohickeys, Nicknacks and Widgets produced.

Application: Industrial Espionage, Continued.

Problem

Suppose our spy observes

$$B = \begin{bmatrix} 14 \\ 12 \\ 7 \end{bmatrix} =$$

Application: Industrial Espionage, Continued.

Problem

Suppose our spy observes

$$B = \begin{bmatrix} 14 \\ 12 \\ 7 \end{bmatrix} = c_D \begin{bmatrix} 10 \\ 5 \\ 2 \end{bmatrix} + c_N \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + c_W \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}.$$

Then what is the number c_D of Doohickeyes produced? How to approach the problem?

Solution

Finding Linear Combinations is the same as solving Linear Systems.

Solution

Finding Linear Combinations is the same as solving Linear Systems. Write down the Augmented Matrix corresponding to the linear combination and find the Reduced Row Echelon Form

$$\left[\begin{array}{ccc|c} 10 & 1 & 3 & 14 \\ 5 & 2 & 5 & 12 \\ 2 & 3 & 2 & 7 \end{array} \right] \simeq$$

Solution

Finding Linear Combinations is the same as solving Linear Systems. Write down the Augmented Matrix corresponding to the linear combination and find the Reduced Row Echelon Form

$$\left[\begin{array}{ccc|c} 10 & 1 & 3 & 14 \\ 5 & 2 & 5 & 12 \\ 2 & 3 & 2 & 7 \end{array} \right] \simeq \left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{array} \right]$$

Solution

Finding Linear Combinations is the same as solving Linear Systems. Write down the Augmented Matrix corresponding to the linear combination and find the Reduced Row Echelon Form

$$\left[\begin{array}{ccc|c} 10 & 1 & 3 & 14 \\ 5 & 2 & 5 & 12 \\ 2 & 3 & 2 & 7 \end{array} \right] \simeq \left[\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{array} \right]$$

So what is c_D ?

Matrices are like Numbers: Matrix Algebra.

Two ways to denote $m \times n$ matrix A (m rows, n column).

- In terms of the **columns** of A :

$$A = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix}$$

Matrices are like Numbers: Matrix Algebra.

Two ways to denote $m \times n$ matrix A (m rows, n column).

- In terms of the **columns** of A :

$$A = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix}$$

- In terms of the **entries** of A :

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

More notation

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- a_{ij} is in the i th row and j th column

More notation

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- a_{ij} is in the i th row and j th column
- \mathbf{a}_j is j^{th} column:

$$\mathbf{a}_j = \begin{bmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{mj} \end{bmatrix}$$

More notation

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

- a_{ij} is in the i th row and j th column
- \mathbf{a}_j is j^{th} column:

$$\mathbf{a}_j = \begin{bmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{mj} \end{bmatrix}$$

- **Main diagonal entries:** $a_{11}, a_{22}, \dots, a_{mm}$ (only care about these when $m = n$)

Even more notation

- **Zero matrix:**

$$0 = \begin{bmatrix} 0 & \cdots & 0 & \cdots & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & \cdots & 0 & \cdots & 0 \\ \vdots & & \vdots & & \vdots \\ 0 & \cdots & 0 & \cdots & 0 \end{bmatrix}$$

Definition

Let $A = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix}$, $B = \begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \cdots & \mathbf{b}_n \end{bmatrix}$ be $m \times n$ -matrices and let r be a scalar.

Definition

Let $A = [\mathbf{a}_1 \ \mathbf{a}_2 \ \cdots \ \mathbf{a}_n]$, $B = [\mathbf{b}_1 \ \mathbf{b}_2 \ \cdots \ \mathbf{b}_n]$ be $m \times n$ -matrices and let r be a scalar. Then

- $A + B$ is defined by

$$A + B = [\mathbf{a}_1 + \mathbf{b}_1 \ \mathbf{a}_2 + \mathbf{b}_2 \ \cdots \ \mathbf{a}_n + \mathbf{b}_n]$$

- Moreover, rA is defined as

$$rA = [r\mathbf{a}_1 \ r\mathbf{a}_2 \ \cdots \ r\mathbf{a}_n]$$

Example

Calculate



$$\begin{bmatrix} 1 & 0 \\ 5 & 2 \end{bmatrix} + \begin{bmatrix} 2 & 3 \\ 3 & 1 \end{bmatrix} =$$

Example

Calculate



$$\begin{bmatrix} 1 & 0 \\ 5 & 2 \end{bmatrix} + \begin{bmatrix} 2 & 3 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 8 & 3 \end{bmatrix}$$

Example

Calculate



$$\begin{bmatrix} 1 & 0 \\ 5 & 2 \end{bmatrix} + \begin{bmatrix} 2 & 3 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 8 & 3 \end{bmatrix}$$



$$10 \begin{bmatrix} 2 & 3 \\ 3 & 1 \end{bmatrix} =$$

Example

Calculate



$$\begin{bmatrix} 1 & 0 \\ 5 & 2 \end{bmatrix} + \begin{bmatrix} 2 & 3 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 8 & 3 \end{bmatrix}$$



$$10 \begin{bmatrix} 2 & 3 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} 20 & 30 \\ 30 & 10 \end{bmatrix}$$

Theorem

Let A , B , and C be matrices of the same size, and let r and s be scalars.

- $A + B = B + A$
- $(A + B) + C = A + (B + C)$
- $A + 0 = A$
- $r(A + B) = rA + rB$
- $(r + s)A = rA + sA$
- $r(sA) = (rs)A$

Theorem

Let A , B , and C be matrices of the same size, and let r and s be scalars.

- $A + B = B + A$
- $(A + B) + C = A + (B + C)$
- $A + 0 = A$
- $r(A + B) = rA + rB$
- $(r + s)A = rA + sA$
- $r(sA) = (rs)A$

Matrices are like Numbers!

How to multiply matrices and vectors

Let \mathbf{x} be a vector, A, B matrices.

- Multiplying B and \mathbf{x} transforms \mathbf{x} into the **vector** $B\mathbf{x}$.

How to multiply matrices and vectors

Let \mathbf{x} be a vector, A, B matrices.

- Multiplying B and \mathbf{x} transforms \mathbf{x} into the **vector** $B\mathbf{x}$.
- In turn, if we multiply A and $B\mathbf{x}$, we transform $B\mathbf{x}$ into $A(B\mathbf{x})$.

How to multiply matrices and vectors

Let \mathbf{x} be a vector, A, B matrices.

- Multiplying B and \mathbf{x} transforms \mathbf{x} into the **vector** $B\mathbf{x}$.
- In turn, if we multiply A and $B\mathbf{x}$, we transform $B\mathbf{x}$ into $A(B\mathbf{x})$.
- So $A(B\mathbf{x})$ is the **composition of two mappings**.

How to multiply matrices and vectors

Let \mathbf{x} be a vector, A, B matrices.

- Multiplying B and \mathbf{x} transforms \mathbf{x} into the **vector** $B\mathbf{x}$.
- In turn, if we multiply A and $B\mathbf{x}$, we transform $B\mathbf{x}$ into $A(B\mathbf{x})$.
- So $A(B\mathbf{x})$ is the **composition of two mappings**.

How to multiply matrices and vectors

Let \mathbf{x} be a vector, A, B matrices.

- Multiplying B and \mathbf{x} transforms \mathbf{x} into the **vector** $B\mathbf{x}$.
- In turn, if we multiply A and $B\mathbf{x}$, we transform $B\mathbf{x}$ into $A(B\mathbf{x})$.
- So $A(B\mathbf{x})$ is the **composition of two mappings**.

Define the product AB so that

$$A(B\mathbf{x}) = (AB)\mathbf{x}$$

Matrix Multiplication

Suppose A is $m \times n$ and B is $n \times p$ where

$$B = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_p] \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_p \end{bmatrix}$$

Matrix Multiplication

Suppose A is $m \times n$ and B is $n \times p$ where

$$B = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_p] \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_p \end{bmatrix}$$

Define

$$B\mathbf{x} = x_1\mathbf{b}_1 + x_2\mathbf{b}_2 + \cdots + x_p\mathbf{b}_p$$

Then

Suppose A is $m \times n$ and B is $n \times p$ where

$$B = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_p] \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_p \end{bmatrix}$$

Define

$$B\mathbf{x} = x_1\mathbf{b}_1 + x_2\mathbf{b}_2 + \cdots + x_n\mathbf{b}_n$$

Then

$$A(B\mathbf{x}) = x_1A\mathbf{b}_1 + x_2A\mathbf{b}_2 + \cdots + x_nA\mathbf{b}_n$$

Example

$$A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Example

$$A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$B\mathbf{x} = B \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$$

Example

$$A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$B\mathbf{x} = B \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $B\mathbf{x} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$:

Example

$$A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$B\mathbf{x} = B \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $B\mathbf{x} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$:

$$A(B\mathbf{x}) = A \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$$

Example

$$A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$B\mathbf{x} = B \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $B\mathbf{x} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$:

$$A(B\mathbf{x}) = A \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix} = (x_1 + 2x_2) \begin{bmatrix} 2 \\ 1 \end{bmatrix} + (x_2) \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Example

$$A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$B\mathbf{x} = B \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $B\mathbf{x} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$:

$$A(B\mathbf{x}) = A \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix} = (x_1 + 2x_2) \begin{bmatrix} 2 \\ 1 \end{bmatrix} + (x_2) \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2x_1 + 4x_2 \\ x_1 + 3x_2 \end{bmatrix}$$

Example

$$A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$B\mathbf{x} = B \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $B\mathbf{x} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$:

$$A(B\mathbf{x}) = A \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix} = (x_1 + 2x_2) \begin{bmatrix} 2 \\ 1 \end{bmatrix} + (x_2) \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2x_1 + 4x_2 \\ x_1 + 3x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $A(B\mathbf{x}) = x_1 A\mathbf{b}_1 + x_2 A\mathbf{b}_2 + \dots + x_n A\mathbf{b}_n$:

Example

$$A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$B\mathbf{x} = B \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $B\mathbf{x} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$:

$$A(B\mathbf{x}) = A \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix} = (x_1 + 2x_2) \begin{bmatrix} 2 \\ 1 \end{bmatrix} + (x_2) \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2x_1 + 4x_2 \\ x_1 + 3x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $A(B\mathbf{x}) = x_1 A\mathbf{b}_1 + x_2 A\mathbf{b}_2 + \dots + x_n A\mathbf{b}_n$:

$$A(B\mathbf{x}) = x_1 A\mathbf{b}_1 + x_2 A\mathbf{b}_2$$

Example

$$A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$B\mathbf{x} = B \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $B\mathbf{x} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$:

$$A(B\mathbf{x}) = A \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix} = (x_1 + 2x_2) \begin{bmatrix} 2 \\ 1 \end{bmatrix} + (x_2) \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2x_1 + 4x_2 \\ x_1 + 3x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $A(B\mathbf{x}) = x_1 A\mathbf{b}_1 + x_2 A\mathbf{b}_2 + \dots + x_n A\mathbf{b}_n$:

$$A(B\mathbf{x}) = x_1 A\mathbf{b}_1 + x_2 A\mathbf{b}_2 = x_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$

Example

$$A = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$B\mathbf{x} = B \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $B\mathbf{x} = \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix}$:

$$A(B\mathbf{x}) = A \begin{bmatrix} x_1 + 2x_2 \\ x_2 \end{bmatrix} = (x_1 + 2x_2) \begin{bmatrix} 2 \\ 1 \end{bmatrix} + (x_2) \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2x_1 + 4x_2 \\ x_1 + 3x_2 \end{bmatrix}$$

Compute $A(B\mathbf{x})$ using $A(B\mathbf{x}) = x_1 A\mathbf{b}_1 + x_2 A\mathbf{b}_2 + \dots + x_n A\mathbf{b}_n$:

$$A(B\mathbf{x}) = x_1 A\mathbf{b}_1 + x_2 A\mathbf{b}_2 = x_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 4 \\ 3 \end{bmatrix} = \begin{bmatrix} 2x_1 + 4x_2 \\ x_1 + 3x_2 \end{bmatrix}$$

Same answer!

Matrix Multiplication

Motto

Matrix Multiplication is Linear Combination!