

Math 415 - Lecture 15

The Four Fundamental Subspaces, the Fundamental Theorem of
Linear Algebra, Linear Transformations

Monday September 28th 2015

Textbook: Chapter 2.4, 2.6

Textbook: Chapter 2.4, 2.6

Suggested Practice Exercise: Chapter 2.4 Exercise 1, 2, 3, 4, 7, 10, 18, 20, 21, 22, 27, 32, 37 Chapter 2.6 Exercise 5, 6, 7, 36, 37

Textbook: Chapter 2.4, 2.6

Suggested Practice Exercise: Chapter 2.4 Exercise 1, 2, 3, 4, 7, 10, 18, 20, 21, 22, 27, 32, 37 Chapter 2.6 Exercise 5, 6, 7, 36, 37

Khan Academy Video: Linear Transformation, Linear Transformations as Matrix Vector Products, Linear Transformation Examples: Rotations in \mathbb{R}^2

Textbook: Chapter 2.4, 2.6

Suggested Practice Exercise: Chapter 2.4 Exercise 1, 2, 3, 4, 7, 10, 18, 20, 21, 22, 27, 32, 37 Chapter 2.6 Exercise 5, 6, 7, 36, 37

Khan Academy Video: Linear Transformation, Linear Transformations as Matrix Vector Products, Linear Transformation Examples: Rotations in \mathbb{R}^2

Strang lectures: Lecture 9: Independence, Basis, and Dimension
Lecture 10: The Four Fundamental Subspaces
Lecture 30: Linear Transformations

* Exam 1 (7-8:15 pm Tuesday September 29):

- * Exam 1 (7-8:15 pm Tuesday September 29):
- * Rooms: look on Moodle.

- * Exam 1 (7-8:15 pm Tuesday September 29):
- * Rooms: look on Moodle.
- * Conflicts: if you have a conflict you should have received an email about it. If not, talk to me after class.

- * Exam 1 (7-8:15 pm Tuesday September 29):
- * Rooms: look on Moodle.
- * Conflicts: if you have a conflict you should have received an email about it. If not, talk to me after class.
- * No Discussion Sections next week.

- * Exam 1 (7-8:15 pm Tuesday September 29):
- * Rooms: look on Moodle.
- * Conflicts: if you have a conflict you should have received an email about it. If not, talk to me after class.
- * No Discussion Sections next week.
- * No Class on Wednesday next week.

- * Exam 1 (7-8:15 pm Tuesday September 29):
- * Rooms: look on Moodle.
- * Conflicts: if you have a conflict you should have received an email about it. If not, talk to me after class.
- * No Discussion Sections next week.
- * No Class on Wednesday next week.
- * The Exam will be part multiple choice. Bring pencils and erasers! Also bring ID.

- * Exam 1 (7-8:15 pm Tuesday September 29):
- * Rooms: look on Moodle.
- * Conflicts: if you have a conflict you should have received an email about it. If not, talk to me after class.
- * No Discussion Sections next week.
- * No Class on Wednesday next week.
- * The Exam will be part multiple choice. Bring pencils and erasers! Also bring ID.
- * The material for the exam covers the lectures upto and including Lecture 12 (last Monday), and this weeks worksheet and quiz.

Review

Basis for the Null Space

- $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a **basis** of V if the vectors span V and are independent.

- $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a **basis** of V if the vectors span V and are independent.
- To find a basis for $\text{Nul}(A)$, solve $A\mathbf{x} = \mathbf{0}$.

- $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a **basis** of V if the vectors span V and are independent.
- To find a basis for $\text{Nul}(A)$, solve $A\mathbf{x} = \mathbf{0}$.

- $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a **basis** of V if the vectors span V and are independent.
- To find a basis for $\text{Nul}(A)$, solve $A\mathbf{x} = \mathbf{0}$.

$$\begin{bmatrix} 3 & 6 & 6 & 3 \\ 6 & 12 & 15 & 0 \end{bmatrix}$$

- $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a **basis** of V if the vectors span V and are independent.
- To find a basis for $\text{Nul}(A)$, solve $A\mathbf{x} = \mathbf{0}$.

$$\begin{bmatrix} 3 & 6 & 6 & 3 \\ 6 & 12 & 15 & 0 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} \boxed{1} & 2 & 0 & 5 \\ 0 & 0 & \boxed{1} & -2 \end{bmatrix}$$

- $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a **basis** of V if the vectors span V and are independent.
- To find a basis for $\text{Nul}(A)$, solve $A\mathbf{x} = \mathbf{0}$.

$$\begin{bmatrix} 3 & 6 & 6 & 3 \\ 6 & 12 & 15 & 0 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} \boxed{1} & 2 & 0 & 5 \\ 0 & 0 & \boxed{1} & -2 \end{bmatrix}$$

$\mathbf{x} =$

- $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a **basis** of V if the vectors span V and are independent.
- To find a basis for $\text{Nul}(A)$, solve $A\mathbf{x} = \mathbf{0}$.

$$\begin{bmatrix} 3 & 6 & 6 & 3 \\ 6 & 12 & 15 & 0 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} \boxed{1} & 2 & 0 & 5 \\ 0 & 0 & \boxed{1} & -2 \end{bmatrix}$$

$$\mathbf{x} = \begin{bmatrix} -2x_2 - 5x_4 \\ x_2 \\ 2x_4 \\ x_4 \end{bmatrix}$$

- $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a **basis** of V if the vectors span V and are independent.
- To find a basis for $Nul(A)$, solve $A\mathbf{x} = \mathbf{0}$.

$$\begin{bmatrix} 3 & 6 & 6 & 3 \\ 6 & 12 & 15 & 0 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} \boxed{1} & 2 & 0 & 5 \\ 0 & 0 & \boxed{1} & -2 \end{bmatrix}$$

$$\mathbf{x} = \begin{bmatrix} -2x_2 - 5x_4 \\ x_2 \\ 2x_4 \\ x_4 \end{bmatrix} = x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -5 \\ 0 \\ 2 \\ 1 \end{bmatrix}$$

So a basis for $Nul(A)$ is

- $\{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ is a **basis** of V if the vectors span V and are independent.
- To find a basis for $\text{Nul}(A)$, solve $A\mathbf{x} = \mathbf{0}$.

$$\begin{bmatrix} 3 & 6 & 6 & 3 \\ 6 & 12 & 15 & 0 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} \boxed{1} & 2 & 0 & 5 \\ 0 & 0 & \boxed{1} & -2 \end{bmatrix}$$

$$\mathbf{x} = \begin{bmatrix} -2x_2 - 5x_4 \\ x_2 \\ 2x_4 \\ x_4 \end{bmatrix} = x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -5 \\ 0 \\ 2 \\ 1 \end{bmatrix}$$

$$\text{So a basis for } \text{Nul}(A) \text{ is } \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -5 \\ 0 \\ 2 \\ 1 \end{bmatrix} \right\}$$

Basis for the Column space.

- To find a basis for $\text{Col}(A)$, take the pivot columns of A .

- To find a basis for $\text{Col}(A)$, take the pivot columns of A .

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix}$$

- To find a basis for $\text{Col}(A)$, take the pivot columns of A .

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \longrightarrow \begin{bmatrix} \boxed{1} & 2 & 0 & 4 \\ 0 & 0 & \boxed{-1} & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

So a basis for $\text{Col}(A)$ is

- To find a basis for $\text{Col}(A)$, take the pivot columns of A .

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \longrightarrow \begin{bmatrix} \boxed{1} & 2 & 0 & 4 \\ 0 & 0 & \boxed{-1} & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

So a basis for $\text{Col}(A)$ is $\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 2 \\ 0 \end{bmatrix} \right\}$

- To find a basis for $\text{Col}(A)$, take the pivot columns of A .

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \longrightarrow \begin{bmatrix} \boxed{1} & 2 & 0 & 4 \\ 0 & 0 & \boxed{-1} & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

So a basis for $\text{Col}(A)$ is $\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 2 \\ 0 \end{bmatrix} \right\}$

Question

Why do we take **columns of A** and not columns of the Echelon form?

The Column spaces of A and U.

Question

Why do we take **columns of A** and not columns of the Echelon form?

Question

Why do we take **columns of A** and not columns of the Echelon form?

- Row operations do **not** preserve the column space.

For example, $\begin{bmatrix} 1 \\ 0 \end{bmatrix} \xrightarrow{R1 \leftrightarrow R2} \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$

- On the other hand, row operations do preserve the null space.
Why?

Question

Why do we take **columns of A** and not columns of the Echelon form?

- Row operations do **not** preserve the column space.

For example, $\begin{bmatrix} 1 \\ 0 \end{bmatrix} \xrightarrow{R1 \leftrightarrow R2} \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$

- On the other hand, row operations do preserve the null space.

Why?

Remember, we can do row operations to solve systems like

$$Ax = \mathbf{0}.$$

Rank and Dimensions

Dimension of Column and Null Space

Definition

The **rank** of a matrix A is the number of pivots it has.

Definition

The **rank** of a matrix A is the number of pivots it has.

Theorem

Rank-Nullity Theorem *Let A be an $m \times n$ matrix of rank r . Then*
 $\dim \text{Col}(A) = r$ Why?

Definition

The **rank** of a matrix A is the number of pivots it has.

Theorem

Rank-Nullity Theorem *Let A be an $m \times n$ matrix of rank r . Then*
 $\dim \text{Col}(A) = r$ Why?

Definition

The **rank** of a matrix A is the number of pivots it has.

Theorem

Rank-Nullity Theorem *Let A be an $m \times n$ matrix of rank r . Then*
 $\dim \text{Col}(A) = r$ Why?

A basis for $\text{Col}(A)$ is given by the pivot columns of A .

Definition

The **rank** of a matrix A is the number of pivots it has.

Theorem

Rank-Nullity Theorem *Let A be an $m \times n$ matrix of rank r . Then*

$\dim \text{Col}(A) = r$ Why?

A basis for $\text{Col}(A)$ is given by the pivot columns of A .

$\dim \text{Nul}(A) = n - r$ is the number of free variables of A . Why?

Definition

The **rank** of a matrix A is the number of pivots it has.

Theorem

Rank-Nullity Theorem *Let A be an $m \times n$ matrix of rank r . Then*

$\dim \text{Col}(A) = r$ Why?

A basis for $\text{Col}(A)$ is given by the pivot columns of A .

$\dim \text{Nul}(A) = n - r$ is the number of free variables of A . Why?

Definition

The **rank** of a matrix A is the number of pivots it has.

Theorem

Rank-Nullity Theorem *Let A be an $m \times n$ matrix of rank r . Then*

$\dim \text{Col}(A) = r$ Why?

A basis for $\text{Col}(A)$ is given by the pivot columns of A .

$\dim \text{Nul}(A) = n - r$ is the number of free variables of A . Why?

In our method for finding a basis for $\text{Nul}(A)$, each free variable corresponds to an element in the basis.

Definition

The **rank** of a matrix A is the number of pivots it has.

Theorem

Rank-Nullity Theorem *Let A be an $m \times n$ matrix of rank r . Then*

$\dim \text{Col}(A) = r$ Why?

A basis for $\text{Col}(A)$ is given by the pivot columns of A .

$\dim \text{Nul}(A) = n - r$ is the number of free variables of A . Why?

In our method for finding a basis for $\text{Nul}(A)$, each free variable corresponds to an element in the basis.

$\dim \text{Col}(A) + \dim \text{Nul}(A) = n$ Why?

Definition

The **rank** of a matrix A is the number of pivots it has.

Theorem

Rank-Nullity Theorem *Let A be an $m \times n$ matrix of rank r . Then*

$\dim \text{Col}(A) = r$ Why?

A basis for $\text{Col}(A)$ is given by the pivot columns of A .

$\dim \text{Nul}(A) = n - r$ is the number of free variables of A . Why?

In our method for finding a basis for $\text{Nul}(A)$, each free variable corresponds to an element in the basis.

$\dim \text{Col}(A) + \dim \text{Nul}(A) = n$ Why?

Definition

The **rank** of a matrix A is the number of pivots it has.

Theorem

Rank-Nullity Theorem *Let A be an $m \times n$ matrix of rank r . Then*
 $\dim \text{Col}(A) = r$ *Why?*

A basis for $\text{Col}(A)$ is given by the pivot columns of A .

$\dim \text{Nul}(A) = n - r$ *is the number of free variables of A . Why?*

In our method for finding a basis for $\text{Nul}(A)$, each free variable corresponds to an element in the basis.

$\dim \text{Col}(A) + \dim \text{Nul}(A) = n$ *Why?*

Each of the n columns of A either contains a pivot or corresponds to a free variable.

The Four Fundamental Subspaces

Two Spaces we know

Let A be a matrix. We already know two fundamental subspaces:

- The **column space** of A and

Let A be a matrix. We already know two fundamental subspaces:

- The **column space** of A and
- The **null space** of A

Let A be a matrix. We already know two fundamental subspaces:

- The **column space** of A and
- The **null space** of A

Let A be a matrix. We already know two fundamental subspaces:

- The **column space** of A and
- The **null space** of A

There are two more!

Row Space and Left Null Space

Definition

- The **row space** of A is the column space of A^T .

Definition

- The **row space** of A is the column space of A^T .
 $\text{Col}(A^T)$ is spanned by the columns of A^T and these are the rows of A (but transposed, to turn into columns!).

Definition

- The **row space** of A is the column space of A^T .
 $\text{Col}(A^T)$ is spanned by the columns of A^T and these are the rows of A (but transposed, to turn into columns!).
- The **left null space** of A is the null space of A^T .

Definition

- The **row space** of A is the column space of A^T .
 $\text{Col}(A^T)$ is spanned by the columns of A^T and these are the rows of A (but transposed, to turn into columns!).
- The **left null space** of A is the null space of A^T .

Definition

- The **row space** of A is the column space of A^T .
 $\text{Col}(A^T)$ is spanned by the columns of A^T and these are the rows of A (but transposed, to turn into columns!).
- The **left null space** of A is the null space of A^T .
 Why is it called the “left” null space?

Definition

- The **row space** of A is the column space of A^T .
 $\text{Col}(A^T)$ is spanned by the columns of A^T and these are the rows of A (but transposed, to turn into columns!).
- The **left null space** of A is the null space of A^T .
 Why is it called the “left” null space?
 Suppose $\mathbf{x} \in \text{Nul}(A^T)$. Thus,

Definition

- The **row space** of A is the column space of A^T .
 $\text{Col}(A^T)$ is spanned by the columns of A^T and these are the rows of A (but transposed, to turn into columns!).
- The **left null space** of A is the null space of A^T .

Why is it called the “left” null space?

Suppose $\mathbf{x} \in \text{Nul}(A^T)$. Thus,

$$\iff A^T \mathbf{x} = \mathbf{0}.$$

Take transposes of both sides:

Definition

- The **row space** of A is the column space of A^T .
 $\text{Col}(A^T)$ is spanned by the columns of A^T and these are the rows of A (but transposed, to turn into columns!).
- The **left null space** of A is the null space of A^T .

Why is it called the “left” null space?

Suppose $\mathbf{x} \in \text{Nul}(A^T)$. Thus,

$$\iff A^T \mathbf{x} = \mathbf{0}. \quad \text{Take transposes of both sides:}$$

$$\iff (A^T \mathbf{x})^T = \mathbf{0}^T. \quad \text{So,}$$

Definition

- The **row space** of A is the column space of A^T .
 $\text{Col}(A^T)$ is spanned by the columns of A^T and these are the rows of A (but transposed, to turn into columns!).
- The **left null space** of A is the null space of A^T .

Why is it called the “left” null space?

Suppose $\mathbf{x} \in \text{Nul}(A^T)$. Thus,

$$\iff A^T \mathbf{x} = \mathbf{0}.$$

Take transposes of both sides:

$$\iff (A^T \mathbf{x})^T = \mathbf{0}^T. \quad \text{So,}$$

$$\iff \mathbf{x}^T A = \mathbf{0}.$$

Definition

- The **row space** of A is the column space of A^T .
 $\text{Col}(A^T)$ is spanned by the columns of A^T and these are the rows of A (but transposed, to turn into columns!).
- The **left null space** of A is the null space of A^T .

Why is it called the “left” null space?

Suppose $\mathbf{x} \in \text{Nul}(A^T)$. Thus,

$$\iff A^T \mathbf{x} = \mathbf{0}. \quad \text{Take transposes of both sides:}$$

$$\iff (A^T \mathbf{x})^T = \mathbf{0}^T. \quad \text{So,}$$

$$\iff \mathbf{x}^T A = \mathbf{0}.$$

Therefore, $\mathbf{x} \in \text{Nul}(A^T) \iff \mathbf{x}^T A = \mathbf{0}.$

Example

Find a basis for $\text{Col}(A)$ and $\text{Col}(A^T)$ if

$$A = \begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix}.$$

Example

Find a basis for $\text{Col}(A)$ and $\text{Col}(A^T)$ if

$$A = \begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix}.$$

Solution. We need to compute an echelon form of A to find a basis for $\text{Col}(A)$.

Example

Find a basis for $\text{Col}(A)$ and $\text{Col}(A^T)$ if

$$A = \begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix}.$$

Solution. We need to compute an echelon form of A to find a basis for $\text{Col}(A)$. Then we might compute an echelon form of A^T to find a basis for $\text{Col}(A^T)$.

Example

Find a basis for $\text{Col}(A)$ and $\text{Col}(A^T)$ if

$$A = \begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix}.$$

Solution. We need to compute an echelon form of A to find a basis for $\text{Col}(A)$. Then we might compute an echelon form of A^T to find a basis for $\text{Col}(A^T)$. However, an echelon form of A will allow us to find a basis for both $\text{Col}(A)$ and $\text{Col}(A^T)$.

Instead of doing twice the work, we only need to find an echelon form of A .

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix}$$

Instead of doing twice the work, we only need to find an echelon form of A .

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \longrightarrow \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & -1 & -5 \\ 0 & 0 & 2 & 10 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Instead of doing twice the work, we only need to find an echelon form of A .

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \longrightarrow \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & -1 & -5 \\ 0 & 0 & 2 & 10 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

We identify the pivot columns:

Instead of doing twice the work, we only need to find an echelon form of A .

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \longrightarrow \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & -1 & -5 \\ 0 & 0 & 2 & 10 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

We identify the pivot columns:

$$\longrightarrow \begin{bmatrix} \boxed{1} & 2 & 0 & 4 \\ 0 & 0 & \boxed{-1} & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = U$$

Instead of doing twice the work, we only need to find an echelon form of A .

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \longrightarrow \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & -1 & -5 \\ 0 & 0 & 2 & 10 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

We identify the pivot columns:

$$\longrightarrow \begin{bmatrix} \boxed{1} & 2 & 0 & 4 \\ 0 & 0 & \boxed{-1} & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = U$$

So $r = 2$ for A and a basis for $\text{Col}(A)$ is

Instead of doing twice the work, we only need to find an echelon form of A .

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 2 & 4 & -1 & 3 \\ 3 & 6 & 2 & 22 \\ 4 & 8 & 0 & 16 \end{bmatrix} \longrightarrow \begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & 0 & -1 & -5 \\ 0 & 0 & 2 & 10 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

We identify the pivot columns:

$$\longrightarrow \begin{bmatrix} \boxed{1} & 2 & 0 & 4 \\ 0 & 0 & \boxed{-1} & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = U$$

So $r = 2$ for A and a basis for $\text{Col}(A)$ is $\left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 2 \\ 0 \end{bmatrix} \right\}$.

Remark

Key idea: The row space is preserved by elementary row operations.

Remark

Key idea: The row space is preserved by elementary row operations.

Remember, $\text{Col}(A) \neq \text{Col}(U)$ because we did row operations.

However, the row spaces are the same!

Remark

Key idea: The row space is preserved by elementary row operations.

Remember, $\text{Col}(A) \neq \text{Col}(U)$ because we did row operations.

However, the row spaces are the same! i.e.

$$\text{Col}(A^T) = \text{Col}(U^T)$$

Remark

Key idea: The row space is preserved by elementary row operations.

Remember, $\text{Col}(A) \neq \text{Col}(U)$ because we did row operations.

However, the row spaces are the same! i.e.

$$\text{Col}(A^T) = \text{Col}(U^T)$$

$$U = \begin{bmatrix} \boxed{1} & 2 & 0 & 4 \\ 0 & 0 & \boxed{-1} & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

Remark

Key idea: The row space is preserved by elementary row operations.

Remember, $\text{Col}(A) \neq \text{Col}(U)$ because we did row operations.

However, the row spaces are the same! i.e.

$$\text{Col}(A^T) = \text{Col}(U^T)$$

$$U = \begin{bmatrix} \boxed{1} & 2 & 0 & 4 \\ 0 & 0 & \boxed{-1} & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad U^T = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 4 & -5 & 0 & 0 \end{bmatrix}$$

Remark

Key idea: The row space is preserved by elementary row operations.

Remember, $\text{Col}(A) \neq \text{Col}(U)$ because we did row operations.

However, the row spaces are the same! i.e.

$$\text{Col}(A^T) = \text{Col}(U^T)$$

$$U = \begin{bmatrix} \boxed{1} & 2 & 0 & 4 \\ 0 & 0 & \boxed{-1} & -5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad U^T = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 4 & -5 & 0 & 0 \end{bmatrix}$$

In particular, a basis for $\text{Col}(A^T)$ is given by $\left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ -1 \\ -5 \end{bmatrix} \right\}$.

Fundamental Theorem of Linear Algebra (Part 1)

Fundamental Theorem of Linear Algebra (Part 1)

Theorem

Let A be an $m \times n$ matrix with rank r .

Theorem

Let A be an $m \times n$ matrix with rank r .

- $\dim \operatorname{Col}(A) = r$

(subspace of \mathbb{R}^m)

Theorem

Let A be an $m \times n$ matrix with rank r .

- $\dim \operatorname{Col}(A) = r$ (subspace of \mathbb{R}^m)
- $\dim \operatorname{Col}(A^T) = r$ (subspace of \mathbb{R}^n)

Theorem

Let A be an $m \times n$ matrix with rank r .

- $\dim \operatorname{Col}(A) = r$ (subspace of \mathbb{R}^m)
- $\dim \operatorname{Col}(A^T) = r$ (subspace of \mathbb{R}^n)
- $\dim \operatorname{Nul}(A) = n - r$ (subspace of \mathbb{R}^n)

Theorem

Let A be an $m \times n$ matrix with rank r .

- $\dim \operatorname{Col}(A) = r$ (subspace of \mathbb{R}^m)
- $\dim \operatorname{Col}(A^T) = r$ (subspace of \mathbb{R}^n)
- $\dim \operatorname{Nul}(A) = n - r$ (subspace of \mathbb{R}^n)
- $\dim \operatorname{Nul}(A^T) = m - r$ (subspace of \mathbb{R}^m)

Remark

The column and row space always have the same dimension. In other words, A and A^T have the same rank. (i.e. same number of pivots). Why?

Remark

The column and row space always have the same dimension. In other words, A and A^T have the same rank. (i.e. same number of pivots). Why?

It's easy to see this for a matrix in echelon form.

$$A = \begin{bmatrix} 2 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 7 \end{bmatrix},$$

Remark

The column and row space always have the same dimension. In other words, A and A^T have the same rank. (i.e. same number of pivots). Why?

It's easy to see this for a matrix in echelon form.

$$A = \begin{bmatrix} 2 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 7 \end{bmatrix}, \quad A^T = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 2 & 7 \end{bmatrix}$$

(3 pivot columns in A , 3 non-zero columns in A^T .)

Remark

The column and row space always have the same dimension. In other words, A and A^T have the same rank. (i.e. same number of pivots). Why?

It's easy to see this for a matrix in echelon form.

$$A = \begin{bmatrix} 2 & 1 & 3 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 7 \end{bmatrix}, \quad A^T = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 2 & 7 \end{bmatrix}$$

(3 pivot columns in A , 3 non-zero columns in A^T .) But it's not as obvious for a random matrix.

Coordinates

Why Bases?

What is the point of having a *basis* for a vector space V ?

What is the point of having a *basis* for a vector space V ?

- **Dimension!** If you have a basis $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ for V , you know that the dimension of V is p , so that you have an idea of the **Size** of V .

What is the point of having a *basis* for a vector space V ?

- **Dimension!** If you have a basis $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ for V , you know that the dimension of V is p , so that you have an idea of the **Size** of V . In particular, if V has dimension 0 V is just the zero vector space.

What is the point of having a *basis* for a vector space V ?

- **Dimension!** If you have a basis $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ for V , you know that the dimension of V is p , so that you have an idea of the **Size** of V . In particular, if V has dimension 0 V is just the zero vector space.
- **Coordinates!** If $w \in V$ and $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ is a basis for V , we can express w in this basis.

What is the point of having a *basis* for a vector space V ?

- **Dimension!** If you have a basis $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ for V , you know that the dimension of V is p , so that you have an idea of the **Size** of V . In particular, if V has dimension 0 V is just the zero vector space.
- **Coordinates!** If $w \in V$ and $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ is a basis for V , we can express w in this basis. This means that we can write (uniquely!)

$$w = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \cdots + c_p \mathbf{v}_p.$$

What is the point of having a *basis* for a vector space V ?

- **Dimension!** If you have a basis $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ for V , you know that the dimension of V is p , so that you have an idea of the **Size** of V . In particular, if V has dimension 0 V is just the zero vector space.
- **Coordinates!** If $w \in V$ and $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ is a basis for V , we can express w in this basis. This means that we can write (uniquely!)

$$w = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p.$$

We call the scalars c_1, c_2, \dots, c_p the *coordinates* of w with respect to the basis \mathcal{B} .

What is the point of having a *basis* for a vector space V ?

- **Dimension!** If you have a basis $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ for V , you know that the dimension of V is p , so that you have an idea of the **Size** of V . In particular, if V has dimension 0 V is just the zero vector space.
- **Coordinates!** If $w \in V$ and $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ is a basis for V , we can express w in this basis. This means that we can write (uniquely!)

$$w = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \cdots + c_p\mathbf{v}_p.$$

We call the scalars c_1, c_2, \dots, c_p the *coordinates* of w with respect to the basis \mathcal{B} .

We are going to organize the coordinates in a convenient package.

Coordinate Vectors

Definition

If $w \in V$ and $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ is a basis for V , the **coordinate vector** of w with respect to the basis \mathcal{B} is

$$w_{\mathcal{B}} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_p \end{bmatrix}, \quad \text{if } w = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \cdots + c_p \mathbf{v}_p.$$

Definition

If $w \in V$ and $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ is a basis for V , the **coordinate vector** of w with respect to the basis \mathcal{B} is

$$w_{\mathcal{B}} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_p \end{bmatrix}, \quad \text{if } w = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \cdots + c_p \mathbf{v}_p.$$

So w is a vector in some vector space, but its coordinate vector is always a column vector in \mathbb{R}^p , if $\dim(V) = p$.

Definition

If $w \in V$ and $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ is a basis for V , the **coordinate vector** of w with respect to the basis \mathcal{B} is

$$w_{\mathcal{B}} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_p \end{bmatrix}, \quad \text{if } w = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \cdots + c_p \mathbf{v}_p.$$

So w is a vector in some vector space, but its coordinate vector is always a column vector in \mathbb{R}^p , if $\dim(V) = p$. Why is the coordinate vector useful?

Definition

If $w \in V$ and $\mathcal{B} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p)$ is a basis for V , the **coordinate vector** of w with respect to the basis \mathcal{B} is

$$w_{\mathcal{B}} = \begin{bmatrix} c_1 \\ c_2 \\ c_3 \\ \vdots \\ c_p \end{bmatrix}, \quad \text{if } w = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \cdots + c_p \mathbf{v}_p.$$

So w is a vector in some vector space, but its coordinate vector is always a column vector in \mathbb{R}^p , if $\dim(V) = p$. Why is the coordinate vector useful? Computations in V can be translated in computations in the familiar vector space \mathbb{R}^p .

Let $V = \mathbb{R}^2$, $\mathcal{B} = (\mathbf{b}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix})$ and $\mathbf{w} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$.

Let $V = \mathbb{R}^2$, $\mathcal{B} = (\mathbf{b}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix})$ and $\mathbf{w} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$. What is the coordinate vector of \mathbf{w} ?

Let $V = \mathbb{R}^2$, $\mathcal{B} = (\mathbf{b}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix})$ and $\mathbf{w} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$. What is the coordinate vector of \mathbf{w} ? Express in the basis:

$$\begin{bmatrix} 3 \\ -1 \end{bmatrix} = c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2 = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} =$$

Let $V = \mathbb{R}^2$, $\mathcal{B} = (\mathbf{b}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix})$ and $\mathbf{w} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$. What is the coordinate vector of \mathbf{w} ? Express in the basis:

$$\begin{bmatrix} 3 \\ -1 \end{bmatrix} = c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2 = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \textcolor{red}{1} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \textcolor{blue}{2} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Let $V = \mathbb{R}^2$, $\mathcal{B} = (\mathbf{b}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix})$ and $\mathbf{w} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$. What is the coordinate vector of \mathbf{w} ? Express in the basis:

$$\begin{bmatrix} 3 \\ -1 \end{bmatrix} = c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2 = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \textcolor{red}{1} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \textcolor{blue}{2} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Therefore

$$\mathbf{w}_{\mathcal{B}} =$$

Let $V = \mathbb{R}^2$, $\mathcal{B} = (\mathbf{b}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix})$ and $\mathbf{w} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$. What is the coordinate vector of \mathbf{w} ? Express in the basis:

$$\begin{bmatrix} 3 \\ -1 \end{bmatrix} = c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2 = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \textcolor{red}{1} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \textcolor{blue}{2} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Therefore

$$\mathbf{w}_{\mathcal{B}} = \begin{bmatrix} \textcolor{red}{1} \\ \textcolor{blue}{2} \end{bmatrix}.$$

Let $V = \mathbb{R}^2$, $\mathcal{B} = (\mathbf{b}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix})$ and $\mathbf{w} = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$. What is the coordinate vector of \mathbf{w} ? Express in the basis:

$$\begin{bmatrix} 3 \\ -1 \end{bmatrix} = c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2 = c_1 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \textcolor{red}{1} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \textcolor{blue}{2} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Therefore

$$\mathbf{w}_{\mathcal{B}} = \begin{bmatrix} \textcolor{red}{1} \\ \textcolor{blue}{2} \end{bmatrix}.$$

Geometrically: this means that to reach \mathbf{w} walk 1 unit along the \mathbf{b}_1 basis vector and 2 units along the \mathbf{b}_2 basis vector.

Example with polynomials

Example with polynomials

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$.

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$. What is the coordinate vector of \mathbf{w} with respect to basis \mathcal{B} ?

Example with polynomials

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$. What is the coordinate vector of \mathbf{w} with respect to basis \mathcal{B} ?

Express \mathbf{w} in terms of the basis:

$$\mathbf{w} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + c_3\mathbf{b}_3 =$$

Example with polynomials

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$. What is the coordinate vector of \mathbf{w} with respect to basis \mathcal{B} ?

Express \mathbf{w} in terms of the basis:

$$\mathbf{w} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + c_3\mathbf{b}_3 = c_11 + c_2t + c_3t^2 =$$

Example with polynomials

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$. What is the coordinate vector of \mathbf{w} with respect to basis \mathcal{B} ?

Express \mathbf{w} in terms of the basis:

$$\mathbf{w} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + c_3\mathbf{b}_3 = c_1 1 + c_2 t + c_3 t^2 = \textcolor{red}{1} + \textcolor{blue}{2}t + \textcolor{green}{3}t^2.$$

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$. What is the coordinate vector of \mathbf{w} with respect to basis \mathcal{B} ?

Express \mathbf{w} in terms of the basis:

$$\mathbf{w} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + c_3\mathbf{b}_3 = c_11 + c_2t + c_3t^2 = \textcolor{red}{1} + \textcolor{blue}{2}t + \textcolor{green}{3}t^2.$$

Hence

$$\mathbf{w}_{\mathcal{B}} =$$

Example with polynomials

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$. What is the coordinate vector of \mathbf{w} with respect to basis \mathcal{B} ?

Express \mathbf{w} in terms of the basis:

$$\mathbf{w} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + c_3\mathbf{b}_3 = c_1 1 + c_2 t + c_3 t^2 = \textcolor{red}{1} + \textcolor{blue}{2}t + \textcolor{green}{3}t^2.$$

Hence

$$\mathbf{w}_{\mathcal{B}} = \begin{bmatrix} \textcolor{red}{1} \\ \textcolor{blue}{2} \\ \textcolor{green}{3} \end{bmatrix}$$

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$. What is the coordinate vector of \mathbf{w} with respect to basis \mathcal{B} ?

Express \mathbf{w} in terms of the basis:

$$\mathbf{w} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + c_3\mathbf{b}_3 = c_1 1 + c_2 t + c_3 t^2 = \textcolor{red}{1} + \textcolor{blue}{2}t + \textcolor{green}{3}t^2.$$

Hence

$$\mathbf{w}_{\mathcal{B}} = \begin{bmatrix} \textcolor{red}{1} \\ \textcolor{blue}{2} \\ \textcolor{green}{3} \end{bmatrix}$$

What if we take another basis?

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$. What is the coordinate vector of \mathbf{w} with respect to basis \mathcal{B} ?

Express \mathbf{w} in terms of the basis:

$$\mathbf{w} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + c_3\mathbf{b}_3 = c_1 1 + c_2 t + c_3 t^2 = \textcolor{red}{1} + \textcolor{blue}{2}t + \textcolor{green}{3}t^2.$$

Hence

$$\mathbf{w}_{\mathcal{B}} = \begin{bmatrix} \textcolor{red}{1} \\ \textcolor{blue}{2} \\ \textcolor{green}{3} \end{bmatrix}$$

What if we take another basis? Say take $\bar{\mathcal{B}} = (t^2, t, 1)$. (Different order!). Then

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$. What is the coordinate vector of \mathbf{w} with respect to basis \mathcal{B} ?

Express \mathbf{w} in terms of the basis:

$$\mathbf{w} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + c_3\mathbf{b}_3 = c_1 1 + c_2 t + c_3 t^2 = \textcolor{red}{1} + \textcolor{blue}{2}t + \textcolor{green}{3}t^2.$$

Hence

$$\mathbf{w}_{\mathcal{B}} = \begin{bmatrix} \textcolor{red}{1} \\ \textcolor{blue}{2} \\ \textcolor{green}{3} \end{bmatrix}$$

What if we take another basis? Say take $\bar{\mathcal{B}} = (t^2, t, 1)$. (Different order!). Then

$$\mathbf{w}_{\bar{\mathcal{B}}} =$$

Let $V = P_2$, the vector space of polynomials of the form $a_0 + a_1t + a_2t^2$. Let $\mathcal{B} = (\mathbf{b}_1 = 1, \mathbf{b}_2 = t, \mathbf{b}_3 = t^2)$ be the obvious basis of P_2 . Let $\mathbf{w} = 1 + 2t + 3t^2$. What is the coordinate vector of \mathbf{w} with respect to basis \mathcal{B} ?

Express \mathbf{w} in terms of the basis:

$$\mathbf{w} = c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + c_3\mathbf{b}_3 = c_11 + c_2t + c_3t^2 = \textcolor{red}{1} + \textcolor{blue}{2}t + \textcolor{green}{3}t^2.$$

Hence

$$\mathbf{w}_{\mathcal{B}} = \begin{bmatrix} \textcolor{red}{1} \\ \textcolor{blue}{2} \\ \textcolor{green}{3} \end{bmatrix}$$

What if we take another basis? Say take $\bar{\mathcal{B}} = (t^2, t, 1)$. (Different order!). Then

$$\mathbf{w}_{\bar{\mathcal{B}}} = \begin{bmatrix} \textcolor{green}{3} \\ \textcolor{blue}{2} \\ \textcolor{red}{1} \end{bmatrix}$$

Standard Coordinate Vectors

Let $V = \mathbb{R}^3$ and let $E = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ be the standard basis. If

$\mathbf{w} = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$ what is the coordinate vector with respect to the standard basis?

Let $V = \mathbb{R}^3$ and let $E = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ be the standard basis. If

$\mathbf{w} = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$ what is the coordinate vector with respect to the standard basis? Express in the basis:

$$\mathbf{w} = c_1 \mathbf{e}_1 + c_2 \mathbf{e}_2 + c_3 \mathbf{e}_3 = c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} =$$

Let $V = \mathbb{R}^3$ and let $E = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ be the standard basis. If

$\mathbf{w} = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$ what is the coordinate vector with respect to the standard basis? Express in the basis:

$$\mathbf{w} = c_1 \mathbf{e}_1 + c_2 \mathbf{e}_2 + c_3 \mathbf{e}_3 = c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 4 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Let $V = \mathbb{R}^3$ and let $E = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ be the standard basis. If

$\mathbf{w} = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$ what is the coordinate vector with respect to the standard basis? Express in the basis:

$$\mathbf{w} = c_1 \mathbf{e}_1 + c_2 \mathbf{e}_2 + c_3 \mathbf{e}_3 = c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 4 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Hence

$$\mathbf{w}_E =$$

Let $V = \mathbb{R}^3$ and let $E = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ be the standard basis. If

$\mathbf{w} = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$ what is the coordinate vector with respect to the standard basis? Express in the basis:

$$\mathbf{w} = c_1 \mathbf{e}_1 + c_2 \mathbf{e}_2 + c_3 \mathbf{e}_3 = c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 4 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Hence

$$\mathbf{w}_E = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix} =$$

Let $V = \mathbb{R}^3$ and let $E = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ be the standard basis. If

$\mathbf{w} = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$ what is the coordinate vector with respect to the standard basis? Express in the basis:

$$\mathbf{w} = c_1 \mathbf{e}_1 + c_2 \mathbf{e}_2 + c_3 \mathbf{e}_3 = c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 4 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Hence

$$\mathbf{w}_E = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix} = \mathbf{w}!$$

Let $V = \mathbb{R}^3$ and let $E = (\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ be the standard basis. If

$\mathbf{w} = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$ what is the coordinate vector with respect to the standard basis? Express in the basis:

$$\mathbf{w} = c_1 \mathbf{e}_1 + c_2 \mathbf{e}_2 + c_3 \mathbf{e}_3 = c_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 4 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Hence

$$\mathbf{w}_E = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix} = \mathbf{w}!$$

So the coordinate vector with respect to the standard basis is just the vector itself!

Linear Transformations

Definition

Let V and W be vector spaces.

Definition

A map $T : V \rightarrow W$ is a **linear transformation** if

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y})$$

for all $\mathbf{x}, \mathbf{y} \in V$ and all $c, d \in \mathbb{R}$.

Definition

Let V and W be vector spaces.

Definition

A map $T : V \rightarrow W$ is a **linear transformation** if

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y})$$

for all $\mathbf{x}, \mathbf{y} \in V$ and all $c, d \in \mathbb{R}$. In other words, a linear transformation respects **addition** and **scaling**.

Definition

Let V and W be vector spaces.

Definition

A map $T : V \rightarrow W$ is a **linear transformation** if

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y})$$

for all $\mathbf{x}, \mathbf{y} \in V$ and all $c, d \in \mathbb{R}$. In other words, a linear transformation respects **addition** and **scaling**.

Remark

It follows immediately that

- $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$
- $T(c\mathbf{x}) = cT(\mathbf{x})$

Definition

Let V and W be vector spaces.

Definition

A map $T : V \rightarrow W$ is a **linear transformation** if

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y})$$

for all $\mathbf{x}, \mathbf{y} \in V$ and all $c, d \in \mathbb{R}$. In other words, a linear transformation respects **addition** and **scaling**.

Remark

It follows immediately that

- $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$
- $T(c\mathbf{x}) = cT(\mathbf{x})$
- $T(\mathbf{0}) = \mathbf{0}$ (because $T(\mathbf{0}) = T(0 \cdot \mathbf{0}) =$

Definition

Let V and W be vector spaces.

Definition

A map $T : V \rightarrow W$ is a **linear transformation** if

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y})$$

for all $\mathbf{x}, \mathbf{y} \in V$ and all $c, d \in \mathbb{R}$. In other words, a linear transformation respects **addition** and **scaling**.

Remark

It follows immediately that

- $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$
- $T(c\mathbf{x}) = cT(\mathbf{x})$
- $T(\mathbf{0}) = \mathbf{0}$ (because $T(\mathbf{0}) = T(0 \cdot \mathbf{0}) = 0 \cdot T(\mathbf{0}) =$

Definition

Let V and W be vector spaces.

Definition

A map $T : V \rightarrow W$ is a **linear transformation** if

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y})$$

for all $\mathbf{x}, \mathbf{y} \in V$ and all $c, d \in \mathbb{R}$. In other words, a linear transformation respects **addition** and **scaling**.

Remark

It follows immediately that

- $T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y})$
- $T(c\mathbf{x}) = cT(\mathbf{x})$
- $T(\mathbf{0}) = \mathbf{0}$ (because $T(\mathbf{0}) = T(0 \cdot \mathbf{0}) = 0 \cdot T(\mathbf{0}) = \mathbf{0}$)

Some examples

First example

Example

Let $V = \mathbb{R}$, $W = \mathbb{R}$. Then the map $f(x) = 3x$ is linear. Why?

First example

Example

Let $V = \mathbb{R}$, $W = \mathbb{R}$. Then the map $f(x) = 3x$ is linear. Why?

If $x, y \in \mathbb{R}$, then

$$f(ax + by) =$$

First example

Example

Let $V = \mathbb{R}$, $W = \mathbb{R}$. Then the map $f(x) = 3x$ is linear. Why?

If $x, y \in \mathbb{R}$, then

$$f(ax + by) = 3(ax + by) =$$

First example

Example

Let $V = \mathbb{R}$, $W = \mathbb{R}$. Then the map $f(x) = 3x$ is linear. Why?

If $x, y \in \mathbb{R}$, then

$$f(ax + by) = 3(ax + by) = a \cdot 3x + b \cdot 3y$$

First example

Example

Let $V = \mathbb{R}$, $W = \mathbb{R}$. Then the map $f(x) = 3x$ is linear. Why?

If $x, y \in \mathbb{R}$, then

$$f(ax + by) = 3(ax + by) = a \cdot 3x + b \cdot 3y = af(x) + bf(y).$$

First example

Example

Let $V = \mathbb{R}$, $W = \mathbb{R}$. Then the map $f(x) = 3x$ is linear. Why?

If $x, y \in \mathbb{R}$, then

$$f(ax + by) = 3(ax + by) = a \cdot 3x + b \cdot 3y = af(x) + bf(y).$$

What about the function $g(x) = 2x - 2$?

First example

Example

Let $V = \mathbb{R}$, $W = \mathbb{R}$. Then the map $f(x) = 3x$ is linear. Why?

If $x, y \in \mathbb{R}$, then

$$f(ax + by) = 3(ax + by) = a \cdot 3x + b \cdot 3y = af(x) + bf(y).$$

What about the function $g(x) = 2x - 2$? Is this a linear transformation?

Matrices are linear transformations!

Example

Let A be an $m \times n$ matrix. Then the map $T(\mathbf{x}) = A\mathbf{x}$ is a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Why?

Matrices are linear transformations!

Example

Let A be an $m \times n$ matrix. Then the map $T(\mathbf{x}) = A\mathbf{x}$ is a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Why? Because matrix multiplication is linear.

Matrices are linear transformations!

Example

Let A be an $m \times n$ matrix. Then the map $T(\mathbf{x}) = A\mathbf{x}$ is a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Why? Because matrix multiplication is linear.

$$A(c\mathbf{x} + d\mathbf{y}) = cA\mathbf{x} + dA\mathbf{y}$$

Matrices are linear transformations!

Example

Let A be an $m \times n$ matrix. Then the map $T(\mathbf{x}) = A\mathbf{x}$ is a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Why? Because matrix multiplication is linear.

$$A(c\mathbf{x} + d\mathbf{y}) = cA\mathbf{x} + dA\mathbf{y}$$

The left-hand side is $T(c\mathbf{x} + d\mathbf{y})$ and the right-hand side is $cT(\mathbf{x}) + dT(\mathbf{y})$.

Stranger example

Example

Let P_n be the vector space of all polynomials of degree at most n .

Stranger example

Example

Let P_n be the vector space of all polynomials of degree at most n . Consider the map $T : P_n \rightarrow P_{n-1}$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

This map is linear! Why?

Stranger example

Example

Let P_n be the vector space of all polynomials of degree at most n . Consider the map $T : P_n \rightarrow P_{n-1}$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

This map is linear! Why?

Because differentiation is linear:

$$\frac{d}{dt} [ap(t) + bq(t)] =$$

Stranger example

Example

Let P_n be the vector space of all polynomials of degree at most n . Consider the map $T : P_n \rightarrow P_{n-1}$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

This map is linear! Why?

Because differentiation is linear:

$$\frac{d}{dt} [ap(t) + bq(t)] = a \frac{d}{dt} p(t) + b \frac{d}{dt} q(t).$$

Stranger example

Example

Let P_n be the vector space of all polynomials of degree at most n . Consider the map $T : P_n \rightarrow P_{n-1}$ given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

This map is linear! Why?

Because differentiation is linear:

$$\frac{d}{dt} [ap(t) + bq(t)] = a \frac{d}{dt} p(t) + b \frac{d}{dt} q(t).$$

The left-hand side is $T(ap(t) + bq(t))$ and the right-hand side is $aT(p(t)) + bT(q(t))$.