

Math 415 - Lecture 34

Discrete dynamical systems, Spectral Theorem

Wednesday November 18th 2015

Textbook reading: Chapter 5.3, Chapter 5.6 p. 297-298

Suggested practice exercises: Chapter 5.3, 2, 3, 4, 7, 8, 9, 10, 12, 14

Strang lecture: Lecture 25: Symmetric Matrices and Positive Definiteness

1 Review

Diagonalization

Suppose that A is an $n \times n$ and has independent eigenvectors $\mathbf{v}_1, \dots, \mathbf{v}_n$. Then A can be **diagonalized** as $A = PDP^{-1}$.

- the columns of P are eigenvectors
- the diagonal matrix D has the eigenvalues on the diagonal

Such a diagonalization is possible if and only if A has enough eigenvectors.

Calculating Powers

If $A = PDP^{-1}$ for some diagonal matrix D , then $A^n = PD^nP^{-1}$ for every n . This is helpful, because calculating powers of diagonal matrices is very easy!

2 Application: Discrete Dynamical Systems

Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time $t = 0$ by a vector $\mathbf{x}_{t=0}$, the **state-vector**. Then you want to know what the state \mathbf{x}_t at arbitrary time t is. How? Assume

$$\mathbf{x}_{t+1} = A\mathbf{x}_t.$$

In other words, time evolution by one time step is given by matrix multiplication by some matrix A . If we start with \mathbf{x}_0 , we get $\mathbf{x}_1 = A\mathbf{x}_0$, $\mathbf{x}_2 = A\mathbf{x}_1 = A(A\mathbf{x}_0) = A^2\mathbf{x}_0$, and more generally, the state of the system at arbitrary time $t = k$ is


$$\mathbf{x}_k = A^k \mathbf{x}_0.$$

So to solve our system we need to be able to calculate high powers of the matrix A .

2.1 Golden ratio and Fibonacci numbers

Example 1. ‘A certain man put a pair of rabbits in a place surrounded by a wall. How many pairs of rabbits can be produced from that pair in a year if it is supposed that every month each pair begets a new pair from which the second month on becomes productive?’ (Liber abbaci, chapter 12, p. 283-4)

Solution.





Fibonacci numbers: $0, 1, 1, 2, 3, 5, 8, 13, 21, 34, \dots$ Did you notice: $\frac{13}{8} = 1.625$, $\frac{21}{13} = 1.615$, $\frac{34}{21} = 1.619, \dots$ The **golden ratio** $\varphi = 1.618\dots$ Where's that from? We just showed that $F_n = \text{round} \left(\frac{1}{\sqrt{5}} \left(\frac{1+\sqrt{5}}{2} \right)^n \right)$. Therefore

$$\lim_{n \rightarrow \infty} \frac{F_{n+1}}{F_n} = \left(\frac{1 + \sqrt{5}}{2} \right).$$

Definition 2. Let A be a $n \times n$ -matrix with eigenvalues $\lambda_1, \dots, \lambda_n$. The discrete dynamical system $\mathbf{x}_{t+1} = A\mathbf{x}_t$ is

- **stable** if all eigenvalues satisfy $|\lambda_i| < 1$,
- **neutrally stable** if some $|\lambda_i| = 1$ and all the other $|\lambda_i| < 1$,
- **unstable** if at least one eigenvalue has $|\lambda_i| > 1$.

Example 3. 1. The discrete dynamical system used to construct the Fibonacci numbers is unstable.

2. If A is a Markov matrix with positive entries, then $\mathbf{x}_{t+1} = A\mathbf{x}_t$ is neutrally stable.

3. If $A = \begin{bmatrix} 0 & 4 \\ 0 & \frac{1}{2} \end{bmatrix}$, is $\mathbf{x}_{t+1} = A\mathbf{x}_t$ stable?

Solution.


3 Spectral Theorem

- Not every matrix A has a basis of eigenvectors 😞
- Special case:


Definition. A is symmetric if $A = A^T$

Theorem 1. *If A is symmetric, then it has an **orthonormal** basis of eigenvectors and all eigenvalues are real!* 😊

If Q is the matrix of eigenvectors, then Q is orthogonal. So, $Q^{-1} = Q^T$. Thus,



Remark. • The converse is also true: If A has an orthogonal basis of eigenvectors, then A is symmetric! Why?





- It is important that if A is symmetric the eigenvalues are always **real**. No complex eigenvalues!

Example 4. Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. Write A as QDQ^T .



Solution.

What does A do to the eigenvectors?




What happens to a vector \mathbf{x} ?

Suppose $\mathbf{x} = 3\mathbf{q}_1 + \mathbf{q}_2$:



Why are symmetric matrices special? Why does spectral theorem work?



Example 5. Let $A = \begin{bmatrix} 3 & -1 & 0 & 0 \\ -1 & 3 & 0 & 0 \\ 0 & 0 & 4 & 1 \\ 0 & 0 & 1 & 4 \end{bmatrix}$. Then

$$A = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 5 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{bmatrix}.$$

Let $\mathbf{x} = \begin{bmatrix} 1 \\ 1 \\ 2 \\ -2 \end{bmatrix}$. Find $A^3\mathbf{x}$.

Solution.

