

# Math 415 - Lecture 34

## Discrete dynamical systems, Spectral Theorem

Wednesday November 18th 2015

Textbook reading: Chapter 5.3, Chapter 5.6 p. 297-298

Textbook reading: Chapter 5.3, Chapter 5.6 p. 297-298

Suggested practice exercises: Chapter 5.3, 2, 3, 4, 7, 8, 9, 10, 12,

14

Textbook reading: Chapter 5.3, Chapter 5.6 p. 297-298

Suggested practice exercises: Chapter 5.3, 2, 3, 4, 7, 8, 9, 10, 12,  
14

Strang lecture: Lecture 25: Symmetric Matrices and Positive  
Definiteness

# Review

## Diagonalization

Suppose that  $A$  is an  $n \times n$  and has independent eigenvectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$ . Then  $A$  can be **diagonalized** as  $A = PDP^{-1}$ .

Such a diagonalization is possible if and only if  $A$  has an eigenbasis.

## Diagonalization

Suppose that  $A$  is an  $n \times n$  and has independent eigenvectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$ . Then  $A$  can be **diagonalized** as  $A = PDP^{-1}$ .

- the columns of  $P$  are eigenvectors

Such a diagonalization is possible if and only if  $A$  has an eigenbasis.

## Diagonalization

Suppose that  $A$  is an  $n \times n$  and has independent eigenvectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$ . Then  $A$  can be **diagonalized** as  $A = PDP^{-1}$ .

- the columns of  $P$  are eigenvectors
- the diagonal matrix  $D$  has the eigenvalues on the diagonal

Such a diagonalization is possible if and only if  $A$  has an eigenbasis.



## Diagonalization

Suppose that  $A$  is an  $n \times n$  and has independent eigenvectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$ . Then  $A$  can be **diagonalized** as  $A = PDP^{-1}$ .

- the columns of  $P$  are eigenvectors
- the diagonal matrix  $D$  has the eigenvalues on the diagonal

Such a diagonalization is possible if and only if  $A$  has an eigenbasis.

## Calculating Powers

If  $A = PDP^{-1}$  for some diagonal matrix  $D$ , then  $A^n = PD^nP^{-1}$  for every  $n$ . This is helpful, because calculating powers of diagonal matrices is very easy!

## Application: Discrete Dynamical Systems

Suppose you want to describe the evolution of some part of the world.

Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time  $t = 0$  by a vector  $\mathbf{x}_{t=0}$ , the **state-vector**.

Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time  $t = 0$  by a vector  $\mathbf{x}_{t=0}$ , the **state-vector**. For a simple system the state vector  $\mathbf{x}_t$  might have 2 components, for a complicated system there might be thousands of components.

Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time  $t = 0$  by a vector  $\mathbf{x}_{t=0}$ , the **state-vector**. For a simple system the state vector  $\mathbf{x}_t$  might have 2 components, for a complicated system there might be thousands of components. Then you want to know what the state  $\mathbf{x}_t$  at arbitrary time  $t$  is.

Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time  $t = 0$  by a vector  $\mathbf{x}_{t=0}$ , the **state-vector**. For a simple system the state vector  $\mathbf{x}_t$  might have 2 components, for a complicated system there might be thousands of components. Then you want to know what the state  $\mathbf{x}_t$  at arbitrary time  $t$  is. How?

Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time  $t = 0$  by a vector  $\mathbf{x}_{t=0}$ , the **state-vector**. For a simple system the state vector  $\mathbf{x}_t$  might have 2 components, for a complicated system there might be thousands of components. Then you want to know what the state  $\mathbf{x}_t$  at arbitrary time  $t$  is. How? Assume

$$\mathbf{x}_{t+1} = A\mathbf{x}_t.$$



Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time  $t = 0$  by a vector  $\mathbf{x}_{t=0}$ , the **state-vector**. For a simple system the state vector  $\mathbf{x}_t$  might have 2 components, for a complicated system there might be thousands of components. Then you want to know what the state  $\mathbf{x}_t$  at arbitrary time  $t$  is. How? Assume

$$\mathbf{x}_{t+1} = A\mathbf{x}_t.$$

In other words, time evolution by one time step is given by matrix multiplication by some matrix  $A$ .

Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time  $t = 0$  by a vector  $\mathbf{x}_{t=0}$ , the **state-vector**. For a simple system the state vector  $\mathbf{x}_t$  might have 2 components, for a complicated system there might be thousands of components. Then you want to know what the state  $\mathbf{x}_t$  at arbitrary time  $t$  is. How? Assume

$$\mathbf{x}_{t+1} = A\mathbf{x}_t.$$

In other words, time evolution by one time step is given by matrix multiplication by some matrix  $A$ . If we start with  $\mathbf{x}_0$ , we get  $\mathbf{x}_1 = A\mathbf{x}_0$ ,  $\mathbf{x}_2 = A\mathbf{x}_1 = A(A\mathbf{x}_0) = A^2\mathbf{x}_0$ , and more generally, the state of the system at arbitrary time  $t = k$  is

Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time  $t = 0$  by a vector  $\mathbf{x}_{t=0}$ , the **state-vector**. For a simple system the state vector  $\mathbf{x}_t$  might have 2 components, for a complicated system there might be thousands of components. Then you want to know what the state  $\mathbf{x}_t$  at arbitrary time  $t$  is. How? Assume

$$\mathbf{x}_{t+1} = A\mathbf{x}_t.$$

In other words, time evolution by one time step is given by matrix multiplication by some matrix  $A$ . If we start with  $\mathbf{x}_0$ , we get  $\mathbf{x}_1 = A\mathbf{x}_0$ ,  $\mathbf{x}_2 = A\mathbf{x}_1 = A(A\mathbf{x}_0) = A^2\mathbf{x}_0$ , and more generally, the state of the system at arbitrary time  $t = k$  is

$$\mathbf{x}_k = A^k\mathbf{x}_0.$$

Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time  $t = 0$  by a vector  $\mathbf{x}_{t=0}$ , the **state-vector**. For a simple system the state vector  $\mathbf{x}_t$  might have 2 components, for a complicated system there might be thousands of components. Then you want to know what the state  $\mathbf{x}_t$  at arbitrary time  $t$  is. How? Assume

$$\mathbf{x}_{t+1} = A\mathbf{x}_t.$$

In other words, time evolution by one time step is given by matrix multiplication by some matrix  $A$ . If we start with  $\mathbf{x}_0$ , we get  $\mathbf{x}_1 = A\mathbf{x}_0$ ,  $\mathbf{x}_2 = A\mathbf{x}_1 = A(A\mathbf{x}_0) = A^2\mathbf{x}_0$ , and more generally, the state of the system at arbitrary time  $t = k$  is

$$\mathbf{x}_k = A^k\mathbf{x}_0.$$

So to solve our system we need to be able to calculate high powers of the matrix  $A$ .

Suppose you want to describe the evolution of some part of the world. Describe the **state** of your part of the world at time  $t = 0$  by a vector  $\mathbf{x}_{t=0}$ , the **state-vector**. For a simple system the state vector  $\mathbf{x}_t$  might have 2 components, for a complicated system there might be thousands of components. Then you want to know what the state  $\mathbf{x}_t$  at arbitrary time  $t$  is. How? Assume

$$\mathbf{x}_{t+1} = A\mathbf{x}_t.$$

In other words, time evolution by one time step is given by matrix multiplication by some matrix  $A$ . If we start with  $\mathbf{x}_0$ , we get  $\mathbf{x}_1 = A\mathbf{x}_0$ ,  $\mathbf{x}_2 = A\mathbf{x}_1 = A(A\mathbf{x}_0) = A^2\mathbf{x}_0$ , and more generally, the state of the system at arbitrary time  $t = k$  is

$$\mathbf{x}_k = A^k\mathbf{x}_0.$$

So to solve our system we need to be able to calculate high powers of the matrix  $A$ . Use eigenbasis of  $A$  for this.

## Golden ratio and Fibonacci numbers

## Example

*'A certain man put a pair of rabbits in a place surrounded by a wall. How many pairs of rabbits can be produced from that pair in a year if it is supposed that every month each pair begets a new pair from which the second month on becomes productive?' (Liber abbaci, chapter 12, p. 283-4)*

## Example

*'A certain man put a pair of rabbits in a place surrounded by a wall. How many pairs of rabbits can be produced from that pair in a year if it is supposed that every month each pair begets a new pair from which the second month on becomes productive?' (Liber abbaci, chapter 12, p. 283-4)*

## Solution

Idea: use discrete dynamical system to produce the Fibonacci numbers.



## Example

*'A certain man put a pair of rabbits in a place surrounded by a wall. How many pairs of rabbits can be produced from that pair in a year if it is supposed that every month each pair begets a new pair from which the second month on becomes productive?' (Liber abbaci, chapter 12, p. 283-4)*

## Solution

Idea: use discrete dynamical system to produce the Fibonacci numbers.

- $F_{n+1} = F_n + F_{n-1}$

## Example

*'A certain man put a pair of rabbits in a place surrounded by a wall. How many pairs of rabbits can be produced from that pair in a year if it is supposed that every month each pair begets a new pair from which the second month on becomes productive?' (Liber abbaci, chapter 12, p. 283-4)*

## Solution

Idea: use discrete dynamical system to produce the Fibonacci numbers.

$$\bullet F_{n+1} = F_n + F_{n-1} \implies \begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} F_n \\ F_{n-1} \end{bmatrix}$$

## Example

*'A certain man put a pair of rabbits in a place surrounded by a wall. How many pairs of rabbits can be produced from that pair in a year if it is supposed that every month each pair begets a new pair from which the second month on becomes productive?' (Liber abbaci, chapter 12, p. 283-4)*

## Solution

Idea: use discrete dynamical system to produce the Fibonacci numbers.

- $F_{n+1} = F_n + F_{n-1} \implies \begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} F_n \\ F_{n-1} \end{bmatrix}$
- Hence  $\begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n \begin{bmatrix} F_1 \\ F_0 \end{bmatrix} \quad \left( \begin{bmatrix} F_1 \\ F_0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right)$

## Example

*'A certain man put a pair of rabbits in a place surrounded by a wall. How many pairs of rabbits can be produced from that pair in a year if it is supposed that every month each pair begets a new pair from which the second month on becomes productive?' (Liber abbaci, chapter 12, p. 283-4)*

## Solution

Idea: use discrete dynamical system to produce the Fibonacci numbers.

- $F_{n+1} = F_n + F_{n-1} \implies \begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} F_n \\ F_{n-1} \end{bmatrix}$
- Hence  $\begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n \begin{bmatrix} F_1 \\ F_0 \end{bmatrix} \quad \left( \begin{bmatrix} F_1 \\ F_0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right)$
- But we know how to compute  $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n$  or  $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n \begin{bmatrix} 1 \\ 0 \end{bmatrix}!$



## Solution

- The characteristic polynomial of  $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  is  $\lambda^2 - \lambda - 1$

## Solution

- The characteristic polynomial of  $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  is  $\lambda^2 - \lambda - 1$
- The eigenvalues are  $\lambda_1 = \frac{1+\sqrt{5}}{2} \approx 1.618$  (the golden mean!) and  $\lambda_2 = \frac{1-\sqrt{5}}{2} \approx -0.618$

## Solution

- The characteristic polynomial of  $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  is  $\lambda^2 - \lambda - 1$
- The eigenvalues are  $\lambda_1 = \frac{1+\sqrt{5}}{2} \approx 1.618$  (the golden mean!) and  $\lambda_2 = \frac{1-\sqrt{5}}{2} \approx -0.618$
- Corresponding eigenvectors:  $\mathbf{v}_1 = \begin{bmatrix} \lambda_1 \\ 1 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} \lambda_2 \\ 1 \end{bmatrix}$



## Solution

- The characteristic polynomial of  $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  is  $\lambda^2 - \lambda - 1$
- The eigenvalues are  $\lambda_1 = \frac{1+\sqrt{5}}{2} \approx 1.618$  (the golden mean!) and  $\lambda_2 = \frac{1-\sqrt{5}}{2} \approx -0.618$
- Corresponding eigenvectors:  $\mathbf{v}_1 = \begin{bmatrix} \lambda_1 \\ 1 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} \lambda_2 \\ 1 \end{bmatrix}$
- Write  $\begin{bmatrix} 1 \\ 0 \end{bmatrix} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2$ . ( $c_1 = \frac{1}{\sqrt{5}}$ ,  $c_2 = -\frac{1}{\sqrt{5}}$ )

## Solution

- The characteristic polynomial of  $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  is  $\lambda^2 - \lambda - 1$
- The eigenvalues are  $\lambda_1 = \frac{1+\sqrt{5}}{2} \approx 1.618$  (the golden mean!) and  $\lambda_2 = \frac{1-\sqrt{5}}{2} \approx -0.618$
- Corresponding eigenvectors:  $\mathbf{v}_1 = \begin{bmatrix} \lambda_1 \\ 1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} \lambda_2 \\ 1 \end{bmatrix}$
- Write  $\begin{bmatrix} 1 \\ 0 \end{bmatrix} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2$ . ( $c_1 = \frac{1}{\sqrt{5}}, c_2 = -\frac{1}{\sqrt{5}}$ )
- $\begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = A^n \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \lambda_1^n c_1 \mathbf{v}_1 + \lambda_2^n c_2 \mathbf{v}_2$

## Solution

- The characteristic polynomial of  $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  is  $\lambda^2 - \lambda - 1$
- The eigenvalues are  $\lambda_1 = \frac{1+\sqrt{5}}{2} \approx 1.618$  (the golden mean!) and  $\lambda_2 = \frac{1-\sqrt{5}}{2} \approx -0.618$
- Corresponding eigenvectors:  $\mathbf{v}_1 = \begin{bmatrix} \lambda_1 \\ 1 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} \lambda_2 \\ 1 \end{bmatrix}$
- Write  $\begin{bmatrix} 1 \\ 0 \end{bmatrix} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2$ . ( $c_1 = \frac{1}{\sqrt{5}}$ ,  $c_2 = -\frac{1}{\sqrt{5}}$ )
- $\begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = A^n \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \lambda_1^n c_1 \mathbf{v}_1 + \lambda_2^n c_2 \mathbf{v}_2$
- Hence,  $F_n = \lambda_1^n c_1 + \lambda_2^n c_2 = \frac{1}{\sqrt{5}} \left[ \left( \frac{1+\sqrt{5}}{2} \right)^n - \left( \frac{1-\sqrt{5}}{2} \right)^n \right]$ .  
That is **Binet's formula**.

## Solution


- The characteristic polynomial of  $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$  is  $\lambda^2 - \lambda - 1$
- The eigenvalues are  $\lambda_1 = \frac{1+\sqrt{5}}{2} \approx 1.618$  (the golden mean!) and  $\lambda_2 = \frac{1-\sqrt{5}}{2} \approx -0.618$
- Corresponding eigenvectors:  $\mathbf{v}_1 = \begin{bmatrix} \lambda_1 \\ 1 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} \lambda_2 \\ 1 \end{bmatrix}$
- Write  $\begin{bmatrix} 1 \\ 0 \end{bmatrix} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2$ . ( $c_1 = \frac{1}{\sqrt{5}}$ ,  $c_2 = -\frac{1}{\sqrt{5}}$ )
- $\begin{bmatrix} F_{n+1} \\ F_n \end{bmatrix} = A^n \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \lambda_1^n c_1 \mathbf{v}_1 + \lambda_2^n c_2 \mathbf{v}_2$
- Hence,  $F_n = \lambda_1^n c_1 + \lambda_2^n c_2 = \frac{1}{\sqrt{5}} \left[ \left( \frac{1+\sqrt{5}}{2} \right)^n - \left( \frac{1-\sqrt{5}}{2} \right)^n \right]$ .  
That is **Binet's formula**.
- but  $|\lambda_2| < 1$ , so  $F_n \approx \lambda_1^n c_1 = \frac{1}{\sqrt{5}} \left( \frac{1+\sqrt{5}}{2} \right)^n$ . In fact,  
 $F_n = \text{round} \left( \frac{1}{\sqrt{5}} \left( \frac{1+\sqrt{5}}{2} \right)^n \right)$ .

## Golden ratio and Fibonacci numbers



Fibonacci numbers: 0, 1, 1, 2, 3, 5, 8, 13, 21, 34, ...

## Golden ratio and Fibonacci numbers



Fibonacci numbers: 0, 1, 1, 2, 3, 5, 8, 13, 21, 34, ...

Did you notice:  $\frac{13}{8} = 1.625$ ,  $\frac{21}{13} = 1.615$ ,  $\frac{34}{21} = 1.619$ , ...

## Golden ratio and Fibonacci numbers



Fibonacci numbers: 0, 1, 1, 2, 3, 5, 8, 13, 21, 34, ...

Did you notice:  $\frac{13}{8} = 1.625$ ,  $\frac{21}{13} = 1.615$ ,  $\frac{34}{21} = 1.619$ , ...

The **golden ratio**  $\varphi = 1.618...$  Where's that from? We just showed that  $F_n = \text{round} \left( \frac{1}{\sqrt{5}} \left( \frac{1+\sqrt{5}}{2} \right)^n \right)$ . Therefore

$$\lim_{n \rightarrow \infty} \frac{F_{n+1}}{F_n} = \left( \frac{1 + \sqrt{5}}{2} \right).$$

## Definition

Let  $A$  be a  $n \times n$ -matrix with eigenvalues  $\lambda_1, \dots, \lambda_n$ . The discrete dynamical system  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  is

- **stable** if all eigenvalues satisfy  $|\lambda_i| < 1$ ,



## Definition

Let  $A$  be a  $n \times n$ -matrix with eigenvalues  $\lambda_1, \dots, \lambda_n$ . The discrete dynamical system  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  is

- **stable** if all eigenvalues satisfy  $|\lambda_i| < 1$ ,
- **neutrally stable** if some  $|\lambda_i| = 1$  and all the other  $|\lambda_i| < 1$ ,

## Definition

Let  $A$  be a  $n \times n$ -matrix with eigenvalues  $\lambda_1, \dots, \lambda_n$ . The discrete dynamical system  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  is

- **stable** if all eigenvalues satisfy  $|\lambda_i| < 1$ ,
- **neutrally stable** if some  $|\lambda_i| = 1$  and all the other  $|\lambda_i| < 1$ ,
- **unstable** if at least one eigenvalue has  $|\lambda_i| > 1$ .

## Definition

Let  $A$  be a  $n \times n$ -matrix with eigenvalues  $\lambda_1, \dots, \lambda_n$ . The discrete dynamical system  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  is

- **stable** if all eigenvalues satisfy  $|\lambda_i| < 1$ ,
- **neutrally stable** if some  $|\lambda_i| = 1$  and all the other  $|\lambda_i| < 1$ ,
- **unstable** if at least one eigenvalue has  $|\lambda_i| > 1$ .

## Example

1. The discrete dynamical system used to construct the Fibonacci numbers is unstable.

## Definition

Let  $A$  be a  $n \times n$ -matrix with eigenvalues  $\lambda_1, \dots, \lambda_n$ . The discrete dynamical system  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  is

- **stable** if all eigenvalues satisfy  $|\lambda_i| < 1$ ,
- **neutrally stable** if some  $|\lambda_i| = 1$  and all the other  $|\lambda_i| < 1$ ,
- **unstable** if at least one eigenvalue has  $|\lambda_i| > 1$ .

## Example

1. The discrete dynamical system used to construct the Fibonacci numbers is unstable.
2. If  $A$  is a Markov matrix with positive entries, then  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  is neutrally stable.

## Definition

Let  $A$  be a  $n \times n$ -matrix with eigenvalues  $\lambda_1, \dots, \lambda_n$ . The discrete dynamical system  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  is

- **stable** if all eigenvalues satisfy  $|\lambda_i| < 1$ ,
- **neutrally stable** if some  $|\lambda_i| = 1$  and all the other  $|\lambda_i| < 1$ ,
- **unstable** if at least one eigenvalue has  $|\lambda_i| > 1$ .

## Example

1. The discrete dynamical system used to construct the Fibonacci numbers is unstable.
2. If  $A$  is a Markov matrix with positive entries, then  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  is neutrally stable.
3. If  $A = \begin{bmatrix} 0 & 4 \\ 0 & \frac{1}{2} \end{bmatrix}$ , is  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  stable?

## Definition

Let  $A$  be a  $n \times n$ -matrix with eigenvalues  $\lambda_1, \dots, \lambda_n$ . The discrete dynamical system  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  is

- **stable** if all eigenvalues satisfy  $|\lambda_i| < 1$ ,
- **neutrally stable** if some  $|\lambda_i| = 1$  and all the other  $|\lambda_i| < 1$ ,
- **unstable** if at least one eigenvalue has  $|\lambda_i| > 1$ .

## Example

1. The discrete dynamical system used to construct the Fibonacci numbers is unstable.
2. If  $A$  is a Markov matrix with positive entries, then  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  is neutrally stable.
3. If  $A = \begin{bmatrix} 0 & 4 \\ 0 & \frac{1}{2} \end{bmatrix}$ , is  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  stable?

## Spectral Theorem

- Not every matrix  $A$  has a basis of eigenvectors



- Not every matrix  $A$  has a basis of eigenvectors 😞

- Not every matrix  $A$  has a basis of eigenvectors
- Special case:



- Not every matrix  $A$  has a basis of eigenvectors
- Special case:



### Definition

$A$  is symmetric if  $A = A^T$

- Not every matrix  $A$  has a basis of eigenvectors
- Special case:



### Definition

$A$  is symmetric if  $A = A^T$

### Theorem

If  $A$  is symmetric, then it has an *orthonormal* basis of eigenvectors  
and all eigenvalues are *real*!

- Not every matrix  $A$  has a basis of eigenvectors
- Special case:



### Definition

$A$  is symmetric if  $A = A^T$

### Theorem

If  $A$  is symmetric, then it has an *orthonormal* basis of eigenvectors

and all eigenvalues are *real!*



- Not every matrix  $A$  has a basis of eigenvectors
- Special case:



### Definition

$A$  is symmetric if  $A = A^T$

### Theorem

If  $A$  is symmetric, then it has an *orthonormal* basis of eigenvectors  
and all eigenvalues are *real!*



If  $Q$  is the matrix of eigenvectors, then  $Q$  is orthogonal.

- Not every matrix  $A$  has a basis of eigenvectors
- Special case:



### Definition

$A$  is symmetric if  $A = A^T$

### Theorem

If  $A$  is symmetric, then it has an *orthonormal* basis of eigenvectors

and all eigenvalues are *real!*



If  $Q$  is the matrix of eigenvectors, then  $Q$  is orthogonal. So,  
 $Q^{-1} =$

- Not every matrix  $A$  has a basis of eigenvectors
- Special case:



### Definition

$A$  is symmetric if  $A = A^T$

### Theorem

If  $A$  is symmetric, then it has an *orthonormal* basis of eigenvectors

and all eigenvalues are *real!*



If  $Q$  is the matrix of eigenvectors, then  $Q$  is orthogonal. So,  
 $Q^{-1} = Q^T$ . Thus,



- Not every matrix  $A$  has a basis of eigenvectors
- Special case:



### Definition

$A$  is symmetric if  $A = A^T$

### Theorem

If  $A$  is symmetric, then it has an *orthonormal* basis of eigenvectors

and all eigenvalues are *real!*



If  $Q$  is the matrix of eigenvectors, then  $Q$  is orthogonal. So,  
 $Q^{-1} = Q^T$ . Thus,

$$A = QDQ^{-1} =$$

- Not every matrix  $A$  has a basis of eigenvectors
- Special case:



### Definition

$A$  is symmetric if  $A = A^T$

### Theorem

If  $A$  is symmetric, then it has an *orthonormal* basis of eigenvectors

and all eigenvalues are *real*!



If  $Q$  is the matrix of eigenvectors, then  $Q$  is orthogonal. So,  
 $Q^{-1} = Q^T$ . Thus,

$$A = QDQ^{-1} = QDQ^T$$

and

$$D = Q^{-1}AQ =$$

- Not every matrix  $A$  has a basis of eigenvectors
- Special case:



### Definition

$A$  is symmetric if  $A = A^T$

### Theorem

If  $A$  is symmetric, then it has an *orthonormal* basis of eigenvectors

and all eigenvalues are *real*!



If  $Q$  is the matrix of eigenvectors, then  $Q$  is orthogonal. So,  
 $Q^{-1} = Q^T$ . Thus,

$$A = QDQ^{-1} = QDQ^T$$

and

$$D = Q^{-1}AQ = Q^T AQ$$

## Remark

- The converse is also true:

## Remark

- The converse is also true: If  $A$  has an orthogonal basis of eigenvectors, then  $A$  is symmetric!

## Remark

- The converse is also true: If  $A$  has an orthogonal basis of eigenvectors, then  $A$  is symmetric!  
Why?

## Remark

- The converse is also true: If  $A$  has an orthogonal basis of eigenvectors, then  $A$  is symmetric!  
Why?
- It is important that if  $A$  is symmetric the eigenvalues are always **real**.

## Remark

- The converse is also true: If  $A$  has an orthogonal basis of eigenvectors, then  $A$  is symmetric!  
Why?
- It is important that if  $A$  is symmetric the eigenvalues are always **real**.  
No complex eigenvalues!



### Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . Write  $A$  as  $QDQ^T$ .

### Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . Write  $A$  as  $QDQ^T$ .

### Solution

### Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . Write  $A$  as  $QDQ^T$ .

### Solution

We've seen this matrix before!

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . Write  $A$  as  $QDQ^T$ .

## Solution

We've seen this matrix before!

**Find eigenvalues:** We have seen that  $A$  has eigenvalues 2 and 4.

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . Write  $A$  as  $QDQ^T$ .

## Solution

We've seen this matrix before!

Find eigenvalues: We have seen that  $A$  has eigenvalues 2 and 4.

Find eigenbasis corresponding to eigenvalues:

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . Write  $A$  as  $QDQ^T$ .

## Solution

We've seen this matrix before!

**Find eigenvalues:** We have seen that  $A$  has eigenvalues 2 and 4.

**Find eigenbasis corresponding to eigenvalues:**

$\lambda_1 = 2$ : We have seen  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . Write  $A$  as  $QDQ^T$ .

## Solution

We've seen this matrix before!

**Find eigenvalues:** We have seen that  $A$  has eigenvalues 2 and 4.

**Find eigenbasis corresponding to eigenvalues:**

$\lambda_1 = 2$ : We have seen  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ . Normalized, we get

$$\mathbf{q}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . Write  $A$  as  $QDQ^T$ .

## Solution

We've seen this matrix before!

**Find eigenvalues:** We have seen that  $A$  has eigenvalues 2 and 4.

**Find eigenbasis corresponding to eigenvalues:**

$\lambda_1 = 2$ : We have seen  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ . Normalized, we get

$$\mathbf{q}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

$\lambda_2 = 4$ : We have seen  $\mathbf{x}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .



## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . Write  $A$  as  $QDQ^T$ .

## Solution

We've seen this matrix before!

**Find eigenvalues:** We have seen that  $A$  has eigenvalues 2 and 4.

**Find eigenbasis corresponding to eigenvalues:**

$\lambda_1 = 2$ : We have seen  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ . Normalized, we get

$$\mathbf{q}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

$\lambda_2 = 4$ : We have seen  $\mathbf{x}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . Normalized, we get

$$\mathbf{q}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$


## Solution (continued)

Write  $D$ :  $D = \begin{bmatrix} 2 & 0 \\ 0 & 4 \end{bmatrix}$



Write  $Q$ :  $Q = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$

Get  $A = QDQ^T$ :  $\begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 4 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$

What does  $A$  do to the eigenvectors?




What does  $A$  do to the eigenvectors?



What happens to a vector  $\mathbf{x}$ ?



What happens to a vector  $\mathbf{x}$ ?

Suppose  $\mathbf{x} = 3\mathbf{q}_1 + \mathbf{q}_2$ :



What happens to a vector  $\mathbf{x}$ ?

Suppose  $\mathbf{x} = 3\mathbf{q}_1 + \mathbf{q}_2$ :



Why are symmetric matrices special?



Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal!

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Let's show  $\mathbf{x} \cdot \mathbf{y} = 0$ :

$$\lambda_1(\mathbf{x} \cdot \mathbf{y}) =$$

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Let's show  $\mathbf{x} \cdot \mathbf{y} = 0$ :

$$\lambda_1(\mathbf{x} \cdot \mathbf{y}) = (\lambda_1\mathbf{x}) \cdot \mathbf{y}$$

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Let's show  $\mathbf{x} \cdot \mathbf{y} = 0$ :

$$\begin{aligned}\lambda_1(\mathbf{x} \cdot \mathbf{y}) &= (\lambda_1\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x}) \cdot \mathbf{y}\end{aligned}$$

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Let's show  $\mathbf{x} \cdot \mathbf{y} = 0$ :

$$\begin{aligned}\lambda_1(\mathbf{x} \cdot \mathbf{y}) &= (\lambda_1\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x})^T \mathbf{y}\end{aligned}$$

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Let's show  $\mathbf{x} \cdot \mathbf{y} = 0$ :

$$\begin{aligned}\lambda_1(\mathbf{x} \cdot \mathbf{y}) &= (\lambda_1\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x})^T \mathbf{y} \\ &= \mathbf{x}^T A^T \mathbf{y}\end{aligned}$$



Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Let's show  $\mathbf{x} \cdot \mathbf{y} = 0$ :

$$\begin{aligned}\lambda_1(\mathbf{x} \cdot \mathbf{y}) &= (\lambda_1\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x})^T \mathbf{y} \\ &= \mathbf{x}^T A^T \mathbf{y} \\ &= \mathbf{x}^T A \mathbf{y} \quad \longleftarrow \text{because } A \text{ is symmetric!}\end{aligned}$$

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Let's show  $\mathbf{x} \cdot \mathbf{y} = 0$ :

$$\begin{aligned}\lambda_1(\mathbf{x} \cdot \mathbf{y}) &= (\lambda_1\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x})^T \mathbf{y} \\ &= \mathbf{x}^T A^T \mathbf{y} \\ &= \mathbf{x}^T A \mathbf{y} \quad \leftarrow \text{because } A \text{ is symmetric!} \\ &= \mathbf{x} \cdot (A\mathbf{y})\end{aligned}$$

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Let's show  $\mathbf{x} \cdot \mathbf{y} = 0$ :

$$\begin{aligned}\lambda_1(\mathbf{x} \cdot \mathbf{y}) &= (\lambda_1\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x})^T \mathbf{y} \\ &= \mathbf{x}^T A^T \mathbf{y} \\ &= \mathbf{x}^T A \mathbf{y} \quad \leftarrow \text{because } A \text{ is symmetric!} \\ &= \mathbf{x} \cdot (A\mathbf{y}) \\ &= \lambda_2(\mathbf{x} \cdot \mathbf{y})\end{aligned}$$

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Let's show  $\mathbf{x} \cdot \mathbf{y} = 0$ :

$$\begin{aligned}\lambda_1(\mathbf{x} \cdot \mathbf{y}) &= (\lambda_1\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x})^T \mathbf{y} \\ &= \mathbf{x}^T A^T \mathbf{y} \\ &= \mathbf{x}^T A \mathbf{y} \quad \leftarrow \text{because } A \text{ is symmetric!} \\ &= \mathbf{x} \cdot (A\mathbf{y}) \\ &= \lambda_2(\mathbf{x} \cdot \mathbf{y})\end{aligned}$$

Since  $\lambda_1 \neq \lambda_2$ , must have  $\mathbf{x} \cdot \mathbf{y} = 0$ !

Why are symmetric matrices special? Why does spectral theorem work? If  $A = A^T$ , and if

$$A\mathbf{x} = \lambda_1\mathbf{x} \text{ and } A\mathbf{y} = \lambda_2\mathbf{y}$$

(for  $\lambda_1 \neq \lambda_2$ ), then  $\mathbf{x}$  and  $\mathbf{y}$  **must** be orthogonal! Why?

Let's show  $\mathbf{x} \cdot \mathbf{y} = 0$ :

$$\begin{aligned}\lambda_1(\mathbf{x} \cdot \mathbf{y}) &= (\lambda_1\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x}) \cdot \mathbf{y} \\ &= (A\mathbf{x})^T \mathbf{y} \\ &= \mathbf{x}^T A^T \mathbf{y} \\ &= \mathbf{x}^T A \mathbf{y} \quad \leftarrow \text{because } A \text{ is symmetric!} \\ &= \mathbf{x} \cdot (A\mathbf{y}) \\ &= \lambda_2(\mathbf{x} \cdot \mathbf{y})\end{aligned}$$

Since  $\lambda_1 \neq \lambda_2$ , must have  $\mathbf{x} \cdot \mathbf{y} = 0$ ! By a similar argument you can show that the eigenvalues of a symmetric matrix **must** be real.

## Example

Let  $A = \begin{bmatrix} 3 & -1 & 0 & 0 \\ -1 & 3 & 0 & 0 \\ 0 & 0 & 4 & 1 \\ 0 & 0 & 1 & 4 \end{bmatrix}$ .

## Example

$$\text{Let } A = \begin{bmatrix} 3 & -1 & 0 & 0 \\ -1 & 3 & 0 & 0 \\ 0 & 0 & 4 & 1 \\ 0 & 0 & 1 & 4 \end{bmatrix}. \text{ Then}$$

$$A = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 5 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{bmatrix}.$$

## Example

$$\text{Let } A = \begin{bmatrix} 3 & -1 & 0 & 0 \\ -1 & 3 & 0 & 0 \\ 0 & 0 & 4 & 1 \\ 0 & 0 & 1 & 4 \end{bmatrix}. \text{ Then}$$

$$A = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 5 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{bmatrix}.$$

$$\text{Let } \mathbf{x} = \begin{bmatrix} 1 \\ 1 \\ 2 \\ -2 \end{bmatrix}.$$



## Example

$$\text{Let } A = \begin{bmatrix} 3 & -1 & 0 & 0 \\ -1 & 3 & 0 & 0 \\ 0 & 0 & 4 & 1 \\ 0 & 0 & 1 & 4 \end{bmatrix}. \text{ Then}$$

$$A = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 5 \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{bmatrix}.$$

$$\text{Let } \mathbf{x} = \begin{bmatrix} 1 \\ 1 \\ 2 \\ -2 \end{bmatrix}. \text{ Find } A^3 \mathbf{x}.$$