

# Math 415 - Lecture 18

## Inner Product and Orthogonality

Wednesday October 7th 2015

# Math 415 - Lecture 18

## Inner Product and Orthogonality

Wednesday October 7th 2015

Textbook reading: Ch 3.1

Textbook reading: Ch 3.1

Suggested practice exercises: 1, 2, 4, 5, 14, 16

Textbook reading: Ch 3.1

Suggested practice exercises: 1, 2, 4, 5, 14, 16

Khan Academy video: Vector Dot Product and Vector Length

Textbook reading: Ch 3.1

Suggested practice exercises: 1, 2, 4, 5, 14, 16

Khan Academy video: Vector Dot Product and Vector Length

Strang lectures: Lecture 30: Linear Transformations / Lecture 14:  
Orthogonality

Textbook reading: Ch 3.1

Suggested practice exercises: 1, 2, 4, 5, 14, 16

Khan Academy video: Vector Dot Product and Vector Length

Strang lectures: Lecture 30: Linear Transformations / Lecture 14:  
Orthogonality

Applications: Information retrieval

## Review



- A **linear map**  $T : V \rightarrow W$  satisfies

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y}).$$

- A **linear map**  $T : V \rightarrow W$  satisfies

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y}).$$

- $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T(\mathbf{x}) = A\mathbf{x}$  is linear.  $A$  is the matrix representing  $T$  in the standard bases.

- A **linear map**  $T : V \rightarrow W$  satisfies

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y}).$$

- $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T(\mathbf{x}) = A\mathbf{x}$  is linear.  $A$  is the matrix representing  $T$  in the standard bases.

- A **linear map**  $T : V \rightarrow W$  satisfies

$$T(c\mathbf{x} + d\mathbf{y}) = cT(\mathbf{x}) + dT(\mathbf{y}).$$

- $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by  $T(\mathbf{x}) = A\mathbf{x}$  is linear.  $A$  is the matrix representing  $T$  in the standard bases.  
For example,  $T(\mathbf{e}_1) = A\mathbf{e}_1 =$  first column of  $A$ .

$$\left( \mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} \right)$$

- Any  $T : V \rightarrow W$  can be represented by a matrix.

**What is the Point?** Why write  $T: V \rightarrow W$  as a matrix?

**What is the Point?** Why write  $T: V \rightarrow W$  as a matrix?

- Replace obscure computations in  $V$  and  $W$  by transparent computations with matrices.

**What is the Point?** Why write  $T: V \rightarrow W$  as a matrix?

- Replace obscure computations in  $V$  and  $W$  by transparent computations with matrices.
- Even if  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  (already have standard coordinates),  $T$  may be simpler in a different coordinate system.

**Summary:** Given  $\mathbf{v}$  in  $V$ , want to calculate  $T(\mathbf{v})$  in  $W$ .



**Summary:** Given  $\mathbf{v}$  in  $V$ , want to calculate  $T(\mathbf{v})$  in  $W$ . Take an input basis  $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$  and an output basis  $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$ .

**Summary:** Given  $\mathbf{v}$  in  $V$ , want to calculate  $T(\mathbf{v})$  in  $W$ . Take an input basis  $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$  and an output basis  $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$ .

- The abstract input vector  $\mathbf{v}$  and the coordinate vector  $\mathbf{v}_{\mathcal{A}}$  determine each other.

**Summary:** Given  $\mathbf{v}$  in  $V$ , want to calculate  $T(\mathbf{v})$  in  $W$ . Take an input basis  $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$  and an output basis  $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$ .

- The abstract input vector  $\mathbf{v}$  and the coordinate vector  $\mathbf{v}_{\mathcal{A}}$  determine each other.
- The abstract output vector  $T(\mathbf{v})$  and the coordinate vector  $T(\mathbf{v})_{\mathcal{B}}$  determine each other.

**Summary:** Given  $\mathbf{v}$  in  $V$ , want to calculate  $T(\mathbf{v})$  in  $W$ . Take an input basis  $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$  and an output basis  $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$ .

- The abstract input vector  $\mathbf{v}$  and the coordinate vector  $\mathbf{v}_{\mathcal{A}}$  determine each other.
- The abstract output vector  $T(\mathbf{v})$  and the coordinate vector  $T(\mathbf{v})_{\mathcal{B}}$  determine each other.
- So we know  $T$  if we know the matrix  $T_{\mathcal{B}\mathcal{A}}$ :

$$T(\mathbf{v})_{\mathcal{B}} = T_{\mathcal{B}\mathcal{A}}\mathbf{v}_{\mathcal{A}}.$$

**Summary:** Given  $\mathbf{v}$  in  $V$ , want to calculate  $T(\mathbf{v})$  in  $W$ . Take an input basis  $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$  and an output basis  $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$ .

- The abstract input vector  $\mathbf{v}$  and the coordinate vector  $\mathbf{v}_{\mathcal{A}}$  determine each other.
- The abstract output vector  $T(\mathbf{v})$  and the coordinate vector  $T(\mathbf{v})_{\mathcal{B}}$  determine each other.
- So we know  $T$  if we know the matrix  $T_{\mathcal{B}\mathcal{A}}$ :

$$T(\mathbf{v})_{\mathcal{B}} = T_{\mathcal{B}\mathcal{A}}\mathbf{v}_{\mathcal{A}}.$$

The output coordinate vector equals the matrix for  $T$  times the input coordinate vector.

**Formula For the Coordinate matrix.** To write  $T: V \rightarrow W$  as a matrix, take an input basis  $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$  and an output basis  $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$ . Then

$$T_{\mathcal{B}\mathcal{A}} = \begin{bmatrix} T(\mathbf{x}_1)_{\mathcal{B}} & T(\mathbf{x}_2)_{\mathcal{B}} & \dots & T(\mathbf{x}_n)_{\mathcal{B}} \end{bmatrix}$$

**Formula For the Coordinate matrix.** To write  $T: V \rightarrow W$  as a matrix, take an input basis  $\mathcal{A} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$  and an output basis  $\mathcal{B} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_m)$ . Then

$$T_{\mathcal{B}\mathcal{A}} = \begin{bmatrix} T(\mathbf{x}_1)_{\mathcal{B}} & T(\mathbf{x}_2)_{\mathcal{B}} & \dots & T(\mathbf{x}_n)_{\mathcal{B}} \end{bmatrix}$$

### Example

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be reflection across the x-y plane,  
 $(x, y, z) \mapsto (x, y, -z)$ .



### Example

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be reflection across the x-y plane,  
 $(x, y, z) \mapsto (x, y, -z)$ .

Determine the matrix representing  $T$  in the basis

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

### Example

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be reflection across the x-y plane,  
 $(x, y, z) \mapsto (x, y, -z)$ .

Determine the matrix representing  $T$  in the basis

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

$T: (x, y, z) \mapsto (x, y, -z)$ . So calculate

$T: (x, y, z) \mapsto (x, y, -z)$ . So calculate

$$T \left( \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right) =$$

$T: (x, y, z) \mapsto (x, y, -z)$ . So calculate

$$T \left( \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right) = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} =$$

$T: (x, y, z) \mapsto (x, y, -z)$ . So calculate

$$T \left( \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right) = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} - 2 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$T: (x, y, z) \mapsto (x, y, -z)$ . So calculate

$$T \left( \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right) = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} - 2 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$T \left( \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right) = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \quad T \left( \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right) = (-1) \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

Hence

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & -1 \end{bmatrix}.$$

### Example

Let  $T : \mathbb{P}_3 \rightarrow \mathbb{P}_2$  be the linear map given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

What's the matrix  $A$  representing  $T$  in the standard bases?



### Example

Let  $T : \mathbb{P}_3 \rightarrow \mathbb{P}_2$  be the linear map given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

What's the matrix  $A$  representing  $T$  in the standard bases?

### Solution

The standard bases are  $\{1, t, t^2, t^3\}$  for  $\mathbb{P}_3$  and  $\{1, t, t^2\}$  for  $\mathbb{P}_2$ .

### Example

Let  $T : \mathbb{P}_3 \rightarrow \mathbb{P}_2$  be the linear map given by

$$T(p(t)) = \frac{d}{dt}p(t).$$

What's the matrix  $A$  representing  $T$  in the standard bases?

### Solution

The standard bases are  $\{1, t, t^2, t^3\}$  for  $\mathbb{P}_3$  and  $\{1, t, t^2\}$  for  $\mathbb{P}_2$ .  
The matrix  $A$  has 4 columns and 3 rows.

- $T(1) = 0$ , so the first column is  $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ .
- $T(t) = 1$ , so the second column is  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ .
- $T(t^2) = 2t$ , so the third column is  $\begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix}$ .
- $T(t^3) = 3t^2$ , so the last (fourth) column is  $\begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix}$ .

- $T(1) = 0$ , so the first column is  $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ .
- $T(t) = 1$ , so the second column is  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ .
- $T(t^2) = 2t$ , so the third column is  $\begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix}$ .
- $T(t^3) = 3t^2$ , so the last (fourth) column is  $\begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix}$ .

- $T(1) = 0$ , so the first column is  $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ .
- $T(t) = 1$ , so the second column is  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ .
- $T(t^2) = 2t$ , so the third column is  $\begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix}$ .
- $T(t^3) = 3t^2$ , so the last (fourth) column is  $\begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix}$ .

- $T(1) = 0$ , so the first column is  $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ .
- $T(t) = 1$ , so the second column is  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ .
- $T(t^2) = 2t$ , so the third column is  $\begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix}$ .
- $T(t^3) = 3t^2$ , so the last (fourth) column is  $\begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix}$ .

- $T(1) = 0$ , so the first column is  $\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ .
- $T(t) = 1$ , so the second column is  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ .
- $T(t^2) = 2t$ , so the third column is  $\begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix}$ .
- $T(t^3) = 3t^2$ , so the last (fourth) column is  $\begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix}$ .

So the matrix  $A$  representing  $T$  in the standard bases is

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}.$$

What is  $Col(A)$  and  $Nul(A)$  for  $A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}$ ?



What is  $\text{Col}(A)$  and  $\text{Nul}(A)$  for  $A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}$ ?

### Solution

$\text{Col}(A) = \mathbb{R}^3$ . Every quadratic polynomial is the derivative of some cubic polynomial.

What is  $Col(A)$  and  $Nul(A)$  for  $A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}$ ?

### Solution

$Col(A) = \mathbb{R}^3$ . Every quadratic polynomial is the derivative of some cubic polynomial.

$$Nul(A) = span \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right\}.$$

What is  $Col(A)$  and  $Nul(A)$  for  $A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}$ ?

### Solution

$Col(A) = \mathbb{R}^3$ . Every quadratic polynomial is the derivative of some cubic polynomial.

$$Nul(A) = span \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right\}.$$

The corresponding polynomial is  $p(t) = 1$ . That makes sense because differentiation kills constant polynomials.

Let's try differentiating  $7t^3 - t + 3$  using the matrix  $A$ .

Let's try differentiating  $7t^3 - t + 3$  using the matrix  $A$ .

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} 3 \\ -1 \\ 0 \\ 7 \end{bmatrix} =$$

Let's try differentiating  $7t^3 - t + 3$  using the matrix  $A$ .

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} 3 \\ -1 \\ 0 \\ 7 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 21 \end{bmatrix}$$

We get  $-1 + 0t + 21t^2$ , which is indeed the derivative of  $7t^3 - t + 3$ .

## Inner Product and Distances

## Definition

The **inner product** (or **dot product**) of  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$  is



## Definition

The **inner product** (or **dot product**) of  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$  is

$$\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \cdots + v_n w_n.$$

### Example

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix} =$$

### Example

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix} = [1 \quad 2 \quad 3] \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix}$$

### Example

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix} = [1 \quad 2 \quad 3] \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix} = 1 - 2 - 6 = -7$$

## Theorem

*Let  $\mathbf{u}, \mathbf{v}$  and  $\mathbf{w}$  be vectors in  $\mathbb{R}^n$ , and let  $c$  be any scalar. Then*

## Theorem

Let  $\mathbf{u}, \mathbf{v}$  and  $\mathbf{w}$  be vectors in  $\mathbb{R}^n$ , and let  $c$  be any scalar. Then

(a)  $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$

## Theorem

Let  $\mathbf{u}, \mathbf{v}$  and  $\mathbf{w}$  be vectors in  $\mathbb{R}^n$ , and let  $c$  be any scalar. Then

(a)  $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$

(b)  $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = \mathbf{u} \cdot \mathbf{w} + \mathbf{v} \cdot \mathbf{w}$

## Theorem

Let  $\mathbf{u}, \mathbf{v}$  and  $\mathbf{w}$  be vectors in  $\mathbb{R}^n$ , and let  $c$  be any scalar. Then

- (a)  $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
- (b)  $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = \mathbf{u} \cdot \mathbf{w} + \mathbf{v} \cdot \mathbf{w}$
- (c)  $(c\mathbf{u}) \cdot \mathbf{v} = c(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \cdot (c\mathbf{v})$



## Theorem

Let  $\mathbf{u}, \mathbf{v}$  and  $\mathbf{w}$  be vectors in  $\mathbb{R}^n$ , and let  $c$  be any scalar. Then

- (a)  $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
- (b)  $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = \mathbf{u} \cdot \mathbf{w} + \mathbf{v} \cdot \mathbf{w}$
- (c)  $(c\mathbf{u}) \cdot \mathbf{v} = c(\mathbf{u} \cdot \mathbf{v}) = \mathbf{u} \cdot (c\mathbf{v})$
- (d)  $\mathbf{u} \cdot \mathbf{u} \geq 0$ , and  $\mathbf{u} \cdot \mathbf{u} = 0$  if and only if  $\mathbf{u} = \mathbf{0}$ .

## Definition

- The **norm** (or **length**) of a vector  $\mathbf{v} \in \mathbb{R}^n$  is

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + \cdots + v_n^2}.$$

- The **distance** between points  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$  is

$$\text{dist}(\mathbf{v}, \mathbf{w}) = \|\mathbf{v} - \mathbf{w}\|.$$

## Example

(a)

$$\left\| \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix} \right\|$$

## Example

(a)

$$\left\| \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix} \right\| = \sqrt{1^2 + (-1)^2 + 3^2}$$

## Example

(a)

$$\left\| \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix} \right\| = \sqrt{1^2 + (-1)^2 + 3^2} = \sqrt{11}$$

## Example

(a)

$$\left\| \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix} \right\| = \sqrt{1^2 + (-1)^2 + 3^2} = \sqrt{11}$$

(b)

$$\text{dist} \left( \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} \right)$$

## Example

(a)

$$\left\| \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix} \right\| = \sqrt{1^2 + (-1)^2 + 3^2} = \sqrt{11}$$

(b)

$$\text{dist} \left( \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} \right) = \left\| \begin{bmatrix} x_1 - x_2 \\ y_1 - y_2 \end{bmatrix} \right\|$$

## Example

(a)

$$\left\| \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix} \right\| = \sqrt{1^2 + (-1)^2 + 3^2} = \sqrt{11}$$

(b)

$$\text{dist} \left( \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} \right) = \left\| \begin{bmatrix} x_1 - x_2 \\ y_1 - y_2 \end{bmatrix} \right\| = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$



## Inner Product and Angles

We can use the dot product to compute angles too.

We can use the dot product to compute angles too.

### Theorem

*If  $\mathbf{v}$  and  $\mathbf{w}$  are linearly independent, they form an angle  $\theta$ , and*


$$\mathbf{v} \cdot \mathbf{w} = \|\mathbf{v}\| \|\mathbf{w}\| \cos \theta$$

We can use the dot product to compute angles too.

### Theorem

*If  $\mathbf{v}$  and  $\mathbf{w}$  are linearly independent, they form an angle  $\theta$ , and*

$$\mathbf{v} \cdot \mathbf{w} = \|\mathbf{v}\| \|\mathbf{w}\| \cos \theta$$



### Example

What is the angle formed in  $\mathbb{R}^3$  between the vectors

$$\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 0 \\ -1 \\ -1 \end{bmatrix}?$$

(A base jumper runs at a cliff at a  $45^\circ$  angle, then jumps straight away from the cliff and  $45^\circ$  downwards; what angle does he turn as he jumps?)

## Solution

$$\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 0 \\ -1 \\ -1 \end{bmatrix}$$

## Solution

$$\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 0 \\ -1 \\ -1 \end{bmatrix}$$

$$\|\mathbf{v}\| = \sqrt{1^2 + 1^2 + 0^2} = \sqrt{2}$$

## Solution

$$\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 0 \\ -1 \\ -1 \end{bmatrix}$$

$$\|\mathbf{v}\| = \sqrt{1^2 + 1^2 + 0^2} = \sqrt{2}$$

$$\|\mathbf{w}\| = \sqrt{0^2 + (-1)^2 + (-1)^2} = \sqrt{2}$$



## Solution

$$\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 0 \\ -1 \\ -1 \end{bmatrix}$$

$$\|\mathbf{v}\| = \sqrt{1^2 + 1^2 + 0^2} = \sqrt{2}$$

$$\|\mathbf{w}\| = \sqrt{0^2 + (-1)^2 + (-1)^2} = \sqrt{2}$$

$$\mathbf{v} \cdot \mathbf{w} = -1$$

## Solution

$$\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 0 \\ -1 \\ -1 \end{bmatrix}$$

$$\|\mathbf{v}\| = \sqrt{1^2 + 1^2 + 0^2} = \sqrt{2}$$

$$\|\mathbf{w}\| = \sqrt{0^2 + (-1)^2 + (-1)^2} = \sqrt{2}$$

$$\mathbf{v} \cdot \mathbf{w} = -1$$

$$\mathbf{v} \cdot \mathbf{w} = \|\mathbf{v}\| \|\mathbf{w}\| \cos \theta$$

## Solution

$$\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 0 \\ -1 \\ -1 \end{bmatrix}$$

$$\|\mathbf{v}\| = \sqrt{1^2 + 1^2 + 0^2} = \sqrt{2}$$

$$\|\mathbf{w}\| = \sqrt{0^2 + (-1)^2 + (-1)^2} = \sqrt{2}$$

$$\mathbf{v} \cdot \mathbf{w} = -1$$

$$\mathbf{v} \cdot \mathbf{w} = \|\mathbf{v}\| \|\mathbf{w}\| \cos \theta$$

$$-1 = 2 \cos \theta \quad \Rightarrow \quad \cos \theta = -\frac{1}{2}$$

## Solution

$$\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{w} = \begin{bmatrix} 0 \\ -1 \\ -1 \end{bmatrix}$$

$$\|\mathbf{v}\| = \sqrt{1^2 + 1^2 + 0^2} = \sqrt{2}$$

$$\|\mathbf{w}\| = \sqrt{0^2 + (-1)^2 + (-1)^2} = \sqrt{2}$$

$$\mathbf{v} \cdot \mathbf{w} = -1$$

$$\mathbf{v} \cdot \mathbf{w} = \|\mathbf{v}\| \|\mathbf{w}\| \cos \theta$$

$$-1 = 2 \cos \theta \quad \Rightarrow \quad \cos \theta = -\frac{1}{2}$$

## Orthogonal vectors

## Definition

$\mathbf{v}$  and  $\mathbf{w}$  in  $\mathbb{R}^n$  are **orthogonal** if

$$\mathbf{v} \cdot \mathbf{w} = 0.$$

### Definition

$\mathbf{v}$  and  $\mathbf{w}$  in  $\mathbb{R}^n$  are **orthogonal** if

$$\mathbf{v} \cdot \mathbf{w} = 0.$$

### Remark


We write  $\mathbf{v} \perp \mathbf{w}$  when  $\mathbf{v}$  and  $\mathbf{w}$  are orthogonal. Nonzero vectors  $\mathbf{v}$ ,  $\mathbf{w}$  are orthogonal if and only if they are perpendicular.






Nonzero vectors  $\mathbf{v}$ ,  $\mathbf{w}$  are orthogonal if and only if they are perpendicular.

Nonzero vectors  $\mathbf{v}$ ,  $\mathbf{w}$  are orthogonal if and only if they are perpendicular. We can derive this from Pythagoras' theorem.




Nonzero vectors  $\mathbf{v}$ ,  $\mathbf{w}$  are orthogonal if and only if they are perpendicular. We can derive this from Pythagoras' theorem.



$$\mathbf{v} \perp \mathbf{w}$$


Nonzero vectors  $\mathbf{v}$ ,  $\mathbf{w}$  are orthogonal if and only if they are perpendicular. We can derive this from Pythagoras' theorem.



$$\mathbf{v} \perp \mathbf{w}$$

$$\iff \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 = \|\mathbf{v} - \mathbf{w}\|^2$$

Nonzero vectors  $\mathbf{v}$ ,  $\mathbf{w}$  are orthogonal if and only if they are perpendicular. We can derive this from Pythagoras' theorem.




$$\mathbf{v} \perp \mathbf{w}$$

$$\iff \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 = \|\mathbf{v} - \mathbf{w}\|^2$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = (\mathbf{v} - \mathbf{w}) \cdot (\mathbf{v} - \mathbf{w})$$

Nonzero vectors  $\mathbf{v}$ ,  $\mathbf{w}$  are orthogonal if and only if they are perpendicular. We can derive this from Pythagoras' theorem.




$$\mathbf{v} \perp \mathbf{w}$$

$$\iff \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 = \|\mathbf{v} - \mathbf{w}\|^2$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = (\mathbf{v} - \mathbf{w}) \cdot (\mathbf{v} - \mathbf{w})$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = \mathbf{v} \cdot \mathbf{v} - 2\mathbf{v} \cdot \mathbf{w} + \mathbf{w} \cdot \mathbf{w}$$

Nonzero vectors  $\mathbf{v}$ ,  $\mathbf{w}$  are orthogonal if and only if they are perpendicular. We can derive this from Pythagoras' theorem.



$$\mathbf{v} \perp \mathbf{w}$$

$$\iff \|\mathbf{v}\|^2 + \|\mathbf{w}\|^2 = \|\mathbf{v} - \mathbf{w}\|^2$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = (\mathbf{v} - \mathbf{w}) \cdot (\mathbf{v} - \mathbf{w})$$

$$\iff \mathbf{v} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{w} = \mathbf{v} \cdot \mathbf{v} - 2\mathbf{v} \cdot \mathbf{w} + \mathbf{w} \cdot \mathbf{w}$$

$$\iff \mathbf{v} \cdot \mathbf{w} = 0$$

## Example

Are the following vectors orthogonal?

(a)  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \end{bmatrix}$



## Example

Are the following vectors orthogonal?

(a)  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \end{bmatrix}$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} -2 \\ 1 \end{bmatrix} = 1 \cdot (-2) + 2 \cdot 1 = 0$$

## Example

Are the following vectors orthogonal?

(a)  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \end{bmatrix}$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} -2 \\ 1 \end{bmatrix} = 1 \cdot (-2) + 2 \cdot 1 = 0$$

So yes, they're orthogonal.

## Example

Are the following vectors orthogonal?

(a)  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \end{bmatrix}$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} -2 \\ 1 \end{bmatrix} = 1 \cdot (-2) + 2 \cdot 1 = 0$$

So yes, they're orthogonal.

(b)  $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix}$

## Example

Are the following vectors orthogonal?

(a)  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \end{bmatrix}$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} -2 \\ 1 \end{bmatrix} = 1 \cdot (-2) + 2 \cdot 1 = 0$$

So yes, they're orthogonal.

(b)  $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix}$

$$\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix} = 1 \cdot (-2) + 2 \cdot 1 + 1 \cdot 1 = 1$$

## Example

Are the following vectors orthogonal?

(a)  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \end{bmatrix}$

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} -2 \\ 1 \end{bmatrix} = 1 \cdot (-2) + 2 \cdot 1 = 0$$

So yes, they're orthogonal.

(b)  $\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix}$

$$\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix} = 1 \cdot (-2) + 2 \cdot 1 + 1 \cdot 1 = 1$$

So no, they're not orthogonal.

### Example

Let  $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ .

### Example

Let  $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ .

Is the set of vectors orthogonal to  $\mathbf{v}$  a subspace of  $\mathbb{R}^3$ ?

### Example

Let  $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ .

Is the set of vectors orthogonal to  $\mathbf{v}$  a subspace of  $\mathbb{R}^3$ ?

$$V = \{\mathbf{x} \in \mathbb{R}^3 : \mathbf{v} \cdot \mathbf{x} = 0\}$$



### Example

Let  $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ .

Is the set of vectors orthogonal to  $\mathbf{v}$  a subspace of  $\mathbb{R}^3$ ?

$$V = \{\mathbf{x} \in \mathbb{R}^3 : \mathbf{v} \cdot \mathbf{x} = 0\}$$

### Solution

### Example

Let  $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ .

Is the set of vectors orthogonal to  $\mathbf{v}$  a subspace of  $\mathbb{R}^3$ ?

$$V = \{\mathbf{x} \in \mathbb{R}^3 : \mathbf{v} \cdot \mathbf{x} = 0\}$$

### Solution

$$\mathbf{v} \cdot \mathbf{x} = 0$$

### Example

Let  $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ .

Is the set of vectors orthogonal to  $\mathbf{v}$  a subspace of  $\mathbb{R}^3$ ?

$$V = \{\mathbf{x} \in \mathbb{R}^3 : \mathbf{v} \cdot \mathbf{x} = 0\}$$

### Solution

$$\mathbf{v} \cdot \mathbf{x} = 0$$

$$\Leftrightarrow \begin{bmatrix} 2 & -1 & 1 \end{bmatrix} \mathbf{x} = 0$$

### Example

Let  $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ .

Is the set of vectors orthogonal to  $\mathbf{v}$  a subspace of  $\mathbb{R}^3$ ?

$$V = \{\mathbf{x} \in \mathbb{R}^3 : \mathbf{v} \cdot \mathbf{x} = 0\}$$

### Solution

$$\mathbf{v} \cdot \mathbf{x} = 0$$

$$\Leftrightarrow \begin{bmatrix} 2 & -1 & 1 \end{bmatrix} \mathbf{x} = 0$$

$V$  is just the null space of the matrix  $\mathbf{v}^T = \begin{bmatrix} 2 & -1 & 1 \end{bmatrix}$ .

### Example

Let  $\mathbf{v} = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$ .

Is the set of vectors orthogonal to  $\mathbf{v}$  a subspace of  $\mathbb{R}^3$ ?

$$V = \{\mathbf{x} \in \mathbb{R}^3 : \mathbf{v} \cdot \mathbf{x} = 0\}$$

### Solution

$$\mathbf{v} \cdot \mathbf{x} = 0$$

$$\Leftrightarrow \begin{bmatrix} 2 & -1 & 1 \end{bmatrix} \mathbf{x} = 0$$

$V$  is just the null space of the matrix  $\mathbf{v}^T = \begin{bmatrix} 2 & -1 & 1 \end{bmatrix}$ . So yes, it is a subspace.

## Definition

If  $V$  is a subspace of  $\mathbb{R}^n$ ,

## Definition

If  $V$  is a subspace of  $\mathbb{R}^n$ , a vector  $\mathbf{x}$  is **orthogonal to**  $V$  if

## Definition

If  $V$  is a subspace of  $\mathbb{R}^n$ , a vector  $\mathbf{x}$  is **orthogonal to**  $V$  if it is orthogonal to every vector in  $V$ .



## Definition

If  $V$  is a subspace of  $\mathbb{R}^n$ , a vector  $\mathbf{x}$  is **orthogonal to**  $V$  if it is orthogonal to every vector in  $V$ .

## Example

Let  $V = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ . Is  $\mathbf{x} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  orthogonal to  $V$ ?

## Definition

If  $V$  is a subspace of  $\mathbb{R}^n$ , a vector  $\mathbf{x}$  is **orthogonal to**  $V$  if it is orthogonal to every vector in  $V$ .

## Example

Let  $V = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ . Is  $\mathbf{x} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  orthogonal to  $V$ ?

## Solution

$$\begin{bmatrix} -1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} a \\ a \end{bmatrix} =$$

## Definition

If  $V$  is a subspace of  $\mathbb{R}^n$ , a vector  $\mathbf{x}$  is **orthogonal to**  $V$  if it is orthogonal to every vector in  $V$ .

## Example

Let  $V = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ . Is  $\mathbf{x} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  orthogonal to  $V$ ?

## Solution

$$\begin{bmatrix} -1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} a \\ a \end{bmatrix} = -a + a =$$

## Definition

If  $V$  is a subspace of  $\mathbb{R}^n$ , a vector  $\mathbf{x}$  is **orthogonal to**  $V$  if it is orthogonal to every vector in  $V$ .

## Example

Let  $V = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ . Is  $\mathbf{x} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  orthogonal to  $V$ ?

## Solution

$$\begin{bmatrix} -1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} a \\ a \end{bmatrix} = -a + a = 0$$

## Definition

If  $V$  is a subspace of  $\mathbb{R}^n$ , a vector  $\mathbf{x}$  is **orthogonal to**  $V$  if it is orthogonal to every vector in  $V$ .

## Example

Let  $V = \text{span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ . Is  $\mathbf{x} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  orthogonal to  $V$ ?

## Solution

$$\begin{bmatrix} -1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} a \\ a \end{bmatrix} = -a + a = 0$$

So yes.