

# Math 415 - Lecture 30

## Eigenvectors and Eigenvalues

Friday November 6th 2015

Textbook reading: Chapter 5.1

Textbook reading: Chapter 5.1

Suggested practice exercises: 12, 20, 21, 22, 36

Textbook reading: Chapter 5.1

Suggested practice exercises: 12, 20, 21, 22, 36

Khan Academy video: Introduction to Eigenvalues and  
Eigenvectors, Proof of formula for determining  
Eigenvalues, Finding Eigenvectors and Eigenspaces  
example

Textbook reading: Chapter 5.1

Suggested practice exercises: 12, 20, 21, 22, 36

Khan Academy video: Introduction to Eigenvalues and  
Eigenvectors, Proof of formula for determining  
Eigenvalues, Finding Eigenvectors and Eigenspaces  
example

Strang lecture: Lecture 21: Eigenvalues and eigenvectors

## Review

## Definition

The **determinant** is characterized by:

## Definition

The **determinant** is characterized by:



## Definition

The **determinant** is characterized by:

- the normalization  $\det I_{n \times n} = 1$ ,

## Definition

The **determinant** is characterized by:

- the normalization  $\det I_{n \times n} = 1$ ,
- and how it is affected by elementary row operations:

## Definition

The **determinant** is characterized by:

- the normalization  $\det I_{n \times n} = 1$ ,
- and how it is affected by elementary row operations:
  - **(Replacement)** Add a multiple of one row to another row.  
*Does not change* the determinant.

## Definition

The **determinant** is characterized by:

- the normalization  $\det I_{n \times n} = 1$ ,
- and how it is affected by elementary row operations:
  - **(Replacement)** Add a multiple of one row to another row.  
*Does not change* the determinant.
  - **(Interchange)** Interchange two rows.  
*Reverses the sign* of the determinant.

## Definition

The **determinant** is characterized by:

- the normalization  $\det I_{n \times n} = 1$ ,
- and how it is affected by elementary row operations:
  - **(Replacement)** Add a multiple of one row to another row.  
*Does not change* the determinant.
  - **(Interchange)** Interchange two rows.  
*Reverses the sign* of the determinant.
  - **(Scaling)** Multiply all entries in a row by  $s$ .  
*Multiplies* the determinant by  $s$ .

## Definition

The **determinant** is characterized by:

- the normalization  $\det I_{n \times n} = 1$ ,
- and how it is affected by elementary row operations:
  - **(Replacement)** Add a multiple of one row to another row.  
*Does not change* the determinant.
  - **(Interchange)** Interchange two rows.  
*Reverses the sign* of the determinant.
  - **(Scaling)** Multiply all entries in a row by  $s$ .  
*Multiplies* the determinant by  $s$ .
- For triangular  $A$  the determinant is just product of the diagonal entries.

## Definition

The **determinant** is characterized by:

- the normalization  $\det I_{n \times n} = 1$ ,
- and how it is affected by elementary row operations:
  - **(Replacement)** Add a multiple of one row to another row.  
*Does not change* the determinant.
  - **(Interchange)** Interchange two rows.  
*Reverses the sign* of the determinant.
  - **(Scaling)** Multiply all entries in a row by  $s$ .  
*Multiplies* the determinant by  $s$ .
- For triangular  $A$  the determinant is just product of the diagonal entries.

## Definition

The **determinant** is characterized by:

- the normalization  $\det I_{n \times n} = 1$ ,
- and how it is affected by elementary row operations:
  - **(Replacement)** Add a multiple of one row to another row.  
*Does not change* the determinant.
  - **(Interchange)** Interchange two rows.  
*Reverses the sign* of the determinant.
  - **(Scaling)** Multiply all entries in a row by  $s$ .  
*Multiplies* the determinant by  $s$ .
- For triangular  $A$  the determinant is just product of the diagonal entries.

This allows us to compute the determinant using just **row operations!**.



## Definition

The **determinant** is characterized by:

- the normalization  $\det I_{n \times n} = 1$ ,
- and how it is affected by elementary row operations:
  - **(Replacement)** Add a multiple of one row to another row.  
*Does not change* the determinant.
  - **(Interchange)** Interchange two rows.  
*Reverses the sign* of the determinant.
  - **(Scaling)** Multiply all entries in a row by  $s$ .  
*Multiplies* the determinant by  $s$ .
- For triangular  $A$  the determinant is just product of the diagonal entries.

This allows us to compute the determinant using just **row operations!**. Bring  $A$  into echelon form = triangular form, keeping track how the determinant changes under the row operations you are using.

- What's **wrong**?!

$$\det(A^{-1}) = \det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc} (da - (-b)(-c)) = 1$$

- What's **wrong**?!

$$\det(A^{-1}) = \det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc} (da - (-b)(-c)) = 1$$

- What's **wrong**?!

$$\det(A^{-1}) = \det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc} (da - (-b)(-c)) = 1$$

The correct calculation is:

$$\det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} =$$

- What's **wrong**?!

$$\det(A^{-1}) = \det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc} (da - (-b)(-c)) = 1$$

The correct calculation is:

$$\det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{(ad - bc)^2} \det \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc}$$

- What's **wrong**?!

$$\det(A^{-1}) = \det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc} (da - (-b)(-c)) = 1$$

The correct calculation is:

$$\det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{(ad - bc)^2} \det \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc}$$

### Example

Suppose  $A$  is a  $3 \times 3$  matrix with  $\det(A) = 5$ . What is  $\det(2A)$ ?

- What's **wrong**?!

$$\det(A^{-1}) = \det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc} (da - (-b)(-c)) = 1$$

The correct calculation is:

$$\det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{(ad - bc)^2} \det \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc}$$

### Example

Suppose  $A$  is a  $3 \times 3$  matrix with  $\det(A) = 5$ . What is  $\det(2A)$ ?

- What's **wrong**?!

$$\det(A^{-1}) = \det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc} (da - (-b)(-c)) = 1$$

The correct calculation is:

$$\det \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{(ad - bc)^2} \det \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{ad - bc}$$

### Example

Suppose  $A$  is a  $3 \times 3$  matrix with  $\det(A) = 5$ . What is  $\det(2A)$ ?

### Solution

*A has three rows. Multiplying all 3 of them produces  $2A$ . Hence,  $\det(2A) = 2^3 \det(A) = 40$ .*



## Eigenvectors and eigenvalues

## Definition

Throughout,  $A$  will be an  $n \times n$  matrix.

Throughout,  $A$  will be an  $n \times n$  matrix.

### Definition

An **eigenvector** of  $A$  is a nonzero  $\mathbf{x}$  such that

$$A\mathbf{x} = \lambda\mathbf{x} \text{ for some scalar } \lambda.$$

The scalar  $\lambda$  is the corresponding **eigenvalue**.

Throughout,  $A$  will be an  $n \times n$  matrix.

### Definition

An **eigenvector** of  $A$  is a nonzero  $\mathbf{x}$  such that

$$A\mathbf{x} = \lambda\mathbf{x} \text{ for some scalar } \lambda.$$

The scalar  $\lambda$  is the corresponding **eigenvalue**.

In words, eigenvectors are those  $\mathbf{x}$ , for which  $A\mathbf{x}$  is parallel to  $\mathbf{x}$ .

Throughout,  $A$  will be an  $n \times n$  matrix.

### Definition

An **eigenvector** of  $A$  is a nonzero  $\mathbf{x}$  such that

$$A\mathbf{x} = \lambda\mathbf{x} \text{ for some scalar } \lambda.$$

The scalar  $\lambda$  is the corresponding **eigenvalue**.

In words, eigenvectors are those  $\mathbf{x}$ , for which  $A\mathbf{x}$  is parallel to  $\mathbf{x}$ .

### Example

Verify that  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ . Is  $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$  an eigenvector?

### Example

Verify that  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ . Is  $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$  an eigenvector?

### Solution

$$A\mathbf{x} = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \end{bmatrix} = -2\mathbf{x}$$




### Example

Verify that  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ . Is  $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$  an eigenvector?

### Solution

$$A\mathbf{x} = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \end{bmatrix} = -2\mathbf{x}$$

Hence,  $\mathbf{x}$  is an eigenvector of  $A$  with eigenvalue  $-2$ .



## Geometric interpretation

## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

## Solution

$$A \begin{bmatrix} x \\ y \end{bmatrix} =$$

## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

## Solution

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ x \end{bmatrix},$$

## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ x \end{bmatrix}$ , i.e. multiplication with  $A$  is reflection through the line  $y = x$ .

## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ x \end{bmatrix}$ , i.e. multiplication with  $A$  is reflection through the line  $y = x$ .

- $A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . So  $\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = 1$ .




## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ x \end{bmatrix}$ , i.e. multiplication with  $A$  is reflection through the line  $y = x$ .

- $A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . So  $\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = 1$ .




## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .

## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ x \end{bmatrix}$ , i.e. multiplication with  $A$  is reflection through the line  $y = x$ .

- $A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . So  $\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = 1$ .
- $A \begin{bmatrix} -1 \\ 1 \end{bmatrix} = -1 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$



## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ .


## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ x \end{bmatrix}$ , i.e. multiplication with  $A$  is reflection through the line  $y = x$ .

- $A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . So  $\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = 1$ .

- $A \begin{bmatrix} -1 \\ 1 \end{bmatrix} = -1 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

So  $\mathbf{x} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = -1$ .



## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ .

## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ .

## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ 0 \end{bmatrix}$ , i.e. multiplication with  $A$  is projection on the  $x$ -axis.

- $A \begin{bmatrix} 1 \\ 0 \end{bmatrix} =$

## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ .

## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ 0 \end{bmatrix}$ , i.e. multiplication with  $A$  is projection on the  $x$ -axis.

- $A \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ .

## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ .

## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ 0 \end{bmatrix}$ , i.e. multiplication with  $A$  is projection on the  $x$ -axis.

- $A \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ . So  $\mathbf{x} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = 1$ .

## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ .

## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ 0 \end{bmatrix}$ , i.e. multiplication with  $A$  is projection on the  $x$ -axis.

- $A \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ . So  $\mathbf{x} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = 1$ .
- $A \begin{bmatrix} 0 \\ 1 \end{bmatrix} =$



## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ .

## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ 0 \end{bmatrix}$ , i.e. multiplication with  $A$  is projection on the  $x$ -axis.

- $A \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ . So  $\mathbf{x} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = 1$ .
- $A \begin{bmatrix} 0 \\ 1 \end{bmatrix} = 0 \begin{bmatrix} 0 \\ 1 \end{bmatrix}$

## Example

Use your geometric understanding to find the eigenvectors and the eigenvalues of  $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ .

## Solution

$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ 0 \end{bmatrix}$ , i.e. multiplication with  $A$  is projection on the  $x$ -axis.

- $A \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 1 \cdot \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ . So  $\mathbf{x} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = 1$ .

- $A \begin{bmatrix} 0 \\ 1 \end{bmatrix} = 0 \begin{bmatrix} 0 \\ 1 \end{bmatrix}$

So  $\mathbf{x} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda = 0$ .

# Summary

# Summary

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero.

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero.



## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ .

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero.

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero.

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero. See the projection example.

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero. See the projection example.

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero. See the projection example.

## Problems



## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero. See the projection example.

## Problems

- \* How to find possible eigenvalues for  $A$ ?

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero. See the projection example.

## Problems

- \* How to find possible eigenvalues for  $A$ ?

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero. See the projection example.

## Problems

- \* How to find possible eigenvalues for  $A$ ? This uses determinants.

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero. See the projection example.

## Problems

- \* How to find possible eigenvalues for  $A$ ? This uses determinants.
- \* How to find eigenvectors?

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero. See the projection example.

## Problems

- \* How to find possible eigenvalues for  $A$ ? This uses determinants.
- \* How to find eigenvectors?

## Summary

- \* Eigenvectors  $\mathbf{x}$  get stretched by eigenvalue  $\lambda$  under multiplication by  $A$ :

$$A\mathbf{x} = \lambda\mathbf{x}.$$

- \* Eigenvectors  $\mathbf{x}$  **CANNOT** be zero. Why?  $A\mathbf{0} = \lambda\mathbf{0}$  for any  $\lambda$ . Not useful!
- \* Eigenvalues  $\lambda$  **CAN** be zero. See the projection example.

## Problems

- \* How to find possible eigenvalues for  $A$ ? This uses determinants.
- \* How to find eigenvectors? This uses null spaces.

## Eigenspaces

## Definition

The **eigenspace** of  $A$  corresponding to  $\lambda$  is the set of all  $\mathbf{x}$  satisfying  $A\mathbf{x} = \lambda\mathbf{x}$ .



## Definition

The **eigenspace** of  $A$  corresponding to  $\lambda$  is the set of all  $\mathbf{x}$  satisfying  $A\mathbf{x} = \lambda\mathbf{x}$ .

## Definition

The **eigenspace** of  $A$  corresponding to  $\lambda$  is the set of all  $\mathbf{x}$  satisfying  $A\mathbf{x} = \lambda\mathbf{x}$ . It consists of all the eigenvectors of  $A$  with eigenvalue  $\lambda$ , and also the zero vector.

## Definition

The **eigenspace** of  $A$  corresponding to  $\lambda$  is the set of all  $\mathbf{x}$  satisfying  $A\mathbf{x} = \lambda\mathbf{x}$ . It consists of all the eigenvectors of  $A$  with eigenvalue  $\lambda$ , and also the zero vector.

## Example

We saw the projection matrix  $P$  of the projection onto a subspace  $V$  has two eigenvalues  $\lambda = 0, 1$ .

## Definition

The **eigenspace** of  $A$  corresponding to  $\lambda$  is the set of all  $\mathbf{x}$  satisfying  $A\mathbf{x} = \lambda\mathbf{x}$ . It consists of all the eigenvectors of  $A$  with eigenvalue  $\lambda$ , and also the zero vector.

## Example

We saw the projection matrix  $P$  of the projection onto a subspace  $V$  has two eigenvalues  $\lambda = 0, 1$ .

- The eigenspace of  $\lambda = 1$  is

## Definition

The **eigenspace** of  $A$  corresponding to  $\lambda$  is the set of all  $\mathbf{x}$  satisfying  $A\mathbf{x} = \lambda\mathbf{x}$ . It consists of all the eigenvectors of  $A$  with eigenvalue  $\lambda$ , and also the zero vector.

## Example

We saw the projection matrix  $P$  of the projection onto a subspace  $V$  has two eigenvalues  $\lambda = 0, 1$ .

- The eigenspace of  $\lambda = 1$  is

## Definition

The **eigenspace** of  $A$  corresponding to  $\lambda$  is the set of all  $\mathbf{x}$  satisfying  $A\mathbf{x} = \lambda\mathbf{x}$ . It consists of all the eigenvectors of  $A$  with eigenvalue  $\lambda$ , and also the zero vector.

## Example

We saw the projection matrix  $P$  of the projection onto a subspace  $V$  has two eigenvalues  $\lambda = 0, 1$ .

- The eigenspace of  $\lambda = 1$  is  $V$ .
- The eigenspace of  $\lambda = 0$  is

## Definition

The **eigenspace** of  $A$  corresponding to  $\lambda$  is the set of all  $\mathbf{x}$  satisfying  $A\mathbf{x} = \lambda\mathbf{x}$ . It consists of all the eigenvectors of  $A$  with eigenvalue  $\lambda$ , and also the zero vector.

## Example

We saw the projection matrix  $P$  of the projection onto a subspace  $V$  has two eigenvalues  $\lambda = 0, 1$ .

- The eigenspace of  $\lambda = 1$  is  $V$ .
- The eigenspace of  $\lambda = 0$  is

## Definition

The **eigenspace** of  $A$  corresponding to  $\lambda$  is the set of all  $\mathbf{x}$  satisfying  $A\mathbf{x} = \lambda\mathbf{x}$ . It consists of all the eigenvectors of  $A$  with eigenvalue  $\lambda$ , and also the zero vector.

## Example

We saw the projection matrix  $P$  of the projection onto a subspace  $V$  has two eigenvalues  $\lambda = 0, 1$ .

- The eigenspace of  $\lambda = 1$  is  $V$ .
- The eigenspace of  $\lambda = 0$  is  $V^\perp$ .



How to solve  $A\mathbf{x} = \lambda\mathbf{x}$

Key observation:

Key observation:  $\mathbf{x} \neq 0$  is an eigenvector means:

Key observation:  $\mathbf{x} \neq \mathbf{0}$  is an eigenvector means:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$\iff A\mathbf{x} - \lambda\mathbf{x} = \mathbf{0}$$

Key observation:  $\mathbf{x} \neq \mathbf{0}$  is an eigenvector means:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$\iff A\mathbf{x} - \lambda\mathbf{x} = \mathbf{0}$$

$$\iff (A - \lambda I)\mathbf{x} = \mathbf{0}$$

Key observation:  $\mathbf{x} \neq \mathbf{0}$  is an eigenvector means:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$\iff A\mathbf{x} - \lambda\mathbf{x} = \mathbf{0}$$

$$\iff (A - \lambda I)\mathbf{x} = \mathbf{0}$$

Key observation:  $\mathbf{x} \neq \mathbf{0}$  is an eigenvector means:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$\iff A\mathbf{x} - \lambda\mathbf{x} = \mathbf{0}$$

$$\iff (A - \lambda I)\mathbf{x} = \mathbf{0}$$

This  $\mathbf{x}$  is a non trivial solution!

Key observation:  $\mathbf{x} \neq \mathbf{0}$  is an eigenvector means:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$\iff A\mathbf{x} - \lambda\mathbf{x} = \mathbf{0}$$

$$\iff (A - \lambda I)\mathbf{x} = \mathbf{0}$$

This  $\mathbf{x}$  is a non trivial solution! This can happen  $\iff$  the square matrix  $A - \lambda I$  is not invertible



Key observation:  $\mathbf{x} \neq \mathbf{0}$  is an eigenvector means:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$\iff A\mathbf{x} - \lambda\mathbf{x} = \mathbf{0}$$

$$\iff (A - \lambda I)\mathbf{x} = \mathbf{0}$$

This  $\mathbf{x}$  is a non trivial solution! This can happen  $\iff$  the square matrix  $A - \lambda I$  is not invertible  $\iff \det(A - \lambda I) = 0$

Key observation:  $\mathbf{x} \neq \mathbf{0}$  is an eigenvector means:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$\iff A\mathbf{x} - \lambda\mathbf{x} = \mathbf{0}$$

$$\iff (A - \lambda I)\mathbf{x} = \mathbf{0}$$

This  $\mathbf{x}$  is a non trivial solution! This can happen  $\iff$  the square matrix  $A - \lambda I$  is not invertible  $\iff \det(A - \lambda I) = 0$

## Recipe

To find the eigenvectors and eigenvalues of  $A$ :

Key observation:  $\mathbf{x} \neq \mathbf{0}$  is an eigenvector means:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$\iff A\mathbf{x} - \lambda\mathbf{x} = \mathbf{0}$$

$$\iff (A - \lambda I)\mathbf{x} = \mathbf{0}$$

This  $\mathbf{x}$  is a non trivial solution! This can happen  $\iff$  the square matrix  $A - \lambda I$  is not invertible  $\iff \det(A - \lambda I) = 0$

## Recipe

To find the eigenvectors and eigenvalues of  $A$ :

- First, find the eigenvalues using

$$\lambda \text{ is an eigenvalue} \iff \det(A - \lambda I) = 0$$

Key observation:  $\mathbf{x} \neq \mathbf{0}$  is an eigenvector means:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$\iff A\mathbf{x} - \lambda\mathbf{x} = \mathbf{0}$$

$$\iff (A - \lambda I)\mathbf{x} = \mathbf{0}$$

This  $\mathbf{x}$  is a non trivial solution! This can happen  $\iff$  the square matrix  $A - \lambda I$  is not invertible  $\iff \det(A - \lambda I) = 0$

## Recipe

To find the eigenvectors and eigenvalues of  $A$ :

- First, find the eigenvalues using  
 $\lambda$  is an eigenvalue  $\iff \det(A - \lambda I) = 0$
- Then, for each eigenvalue  $\lambda$ , find the corresponding eigenvectors by solving  $(A - \lambda I)\mathbf{x} = \mathbf{0}$ .

Key observation:  $\mathbf{x} \neq \mathbf{0}$  is an eigenvector means:

$$A\mathbf{x} = \lambda\mathbf{x}$$

$$\iff A\mathbf{x} - \lambda\mathbf{x} = \mathbf{0}$$

$$\iff (A - \lambda I)\mathbf{x} = \mathbf{0}$$

This  $\mathbf{x}$  is a non trivial solution! This can happen  $\iff$  the square matrix  $A - \lambda I$  is not invertible  $\iff \det(A - \lambda I) = 0$

## Recipe

To find the eigenvectors and eigenvalues of  $A$ :

- First, find the eigenvalues using  
 $\lambda$  is an eigenvalue  $\iff \det(A - \lambda I) = 0$
- Then, for each eigenvalue  $\lambda$ , find the corresponding eigenvectors by solving  $(A - \lambda I)\mathbf{x} = \mathbf{0}$ . So you need to find the null space  $\text{Nul}(A - \lambda I)$ .

## The characteristic polynomial

## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

## Solution

- $A - \lambda I = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{bmatrix}$



## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

## Solution

- $A - \lambda I = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{bmatrix}$
- $\det(A - \lambda I) = \begin{vmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{vmatrix} = (3 - \lambda)^2 - 1$

## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

## Solution

- $A - \lambda I = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{bmatrix}$
- $\det(A - \lambda I) = \begin{vmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{vmatrix} = (3 - \lambda)^2 - 1$ 
$$= \lambda^2 - 6\lambda + 8 = 0$$

## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

## Solution

- $A - \lambda I = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{bmatrix}$
- $\det(A - \lambda I) = \begin{vmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{vmatrix} = (3 - \lambda)^2 - 1$   

$$= \lambda^2 - 6\lambda + 8 = 0 \implies \lambda_1 = 2, \lambda_2 = 4$$

## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

## Solution

- $A - \lambda I = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{bmatrix}$
- $\det(A - \lambda I) = \begin{vmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{vmatrix} = (3 - \lambda)^2 - 1$   

$$= \lambda^2 - 6\lambda + 8 = 0 \implies \lambda_1 = 2, \lambda_2 = 4$$

This is the **characteristic polynomial** of  $A$ .

## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$$

## Solution

- $A - \lambda I = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{bmatrix}$
- $\det(A - \lambda I) = \begin{vmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{vmatrix} = (3 - \lambda)^2 - 1$ 
$$= \lambda^2 - 6\lambda + 8 = 0 \implies \lambda_1 = 2, \lambda_2 = 4$$

This is the **characteristic polynomial** of  $A$ . Its roots are the eigenvalues of  $A$ .

## Solution (continued)

- Next, find the eigenvectors with eigenvalue  $\lambda_1 = 2$ :

## Solution (continued)

- Next, find the eigenvectors with eigenvalue  $\lambda_1 = 2$ :

$$A - \lambda_1 I =$$

## Solution (continued)

- Next, find the eigenvectors with eigenvalue  $\lambda_1 = 2$ :

$$A - \lambda_1 I = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad \left( A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \right)$$



## Solution (continued)

- Next, find the eigenvectors with eigenvalue  $\lambda_1 = 2$ :

$$A - \lambda_1 I = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad \left( A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \right)$$

Solutions to  $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \mathbf{0}$  have basis  $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ .

## Solution (continued)

- Next, find the eigenvectors with eigenvalue  $\lambda_1 = 2$ :

$$A - \lambda_1 I = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad \left( A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \right)$$

Solutions to  $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \mathbf{0}$  have basis  $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ .

So:  $\mathbf{x}_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda_1 = 2$ .

## Solution (continued)

- Next, find the eigenvectors with eigenvalue  $\lambda_1 = 2$ :

$$A - \lambda_1 I = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad \left( A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \right)$$

Solutions to  $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \mathbf{0}$  have basis  $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ .

So:  $\mathbf{x}_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda_1 = 2$ .

All other eigenvectors with eigenvalue  $\lambda = 2$  are multiples of  $\mathbf{x}_1$ .

## Solution (continued)

- Next, find the eigenvectors with eigenvalue  $\lambda_1 = 2$ :

$$A - \lambda_1 I = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad \left( A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \right)$$

Solutions to  $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \mathbf{x} = \mathbf{0}$  have basis  $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ .

So:  $\mathbf{x}_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda_1 = 2$ .

All other eigenvectors with eigenvalue  $\lambda = 2$  are multiples of  $\mathbf{x}_1$ .

$\text{Span} \left\{ \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$  is the **eigenspace** for the eigenvalue  $\lambda = 2$ .

## Solution (continued)

- Find the eigenvectors with eigenvalue  $\lambda_2 = 4$ :

## Solution (continued)

- Find the eigenvectors with eigenvalue  $\lambda_2 = 4$ :

$$A - \lambda_2 I =$$

## Solution (continued)

- Find the eigenvectors with eigenvalue  $\lambda_2 = 4$ :

$$A - \lambda_2 I = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \quad \left( A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \right)$$

## Solution (continued)

- Find the eigenvectors with eigenvalue  $\lambda_2 = 4$ :

$$A - \lambda_2 I = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \quad \left( A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \right)$$

Solutions to  $\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0}$  have basis  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .



## Solution (continued)

- Find the eigenvectors with eigenvalue  $\lambda_2 = 4$ :

$$A - \lambda_2 I = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \quad \left( A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \right)$$

Solutions to  $\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0}$  have basis  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .

So:  $\mathbf{x}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda_2 = 4$ .

## Solution (continued)

- Find the eigenvectors with eigenvalue  $\lambda_2 = 4$ :

$$A - \lambda_2 I = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \quad \left( A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \right)$$

Solutions to  $\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0}$  have basis  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .

So:  $\mathbf{x}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda_2 = 4$ .

The eigenspace of  $\lambda = 4$  is  $\text{Span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ .

## Solution (continued)

- Find the eigenvectors with eigenvalue  $\lambda_2 = 4$ :

$$A - \lambda_2 I = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \quad \left( A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix} \right)$$

Solutions to  $\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0}$  have basis  $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .

So:  $\mathbf{x}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector with eigenvalue  $\lambda_2 = 4$ .

The eigenspace of  $\lambda = 4$  is  $\text{Span} \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$ .



## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$$

## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$$

## Solution

- The characteristic polynomial is:

$$\det(A - \lambda I) = \begin{vmatrix} 3 - \lambda & 2 & 3 \\ 0 & 6 - \lambda & 10 \\ 0 & 0 & 2 - \lambda \end{vmatrix} = (3 - \lambda)(6 - \lambda)(2 - \lambda)$$

## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$$

## Solution

- The characteristic polynomial is:

$$\det(A - \lambda I) = \begin{vmatrix} 3 - \lambda & 2 & 3 \\ 0 & 6 - \lambda & 10 \\ 0 & 0 & 2 - \lambda \end{vmatrix} = (3 - \lambda)(6 - \lambda)(2 - \lambda)$$

- $A$  has eigenvalues 2, 3, 6.

## Example

Find the eigenvectors and eigenvalues of

$$A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$$

## Solution

- The characteristic polynomial is:

$$\det(A - \lambda I) = \begin{vmatrix} 3 - \lambda & 2 & 3 \\ 0 & 6 - \lambda & 10 \\ 0 & 0 & 2 - \lambda \end{vmatrix} = (3 - \lambda)(6 - \lambda)(2 - \lambda)$$

- $A$  has eigenvalues 2, 3, 6.

The eigenvalues of a triangular matrix are its diagonal entries.



## Solution (continued)

- $\lambda_1 = 2$ :

$$(A - \lambda_1 I)\mathbf{x} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 10 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

## Solution (continued)

- $\lambda_1 = 2$ :

$$(A - \lambda_1 I)\mathbf{x} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 10 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 2 \\ -5/2 \\ 1 \end{bmatrix}$$

## Solution (continued)

- $\lambda_1 = 2$ :

$$(A - \lambda_1 I)\mathbf{x} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 10 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 2 \\ -5/2 \\ 1 \end{bmatrix}$$

- $\lambda_2 = 3$ :

$$(A - \lambda_2 I)\mathbf{x} = \begin{bmatrix} 0 & 2 & 3 \\ 0 & 3 & 10 \\ 0 & 0 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

## Solution (continued)

- $\lambda_1 = 2$ :

$$(A - \lambda_1 I)\mathbf{x} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 10 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 2 \\ -5/2 \\ 1 \end{bmatrix}$$

- $\lambda_2 = 3$ :

$$(A - \lambda_2 I)\mathbf{x} = \begin{bmatrix} 0 & 2 & 3 \\ 0 & 3 & 10 \\ 0 & 0 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_2 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

## Solution (continued)

- $\lambda_1 = 2$ :

$$(A - \lambda_1 I)\mathbf{x} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 10 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 2 \\ -5/2 \\ 1 \end{bmatrix}$$

- $\lambda_2 = 3$ :

$$(A - \lambda_2 I)\mathbf{x} = \begin{bmatrix} 0 & 2 & 3 \\ 0 & 3 & 10 \\ 0 & 0 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_2 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- $\lambda_3 = 6$ :

$$(A - \lambda_3 I)\mathbf{x} = \begin{bmatrix} -3 & 2 & 3 \\ 0 & 0 & 10 \\ 0 & 0 & -4 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

## Solution (continued)

- $\lambda_1 = 2$ :

$$(A - \lambda_1 I)\mathbf{x} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 10 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 2 \\ -5/2 \\ 1 \end{bmatrix}$$

- $\lambda_2 = 3$ :

$$(A - \lambda_2 I)\mathbf{x} = \begin{bmatrix} 0 & 2 & 3 \\ 0 & 3 & 10 \\ 0 & 0 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_2 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- $\lambda_3 = 6$ :

$$(A - \lambda_3 I)\mathbf{x} = \begin{bmatrix} -3 & 2 & 3 \\ 0 & 0 & 10 \\ 0 & 0 & -4 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_3 = \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}$$

## Solution (continued)

- Each of those matrices had a one-dimensional null space.

## Solution (continued)

- Each of those matrices had a one-dimensional null space. So our eigenvectors are not unique.



## Solution (continued)

- Each of those matrices had a one-dimensional null space. So our eigenvectors are not unique. They are unique up to scaling.

## Solution (continued)

- Each of those matrices had a one-dimensional null space. So our eigenvectors are not unique. They are unique up to scaling.

- In summary,  $A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$  has eigenvalues 2, 3, 6

## Solution (continued)

- Each of those matrices had a one-dimensional null space. So our eigenvectors are not unique. They are unique up to scaling.

- In summary,  $A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$  has eigenvalues 2, 3, 6 with corresponding eigenvectors

$$\begin{bmatrix} 2 \\ -5/2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2/3 \\ 1 \\ 0 \end{bmatrix}.$$

## Solution (continued)

- Each of those matrices had a one-dimensional null space. So our eigenvectors are not unique. They are unique up to scaling.
- In summary,  $A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$  has eigenvalues 2, 3, 6 with corresponding eigenvectors

$$\begin{bmatrix} 2 \\ -5/2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2/3 \\ 1 \\ 0 \end{bmatrix}.$$

These three vectors are independent. By the next result, this is always so.



## Theorem

*If  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are eigenvectors of  $A$  corresponding to different eigenvalues, then they are independent.*

## Theorem

*If  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are eigenvectors of  $A$  corresponding to different eigenvalues, then they are independent.*

## Proof.

Suppose, for contradiction, that  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are dependent.

## Theorem

*If  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are eigenvectors of  $A$  corresponding to different eigenvalues, then they are independent.*

## Proof.

Suppose, for contradiction, that  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are dependent. By kicking out some vectors, we may assume that there is (up to multiples) only one linear relation:  $c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m = \mathbf{0}$ .



## Theorem

*If  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are eigenvectors of  $A$  corresponding to different eigenvalues, then they are independent.*

## Proof.

Suppose, for contradiction, that  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are dependent.

By kicking out some vectors, we may assume that there is (up to multiples) only one linear relation:  $c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m = \mathbf{0}$ .

In other words, the matrix with columns  $\mathbf{x}_1, \dots, \mathbf{x}_m$  has one-dimensional null space.

## Theorem

*If  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are eigenvectors of  $A$  corresponding to different eigenvalues, then they are independent.*

## Proof.

Suppose, for contradiction, that  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are dependent.

By kicking out some vectors, we may assume that there is (up to multiples) only one linear relation:  $c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m = \mathbf{0}$ .

In other words, the matrix with columns  $\mathbf{x}_1, \dots, \mathbf{x}_m$  has one-dimensional null space. Now multiply this relation with  $A$ :

$$A(c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m) = c_1\lambda_1\mathbf{x}_1 + \dots + c_m\lambda_m\mathbf{x}_m = \mathbf{0}$$

This is a second independent relation!

## Theorem

*If  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are eigenvectors of  $A$  corresponding to different eigenvalues, then they are independent.*

## Proof.

Suppose, for contradiction, that  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are dependent.

By kicking out some vectors, we may assume that there is (up to multiples) only one linear relation:  $c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m = \mathbf{0}$ .

In other words, the matrix with columns  $\mathbf{x}_1, \dots, \mathbf{x}_m$  has one-dimensional null space. Now multiply this relation with  $A$ :

$$A(c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m) = c_1\lambda_1\mathbf{x}_1 + \dots + c_m\lambda_m\mathbf{x}_m = \mathbf{0}$$

This is a second independent relation! In other words, two independent vectors living in a one-dimensional vector space.

## Theorem

*If  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are eigenvectors of  $A$  corresponding to different eigenvalues, then they are independent.*

## Proof.

Suppose, for contradiction, that  $\mathbf{x}_1, \dots, \mathbf{x}_m$  are dependent.

By kicking out some vectors, we may assume that there is (up to multiples) only one linear relation:  $c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m = \mathbf{0}$ .

In other words, the matrix with columns  $\mathbf{x}_1, \dots, \mathbf{x}_m$  has one-dimensional null space. Now multiply this relation with  $A$ :

$$A(c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m) = c_1\lambda_1\mathbf{x}_1 + \dots + c_m\lambda_m\mathbf{x}_m = \mathbf{0}$$

This is a second independent relation! In other words, two independent vectors living in a one-dimensional vector space.

Contradiction. □

## Relations between eigenvalues

## Product of Eigenvalues

If  $A$  is  $n \times n$  get in principle  $n$  eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ .

If  $A$  is  $n \times n$  get in principle  $n$  eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ . How are these eigenvalues related?



If  $A$  is  $n \times n$  get in principle  $n$  eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ . How are these eigenvalues related?

### Theorem

*The product of eigenvalues  $\lambda_1 \lambda_2 \dots \lambda_n$  is equal to the determinant of  $A$ .*

If  $A$  is  $n \times n$  get in principle  $n$  eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ . How are these eigenvalues related?

### Theorem

*The product of eigenvalues  $\lambda_1 \lambda_2 \dots \lambda_n$  is equal to the determinant of  $A$ .*

### Proof.

The characteristic polynomial  $p(\lambda) = \det(A - \lambda I)$  has constant term  $\det(A)$ .

If  $A$  is  $n \times n$  get in principle  $n$  eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ . How are these eigenvalues related?

### Theorem

*The product of eigenvalues  $\lambda_1 \lambda_2 \dots \lambda_n$  is equal to the determinant of  $A$ .*

### Proof.

The characteristic polynomial  $p(\lambda) = \det(A - \lambda I)$  has constant term  $\det(A)$ . On the other hand  $p(\lambda)$  factors, because the roots are the eigenvalues we get  $p(\lambda) = (\pm 1)(\lambda - \lambda_1)(\lambda - \lambda_2) \dots (\lambda - \lambda_n)$ , which has constant term  $\lambda_1 \lambda_2 \dots \lambda_n$ . □

## Product of Eigenvalues

If  $A$  is  $n \times n$  get in principle  $n$  eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ . How are these eigenvalues related?

## Theorem

*The product of eigenvalues  $\lambda_1 \lambda_2 \dots \lambda_n$  is equal to the determinant of  $A$ .*

## Proof.

The characteristic polynomial  $p(\lambda) = \det(A - \lambda I)$  has constant term  $\det(A)$ . On the other hand  $p(\lambda)$  factors, because the roots are the eigenvalues we get  $p(\lambda) = (\pm 1)(\lambda - \lambda_1)(\lambda - \lambda_2) \dots (\lambda - \lambda_n)$ , which has constant term  $\lambda_1 \lambda_2 \dots \lambda_n$ . □

## Example

Let  $A = \begin{bmatrix} \lambda_1 & b \\ 0 & \lambda_2 \end{bmatrix}$ . Then the eigenvalues are  $\lambda_1, \lambda_2$  and  $\det(A) = \lambda_1 \lambda_2$ .

## Sum of Eigenvalues

What other relations are there between the eigenvalues?

What other relations are there between the eigenvalues?

### Definition

Let  $A = \begin{bmatrix} a_{11} & a_{12} & \cdots \\ a_{21} & a_{22} & \cdots \\ \vdots & \vdots & \ddots \end{bmatrix}$  be  $n \times n$ . Then the **TRACE** of  $A$  is the sum of the diagonal entries:  $\text{Tr}(A) = a_{11} + a_{22} + \cdots + a_{nn}$ .

What other relations are there between the eigenvalues?

### Definition

Let  $A = \begin{bmatrix} a_{11} & a_{12} & \cdots \\ a_{21} & a_{22} & \cdots \\ \vdots & \vdots & \ddots \end{bmatrix}$  be  $n \times n$ . Then the **TRACE** of  $A$  is the sum of the diagonal entries:  $\text{Tr}(A) = a_{11} + a_{22} + \cdots + a_{nn}$ .

### Theorem

Let  $A$  be  $n \times n$ . Then the trace of  $A$  is the **sum** of eigenvalues:

$$\text{Tr}(A) = \lambda_1 + \lambda_2 + \cdots + \lambda_n$$



### Example

Let  $A = \begin{bmatrix} \lambda_1 & b \\ 0 & \lambda_2 \end{bmatrix}$ . What are the eigenvalues and what is  $\text{Tr}(A)$ ?

### Example

Let  $A = \begin{bmatrix} \lambda_1 & b \\ 0 & \lambda_2 \end{bmatrix}$ . What are the eigenvalues and what is  $\text{Tr}(A)$ ?

### Solution

The eigenvalues are  $\lambda_1, \lambda_2$  and  $\text{Tr}(A) = \lambda_1 + \lambda_2$ .

## The Characteristic Polynomial for $2 \times 2$

◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻

The Characteristic Polynomial for  $2 \times 2$ 

$2 \times 2$  matrices are easy.

### Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

The Characteristic Polynomial for  $2 \times 2$ 

$2 \times 2$  matrices are easy.

## Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

## The Characteristic Polynomial for $2 \times 2$

$2 \times 2$  matrices are easy.

### Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

### Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

### Solution

$$\text{Tr}(A) =$$

# The Characteristic Polynomial for $2 \times 2$

$2 \times 2$  matrices are easy.

## Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

## Solution

$$\text{Tr}(A) = 6,$$



# The Characteristic Polynomial for $2 \times 2$

$2 \times 2$  matrices are easy.

## Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

## Solution

$$\text{Tr}(A) = 6, \det(A) =$$

The Characteristic Polynomial for  $2 \times 2$ 

$2 \times 2$  matrices are easy.

## Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

## Solution

$$\text{Tr}(A) = 6, \det(A) = 8,$$

The Characteristic Polynomial for  $2 \times 2$ 

$2 \times 2$  matrices are easy.

## Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

## Solution

$\text{Tr}(A) = 6$ ,  $\det(A) = 8$ , so  $p(\lambda) =$

# The Characteristic Polynomial for $2 \times 2$

$2 \times 2$  matrices are easy.

## Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

## Solution

$\text{Tr}(A) = 6$ ,  $\det(A) = 8$ , so  $p(\lambda) = \lambda^2 - 6\lambda + 8$ .

# The Characteristic Polynomial for $2 \times 2$

$2 \times 2$  matrices are easy.

## Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

## Solution

$\text{Tr}(A) = 6$ ,  $\det(A) = 8$ , so  $p(\lambda) = \lambda^2 - 6\lambda + 8$ . Also in terms of eigenvalues  $\text{Tr}(A) =$

The Characteristic Polynomial for  $2 \times 2$ 

$2 \times 2$  matrices are easy.

### Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

### Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

### Solution

$\text{Tr}(A) = 6$ ,  $\det(A) = 8$ , so  $p(\lambda) = \lambda^2 - 6\lambda + 8$ . Also in terms of eigenvalues  $\text{Tr}(A) = \lambda_1 + \lambda_2$

The Characteristic Polynomial for  $2 \times 2$ 

$2 \times 2$  matrices are easy.

### Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

### Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

### Solution

$\text{Tr}(A) = 6$ ,  $\det(A) = 8$ , so  $p(\lambda) = \lambda^2 - 6\lambda + 8$ . Also in terms of eigenvalues  $\text{Tr}(A) = \lambda_1 + \lambda_2$  and  $\det(A) =$

The Characteristic Polynomial for  $2 \times 2$ 

$2 \times 2$  matrices are easy.

### Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

### Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

### Solution

$\text{Tr}(A) = 6$ ,  $\det(A) = 8$ , so  $p(\lambda) = \lambda^2 - 6\lambda + 8$ . Also in terms of eigenvalues  $\text{Tr}(A) = \lambda_1 + \lambda_2$  and  $\det(A) = \lambda_1\lambda_2$ .



# The Characteristic Polynomial for $2 \times 2$

$2 \times 2$  matrices are easy.

## Theorem

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

## Example

Let  $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ . What are the eigenvalues and what is the characteristic polynomial?

## Solution

$\text{Tr}(A) = 6$ ,  $\det(A) = 8$ , so  $p(\lambda) = \lambda^2 - 6\lambda + 8$ . Also in terms of eigenvalues  $\text{Tr}(A) = \lambda_1 + \lambda_2$  and  $\det(A) = \lambda_1\lambda_2$ . So  $\lambda_1 = 2$ ,  $\lambda_2 = 4$

## Practice problems

## Example

Find the eigenvectors and eigenvalues of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .

### Example

Find the eigenvectors and eigenvalues of  $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$ .

### Example

What are the eigenvalues of  $A = \begin{bmatrix} 2 & 0 & 0 & 0 \\ -1 & 3 & 0 & 0 \\ -1 & 1 & 3 & 0 \\ 0 & 1 & 2 & 4 \end{bmatrix}$ .

No calculations!