

Math 415 - Lecture 36

Minima, maxima and saddle points, Constrained Optimization

Wednesday December 2nd 2015

Textbook reading: Chapter 6.1

Textbook reading: Chapter 6.1

Suggested practice exercises: Chapter 6.1, # 1, 16

Textbook reading: Chapter 6.1

Suggested practice exercises: Chapter 6.1, # 1, 16

Strang lecture: Lecture 27: Positive definite matrices and minima

Review

Spectral theorem:

- A is a **symmetric** matrix if $A^T = A$.

Spectral theorem:

- A is a **symmetric** matrix if $A^T = A$.
- Any $n \times n$ symmetric matrix A has n **real eigenvalues** and an **orthonormal eigenbasis** $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$.

Spectral theorem:

- A is a **symmetric** matrix if $A^T = A$.
- Any $n \times n$ symmetric matrix A has n **real eigenvalues** and an **orthonormal eigenbasis** $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$.
- So, we can write $A = QDQ^T$ where

$$D = \underbrace{\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \quad \text{and} \quad Q = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{q}_1 & \cdots & \mathbf{q}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}}$$

Spectral theorem:

- A is a **symmetric** matrix if $A^T = A$.
- Any $n \times n$ symmetric matrix A has n **real eigenvalues** and an **orthonormal eigenbasis** $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$.
- So, we can write $A = QDQ^T$ where

$$D = \underbrace{\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \quad \text{and} \quad Q = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{q}_1 & \cdots & \mathbf{q}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}}$$

- A is called **positive definite** if $\mathbf{x}^T A \mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$.

Spectral theorem:

- A is a **symmetric** matrix if $A^T = A$.
- Any $n \times n$ symmetric matrix A has n **real eigenvalues** and an **orthonormal eigenbasis** $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$.
- So, we can write $A = QDQ^T$ where

$$D = \underbrace{\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \quad \text{and} \quad Q = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{q}_1 & \cdots & \mathbf{q}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}}$$

- A is called **positive definite** if $\mathbf{x}^T A \mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$.
- a function of the form $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ is called a **quadratic form**.

Spectral theorem:

- A is a **symmetric** matrix if $A^T = A$.
- Any $n \times n$ symmetric matrix A has n **real eigenvalues** and an **orthonormal eigenbasis** $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$.
- So, we can write $A = QDQ^T$ where

$$D = \underbrace{\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \quad \text{and} \quad Q = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{q}_1 & \cdots & \mathbf{q}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}}$$

- A is called **positive definite** if $\mathbf{x}^T A \mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$.
- a function of the form $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ is called a **quadratic form**.
- Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A . Then

Spectral theorem:

- A is a **symmetric** matrix if $A^T = A$.
- Any $n \times n$ symmetric matrix A has n **real eigenvalues** and an **orthonormal eigenbasis** $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$.
- So, we can write $A = QDQ^T$ where

$$D = \underbrace{\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \quad \text{and} \quad Q = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{q}_1 & \cdots & \mathbf{q}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}}$$

- A is called **positive definite** if $\mathbf{x}^T A \mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$.
- a function of the form $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ is called a **quadratic form**.
- Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A . Then
 - ① If all $\lambda_i > 0$, then A is positive definite,

Spectral theorem:

- A is a **symmetric** matrix if $A^T = A$.
- Any $n \times n$ symmetric matrix A has n **real eigenvalues** and an **orthonormal eigenbasis** $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$.
- So, we can write $A = QDQ^T$ where

$$D = \underbrace{\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \quad \text{and} \quad Q = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{q}_1 & \cdots & \mathbf{q}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}}$$

- A is called **positive definite** if $\mathbf{x}^T A \mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$.
- a function of the form $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ is called a **quadratic form**.
- Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A . Then
 - 1 If all $\lambda_i > 0$, then A is positive definite,
 - 2 If all $\lambda_i < 0$, then $\mathbf{x}^T A \mathbf{x} < 0$ for all $\mathbf{x} \neq \mathbf{0}$

Spectral theorem:

- A is a **symmetric** matrix if $A^T = A$.
- Any $n \times n$ symmetric matrix A has n **real eigenvalues** and an **orthonormal eigenbasis** $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$.
- So, we can write $A = QDQ^T$ where

$$D = \underbrace{\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}}_{\text{matrix of eigenvalues}} \quad \text{and} \quad Q = \underbrace{\begin{bmatrix} | & & | \\ \mathbf{q}_1 & \cdots & \mathbf{q}_n \\ | & & | \end{bmatrix}}_{\text{matrix of eigenvectors}}$$

- A is called **positive definite** if $\mathbf{x}^T A \mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$.
- a function of the form $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ is called a **quadratic form**.
- Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A . Then
 - ① If all $\lambda_i > 0$, then A is positive definite,
 - ② If all $\lambda_i < 0$, then $\mathbf{x}^T A \mathbf{x} < 0$ for all $\mathbf{x} \neq \mathbf{0}$
 - ③ If some $\lambda_i > 0$, some $\lambda_j < 0$, $\mathbf{x}^T A \mathbf{x}$ will have both positive and negative values.

oooooooo

2nd derivative test

●○○○○○○○

2nd derivative test

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable, the **Hessian** matrix of f

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial^2 x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_1 \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(\mathbf{0}) \\ \frac{\partial^2 f}{\partial x_2 \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial^2 x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n}(\mathbf{0}) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_n \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial^2 x_n}(\mathbf{0}) \end{bmatrix}$$

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable, the **Hessian** matrix of f

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial^2 x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_1 \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(\mathbf{0}) \\ \frac{\partial^2 f}{\partial x_2 \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial^2 x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n}(\mathbf{0}) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_n \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial^2 x_n}(\mathbf{0}) \end{bmatrix}$$

Idea

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable and

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable, the **Hessian** matrix of f

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial^2 x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_1 \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(\mathbf{0}) \\ \frac{\partial^2 f}{\partial x_2 \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial^2 x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n}(\mathbf{0}) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_n \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial^2 x_n}(\mathbf{0}) \end{bmatrix}$$

Idea

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable and

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable, the **Hessian** matrix of f

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial^2 x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_1 \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(\mathbf{0}) \\ \frac{\partial^2 f}{\partial x_2 \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial^2 x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n}(\mathbf{0}) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_n \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial^2 x_n}(\mathbf{0}) \end{bmatrix}$$

Idea

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable and $\mathbf{0}$ is a critical point, then

Definition

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable, the **Hessian** matrix of f

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial^2 x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_1 \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n}(\mathbf{0}) \\ \frac{\partial^2 f}{\partial x_2 \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial^2 x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n}(\mathbf{0}) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1}(\mathbf{0}) & \frac{\partial^2 f}{\partial x_n \partial x_2}(\mathbf{0}) & \cdots & \frac{\partial^2 f}{\partial^2 x_n}(\mathbf{0}) \end{bmatrix}$$

Idea

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is twice differentiable and $\mathbf{0}$ is a critical point, then $f(\mathbf{x}) \approx f(\mathbf{0}) + \frac{1}{2}\mathbf{x} \cdot H\mathbf{x}$.

- H is always symmetric

2nd derivative test

- H is always symmetric
- We're approximating $f(\mathbf{x})$ by $f(\mathbf{0})$ plus a **quadratic** function,

2nd derivative test

- H is always symmetric
- We're approximating $f(\mathbf{x})$ by $f(\mathbf{0})$ plus a quadratic function,

- H is always symmetric
- We're approximating $f(\mathbf{x})$ by $f(\mathbf{0})$ plus a **quadratic** function,
 $\frac{1}{2}\mathbf{x} \cdot H\mathbf{x}$!

2nd derivative test

- H is always symmetric
- We're approximating $f(\mathbf{x})$ by $f(\mathbf{0})$ plus a **quadratic** function,
 $\frac{1}{2}\mathbf{x} \cdot H\mathbf{x}$!
- We understand $q(\mathbf{x}) = \frac{1}{2}\mathbf{x} \cdot H\mathbf{x} \implies$

- H is always symmetric
- We're approximating $f(\mathbf{x})$ by $f(\mathbf{0})$ plus a **quadratic** function,
 $\frac{1}{2}\mathbf{x} \cdot H\mathbf{x}$!
- We understand $q(\mathbf{x}) = \frac{1}{2}\mathbf{x} \cdot H\mathbf{x} \implies$

- H is always symmetric
- We're approximating $f(\mathbf{x})$ by $f(\mathbf{0})$ plus a **quadratic** function, $\frac{1}{2}\mathbf{x} \cdot H\mathbf{x}$!
- We understand $q(\mathbf{x}) = \frac{1}{2}\mathbf{x} \cdot H\mathbf{x} \implies$ we understand if $\mathbf{0}$ is a max, min or neither for f !

- H is always symmetric
- We're approximating $f(\mathbf{x})$ by $f(\mathbf{0})$ plus a **quadratic** function, $\frac{1}{2}\mathbf{x} \cdot H\mathbf{x}$!
- We understand $q(\mathbf{x}) = \frac{1}{2}\mathbf{x} \cdot H\mathbf{x} \implies$ we understand if $\mathbf{0}$ is a max, min or neither for f !
- Turns out:

- H is always symmetric
- We're approximating $f(\mathbf{x})$ by $f(\mathbf{0})$ plus a **quadratic** function, $\frac{1}{2}\mathbf{x} \cdot H\mathbf{x}$!
- We understand $q(\mathbf{x}) = \frac{1}{2}\mathbf{x} \cdot H\mathbf{x} \implies$ we understand if $\mathbf{0}$ is a max, min or neither for f !
- Turns out:

- H is always symmetric
- We're approximating $f(\mathbf{x})$ by $f(\mathbf{0})$ plus a **quadratic** function, $\frac{1}{2}\mathbf{x} \cdot H\mathbf{x}$!
- We understand $q(\mathbf{x}) = \frac{1}{2}\mathbf{x} \cdot H\mathbf{x} \implies$ we understand if $\mathbf{0}$ is a max, min or neither for f !
- Turns out: $q(\mathbf{x})$ is determined by eigenvectors and eigenvalues of H !

Theorem (2nd derivative test)

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$, then

Theorem (2nd derivative test)

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$, then

- 1 If all eigenvalues of H are *positive*, then $\mathbf{0}$ is a local *min*.
 H is positive-definite, graph is a bowl.

Theorem (2nd derivative test)

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$, then

- 1 If all eigenvalues of H are **positive**, then $\mathbf{0}$ is a local **min**.
 H is positive-definite, graph is a bowl.
- 2 If all eigenvalues of H are **negative**, then $\mathbf{0}$ is a local **max**.
 H is negative-definite, graph is a dome.

Theorem (2nd derivative test)

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$, then

- 1 If all eigenvalues of H are **positive**, then $\mathbf{0}$ is a local **min**.
 H is positive-definite, graph is a bowl.
- 2 If all eigenvalues of H are **negative**, then $\mathbf{0}$ is a local **max**.
 H is negative-definite, graph is a dome.
- 3 If one eigenvalue of H is **positive** and one is **negative**, then $\mathbf{0}$ is **neither** a max nor a min.
 H is indefinite, graph is a saddle

Theorem (2nd derivative test)

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$, then

- 1 If all eigenvalues of H are **positive**, then $\mathbf{0}$ is a local **min**.
 H is positive-definite, graph is a bowl.
- 2 If all eigenvalues of H are **negative**, then $\mathbf{0}$ is a local **max**.
 H is negative-definite, graph is a dome.
- 3 If one eigenvalue of H is **positive** and one is **negative**, then $\mathbf{0}$ is **neither** a max nor a min.
 H is indefinite, graph is a saddle
- 4 Otherwise (e.g. all eigenvalues positive or zero), no information!

Theorem (2nd derivative test)

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$, then

- 1 If all eigenvalues of H are **positive**, then $\mathbf{0}$ is a local **min**.
 H is positive-definite, graph is a bowl.
- 2 If all eigenvalues of H are **negative**, then $\mathbf{0}$ is a local **max**.
 H is negative-definite, graph is a dome.
- 3 If one eigenvalue of H is **positive** and one is **negative**, then $\mathbf{0}$ is **neither** a max nor a min.
 H is indefinite, graph is a saddle
- 4 Otherwise (e.g. all eigenvalues positive or zero), no information!

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian

$$H = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}.$$

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

(An example of such a function is $f(x, y) = \frac{1}{2}x^2 + 2xy + \frac{1}{2}y^2$).

○○○○○●○○

Solution

Solution

We showed that H has eigenvalues 3 and -1. So

Solution

We showed that H has eigenvalues 3 and -1. So f has a **saddle point** at $\mathbf{0}$.

2nd derivative test




Figure : Graph of the function $f(x, y) = \frac{1}{2}x^2 + 2xy + \frac{1}{2}y^2$

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian

$$H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}.$$

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Solution

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Solution

Eigenvalues: Sum $\lambda_1 + \lambda_2 = \text{Tr}(H)$

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Solution

Eigenvalues: Sum $\lambda_1 + \lambda_2 = \text{Tr}(H) = 4$

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Solution

Eigenvalues: Sum $\lambda_1 + \lambda_2 = \text{Tr}(H) = 4$
Product $\lambda_1 \lambda_2 = \det(H)$

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Solution

Eigenvalues: Sum $\lambda_1 + \lambda_2 = \text{Tr}(H) = 4$
Product $\lambda_1 \lambda_2 = \det(H) = 2$.

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Solution

Eigenvalues: Sum $\lambda_1 + \lambda_2 = \text{Tr}(H) = 4$
Product $\lambda_1 \lambda_2 = \det(H) = 2$.
So, λ_1, λ_2 must be positive!

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Solution

Eigenvalues: Sum $\lambda_1 + \lambda_2 = \text{Tr}(H) = 4$

Product $\lambda_1 \lambda_2 = \det(H) = 2$.

So, λ_1, λ_2 must be positive!

(positive product \implies both positive or both negative.)

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Solution

Eigenvalues: Sum $\lambda_1 + \lambda_2 = \text{Tr}(H) = 4$

Product $\lambda_1 \lambda_2 = \det(H) = 2$.

So, λ_1, λ_2 must be positive!

(positive product \implies both positive or both negative.

positive sum \implies both positive.)

Example

Suppose $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ has a critical point at $\mathbf{0}$ and has Hessian $H = \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$. Does f have local max, min or neither at $\mathbf{0}$?

Solution

Eigenvalues: Sum $\lambda_1 + \lambda_2 = \text{Tr}(H) = 4$

Product $\lambda_1 \lambda_2 = \det(H) = 2$.

So, λ_1, λ_2 must be positive!

(positive product \implies both positive or both negative.

positive sum \implies both positive.)

2nd derivative test says: $f(\mathbf{0})$ is local **min**.

2nd derivative test




Figure : Graph of the function $f(x, y) = \frac{1}{2}x^2 - xy + \frac{3}{2}y^2$

Constrained optimization

Problem: Given a quadratic form $q(\mathbf{x})$, find the maximum or minimum value $q(\mathbf{x})$ for \mathbf{x} in some specified set. Typically, the problem can be arranged such that \mathbf{x} varies over the set of vectors with $\mathbf{x}^T \mathbf{x} = 1$.

Example

Let $A = \begin{bmatrix} 9 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 3 \end{bmatrix}$. Find the maximum and minimum values of $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$.

Problem: Given a quadratic form $q(\mathbf{x})$, find the maximum or minimum value $q(\mathbf{x})$ for \mathbf{x} in some specified set. Typically, the problem can be arranged such that \mathbf{x} varies over the set of vectors with $\mathbf{x}^T \mathbf{x} = 1$.

Example

Let $A = \begin{bmatrix} 9 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 3 \end{bmatrix}$. Find the maximum and minimum values of $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$.

Solution

The quadratic form is $q(x_1, x_2, x_3) = 9x_1^2 + 4x_2^2 + 3x_3^2$. We are interested in the maximal value for q when (x_1, x_2, x_3) is such that $x_1^2 + x_2^2 + x_3^2 = 1$.

Problem: Given a quadratic form $q(\mathbf{x})$, find the maximum or minimum value $q(\mathbf{x})$ for \mathbf{x} in some specified set. Typically, the problem can be arranged such that \mathbf{x} varies over the set of vectors with $\mathbf{x}^T \mathbf{x} = 1$.

Example

Let $A = \begin{bmatrix} 9 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 3 \end{bmatrix}$. Find the maximum and minimum values of $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$.

Solution

The quadratic form is $q(x_1, x_2, x_3) = 9x_1^2 + 4x_2^2 + 3x_3^2$. We are interested in the maximal value for q when (x_1, x_2, x_3) is such that $x_1^2 + x_2^2 + x_3^2 = 1$. Now we can give an upper bound for q : we obviously have

$$q(\mathbf{x}) \leq 9x_1^2 + 9x_2^2 + 9x_3^2 = 9$$

Solution

So $q(\mathbf{x})$ can not be bigger than 9, for any \mathbf{x} . Can we get $q(\mathbf{x}) = 9$ for some \mathbf{x} ?

Solution

So $q(\mathbf{x})$ can not be bigger than 9, for any \mathbf{x} . Can we get $q(\mathbf{x}) = 9$ for some \mathbf{x} ?

Solution

So $q(\mathbf{x})$ can not be bigger than 9, for any \mathbf{x} . Can we get $q(\mathbf{x}) = 9$ for some \mathbf{x} ? Obviously for $\mathbf{x} = (1, 0, 0)$ we achieve the upper bound, so 9 is the maximum value for q (under this constraint.)

Solution

So $q(\mathbf{x})$ can not be bigger than 9, for any \mathbf{x} . Can we get $q(\mathbf{x}) = 9$ for some \mathbf{x} ? Obviously for $\mathbf{x} = (1, 0, 0)$ we achieve the upper bound, so 9 is the maximum value for q (under this constraint.) What is a lower bound?

Solution

So $q(\mathbf{x})$ can not be bigger than 9, for any \mathbf{x} . Can we get $q(\mathbf{x}) = 9$ for some \mathbf{x} ? Obviously for $\mathbf{x} = (1, 0, 0)$ we achieve the upper bound, so 9 is the maximum value for q (under this constraint.) What is a lower bound? For which \mathbf{x} is the lower bound achieved?

What if A is not diagonal?

What if A is not diagonal?

Theorem

Let A be a symmetric matrix and let λ_m be the least eigenvalue and λ_M be the greatest eigenvalue of A . Then

$$\lambda_m = \min\{\mathbf{x}^T A \mathbf{x} : \mathbf{x}^T \mathbf{x} = 1\},$$

moreover if \mathbf{u}_m is a unit eigenvector corresponding to λ_m , then $\mathbf{u}_m^T A \mathbf{u}_m = \lambda_m$.

In addition,

$$\lambda_M = \max\{\mathbf{x}^T A \mathbf{x} : \mathbf{x}^T \mathbf{x} = 1\},$$

moreover if \mathbf{u}_M is a unit eigenvector corresponding to λ_M , then $\mathbf{u}_M^T A \mathbf{u}_M = \lambda_M$.

What if A is not diagonal?

Theorem

Let A be a symmetric matrix and let λ_m be the least eigenvalue and λ_M be the greatest eigenvalue of A . Then

$$\lambda_m = \min\{\mathbf{x}^T A \mathbf{x} : \mathbf{x}^T \mathbf{x} = 1\},$$

moreover if \mathbf{u}_m is a unit eigenvector corresponding to λ_m , then $\mathbf{u}_m^T A \mathbf{u}_m = \lambda_m$.

In addition,

$$\lambda_M = \max\{\mathbf{x}^T A \mathbf{x} : \mathbf{x}^T \mathbf{x} = 1\},$$

moreover if \mathbf{u}_M is a unit eigenvector corresponding to λ_M , then $\mathbf{u}_M^T A \mathbf{u}_M = \lambda_M$.

Proof.

We know by the spectral theorem that $A = QDQ^T$, and so we can write $q(\mathbf{x}) = \mathbf{x}^T QDQ^T \mathbf{x} = u^T D u = \lambda_M u_1^2 + \cdots + \lambda_m u_m^2$, where $u = Q^T \mathbf{x}$.

Proof.

We know by the spectral theorem that $A = QDQ^T$, and so we can write $q(\mathbf{x}) = \mathbf{x}^T QDQ^T \mathbf{x} = u^T D u = \lambda_M u_1^2 + \cdots + \lambda_m u_m^2$, where $u = Q^T \mathbf{x}$.

Proof.

We know by the spectral theorem that $A = QDQ^T$, and so we can write $q(\mathbf{x}) = \mathbf{x}^T QDQ^T \mathbf{x} = u^T D u = \lambda_M u_1^2 + \cdots + \lambda_m u_m^2$, where $u = Q^T \mathbf{x}$. As before we see that the largest eigenvalue λ_M is the upper bound for q , achieved for $u = (1, 0, \dots, 0)$ or $\mathbf{x} = Qu$, the normalized eigenvector corresponding to λ_M .

Proof.

We know by the spectral theorem that $A = QDQ^T$, and so we can write $q(\mathbf{x}) = \mathbf{x}^T QDQ^T \mathbf{x} = u^T Du = \lambda_M u_1^2 + \cdots + \lambda_m u_m^2$, where $u = Q^T \mathbf{x}$. As before we see that the largest eigenvalue λ_M is the upper bound for q , achieved for $u = (1, 0, \dots, 0)$ or $\mathbf{x} = Qu$, the normalized eigenvector corresponding to λ_M . Same argument for λ_m . □

Example

Let $A = \begin{bmatrix} 3 & 2 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 4 \end{bmatrix}$. Find the maximum and minimum values of $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$.

Solution

We first find eigenvectors and eigenvalues for A .

Example

Let $A = \begin{bmatrix} 3 & 2 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 4 \end{bmatrix}$. Find the maximum and minimum values of $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$.

Solution

We first find eigenvectors and eigenvalues for A . Let us ask Wolfram Alpha: $\det(A)$, Eigenvalues and eigenvectors .

Example

Let $A = \begin{bmatrix} 3 & 2 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 4 \end{bmatrix}$. Find the maximum and minimum values of $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$.

Solution

We first find eigenvectors and eigenvalues for A . Let us ask Wolfram Alpha: [det\(A\)](#), [Eigenvalues and eigenvectors](#). So $\lambda = 6, 3, 1$, with eigenvectors

$$v_1 = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \quad v_2 = \frac{1}{\sqrt{6}} \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}, \quad \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Example

Let $A = \begin{bmatrix} 3 & 2 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 4 \end{bmatrix}$. Find the maximum and minimum values of $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$.

Solution

We first find eigenvectors and eigenvalues for A . Let us ask Wolfram Alpha: [det\(A\)](#), [Eigenvalues and eigenvectors](#). So $\lambda = 6, 3, 1$, with eigenvectors

$$v_1 = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \quad v_2 = \frac{1}{\sqrt{6}} \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}, \quad \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Then $q(x)$ has maximum value 6, and $q(v_1) = v_1^T A v_1 = 6 \|v_1\|^2 = 6$.

Example

Let $A = \begin{bmatrix} 3 & 2 & 1 \\ 2 & 3 & 1 \\ 1 & 1 & 4 \end{bmatrix}$. Find the maximum and minimum values of $q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x}$ subject to the constraint $\mathbf{x}^T \mathbf{x} = 1$.

Solution

We first find eigenvectors and eigenvalues for A . Let us ask Wolfram Alpha: [det\(A\)](#), [Eigenvalues and eigenvectors](#). So $\lambda = 6, 3, 1$, with eigenvectors

$$\mathbf{v}_1 = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \quad \mathbf{v}_2 = \frac{1}{\sqrt{6}} \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}, \quad \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Then $q(\mathbf{x})$ has maximum value 6, and

$q(\mathbf{v}_1) = \mathbf{v}_1^T A \mathbf{v}_1 = 6 \|\mathbf{v}_1\|^2 = 6$. The minimal value is 1 and

$q(\mathbf{v}_3) = \mathbf{v}_3^T A \mathbf{v}_3 = 1 \|\mathbf{v}_3\|^2 = 1$.