

Math 415 - Lecture 31

Markov matrices and Google

Monday November 9th 2015

Textbook reading: Chapter 5.3

Textbook reading: Chapter 5.3

Suggested practice exercises: Chapter 5.3: 8, 9, 12, 14, 10.

Textbook reading: Chapter 5.3

Suggested practice exercises: Chapter 5.3: 8, 9, 12, 14, 10.

Khan Academy video: Finding Eigenvectors and Eigenspaces
example

Textbook reading: Chapter 5.3

Suggested practice exercises: Chapter 5.3: 8, 9, 12, 14, 10.

Khan Academy video: Finding Eigenvectors and Eigenspaces
example

Strang lecture: Lecture 21: Eigenvalues and eigenvectors Lecture
24: Markov Matrices and Fourier Series.

SOME DATES.

SOME DATES.

SOME DATES.

- * Friday November 13th: No class.

SOME DATES.

- * Friday November 13th: No class.

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.
- * Friday November 20th: No class.

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.
- * Friday November 20th: No class.

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.
- * Friday November 20th: No class.
- * November 23-27th Thanksgiving break: no class.

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.
- * Friday November 20th: No class.
- * November 23-27th Thanksgiving break: no class.

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.
- * Friday November 20th: No class.
- * November 23-27th Thanksgiving break: no class.
- * Wednesday December 9th: last day of class

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.
- * Friday November 20th: No class.
- * November 23-27th Thanksgiving break: no class.
- * Wednesday December 9th: last day of class

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.
- * Friday November 20th: No class.
- * November 23-27th Thanksgiving break: no class.
- * Wednesday December 9th: last day of class
- * Thursday December 17th: Final Exam.

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.
- * Friday November 20th: No class.
- * November 23-27th Thanksgiving break: no class.
- * Wednesday December 9th: last day of class
- * Thursday December 17th: Final Exam.

SOME DATES.

- * Friday November 13th: No class.
- * Next week there will be discussion sections, but no quiz. Prepare for the midterm!
- * Thursday November 19th, 7-8:15PM: Midterm 3.
- * Friday November 20th: No class.
- * November 23-27th Thanksgiving break: no class.
- * Wednesday December 9th: last day of class
- * Thursday December 17th: Final Exam.

Review

Properties of eigenvectors and eigenvalues

- If $A\mathbf{x} = \lambda\mathbf{x}$ then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .

- If $A\mathbf{x} = \lambda\mathbf{x}$ then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .

- If $A\mathbf{x} = \lambda\mathbf{x}$ then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
All eigenvectors (plus $\mathbf{0}$) with eigenvalue λ form **eigenspace** of λ .

- If $A\mathbf{x} = \lambda\mathbf{x}$ then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
All eigenvectors (plus $\mathbf{0}$) with eigenvalue λ form **eigenspace** of λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

- If $A\mathbf{x} = \lambda\mathbf{x}$ then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
All eigenvectors (plus $\mathbf{0}$) with eigenvalue λ form **eigenspace** of λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

- If $A\mathbf{x} = \lambda\mathbf{x}$ then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
All eigenvectors (plus $\mathbf{0}$) with eigenvalue λ form **eigenspace** of λ .
- λ is an eigenvalue of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

- If $A\mathbf{x} = \lambda\mathbf{x}$ then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
All eigenvectors (plus $\mathbf{0}$) with eigenvalue λ form **eigenspace** of λ .
- λ is an eigenvalue of $A \iff \det(A - \lambda I) = 0$.

characteristic polynomial

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

By the way: this means that the eigenspace of λ is just $\text{Nul}(A - \lambda I)$.

- If $A\mathbf{x} = \lambda\mathbf{x}$ then \mathbf{x} is an **eigenvector** of A with **eigenvalue** λ .
All eigenvectors (plus $\mathbf{0}$) with eigenvalue λ form **eigenspace** of λ .
- λ is an eigenvalue of $A \iff \det(A - \lambda I) = 0$.

characteristic polynomial

Why? Because $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$.

By the way: this means that the eigenspace of λ is just $\text{Nul}(A - \lambda I)$.

- E.g. if $A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$ then
 $\det(A - \lambda I) = (3 - \lambda)(6 - \lambda)(2 - \lambda)$.

If $A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$ then the eigenvalues are 2, 3, 6

If $A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$ then the eigenvalues are 2, 3, 6 with
corresponding eigenvectors

$$\begin{bmatrix} 2 \\ -5/2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2/3 \\ 1 \\ 0 \end{bmatrix}.$$

If $A = \begin{bmatrix} 3 & 2 & 3 \\ 0 & 6 & 10 \\ 0 & 0 & 2 \end{bmatrix}$ then the eigenvalues are 2, 3, 6 with corresponding eigenvectors

$$\begin{bmatrix} 2 \\ -5/2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2/3 \\ 1 \\ 0 \end{bmatrix}.$$

These three vectors are independent. By the next result, this is always so.

◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻

If $\mathbf{x}_1, \dots, \mathbf{x}_m$ are eigenvectors of A corresponding to different eigenvalues, then they are independent.

Theorem

If $\mathbf{x}_1, \dots, \mathbf{x}_m$ are eigenvectors of A corresponding to different eigenvalues, then they are independent.

Proof.

Suppose, for contradiction, that $\mathbf{x}_1, \dots, \mathbf{x}_m$ are dependent.

Theorem

If $\mathbf{x}_1, \dots, \mathbf{x}_m$ are eigenvectors of A corresponding to different eigenvalues, then they are independent.

Proof.

Suppose, for contradiction, that $\mathbf{x}_1, \dots, \mathbf{x}_m$ are dependent.

By kicking out some vectors, we may assume that there is (up to multiples) only one linear relation: $c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m = \mathbf{0}$.

Theorem

If $\mathbf{x}_1, \dots, \mathbf{x}_m$ are eigenvectors of A corresponding to different eigenvalues, then they are independent.

Proof.

Suppose, for contradiction, that $\mathbf{x}_1, \dots, \mathbf{x}_m$ are dependent.

By kicking out some vectors, we may assume that there is (up to multiples) only one linear relation: $c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m = \mathbf{0}$.

In other words, the matrix with columns $\mathbf{x}_1, \dots, \mathbf{x}_m$ has one-dimensional null space.

Theorem

If $\mathbf{x}_1, \dots, \mathbf{x}_m$ are eigenvectors of A corresponding to different eigenvalues, then they are independent.

Proof.

Suppose, for contradiction, that $\mathbf{x}_1, \dots, \mathbf{x}_m$ are dependent.

By kicking out some vectors, we may assume that there is (up to multiples) only one linear relation: $c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m = \mathbf{0}$.

In other words, the matrix with columns $\mathbf{x}_1, \dots, \mathbf{x}_m$ has one-dimensional null space. Now multiply this relation with A :

$$A(c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m) = c_1\lambda_1\mathbf{x}_1 + \dots + c_m\lambda_m\mathbf{x}_m = \mathbf{0}$$

This is a second independent relation!

Theorem

If $\mathbf{x}_1, \dots, \mathbf{x}_m$ are eigenvectors of A corresponding to different eigenvalues, then they are independent.

Proof.

Suppose, for contradiction, that $\mathbf{x}_1, \dots, \mathbf{x}_m$ are dependent.

By kicking out some vectors, we may assume that there is (up to multiples) only one linear relation: $c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m = \mathbf{0}$.

In other words, the matrix with columns $\mathbf{x}_1, \dots, \mathbf{x}_m$ has one-dimensional null space. Now multiply this relation with A :

$$A(c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m) = c_1\lambda_1\mathbf{x}_1 + \dots + c_m\lambda_m\mathbf{x}_m = \mathbf{0}$$

This is a second independent relation! In other words, two independent vectors living in a one-dimensional vector space.

Theorem

If $\mathbf{x}_1, \dots, \mathbf{x}_m$ are eigenvectors of A corresponding to different eigenvalues, then they are independent.

Proof.

Suppose, for contradiction, that $\mathbf{x}_1, \dots, \mathbf{x}_m$ are dependent.

By kicking out some vectors, we may assume that there is (up to multiples) only one linear relation: $c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m = \mathbf{0}$.

In other words, the matrix with columns $\mathbf{x}_1, \dots, \mathbf{x}_m$ has one-dimensional null space. Now multiply this relation with A :

$$A(c_1\mathbf{x}_1 + \dots + c_m\mathbf{x}_m) = c_1\lambda_1\mathbf{x}_1 + \dots + c_m\lambda_m\mathbf{x}_m = \mathbf{0}$$

This is a second independent relation! In other words, two independent vectors living in a one-dimensional vector space.

Contradiction. □

Product of Eigenvalues

If A is $n \times n$ get in principle n eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$.

If A is $n \times n$ get in principle n eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. How are these eigenvalues related?

If A is $n \times n$ get in principle n eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. How are these eigenvalues related?

Theorem

The product of eigenvalues $\lambda_1 \lambda_2 \dots \lambda_n$ is equal to the determinant of A .

If A is $n \times n$ get in principle n eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. How are these eigenvalues related?

Theorem

The product of eigenvalues $\lambda_1 \lambda_2 \dots \lambda_n$ is equal to the determinant of A .

Proof.

The characteristic polynomial $p(\lambda) = \det(A - \lambda I)$ has constant term $\det(A)$.

If A is $n \times n$ get in principle n eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. How are these eigenvalues related?

Theorem

The product of eigenvalues $\lambda_1 \lambda_2 \dots \lambda_n$ is equal to the determinant of A .

Proof.

The characteristic polynomial $p(\lambda) = \det(A - \lambda I)$ has constant term $\det(A)$. On the other hand $p(\lambda)$ factors, because the roots are the eigenvalues we get $p(\lambda) = (\pm 1)(\lambda - \lambda_1)(\lambda - \lambda_2) \dots (\lambda - \lambda_n)$, which has constant term $\lambda_1 \lambda_2 \dots \lambda_n$. \square

If A is $n \times n$ get in principle n eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. How are these eigenvalues related?

Theorem

The product of eigenvalues $\lambda_1 \lambda_2 \dots \lambda_n$ is equal to the determinant of A .

Proof.

The characteristic polynomial $p(\lambda) = \det(A - \lambda I)$ has constant term $\det(A)$. On the other hand $p(\lambda)$ factors, because the roots are the eigenvalues we get $p(\lambda) = (\pm 1)(\lambda - \lambda_1)(\lambda - \lambda_2) \dots (\lambda - \lambda_n)$, which has constant term $\lambda_1 \lambda_2 \dots \lambda_n$. \square

Example

Let $A = \begin{bmatrix} \lambda_1 & b \\ 0 & \lambda_2 \end{bmatrix}$. Then the eigenvalues are λ_1, λ_2 and $\det(A) = \lambda_1 \lambda_2$.

Sum of Eigenvalues

What other relations are there between the eigenvalues?

What other relations are there between the eigenvalues?

Definition

Let $A = \begin{bmatrix} a_{11} & a_{12} & \cdots \\ a_{21} & a_{22} & \cdots \\ \vdots & \vdots & \ddots \end{bmatrix}$ be $n \times n$. Then the **TRACE** of A is the sum of the diagonal entries: $\text{Tr}(A) = a_{11} + a_{22} + \cdots + a_{nn}$.

What other relations are there between the eigenvalues?

Definition

Let $A = \begin{bmatrix} a_{11} & a_{12} & \cdots \\ a_{21} & a_{22} & \cdots \\ \vdots & \vdots & \ddots \end{bmatrix}$ be $n \times n$. Then the **TRACE** of A is the sum of the diagonal entries: $\text{Tr}(A) = a_{11} + a_{22} + \cdots + a_{nn}$.

Theorem

*Let A be $n \times n$. Then the trace of A is the **sum** of eigenvalues:*

$$\text{Tr}(A) = \lambda_1 + \lambda_2 + \cdots + \lambda_n$$

Example

Let $A = \begin{bmatrix} \lambda_1 & b \\ 0 & \lambda_2 \end{bmatrix}$. What are the eigenvalues and what is $\text{Tr}(A)$?

Example

Let $A = \begin{bmatrix} \lambda_1 & b \\ 0 & \lambda_2 \end{bmatrix}$. What are the eigenvalues and what is $\text{Tr}(A)$?

Solution

The eigenvalues are λ_1, λ_2 and $\text{Tr}(A) = \lambda_1 + \lambda_2$.

◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻

2×2 matrices are easy.

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$$\text{Tr}(A) =$$

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$$\text{Tr}(A) = 6,$$

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$$\text{Tr}(A) = 6, \det(A) =$$

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$$\text{Tr}(A) = 6, \det(A) = 8,$$

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$\text{Tr}(A) = 6$, $\det(A) = 8$, so $p(\lambda) =$

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$\text{Tr}(A) = 6$, $\det(A) = 8$, so $p(\lambda) = \lambda^2 - 6\lambda + 8$.

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$\text{Tr}(A) = 6$, $\det(A) = 8$, so $p(\lambda) = \lambda^2 - 6\lambda + 8$. Also in terms of eigenvalues $\text{Tr}(A) =$

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$\text{Tr}(A) = 6$, $\det(A) = 8$, so $p(\lambda) = \lambda^2 - 6\lambda + 8$. Also in terms of eigenvalues $\text{Tr}(A) = \lambda_1 + \lambda_2$

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$\text{Tr}(A) = 6$, $\det(A) = 8$, so $p(\lambda) = \lambda^2 - 6\lambda + 8$. Also in terms of eigenvalues $\text{Tr}(A) = \lambda_1 + \lambda_2$ and $\det(A) =$

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$\text{Tr}(A) = 6$, $\det(A) = 8$, so $p(\lambda) = \lambda^2 - 6\lambda + 8$. Also in terms of eigenvalues $\text{Tr}(A) = \lambda_1 + \lambda_2$ and $\det(A) = \lambda_1\lambda_2$.

The Characteristic Polynomial for 2×2

2×2 matrices are easy.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the characteristic polynomial is

$$p(\lambda) = \lambda^2 - (a + d)\lambda + ad - bc = \lambda^2 - \text{Tr}(A)\lambda + \det(A).$$

Example

Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$. What are the eigenvalues and what is the characteristic polynomial?

Solution

$\text{Tr}(A) = 6$, $\det(A) = 8$, so $p(\lambda) = \lambda^2 - 6\lambda + 8$. Also in terms of eigenvalues $\text{Tr}(A) = \lambda_1 + \lambda_2$ and $\det(A) = \lambda_1\lambda_2$. So $\lambda_1 = 2, \lambda_2 = 4$

Practice problems

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$.

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 0 & -2 \\ -4 & 2 \end{bmatrix}$.

Example

What are the eigenvalues of $A = \begin{bmatrix} 2 & 0 & 0 & 0 \\ -1 & 3 & 0 & 0 \\ -1 & 1 & 3 & 0 \\ 0 & 1 & 2 & 4 \end{bmatrix}$.

No calculations!

Example

Find the eigenvalues of A as well as a basis for the corresponding eigenspaces, where

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}.$$

Solution

- The characteristic polynomial is:

Example

Find the eigenvalues of A as well as a basis for the corresponding eigenspaces, where

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}.$$

Solution

- The characteristic polynomial is:

Example

Find the eigenvalues of A as well as a basis for the corresponding eigenspaces, where

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}.$$

Solution

- The characteristic polynomial is:

Example

Find the eigenvalues of A as well as a basis for the corresponding eigenspaces, where

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}.$$

Solution

- The characteristic polynomial is:

$$\det(A - \lambda I) = \begin{vmatrix} 2 - \lambda & 0 & 0 \\ -1 & 3 - \lambda & 1 \\ -1 & 1 & 3 - \lambda \end{vmatrix}$$

Example

Find the eigenvalues of A as well as a basis for the corresponding eigenspaces, where

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}.$$

Solution

- The characteristic polynomial is:

$$\begin{aligned} \det(A - \lambda I) &= \begin{vmatrix} 2 - \lambda & 0 & 0 \\ -1 & 3 - \lambda & 1 \\ -1 & 1 & 3 - \lambda \end{vmatrix} \\ &= (2 - \lambda) \begin{vmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{vmatrix} \end{aligned}$$

Example

Find the eigenvalues of A as well as a basis for the corresponding eigenspaces, where

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}.$$

Solution

- The characteristic polynomial is:

$$\begin{aligned} \det(A - \lambda I) &= \begin{vmatrix} 2 - \lambda & 0 & 0 \\ -1 & 3 - \lambda & 1 \\ -1 & 1 & 3 - \lambda \end{vmatrix} \\ &= (2 - \lambda) \begin{vmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{vmatrix} \\ &= (2 - \lambda)[(3 - \lambda)^2 - 1] \end{aligned}$$

Example

Find the eigenvalues of A as well as a basis for the corresponding eigenspaces, where

$$A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}.$$

Solution

- The characteristic polynomial is:

$$\begin{aligned} \det(A - \lambda I) &= \begin{vmatrix} 2 - \lambda & 0 & 0 \\ -1 & 3 - \lambda & 1 \\ -1 & 1 & 3 - \lambda \end{vmatrix} \\ &= (2 - \lambda) \begin{vmatrix} 3 - \lambda & 1 \\ 1 & 3 - \lambda \end{vmatrix} \\ &= (2 - \lambda)[(3 - \lambda)^2 - 1] \\ &= (2 - \lambda)(\lambda - 2)(\lambda - 4) \end{aligned}$$

Solution (continued)

- A has eigenvalues 2, 2, 4 $\left(A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \right)$

Solution (continued)

- A has eigenvalues 2, 2, 4 $\left(A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \right)$

Since $\lambda = 2$ is a double root, it has **(algebraic) multiplicity 2**.

Solution (continued)

- A has eigenvalues 2, 2, 4 $\left(A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \right)$

Since $\lambda = 2$ is a double root, it has **(algebraic) multiplicity 2**.

- $\lambda_1 = 2$:

Solution (continued)

- A has eigenvalues 2, 2, 4 $\left(A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \right)$

Since $\lambda = 2$ is a double root, it has **(algebraic) multiplicity 2**.

- $\lambda_1 = 2$:

$$(A - \lambda_1 I)\mathbf{x} = \begin{bmatrix} 0 & 0 & 0 \\ -1 & 1 & 1 \\ -1 & 1 & 1 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

Solution (continued)

- A has eigenvalues 2, 2, 4 $\left(A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \right)$

Since $\lambda = 2$ is a double root, it has **(algebraic) multiplicity 2**.

- $\lambda_1 = 2$:

$$(A - \lambda_1 I)\mathbf{x} = \begin{bmatrix} 0 & 0 & 0 \\ -1 & 1 & 1 \\ -1 & 1 & 1 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

Two independent solutions: $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$, $\mathbf{x}_2 = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$

Solution (continued)

- A has eigenvalues 2, 2, 4 $\left(A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \right)$

Since $\lambda = 2$ is a double root, it has **(algebraic) multiplicity 2**.

- $\lambda_1 = 2$:

$$(A - \lambda_1 I)\mathbf{x} = \begin{bmatrix} 0 & 0 & 0 \\ -1 & 1 & 1 \\ -1 & 1 & 1 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

Two independent solutions: $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$, $\mathbf{x}_2 = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$

In other words, the eigenspace for $\lambda_1 = 2$ is

Solution (continued)

- A has eigenvalues 2, 2, 4 $\left(A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \right)$

Since $\lambda = 2$ is a double root, it has **(algebraic) multiplicity 2**.

- $\lambda_1 = 2$:

$$(A - \lambda_1 I)\mathbf{x} = \begin{bmatrix} 0 & 0 & 0 \\ -1 & 1 & 1 \\ -1 & 1 & 1 \end{bmatrix} \mathbf{x} = \mathbf{0}$$

Two independent solutions: $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$, $\mathbf{x}_2 = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$

In other words, the eigenspace for $\lambda_1 = 2$ is

$$\text{Span} \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \right\}.$$

Solution (continued)

- $\lambda_2 = 4$:

Solution (continued)

- $\lambda_2 = 4$: $\left(A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \right)$

$$(A - \lambda_2 I)\mathbf{x} = \begin{bmatrix} -2 & 0 & 0 \\ -1 & -1 & 1 \\ -1 & 1 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0} \Rightarrow$$

Solution (continued)

- $\lambda_2 = 4$: $\left(A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \right)$

$$(A - \lambda_2 I)\mathbf{x} = \begin{bmatrix} -2 & 0 & 0 \\ -1 & -1 & 1 \\ -1 & 1 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_3 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

Solution (continued)

- $\lambda_2 = 4$:
$$\left(A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix} \right)$$

$$(A - \lambda_2 I)\mathbf{x} = \begin{bmatrix} -2 & 0 & 0 \\ -1 & -1 & 1 \\ -1 & 1 & -1 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_3 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

- In summary, A has eigenvalues 2 and 4:

- eigenspace for $\lambda_1 = 2$ has basis $\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix},$
- eigenspace for $\lambda_2 = 4$ has basis $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$

Markov matrices

Definition

An $n \times n$ matrix A is **Markov matrix** if it has non negative entries, and the entries in each column add to 1.

Definition

An $n \times n$ matrices A is **Markov matrix** if has non negative entries, and the entries in each column add to 1.

Theorem

Let A be a Markov matrix. Then

- (i) *1 is an eigenvalue of A and any other eigenvalue λ satisfies $|\lambda| \leq 1$.*

Definition

An $n \times n$ matrices A is **Markov matrix** if has non negative entries, and the entries in each column add to 1.

Theorem

Let A be a Markov matrix. Then

- (i) *1 is an eigenvalue of A and any other eigenvalue λ satisfies $|\lambda| \leq 1$.*
- (ii) *If A has only positive entries, then any other eigenvalue satisfies $|\lambda| < 1$.*

Definition

An $n \times n$ matrices A is **Markov matrix** if has non negative entries, and the entries in each column add to 1.

Theorem

Let A be a Markov matrix. Then

- (i) 1 is an eigenvalue of A and any other eigenvalue λ satisfies $|\lambda| \leq 1$.
- (ii) If A has only positive entries, then any other eigenvalue satisfies $|\lambda| < 1$.

Example

Let A be

$$\begin{bmatrix} .9 & .2 \\ .1 & .8 \end{bmatrix}.$$

Is A a Markov matrix?

Theorem

Let A be an $n \times n$ -Markov matrix with only positive entries and let $\mathbf{v} \in \mathbb{R}^n$. Then

$$\mathbf{v}_\infty := \lim_{k \rightarrow \infty} A^k \mathbf{v} \text{ exists,}$$

and $A\mathbf{v}_\infty = \mathbf{v}_\infty$. In this case \mathbf{v}_∞ is often called the **steady state**.

Theorem

Let A be an $n \times n$ -Markov matrix with only positive entries and let $\mathbf{v} \in \mathbb{R}^n$. Then

$$\mathbf{v}_\infty := \lim_{k \rightarrow \infty} A^k \mathbf{v} \text{ exists,}$$

and $A\mathbf{v}_\infty = \mathbf{v}_\infty$. In this case \mathbf{v}_∞ is often called the **steady state**.

Proof.

If \mathbf{x} is any vector and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is an eigenbasis for a Markov matrix ($A\mathbf{v}_1 = 1\mathbf{v}_1$):

Theorem

Let A be an $n \times n$ -Markov matrix with only positive entries and let $\mathbf{v} \in \mathbb{R}^n$. Then

$$\mathbf{v}_\infty := \lim_{k \rightarrow \infty} A^k \mathbf{v} \text{ exists,}$$

and $A\mathbf{v}_\infty = \mathbf{v}_\infty$. In this case \mathbf{v}_∞ is often called the **steady state**.

Proof.

If \mathbf{x} is any vector and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is an eigenbasis for a Markov matrix ($A\mathbf{v}_1 = 1\mathbf{v}_1$):

$$\mathbf{x} = c_1 \mathbf{v}_1 + \dots + c_n \mathbf{v}_n,$$

then

Theorem

Let A be an $n \times n$ -Markov matrix with only positive entries and let $\mathbf{v} \in \mathbb{R}^n$. Then

$$\mathbf{v}_\infty := \lim_{k \rightarrow \infty} A^k \mathbf{v} \text{ exists,}$$

and $A\mathbf{v}_\infty = \mathbf{v}_\infty$. In this case \mathbf{v}_∞ is often called the **steady state**.

Proof.

If \mathbf{x} is any vector and $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ is an eigenbasis for a Markov matrix ($A\mathbf{v}_1 = 1\mathbf{v}_1$):

$$\mathbf{x} = c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n,$$

then

$$A^k \mathbf{x} = c_1 \lambda_1^k \mathbf{v}_1 + \dots + c_n \lambda_n^k \mathbf{v}_n \rightarrow c_1 \mathbf{v}_1,$$

if the eigenspace of $\lambda = 1$ is 1-dimensional.



Example

Consider a fixed population of people with or without job.

Example

Consider a fixed population of people with or without job. Suppose that each year, 50% of those unemployed find a job while 10% of those employed lose their job.

Example

Consider a fixed population of people with or without job. Suppose that each year, 50% of those unemployed find a job while 10% of those employed lose their job.


What is the unemployment rate in the long term equilibrium?

Example

Consider a fixed population of people with or without job. Suppose that each year, 50% of those unemployed find a job while 10% of those employed lose their job.

What is the unemployment rate in the long term equilibrium?

Solution



Example

Consider a fixed population of people with or without job. Suppose that each year, 50% of those unemployed find a job while 10% of those employed lose their job.

What is the unemployment rate in the long term equilibrium?

Solution



x_t : proportion of population employed at time t (in years)


y_t : proportion of population unemployed at time t

Example

Consider a fixed population of people with or without job. Suppose that each year, 50% of those unemployed find a job while 10% of those employed lose their job.

What is the unemployment rate in the long term equilibrium?

Solution



x_t : proportion of population employed at time t (in years)

y_t : proportion of population unemployed at time t

$$\begin{bmatrix} x_{t+1} \\ y_{t+1} \end{bmatrix} = \begin{bmatrix} 0.9x_t + 0.5y_t \\ 0.1x_t + 0.5y_t \end{bmatrix} = \begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix} \begin{bmatrix} x_t \\ y_t \end{bmatrix}$$

Solution (continued)

The matrix $\begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix}$ is a **Markov matrix**. Its columns add to 1 and it has no negative entries.

Solution (continued)

The matrix $\begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix}$ is a **Markov matrix**. Its columns add to 1 and it has no negative entries.

$$\begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix} \text{ is an equilibrium if } \begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix} = \begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix} \begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix}.$$

Solution (continued)

The matrix $\begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix}$ is a **Markov matrix**. Its columns add to 1 and it has no negative entries.

$$\begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix} \text{ is an equilibrium if } \begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix} = \begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix} \begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix}.$$

In other words, $\begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix}$ is an eigenvector with eigenvalue 1.

Solution (continued)

The matrix $\begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix}$ is a **Markov matrix**. Its columns add to 1 and it has no negative entries.

$$\begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix} \text{ is an equilibrium if } \begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix} = \begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix} \begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix}.$$

In other words, $\begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix}$ is an eigenvector with eigenvalue 1.

$$\text{Eigenspace of } \lambda = 1 : \text{Nul} \left(\begin{bmatrix} -0.1 & 0.5 \\ 0.1 & -0.5 \end{bmatrix} \right) = \text{Span} \left\{ \begin{bmatrix} 5 \\ 1 \end{bmatrix} \right\}$$

Solution (continued)

The matrix $\begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix}$ is a **Markov matrix**. Its columns add to 1 and it has no negative entries.

$$\begin{bmatrix} x_\infty \\ y_\infty \end{bmatrix} \text{ is an equilibrium if } \begin{bmatrix} x_\infty \\ y_\infty \end{bmatrix} = \begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix} \begin{bmatrix} x_\infty \\ y_\infty \end{bmatrix}.$$

In other words, $\begin{bmatrix} x_\infty \\ y_\infty \end{bmatrix}$ is an eigenvector with eigenvalue 1.

$$\text{Eigenspace of } \lambda = 1 : \text{Nul} \left(\begin{bmatrix} -0.1 & 0.5 \\ 0.1 & -0.5 \end{bmatrix} \right) = \text{Span} \left\{ \begin{bmatrix} 5 \\ 1 \end{bmatrix} \right\}$$

$$\text{Since } x_\infty + y_\infty = 1, \text{ we conclude that } \begin{bmatrix} x_\infty \\ y_\infty \end{bmatrix} = \begin{bmatrix} 5/6 \\ 1/6 \end{bmatrix}.$$

Solution (continued)

The matrix $\begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix}$ is a **Markov matrix**. Its columns add to 1 and it has no negative entries.

$$\begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix} \text{ is an equilibrium if } \begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix} = \begin{bmatrix} 0.9 & 0.5 \\ 0.1 & 0.5 \end{bmatrix} \begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix}.$$

In other words, $\begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix}$ is an eigenvector with eigenvalue 1.

$$\text{Eigenspace of } \lambda = 1 : \text{Nul} \left(\begin{bmatrix} -0.1 & 0.5 \\ 0.1 & -0.5 \end{bmatrix} \right) = \text{Span} \left\{ \begin{bmatrix} 5 \\ 1 \end{bmatrix} \right\}$$

Since $x_{\infty} + y_{\infty} = 1$, we conclude that $\begin{bmatrix} x_{\infty} \\ y_{\infty} \end{bmatrix} = \begin{bmatrix} 5/6 \\ 1/6 \end{bmatrix}$.

Hence, the unemployment rate in the long term equilibrium is $1/6$


Page rank (or: the 25000000000 \$ eigenvector)

Google's success is based on an algorithm to rank webpages, the **Page rank**, named after Google founder Larry Page. The idea is to determine how likely it is that a web user randomly gets to a given webpage. The webpages are ranked by these probabilities.


Google's success is based on an algorithm to rank webpages, the **Page rank**, named after Google founder Larry Page. The idea is to determine how likely it is that a web user randomly gets to a given webpage. The webpages are ranked by these probabilities. Suppose the internet consisted of the only four webpages A, B, C, D linked as in the following graph.

Google's success is based on an algorithm to rank webpages, the **Page rank**, named after Google founder Larry Page. The idea is to determine how likely it is that a web user randomly gets to a given webpage. The webpages are ranked by these probabilities. Suppose the internet consisted of the only four webpages A, B, C, D linked as in the following graph.

Imagine a surfer following these links at random. For the probability $PR_n(A)$ that she is at A (after n steps), we need to think about how she could have reached A . We add:




Google's success is based on an algorithm to rank webpages, the **Page rank**, named after Google founder Larry Page. The idea is to determine how likely it is that a web user randomly gets to a given webpage. The webpages are ranked by these probabilities. Suppose the internet consisted of the only four webpages A, B, C, D linked as in the following graph.



Imagine a surfer following these links at random. For the probability $PR_n(A)$ that she is at A (after n steps), we need to think about how she could have reached A . We add:

- the probability that she was at B (at exactly one step before), and left for A , (that's $PR_{n-1}(B) \cdot \frac{1}{2}$)


Google's success is based on an algorithm to rank webpages, the **Page rank**, named after Google founder Larry Page. The idea is to determine how likely it is that a web user randomly gets to a given webpage. The webpages are ranked by these probabilities. Suppose the internet consisted of the only four webpages A, B, C, D linked as in the following graph.



Imagine a surfer following these links at random. For the probability $PR_n(A)$ that she is at A (after n steps), we need to think about how she could have reached A . We add:

- the probability that she was at B (at exactly one step before), and left for A , (that's $PR_{n-1}(B) \cdot \frac{1}{2}$)
- the probability that she was at C , and left for A ,


Google's success is based on an algorithm to rank webpages, the **Page rank**, named after Google founder Larry Page. The idea is to determine how likely it is that a web user randomly gets to a given webpage. The webpages are ranked by these probabilities. Suppose the internet consisted of the only four webpages A, B, C, D linked as in the following graph.



Imagine a surfer following these links at random. For the probability $PR_n(A)$ that she is at A (after n steps), we need to think about how she could have reached A . We add:

- the probability that she was at B (at exactly one step before), and left for A , (that's $PR_{n-1}(B) \cdot \frac{1}{2}$)
- the probability that she was at C , and left for A ,
- the probability that she was at D , and left for A .

Google's success is based on an algorithm to rank webpages, the **Page rank**, named after Google founder Larry Page. The idea is to determine how likely it is that a web user randomly gets to a given webpage. The webpages are ranked by these probabilities. Suppose the internet consisted of the only four webpages A, B, C, D linked as in the following graph.



Imagine a surfer following these links at random. For the probability $PR_n(A)$ that she is at A (after n steps), we need to think about how she could have reached A . We add:

- the probability that she was at B (at exactly one step before), and left for A , (that's $PR_{n-1}(B) \cdot \frac{1}{2}$)
- the probability that she was at C , and left for A ,
- the probability that she was at D , and left for A .

$$\text{Hence: } PR_n(A) = PR_{n-1}(B) \cdot \frac{1}{2} + PR_{n-1}(C) \cdot \frac{1}{1} + PR_{n-1}(D) \cdot \frac{0}{1}.$$

Hence: $PR_n(A) = PR_{n-1}(B) \cdot \frac{1}{2} + PR_{n-1}(C) \cdot \frac{1}{1} + PR_{n-1}(D) \cdot \frac{0}{1}$.

Encode the probabilities at step n in a state vector with four

entries.

$$\begin{bmatrix} PR_n(A) \\ PR_n(B) \\ PR_n(C) \\ PR_n(D) \end{bmatrix} = \begin{bmatrix} 0 & \frac{1}{2} & 1 & 0 \\ \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{3} & 0 & 0 & 1 \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 \end{bmatrix} \begin{bmatrix} PR_{n-1}(A) \\ PR_{n-1}(B) \\ PR_{n-1}(C) \\ PR_{n-1}(D) \end{bmatrix}$$

Hence: $PR_n(A) = PR_{n-1}(B) \cdot \frac{1}{2} + PR_{n-1}(C) \cdot \frac{1}{1} + PR_{n-1}(D) \cdot \frac{0}{1}$.

Encode the probabilities at step n in a state vector with four

entries.

$$\begin{bmatrix} PR_n(A) \\ PR_n(B) \\ PR_n(C) \\ PR_n(D) \end{bmatrix} = \begin{bmatrix} 0 & \frac{1}{2} & 1 & 0 \\ \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{3} & 0 & 0 & 1 \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 \end{bmatrix} \begin{bmatrix} PR_{n-1}(A) \\ PR_{n-1}(B) \\ PR_{n-1}(C) \\ PR_{n-1}(D) \end{bmatrix}$$

Definition

The **PageRank vector** is the long-term equilibrium.

It is an eigenvector of the Markov matrix with eigenvalue 1.

Example (continued)

Let's call the Markov matrix with the probabilities T :

Example (continued)

Let's call the Markov matrix with the probabilities T :

Example (continued)

Let's call the Markov matrix with the probabilities T :

$$\bullet \quad T - I = \begin{bmatrix} -1 & \frac{1}{2} & 1 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & \frac{1}{2} & 0 & -1 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -\frac{2}{3} \\ 0 & 0 & 1 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Example (continued)

Let's call the Markov matrix with the probabilities T :

$$\bullet T - I = \begin{bmatrix} -1 & \frac{1}{2} & 1 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & \frac{1}{2} & 0 & -1 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -\frac{2}{3} \\ 0 & 0 & 1 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Example (continued)

Let's call the Markov matrix with the probabilities T :

$$\bullet \quad T - I = \begin{bmatrix} -1 & \frac{1}{2} & 1 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & \frac{1}{2} & 0 & -1 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -\frac{2}{3} \\ 0 & 0 & 1 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

\implies eigenspace of $\lambda = 1$ is spanned by $\begin{bmatrix} 2 \\ 2 \\ 5 \\ 3 \\ 1 \end{bmatrix}$.

Example (continued)

Let's call the Markov matrix with the probabilities T :

$$\bullet \quad T - I = \begin{bmatrix} -1 & \frac{1}{2} & 1 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & \frac{1}{2} & 0 & -1 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -\frac{2}{3} \\ 0 & 0 & 1 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

\implies eigenspace of $\lambda = 1$ is spanned by $\begin{bmatrix} 2 \\ 2 \\ 5 \\ 3 \\ 1 \end{bmatrix}$.

- Now we need to make the entries add up to 1.

Example (continued)

Let's call the Markov matrix with the probabilities T :

$$\bullet \quad T - I = \begin{bmatrix} -1 & \frac{1}{2} & 1 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & \frac{1}{2} & 0 & -1 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -\frac{2}{3} \\ 0 & 0 & 1 & -\frac{5}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

\implies eigenspace of $\lambda = 1$ is spanned by $\begin{bmatrix} 2 \\ 2 \\ 5 \\ 3 \\ 1 \end{bmatrix}$.

- Now we need to make the entries add up to 1.

Example (continued)

Let's call the Markov matrix with the probabilities T :

$$\bullet \quad T - I = \begin{bmatrix} -1 & \frac{1}{2} & 1 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & \frac{1}{2} & 0 & -1 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -\frac{2}{3} \\ 0 & 0 & 1 & -\frac{2}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

\implies eigenspace of $\lambda = 1$ is spanned by $\begin{bmatrix} 2 \\ 2 \\ 3 \\ 3 \\ 1 \end{bmatrix}$.

- Now we need to make the entries add up to 1.

$$\begin{bmatrix} PR(A) \\ PR(B) \\ PR(C) \\ PR(D) \end{bmatrix} = \frac{3}{16} \begin{bmatrix} 2 \\ 2 \\ 3 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.125 \\ 0.313 \\ 0.188 \end{bmatrix}.$$

Example (continued)

Let's call the Markov matrix with the probabilities T :

$$\bullet \quad T - I = \begin{bmatrix} -1 & \frac{1}{2} & 1 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & \frac{1}{2} & 0 & -1 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -\frac{2}{3} \\ 0 & 0 & 1 & -\frac{2}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

\Rightarrow eigenspace of $\lambda = 1$ is spanned by $\begin{bmatrix} 2 \\ 2 \\ 3 \\ 3 \\ 1 \end{bmatrix}$.

- Now we need to make the entries add up to 1.

$$\begin{bmatrix} PR(A) \\ PR(B) \\ PR(C) \\ PR(D) \end{bmatrix} = \frac{3}{16} \begin{bmatrix} 2 \\ 2 \\ 3 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.125 \\ 0.313 \\ 0.188 \end{bmatrix}.$$

This is the PageRank vector.

Example (continued)

Let's call the Markov matrix with the probabilities T :

$$\bullet \quad T - I = \begin{bmatrix} -1 & \frac{1}{2} & 1 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 \\ 0 & \frac{1}{2} & 0 & -1 \end{bmatrix} \xrightarrow{\text{RREF}} \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & -\frac{2}{3} \\ 0 & 0 & 1 & -\frac{2}{3} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

\Rightarrow eigenspace of $\lambda = 1$ is spanned by $\begin{bmatrix} 2 \\ 2 \\ 3 \\ 3 \\ 1 \end{bmatrix}$.

- Now we need to make the entries add up to 1.

$$\begin{bmatrix} PR(A) \\ PR(B) \\ PR(C) \\ PR(D) \end{bmatrix} = \frac{3}{16} \begin{bmatrix} 2 \\ 2 \\ 3 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.125 \\ 0.313 \\ 0.188 \end{bmatrix}.$$

This is the PageRank vector.

- The corresponding ranking of the webpages is A, C, D, B .

Remark

In practical situations the system might be too large for finding the eigenvalues by row operations.

- Google reports having met 60 trillion webpages.
Google's search index is over 100,000,000 gigabytes.
Number of Google's servers is secret: about 2,500,000
More than 1,000,000,000 websites (i.e. hostnames; about 75% not active)
- Thus we have a gigantic but very sparse matrix.

Remark

In practical situations the system might be too large for finding the eigenvalues by row operations.

- Google reports having met 60 trillion webpages.
Google's search index is over 100,000,000 gigabytes.
Number of Google's servers is secret: about 2,500,000
More than 1,000,000,000 websites (i.e. hostnames; about 75% not active)
- Thus we have a gigantic but very sparse matrix.

An alternative to row operations is the **power method** (see Theorem 2):

Remark

In practical situations the system might be too large for finding the eigenvalues by row operations.

- Google reports having met 60 trillion webpages.
Google's search index is over 100,000,000 gigabytes.
Number of Google's servers is secret: about 2,500,000
More than 1,000,000,000 websites (i.e. hostnames; about 75% not active)
- Thus we have a gigantic but very sparse matrix.

An alternative to row operations is the **power method** (see Theorem 2):

Power method

If T is an (acyclic and irreducible) Markov matrix, then for any \mathbf{v}_0 the vectors $T^n \mathbf{v}_0$ converge to an eigenvector with eigenvalue 1

Here:

$$T = \begin{bmatrix} 0 & \frac{1}{2} & 1 & 0 \\ \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{3} & 0 & 0 & 1 \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 \end{bmatrix}.$$

Here:

$$T = \begin{bmatrix} 0 & \frac{1}{2} & 1 & 0 \\ \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{3} & 0 & 0 & 1 \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 \end{bmatrix}.$$

Start with an arbitrary state vector, hit it with powers of T .

Here:

$$T = \begin{bmatrix} 0 & \frac{1}{2} & 1 & 0 \\ \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{3} & 0 & 0 & 1 \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 \end{bmatrix}.$$

Start with an arbitrary state vector, hit it with powers of T .

$$\begin{pmatrix} \begin{bmatrix} PR(A) \\ PR(B) \\ PR(C) \\ PR(D) \end{bmatrix} \\ = \\ \begin{bmatrix} 0.375 \\ 0.125 \\ 0.313 \\ 0.188 \end{bmatrix} \end{pmatrix},$$

Here:

$$T = \begin{bmatrix} 0 & \frac{1}{2} & 1 & 0 \\ \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{3} & 0 & 0 & 1 \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 \end{bmatrix}.$$

Start with an arbitrary state vector, hit it with powers of T .

$$\left(\begin{bmatrix} PR(A) \\ PR(B) \\ PR(C) \\ PR(D) \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.125 \\ 0.313 \\ 0.188 \end{bmatrix} \right), T \begin{bmatrix} 1/4 \\ 1/4 \\ 1/4 \\ 1/4 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.083 \\ 0.333 \\ 0.208 \end{bmatrix}$$

.

Here:

$$T = \begin{bmatrix} 0 & \frac{1}{2} & 1 & 0 \\ \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{3} & 0 & 0 & 1 \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 \end{bmatrix}.$$

Start with an arbitrary state vector, hit it with powers of T .

$$\begin{pmatrix} PR(A) \\ PR(B) \\ PR(C) \\ PR(D) \end{pmatrix} = \begin{bmatrix} 0.375 \\ 0.125 \\ 0.313 \\ 0.188 \end{bmatrix}, \quad T \begin{bmatrix} 1/4 \\ 1/4 \\ 1/4 \\ 1/4 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.083 \\ 0.333 \\ 0.208 \end{bmatrix}$$

. Note that the ranking of the webpages is already A, C, D, B if we stop here.

$$T \begin{bmatrix} 1/4 \\ 1/4 \\ 1/4 \\ 1/4 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.083 \\ 0.333 \\ 0.208 \end{bmatrix}, \quad T^2 \begin{bmatrix} 1/4 \\ 1/4 \\ 1/4 \\ 1/4 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.125 \\ 0.333 \\ 0.167 \end{bmatrix},$$

Here:

$$T = \begin{bmatrix} 0 & \frac{1}{2} & 1 & 0 \\ \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{3} & 0 & 0 & 1 \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 \end{bmatrix}.$$

Start with an arbitrary state vector, hit it with powers of T .

$$\begin{pmatrix} PR(A) \\ PR(B) \\ PR(C) \\ PR(D) \end{pmatrix} = \begin{bmatrix} 0.375 \\ 0.125 \\ 0.313 \\ 0.188 \end{bmatrix}, \quad T \begin{bmatrix} 1/4 \\ 1/4 \\ 1/4 \\ 1/4 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.083 \\ 0.333 \\ 0.208 \end{bmatrix}$$

. Note that the ranking of the webpages is already A, C, D, B if we stop here.

$$T \begin{bmatrix} 1/4 \\ 1/4 \\ 1/4 \\ 1/4 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.083 \\ 0.333 \\ 0.208 \end{bmatrix}, \quad T^2 \begin{bmatrix} 1/4 \\ 1/4 \\ 1/4 \\ 1/4 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.125 \\ 0.333 \\ 0.167 \end{bmatrix}, \quad T^3 \begin{bmatrix} 1/4 \\ 1/4 \\ 1/4 \\ 1/4 \end{bmatrix} = \begin{bmatrix} 0.375 \\ 0.167 \\ 0.333 \\ 0.125 \end{bmatrix}$$

Remark

Remark

- If all entries of T are positive (no zero entries!), then the power method is guaranteed to work.

Remark

- If all entries of T are positive (no zero entries!), then the power method is guaranteed to work.
- In the context of PageRank, we can make sure that this is the case by replacing T with

$$(1 - p) \cdot \begin{bmatrix} 0 & \frac{1}{2} & 1 & 0 \\ \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{3} & 0 & 0 & 1 \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 \end{bmatrix} + p \cdot \begin{bmatrix} \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \end{bmatrix}.$$

Just to make sure: still a Markov matrix, now with positive entries
Google used to use $p = 0.15$.

Practice problems

Problem

Can you see why 1 is an eigenvalue for every Markov matrix?

Problem

Can you see why 1 is an eigenvalue for every Markov matrix?

Problem (just for fun)

The real web contains pages which have no outgoing links. In that case, our random surfer would get “stuck” (the transition matrix is not a Markov matrix). Do you have an idea how to deal with this issue?