

# Math 415 - Lecture 23

## Projections on subspaces

Monday October 19th 2015

Textbook reading: Chapter 3.2, 3.3, 3.4

Textbook reading: Chapter 3.2, 3.3, 3.4

Suggested practice exercises: Chapter 3.2 Exercise 17, 18, 24,  
Chapter 3.4 Exercise 2, 3 and see exercise at the end  
of this notes

**Textbook reading:** Chapter 3.2, 3.3, 3.4

**Suggested practice exercises:** Chapter 3.2 Exercise 17, 18, 24,  
Chapter 3.4 Exercise 2, 3 and see exercise at the end  
of this notes

**Khan Academy video:** Projections onto subspaces, Visualizing a  
projection onto a plane, Projection is closest vector in  
subspace

## Review

- **Orthogonal projection** of  $x$  onto  $y$ :

$$\hat{x} = \frac{x \cdot y}{y \cdot y} y$$

- **Orthogonal projection** of  $x$  onto  $y$ :

$$\hat{x} = \frac{x \cdot y}{y \cdot y} y$$

- **Orthogonal projection** of  $\mathbf{x}$  onto  $\mathbf{y}$ :

$$\hat{\mathbf{x}} = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}} \mathbf{y}$$

“Error”  $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$  is orthogonal to  $\mathbf{y}$ .




- **Orthogonal projection** of  $x$  onto  $y$ :

$$\hat{\mathbf{x}} = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}} \mathbf{y}$$

“Error”  $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$  is orthogonal to  $\mathbf{y}$ .

- If  $\mathbf{y}_1, \dots, \mathbf{y}_n$  is an **orthogonal basis** of  $V$ ,  $\mathbf{x}$  and  $\mathbf{x}$  is in  $V$ , then  $\mathbf{x} = c_1\mathbf{y}_1 + \dots + c_n\mathbf{y}_n$  with  $c_j = \frac{\mathbf{x} \cdot \mathbf{y}_j}{\mathbf{y}_j \cdot \mathbf{y}_j}$ .




- **Orthogonal projection** of  $\mathbf{x}$  onto  $\mathbf{y}$ :

$$\hat{\mathbf{x}} = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}} \mathbf{y}$$

“Error”  $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$  is orthogonal to  $\mathbf{y}$ .

- If  $\mathbf{y}_1, \dots, \mathbf{y}_n$  is an **orthogonal basis** of  $V$ ,  $\mathbf{x}$  and  $\mathbf{x}$  is in  $V$ , then  $\mathbf{x} = c_1\mathbf{y}_1 + \dots + c_n\mathbf{y}_n$  with  $c_j = \frac{\mathbf{x} \cdot \mathbf{y}_j}{\mathbf{y}_j \cdot \mathbf{y}_j}$ .



### Remark


$\mathbf{x}$  decomposes as the sum of its projections onto each vector in the orthogonal basis.

- **Orthogonal projection** of  $\mathbf{x}$  onto  $\mathbf{y}$ :

$$\hat{\mathbf{x}} = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}} \mathbf{y}$$

“Error”  $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$  is orthogonal to  $\mathbf{y}$ .

- If  $\mathbf{y}_1, \dots, \mathbf{y}_n$  is an **orthogonal basis** of  $V$ ,  $\mathbf{x}$  and  $\mathbf{x}$  is in  $V$ , then  $\mathbf{x} = c_1\mathbf{y}_1 + \dots + c_n\mathbf{y}_n$  with  $c_j = \frac{\mathbf{x} \cdot \mathbf{y}_j}{\mathbf{y}_j \cdot \mathbf{y}_j}$ .



### Remark


$\mathbf{x}$  decomposes as the sum of its projections onto each vector in the orthogonal basis.

- **Orthogonal projection** of  $\mathbf{x}$  onto  $\mathbf{y}$ :

$$\hat{\mathbf{x}} = \frac{\mathbf{x} \cdot \mathbf{y}}{\mathbf{y} \cdot \mathbf{y}} \mathbf{y}$$

“Error”  $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$  is orthogonal to  $\mathbf{y}$ .

- If  $\mathbf{y}_1, \dots, \mathbf{y}_n$  is an **orthogonal basis** of  $V$ ,  $\mathbf{x}$  and  $\mathbf{x}$  is in  $V$ , then  $\mathbf{x} = c_1\mathbf{y}_1 + \dots + c_n\mathbf{y}_n$  with  $c_j = \frac{\mathbf{x} \cdot \mathbf{y}_j}{\mathbf{y}_j \cdot \mathbf{y}_j}$ .



### Remark

$\mathbf{x}$  decomposes as the sum of its projections onto each vector in the orthogonal basis.

### Remark

The formulas simplify when you project on **unit** vectors: all denominators are then 1.

## Example

Express  $\underbrace{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}}_x$  in terms of the basis  $\underbrace{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}_{y_1}, \underbrace{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}_{y_2}, \underbrace{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}_{y_3}$

### Example

Express  $\underbrace{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}}_x$  in terms of the basis  $\underbrace{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}_{y_1}, \underbrace{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}_{y_2}, \underbrace{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}_{y_3}$

**Solution.**

### Example

Express  $\underbrace{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}}_{\mathbf{x}}$  in terms of the basis  $\underbrace{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}_{\mathbf{y}_1}, \underbrace{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}_{\mathbf{y}_2}, \underbrace{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}_{\mathbf{y}_3}$

### Solution.

Notice that  $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3$  is an orthogonal basis of  $\mathbb{R}^3$ .

$$\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$



$$\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} = c_1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$= \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\underbrace{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}_{\text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_1}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} +$$

$$\begin{aligned}
 \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} &= c_1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
 &= \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}}_{\text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_1} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}}_{\text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} +
 \end{aligned}$$

$$\begin{aligned}
 \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} &= c_1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
 &= \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
 &\quad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_1 \qquad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_2 \qquad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_3
 \end{aligned}$$

$$\begin{aligned}
 \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} &= c_1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
 &= \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
 &\quad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_1 \qquad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_2 \qquad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_3 \\
 &= \frac{1}{2} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} +
 \end{aligned}$$

$$\begin{aligned}
 \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} &= c_1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
 &= \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
 &\quad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_1 \qquad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_2 \qquad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_3 \\
 &= \frac{1}{2} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} + \frac{3}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} +
 \end{aligned}$$

$$\begin{aligned}
 \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} &= c_1 \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + c_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
 &= \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + \underbrace{\frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}}} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \\
 &\quad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_1 \qquad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_2 \qquad \text{projection of } \mathbf{x} \text{ onto } \mathbf{y}_3 \\
 &= \frac{1}{2} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} + \frac{3}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}
 \end{aligned}$$

## Orthogonal projection on subspaces

## Projecting onto a subspace



## Theorem

Let  $W$  be a subspace of  $\mathbb{R}^n$ . Then, each  $\mathbf{x}$  in  $\mathbb{R}^n$  can be *uniquely* written as

$$\mathbf{x} = \underbrace{\hat{\mathbf{x}}}_{\text{in } W} + \underbrace{\mathbf{x}^\perp}_{\text{in } W^\perp}$$

## Theorem


Let  $W$  be a subspace of  $\mathbb{R}^n$ . Then, each  $\mathbf{x}$  in  $\mathbb{R}^n$  can be *uniquely* written as

$$\mathbf{x} = \underbrace{\hat{\mathbf{x}}}_{\text{in } W} + \underbrace{\mathbf{x}^\perp}_{\text{in } W^\perp}$$

## Theorem

Let  $W$  be a subspace of  $\mathbb{R}^n$ . Then, each  $\mathbf{x}$  in  $\mathbb{R}^n$  can be **uniquely** written as


$$\mathbf{x} = \underbrace{\hat{\mathbf{x}}}_{\text{in } W} + \underbrace{\mathbf{x}^\perp}_{\text{in } W^\perp}$$



## Theorem


Let  $W$  be a subspace of  $\mathbb{R}^n$ . Then, each  $\mathbf{x}$  in  $\mathbb{R}^n$  can be **uniquely** written as

$$\mathbf{x} = \underbrace{\hat{\mathbf{x}}}_{\text{in } W} + \underbrace{\mathbf{x}^\perp}_{\text{in } W^\perp}$$




$\hat{\mathbf{x}}$  is the **orthogonal projection** of  $\mathbf{x}$  onto  $W$ .

## Projecting onto a subspace




## Projecting onto a subspace




- $\hat{\mathbf{x}}$  is the point in  $W$  closest to  $\mathbf{x}$ .

## Projecting onto a subspace



- $\hat{\mathbf{x}}$  is the point in  $W$  closest to  $\mathbf{x}$ .


## Projecting onto a subspace



- $\hat{\mathbf{x}}$  is the point in  $W$  closest to  $\mathbf{x}$ . For any other  $\mathbf{y}$  in  $W$ ,  $\text{dist}(\mathbf{x}, \hat{\mathbf{x}}) < \text{dist}(\mathbf{x}, \mathbf{y})$ .




## Projecting onto a subspace




- $\hat{\mathbf{x}}$  is the point in  $W$  closest to  $\mathbf{x}$ . For any other  $\mathbf{y}$  in  $W$ ,  $\text{dist}(\mathbf{x}, \hat{\mathbf{x}}) < \text{dist}(\mathbf{x}, \mathbf{y})$ .
- If  $\mathbf{v}_1, \dots, \mathbf{v}_m$  is an orthogonal basis of  $W$ , then

## Projecting onto a subspace



- $\hat{\mathbf{x}}$  is the point in  $W$  closest to  $\mathbf{x}$ . For any other  $\mathbf{y}$  in  $W$ ,  $\text{dist}(\mathbf{x}, \hat{\mathbf{x}}) < \text{dist}(\mathbf{x}, \mathbf{y})$ .
- If  $\mathbf{v}_1, \dots, \mathbf{v}_m$  is an orthogonal basis of  $W$ , then


## Projecting onto a subspace



- $\hat{\mathbf{x}}$  is the point in  $W$  closest to  $\mathbf{x}$ . For any other  $\mathbf{y}$  in  $W$ ,  $\text{dist}(\mathbf{x}, \hat{\mathbf{x}}) < \text{dist}(\mathbf{x}, \mathbf{y})$ .
- If  $\mathbf{v}_1, \dots, \mathbf{v}_m$  is an orthogonal basis of  $W$ , then

$$\hat{\mathbf{x}} = \left( \frac{\mathbf{x} \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \right) \mathbf{v}_1 + \dots + \left( \frac{\mathbf{x} \cdot \mathbf{v}_m}{\mathbf{v}_m \cdot \mathbf{v}_m} \right) \mathbf{v}_m$$

## Projecting onto a subspace




- $\hat{\mathbf{x}}$  is the point in  $W$  closest to  $\mathbf{x}$ . For any other  $\mathbf{y}$  in  $W$ ,  $\text{dist}(\mathbf{x}, \hat{\mathbf{x}}) < \text{dist}(\mathbf{x}, \mathbf{y})$ .
- If  $\mathbf{v}_1, \dots, \mathbf{v}_m$  is an orthogonal basis of  $W$ , then

$$\hat{\mathbf{x}} = \left( \frac{\mathbf{x} \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \right) \mathbf{v}_1 + \dots + \left( \frac{\mathbf{x} \cdot \mathbf{v}_m}{\mathbf{v}_m \cdot \mathbf{v}_m} \right) \mathbf{v}_m$$

Once  $\hat{\mathbf{x}}$  is determined,  $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$ .

## Projecting onto a subspace



- $\hat{\mathbf{x}}$  is the point in  $W$  closest to  $\mathbf{x}$ . For any other  $\mathbf{y}$  in  $W$ ,  $\text{dist}(\mathbf{x}, \hat{\mathbf{x}}) < \text{dist}(\mathbf{x}, \mathbf{y})$ .
- If  $\mathbf{v}_1, \dots, \mathbf{v}_m$  is an orthogonal basis of  $W$ , then

$$\hat{\mathbf{x}} = \left( \frac{\mathbf{x} \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \right) \mathbf{v}_1 + \dots + \left( \frac{\mathbf{x} \cdot \mathbf{v}_m}{\mathbf{v}_m \cdot \mathbf{v}_m} \right) \mathbf{v}_m$$

Once  $\hat{\mathbf{x}}$  is determined,  $\mathbf{x}^\perp = \mathbf{x} - \hat{\mathbf{x}}$ .

(This is also the orthogonal projection of  $\mathbf{x}$  onto  $W^\perp$ .)

## Example

Let  $W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ , and  $\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$ .

- Find the orthogonal projection of  $\mathbf{x}$  onto  $W$ . (Or: find the vector in  $W$  which is closest to  $\mathbf{x}$ )
- Write  $\mathbf{x}$  as a vector in  $W$  plus a vector orthogonal to  $W$ .

## Example

Let  $W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ , and  $\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$ .

- Find the orthogonal projection of  $\mathbf{x}$  onto  $W$ . (Or: find the vector in  $W$  which is closest to  $\mathbf{x}$ )
- Write  $\mathbf{x}$  as a vector in  $W$  plus a vector orthogonal to  $W$ .

**Solution.**

### Example

Let  $W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ , and  $\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$ .

- Find the orthogonal projection of  $\mathbf{x}$  onto  $W$ . (Or: find the vector in  $W$  which is closest to  $\mathbf{x}$ )
- Write  $\mathbf{x}$  as a vector in  $W$  plus a vector orthogonal to  $W$ .

**Solution.**

Note that  $\mathbf{w}_1 = \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}$  and  $\mathbf{w}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$  are an orthogonal basis for  $W$ .



## Example

Let  $W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ , and  $\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$ .

- Find the orthogonal projection of  $\mathbf{x}$  onto  $W$ . (Or: find the vector in  $W$  which is closest to  $\mathbf{x}$ )
- Write  $\mathbf{x}$  as a vector in  $W$  plus a vector orthogonal to  $W$ .

### Solution.

Note that  $\mathbf{w}_1 = \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}$  and  $\mathbf{w}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$  are an orthogonal basis for  $W$ .

(We will soon learn how to construct orthogonal bases ourselves).

Hence, the orthogonal projection of  $\mathbf{x}$  onto  $W$  is:

Hence, the orthogonal projection of  $\mathbf{x}$  onto  $W$  is:

$$\hat{\mathbf{x}} = \frac{\mathbf{x} \cdot \mathbf{w}_1}{\mathbf{w}_1 \cdot \mathbf{w}_1} \mathbf{w}_1 + \frac{\mathbf{x} \cdot \mathbf{w}_2}{\mathbf{w}_2 \cdot \mathbf{w}_2} \mathbf{w}_2$$

Hence, the orthogonal projection of  $\mathbf{x}$  onto  $W$  is:

$$\begin{aligned}\hat{\mathbf{x}} &= \frac{\mathbf{x} \cdot \mathbf{w}_1}{\mathbf{w}_1 \cdot \mathbf{w}_1} \mathbf{w}_1 + \frac{\mathbf{x} \cdot \mathbf{w}_2}{\mathbf{w}_2 \cdot \mathbf{w}_2} \mathbf{w}_2 \\ &= \frac{\begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}\end{aligned}$$

Hence, the orthogonal projection of  $\mathbf{x}$  onto  $W$  is:

$$\begin{aligned}
 \hat{\mathbf{x}} &= \frac{\mathbf{x} \cdot \mathbf{w}_1}{\mathbf{w}_1 \cdot \mathbf{w}_1} \mathbf{w}_1 + \frac{\mathbf{x} \cdot \mathbf{w}_2}{\mathbf{w}_2 \cdot \mathbf{w}_2} \mathbf{w}_2 \\
 &= \frac{\begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \\
 &= \frac{10}{10} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} =
 \end{aligned}$$

Hence, the orthogonal projection of  $\mathbf{x}$  onto  $W$  is:

$$\begin{aligned}
 \hat{\mathbf{x}} &= \frac{\mathbf{x} \cdot \mathbf{w}_1}{\mathbf{w}_1 \cdot \mathbf{w}_1} \mathbf{w}_1 + \frac{\mathbf{x} \cdot \mathbf{w}_2}{\mathbf{w}_2 \cdot \mathbf{w}_2} \mathbf{w}_2 \\
 &= \frac{\begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \\
 &= \frac{10}{10} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix}
 \end{aligned}$$

Hence, the orthogonal projection of  $\mathbf{x}$  onto  $W$  is:

$$\begin{aligned}
 \hat{\mathbf{x}} &= \frac{\mathbf{x} \cdot \mathbf{w}_1}{\mathbf{w}_1 \cdot \mathbf{w}_1} \mathbf{w}_1 + \frac{\mathbf{x} \cdot \mathbf{w}_2}{\mathbf{w}_2 \cdot \mathbf{w}_2} \mathbf{w}_2 \\
 &= \frac{\begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \\
 &= \frac{10}{10} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix}
 \end{aligned}$$

### Warning

This calculation only works for **orthogonal**  $\mathbf{w}_1, \mathbf{w}_2$ !

$\hat{\mathbf{x}}$  is the vector in  $W$  which best approximates  $\mathbf{x}$ .



$\hat{\mathbf{x}}$  is the vector in  $W$  which best approximates  $\mathbf{x}$ .

Orthogonal projection of  $\mathbf{x}$  onto the orthogonal complement of  $W$ :

$\hat{\mathbf{x}}$  is the vector in  $W$  which best approximates  $\mathbf{x}$ .

Orthogonal projection of  $\mathbf{x}$  onto the orthogonal complement of  $W$ :

$$\mathbf{x}^\perp = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} - \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix}$$

$\hat{\mathbf{x}}$  is the vector in  $W$  which best approximates  $\mathbf{x}$ .

Orthogonal projection of  $\mathbf{x}$  onto the orthogonal complement of  $W$ :

$$\mathbf{x}^\perp = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} - \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} -3 \\ 0 \\ 9 \end{bmatrix}$$

$\hat{\mathbf{x}}$  is the vector in  $W$  which best approximates  $\mathbf{x}$ .

Orthogonal projection of  $\mathbf{x}$  onto the orthogonal complement of  $W$ :

$$\mathbf{x}^\perp = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} - \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} -3 \\ 0 \\ 9 \end{bmatrix}$$

Hence,

$$\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} = \underbrace{\begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix}}_{\text{in } W} + \underbrace{\begin{bmatrix} -3 \\ 0 \\ 9 \end{bmatrix}}_{\text{in } W^\perp}$$

$\hat{\mathbf{x}}$  is the vector in  $W$  which best approximates  $\mathbf{x}$ .

Orthogonal projection of  $\mathbf{x}$  onto the orthogonal complement of  $W$ :

$$\mathbf{x}^\perp = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} - \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} -3 \\ 0 \\ 9 \end{bmatrix}$$

Hence,

$$\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} = \underbrace{\begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix}}_{\text{in } W} + \underbrace{\begin{bmatrix} -3 \\ 0 \\ 9 \end{bmatrix}}_{\text{in } W^\perp}$$

Indeed,  $\begin{bmatrix} -3 \\ 0 \\ 9 \end{bmatrix}$  is orthogonal to  $\mathbf{w}_1 = \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}$  and  $\mathbf{w}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ .

## The matrix of a projection

## Definition

Let  $\mathbf{v}_1, \dots, \mathbf{v}_m$  be an orthogonal basis of  $W$ , a subspace of  $\mathbb{R}^n$ .

## Definition

Let  $\mathbf{v}_1, \dots, \mathbf{v}_m$  be an orthogonal basis of  $W$ , a subspace of  $\mathbb{R}^n$ .



## Definition

Let  $\mathbf{v}_1, \dots, \mathbf{v}_m$  be an orthogonal basis of  $W$ , a subspace of  $\mathbb{R}^n$ . Note that the projection map  $\pi_W : \mathbb{R}^n \rightarrow \mathbb{R}^n$  that sends  $\mathbf{x}$  to

$$\hat{\mathbf{x}} = \left( \frac{\mathbf{x} \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \right) \mathbf{v}_1 + \dots + \left( \frac{\mathbf{x} \cdot \mathbf{v}_m}{\mathbf{v}_m \cdot \mathbf{v}_m} \right) \mathbf{v}_m$$

is a linear map.

## Definition

Let  $\mathbf{v}_1, \dots, \mathbf{v}_m$  be an orthogonal basis of  $W$ , a subspace of  $\mathbb{R}^n$ . Note that the projection map  $\pi_W : \mathbb{R}^n \rightarrow \mathbb{R}^n$  that sends  $\mathbf{x}$  to

$$\hat{\mathbf{x}} = \left( \frac{\mathbf{x} \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \right) \mathbf{v}_1 + \dots + \left( \frac{\mathbf{x} \cdot \mathbf{v}_m}{\mathbf{v}_m \cdot \mathbf{v}_m} \right) \mathbf{v}_m$$

is a linear map. The matrix  $P$  representing  $\pi_W$  with respect to the standard basis is the **projection matrix**.

### Example

Find the projection matrix  $P$  for the orthogonal projection onto

$$W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$$

in  $\mathbb{R}^3$ .

### Example

Find the projection matrix  $P$  for the orthogonal projection onto

$$W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$$

in  $\mathbb{R}^3$ .

**Solution.**

### Example

Find the projection matrix  $P$  for the orthogonal projection onto

$$W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$$

in  $\mathbb{R}^3$ .

**Solution.** Standard basis:  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ .

The first column of  $P$  encodes the projection of  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ :

The first column of  $P$  encodes the projection of  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ :

$$\frac{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

The first column of  $P$  encodes the projection of  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ :

$$\frac{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \frac{3}{10} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}.$$



The first column of  $P$  encodes the projection of  $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ :

$$\frac{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \frac{3}{10} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}.$$

Hence  $P = \begin{bmatrix} \frac{9}{10} & * & * \\ 0 & * & * \\ \frac{3}{10} & * & * \end{bmatrix}.$

The second column of  $P$  encodes the projection of  $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ :

The second column of  $P$  encodes the projection of  $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ :

$$\frac{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

The second column of  $P$  encodes the projection of  $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ :

$$\frac{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}.$$

The second column of  $P$  encodes the projection of  $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ :

$$\frac{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}.$$

Hence  $P = \begin{bmatrix} \frac{9}{10} & 0 & * \\ 0 & 1 & * \\ \frac{3}{10} & 0 & * \end{bmatrix}.$

The third column of  $P$  encodes the projection of  $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ :

The third column of  $P$  encodes the projection of  $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ :

$$\frac{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

The third column of  $P$  encodes the projection of  $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ :

$$\frac{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \frac{1}{10} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}.$$



The third column of  $P$  encodes the projection of  $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ :

$$\frac{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}}{\begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix} + \frac{\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \frac{1}{10} \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}.$$

Hence  $P = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix}.$

Let's do the earlier example again using the matrix  $P$ .

Let's do the earlier example again using the matrix  $P$ .

### Example

Let  $W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ , and  $\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$ . Find the orthogonal projection of  $\mathbf{x}$  onto  $W$ .

Let's do the earlier example again using the matrix  $P$ .

### Example

Let  $W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ , and  $\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$ . Find the orthogonal projection of  $\mathbf{x}$  onto  $W$ .

**Solution.**

Let's do the earlier example again using the matrix  $P$ .

### Example

Let  $W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ , and  $\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$ . Find the orthogonal projection of  $\mathbf{x}$  onto  $W$ .

**Solution.**

$$\hat{\mathbf{x}} = P\mathbf{x}$$

Let's do the earlier example again using the matrix  $P$ .

### Example

Let  $W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ , and  $\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$ . Find the orthogonal projection of  $\mathbf{x}$  onto  $W$ .

**Solution.**

$$\hat{\mathbf{x}} = P\mathbf{x} = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix} \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$$

Let's do the earlier example again using the matrix  $P$ .

### Example

Let  $W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ , and  $\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$ . Find the orthogonal projection of  $\mathbf{x}$  onto  $W$ .

**Solution.**

$$\hat{\mathbf{x}} = P\mathbf{x} = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix} \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix},$$

Let's do the earlier example again using the matrix  $P$ .

### Example

Let  $W = \text{span} \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$ , and  $\mathbf{x} = \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix}$ . Find the orthogonal projection of  $\mathbf{x}$  onto  $W$ .

**Solution.**

$$\hat{\mathbf{x}} = P\mathbf{x} = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix} \begin{bmatrix} 0 \\ 3 \\ 10 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ 1 \end{bmatrix},$$

as in the previous example.



## Example

Compute  $P^2$  when

$$P = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix}.$$

Explain why the answer makes sense.

## Example

Compute  $P^2$  when

$$P = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix}.$$

Explain why the answer makes sense.

**Solution.**

## Example

Compute  $P^2$  when

$$P = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix}.$$

Explain why the answer makes sense.

**Solution.**

$$\begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix} \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix} = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix}$$

## Example

Compute  $P^2$  when

$$P = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix}.$$

Explain why the answer makes sense.

**Solution.**

$$\begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix} \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix} = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix}$$

$$P^2 = P$$

## Example

Compute  $P^2$  when

$$P = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix}.$$

Explain why the answer makes sense.

**Solution.**

$$\begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix} \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix} = \begin{bmatrix} \frac{9}{10} & 0 & \frac{3}{10} \\ 0 & 1 & 0 \\ \frac{3}{10} & 0 & \frac{1}{10} \end{bmatrix}$$
$$P^2 = P$$

Once we have projected down onto  $W$ , projecting onto  $W$  again does not change anything!

## Practice problems

## Practice problems

## Example

Find the closest point to  $\mathbf{x}$  in  $\text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  where

$$\mathbf{x} = \begin{bmatrix} 2 \\ 4 \\ 0 \\ -2 \end{bmatrix}, \mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$



### Example

Find the closest point to  $\mathbf{x}$  in  $\text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  where

$$\mathbf{x} = \begin{bmatrix} 2 \\ 4 \\ 0 \\ -2 \end{bmatrix}, \mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

**Solution.**

$$\hat{\mathbf{x}} =$$

## Example

Find the closest point to  $\mathbf{x}$  in  $\text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  where

$$\mathbf{x} = \begin{bmatrix} 2 \\ 4 \\ 0 \\ -2 \end{bmatrix}, \mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

**Solution.**

$$\hat{\mathbf{x}} = \frac{6}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} + \frac{-2}{2} \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

## Example

Find the closest point to  $\mathbf{x}$  in  $\text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  where

$$\mathbf{x} = \begin{bmatrix} 2 \\ 4 \\ 0 \\ -2 \end{bmatrix}, \mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

**Solution.**

$$\hat{\mathbf{x}} = \frac{6}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} + \frac{-2}{2} \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ -1 \\ -1 \end{bmatrix}$$

### Example

If  $P$  is the projection matrix for projecting on  $W$ , what is the projection matrix  $Q$  for projecting on  $W^\perp$ ?

### Example

If  $P$  is the projection matrix for projecting on  $W$ , what is the projection matrix  $Q$  for projecting on  $W^\perp$ ?

**Solution.**  $Q = 1 - P$ !

### Example

Let  $P$  be the projection matrix for projecting on  $W$ , and let  $\mathbf{x}$  be some vector.

### Example

Let  $P$  be the projection matrix for projecting on  $W$ , and let  $\mathbf{x}$  be some vector.

- Suppose  $P\mathbf{x} = \mathbf{x}$ . What can you say about  $\mathbf{x}$ ?

### Example

Let  $P$  be the projection matrix for projecting on  $W$ , and let  $\mathbf{x}$  be some vector.

- Suppose  $P\mathbf{x} = \mathbf{x}$ . What can you say about  $\mathbf{x}$ ?
- Suppose  $P\mathbf{x} = 0$ . What can you say about  $\mathbf{x}$ ?



### Example

Let  $P$  be the projection matrix for projecting on  $W$ , and let  $\mathbf{x}$  be some vector.

- Suppose  $P\mathbf{x} = \mathbf{x}$ . What can you say about  $\mathbf{x}$ ?
- Suppose  $P\mathbf{x} = 0$ . What can you say about  $\mathbf{x}$ ?