

# Math 415 - Lecture 10

## Span is a subspace, Null Space

Wednesday September 16th 2015

Textbook: Chapter 2.1, 2.2.

Textbook: Chapter 2.1, 2.2.

Suggested practice exercises: Chapter 2.1: 3, 21, 28. Chapter 2.2:  
33 and additional exercises in this lecture note.

**Textbook:** Chapter 2.1, 2.2.

**Suggested practice exercises:** Chapter 2.1: 3, 21, 28. Chapter 2.2: 33 and additional exercises in this lecture note.

**Khan Academy videos:** Linear Subspaces, Introduction to the Null Space of a Matrix, Calculating the Null Space of a Matrix

## Review of vector space and subspace

- A **vector space** is a set of vectors which can be **added** and **scaled** (without leaving the space!); subject to the “usual” rules.

- A **vector space** is a set of vectors which can be **added** and **scaled** (without leaving the space!); subject to the “usual” rules.
- The set of all polynomials of degree **up to 2** is a vector space. Why?

- A **vector space** is a set of vectors which can be **added** and **scaled** (without leaving the space!); subject to the “usual” rules.
- The set of all polynomials of degree **up to 2** is a vector space. Why?

$$\begin{aligned}[a_0 + a_1 t + a_2 t^2] + [b_0 + b_1 t + b_2 t^2] &= \\ &= [(a_0 + b_0) + (a_1 + b_1)t + (a_2 + b_2)t^2] \\ r[a_0 + a_1 t + a_2 t^2] &= [(ra_0) + (ra_1)t + (ra_2)t^2]\end{aligned}$$



- A **vector space** is a set of vectors which can be **added** and **scaled** (without leaving the space!); subject to the “usual” rules.
- The set of all polynomials of degree **up to 2** is a vector space. Why?

$$\begin{aligned}[a_0 + a_1 t + a_2 t^2] + [b_0 + b_1 t + b_2 t^2] &= \\ &= [(a_0 + b_0) + (a_1 + b_1)t + (a_2 + b_2)t^2] \\ r[a_0 + a_1 t + a_2 t^2] &= [(ra_0) + (ra_1)t + (ra_2)t^2]\end{aligned}$$

Note how it “works” just like  $\mathbb{R}^3$ .

- A **vector space** is a set of vectors which can be **added** and **scaled** (without leaving the space!); subject to the “usual” rules.
- The set of all polynomials of degree **up to 2** is a vector space. Why?

$$\begin{aligned}[a_0 + a_1t + a_2t^2] + [b_0 + b_1t + b_2t^2] &= \\ &[(a_0 + b_0) + (a_1 + b_1)t + (a_2 + b_2)t^2] \\ r[a_0 + a_1t + a_2t^2] &= [(ra_0) + (ra_1)t + (ra_2)t^2]\end{aligned}$$

Note how it “works” just like  $\mathbb{R}^3$ .

- The set of all polynomials of degree **exactly 2** is **not** a vector space. Why?

- A **vector space** is a set of vectors which can be **added** and **scaled** (without leaving the space!); subject to the “usual” rules.
- The set of all polynomials of degree **up to 2** is a vector space. Why?

$$\begin{aligned}[a_0 + a_1t + a_2t^2] + [b_0 + b_1t + b_2t^2] &= \\ &= [(a_0 + b_0) + (a_1 + b_1)t + (a_2 + b_2)t^2] \\ r[a_0 + a_1t + a_2t^2] &= [(ra_0) + (ra_1)t + (ra_2)t^2]\end{aligned}$$

Note how it “works” just like  $\mathbb{R}^3$ .

- The set of all polynomials of degree **exactly 2** is **not** a vector space. Why?

$$\underbrace{[1 + 4t + t^2]}_{\text{degree 2}} + \underbrace{[3 - t - t^2]}_{\text{degree 2}} =$$

- A **vector space** is a set of vectors which can be **added** and **scaled** (without leaving the space!); subject to the “usual” rules.
- The set of all polynomials of degree **up to 2** is a vector space. Why?

$$\begin{aligned}[a_0 + a_1t + a_2t^2] + [b_0 + b_1t + b_2t^2] &= \\ &= [(a_0 + b_0) + (a_1 + b_1)t + (a_2 + b_2)t^2] \\ r[a_0 + a_1t + a_2t^2] &= [(ra_0) + (ra_1)t + (ra_2)t^2]\end{aligned}$$

Note how it “works” just like  $\mathbb{R}^3$ .

- The set of all polynomials of degree **exactly 2** is **not** a vector space. Why?

$$\underbrace{[1 + 4t + t^2]}_{\text{degree 2}} + \underbrace{[3 - t - t^2]}_{\text{degree 2}} = \underbrace{[4 + 3t]}_{\text{NOT degree 2}}$$

- **Easy test:** Is the zero vector in the set? (If not, then it's **not** a vector space.)

- **Easy test:** Is the zero vector in the set? (If not, then it's **not** a vector space.)

### Example

Let  $V$  be the set of all function  $f : \mathbb{R} \rightarrow \mathbb{R}$ . Is  $V$  a vector space?

- **Easy test:** Is the zero vector in the set? (If not, then it's **not** a vector space.)

### Example

Let  $V$  be the set of all function  $f : \mathbb{R} \rightarrow \mathbb{R}$ . Is  $V$  a vector space?

### Solution

Yes!

- **Easy test:** Is the zero vector in the set? (If not, then it's **not** a vector space.)

### Example

Let  $V$  be the set of all function  $f : \mathbb{R} \rightarrow \mathbb{R}$ . Is  $V$  a vector space?

### Solution

Yes! Adding of functions  $f$  and  $g$ :

$$f(x) + g(x) = (f + g)(x)$$

so  $f(x) + g(x)$  is in  $V$ .



- **Easy test:** Is the zero vector in the set? (If not, then it's **not** a vector space.)

### Example

Let  $V$  be the set of all function  $f : \mathbb{R} \rightarrow \mathbb{R}$ . Is  $V$  a vector space?

### Solution

Yes! Adding of functions  $f$  and  $g$ :

$$f(x) + g(x) = (f + g)(x)$$

so  $f(x) + g(x)$  is in  $V$ .

Note that, once more, this definition is “component-wise”.

- **Easy test:** Is the zero vector in the set? (If not, then it's **not** a vector space.)

### Example

Let  $V$  be the set of all function  $f : \mathbb{R} \rightarrow \mathbb{R}$ . Is  $V$  a vector space?

### Solution

Yes! Adding of functions  $f$  and  $g$ :

$$f(x) + g(x) = (f + g)(x)$$

so  $f(x) + g(x)$  is in  $V$ .

Note that, once more, this definition is “component-wise”. Scalar multiplication works the same way.

# Subspaces

## Definition

A **subspace** of a vector space  $V$  is a subset  $H$  of  $V$  that has three properties:

# Subspaces

## Definition

A **subspace** of a vector space  $V$  is a subset  $H$  of  $V$  that has three properties:

1. The zero vector of  $V$  is in  $H$ .

# Subspaces

## Definition

A **subspace** of a vector space  $V$  is a subset  $H$  of  $V$  that has three properties:

1. The zero vector of  $V$  is in  $H$ .
2. For each  $\mathbf{u}$  and  $\mathbf{v}$  are in  $H$ ,  $\mathbf{u} + \mathbf{v}$  is in  $H$ .  
(In this case we say  $H$  is closed under vector addition.)

# Subspaces

## Definition

A **subspace** of a vector space  $V$  is a subset  $H$  of  $V$  that has three properties:

1. The zero vector of  $V$  is in  $H$ .
2. For each  $\mathbf{u}$  and  $\mathbf{v}$  are in  $H$ ,  $\mathbf{u} + \mathbf{v}$  is in  $H$ .  
(In this case we say  $H$  is closed under vector addition.)
3. For each  $\mathbf{u}$  in  $H$  and each scalar  $c$ ,  $c\mathbf{u}$  is in  $H$ .  
(In this case we say  $H$  is closed under scalar multiplication.)

## Problem

*Find as many subspaces in  $\mathbb{R}^2$  as you can.*

## A Shortcut for Determining Subspaces



# A Shortcut for Determining Subspaces

## Definition

Recall that  $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$  is the collection of all vectors that can be written as

$$x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \cdots + x_p\mathbf{v}_p,$$

where  $x_1, x_2, \dots, x_p$  are scalars.

# A Shortcut for Determining Subspaces

## Definition

Recall that  $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$  is the collection of all vectors that can be written as

$$x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \cdots + x_p\mathbf{v}_p,$$

where  $x_1, x_2, \dots, x_p$  are scalars.

## Theorem

*If  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p$  are in a vector space  $V$ , then  $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_p\}$  is a subspace of  $V$ .*

# A Shortcut for Determining Subspaces

## Example

Is  $V = \left\{ \begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ? Why or why not?

# A Shortcut for Determining Subspaces

## Example

Is  $V = \left\{ \begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ? Why or why not?

## Solution

Write vectors in  $V$  as:

$$\begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} = \begin{bmatrix} a \\ 2a \end{bmatrix} + \begin{bmatrix} 2b \\ -3b \end{bmatrix}$$

# A Shortcut for Determining Subspaces

## Example

Is  $V = \left\{ \begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ? Why or why not?

## Solution

Write vectors in  $V$  as:

$$\begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} = \begin{bmatrix} a \\ 2a \end{bmatrix} + \begin{bmatrix} 2b \\ -3b \end{bmatrix} = a \begin{bmatrix} 1 \\ 2 \end{bmatrix} + b \begin{bmatrix} 2 \\ -3 \end{bmatrix}.$$

# A Shortcut for Determining Subspaces

## Example

Is  $V = \left\{ \begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ? Why or why not?

## Solution

Write vectors in  $V$  as:

$$\begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} = \begin{bmatrix} a \\ 2a \end{bmatrix} + \begin{bmatrix} 2b \\ -3b \end{bmatrix} = a \begin{bmatrix} 1 \\ 2 \end{bmatrix} + b \begin{bmatrix} 2 \\ -3 \end{bmatrix}.$$

So  $V = \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  where

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix},$$

# A Shortcut for Determining Subspaces

## Example

Is  $V = \left\{ \begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ? Why or why not?

## Solution

Write vectors in  $V$  as:

$$\begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} = \begin{bmatrix} a \\ 2a \end{bmatrix} + \begin{bmatrix} 2b \\ -3b \end{bmatrix} = a \begin{bmatrix} 1 \\ 2 \end{bmatrix} + b \begin{bmatrix} 2 \\ -3 \end{bmatrix}.$$

So  $V = \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  where

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 2 \\ -3 \end{bmatrix}$$

# A Shortcut for Determining Subspaces

## Example

Is  $V = \left\{ \begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^2$ ? Why or why not?

## Solution

Write vectors in  $V$  as:

$$\begin{bmatrix} a + 2b \\ 2a - 3b \end{bmatrix} = \begin{bmatrix} a \\ 2a \end{bmatrix} + \begin{bmatrix} 2b \\ -3b \end{bmatrix} = a \begin{bmatrix} 1 \\ 2 \end{bmatrix} + b \begin{bmatrix} 2 \\ -3 \end{bmatrix}.$$

So  $V = \text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  where

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 2 \\ -3 \end{bmatrix}$$

and therefore  $V$  is a subspace of  $\mathbb{R}^2$  by the previous theorem.



### Example

Is  $H = \left\{ \begin{bmatrix} a + 2b \\ a + 1 \\ a \end{bmatrix} : a, b \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^3$ ? Why or why not?

### Example

Is  $H = \left\{ \begin{bmatrix} a + 2b \\ a + 1 \\ a \end{bmatrix} : a, b \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^3$ ? Why or why not?

### Solution

No!  $H$  does not contain the zero vector.

### Example

Is  $H = \left\{ \begin{bmatrix} a + 2b \\ a + 1 \\ a \end{bmatrix} : a, b \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^3$ ? Why or why not?

### Solution

No!  $H$  does not contain the zero vector. In other words,

$$\begin{bmatrix} a + 2b \\ a + 1 \\ a \end{bmatrix} = a \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + b \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

cannot equal the zero vector for any choice of  $a$  or  $b$ .

### Example

Is the set  $H$  of all matrices of the form  $\begin{bmatrix} 2a & b \\ 3a + b & 3b \end{bmatrix}$  a subspace of  $M_{2 \times 2}$ ?

### Example

Is the set  $H$  of all matrices of the form  $\begin{bmatrix} 2a & b \\ 3a+b & 3b \end{bmatrix}$  a subspace of  $M_{2 \times 2}$ ?

### Solution

Yes!

$$H = \text{span} \left\{ \begin{bmatrix} 2 & 0 \\ 3 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 3 \end{bmatrix} \right\}.$$

Since  $H$  can be written as a span, it's a subspace of  $M_{2 \times 2}$ .

# Additional Problems

## Problem

*Determine which of the following sets are subspaces and give reasons:*

# Additional Problems

## Problem

*Determine which of the following sets are subspaces and give reasons:*

1.  $W_1 = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a - 2b = c, 4a + 2c = 1 \right\}.$

# Additional Problems

## Problem

*Determine which of the following sets are subspaces and give reasons:*

1.  $W_1 = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a - 2b = c, 4a + 2c = 1 \right\}.$



# Additional Problems

## Problem

*Determine which of the following sets are subspaces and give reasons:*

$$1. W_1 = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a - 2b = c, 4a + 2c = 1 \right\}.$$

$$2. W_2 = \left\{ \begin{bmatrix} a - b \\ c \\ a + c \\ a - 2b - c \end{bmatrix} : a, b, c \in \mathbb{R} \right\}.$$

# Additional Problems

## Problem

*Determine which of the following sets are subspaces and give reasons:*

$$1. W_1 = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a - 2b = c, 4a + 2c = 1 \right\}.$$

$$2. W_2 = \left\{ \begin{bmatrix} a - b \\ c \\ a + c \\ a - 2b - c \end{bmatrix} : a, b, c \in \mathbb{R} \right\}.$$

# Additional Problems

## Problem

*Determine which of the following sets are subspaces and give reasons:*

$$1. W_1 = \left\{ \begin{bmatrix} a \\ b \\ c \end{bmatrix} : a - 2b = c, 4a + 2c = 1 \right\}.$$

$$2. W_2 = \left\{ \begin{bmatrix} a - b \\ c \\ a + c \\ a - 2b - c \end{bmatrix} : a, b, c \in \mathbb{R} \right\}.$$

$$3. W_3 = \left\{ \begin{bmatrix} a \\ b \end{bmatrix} : a \cdot b \geq 0 \right\}.$$

## Null Spaces

# Null Spaces

## Definition

The **nullspace** of an  $m \times n$  matrix  $A$ , written as  $\text{Nul}(A)$ , is the set of all solutions to the homogeneous equation  $A\mathbf{x} = \mathbf{0}$ .

# Null Spaces

## Definition

The **nullspace** of an  $m \times n$  matrix  $A$ , written as  $\text{Nul}(A)$ , is the set of all solutions to the homogeneous equation  $A\mathbf{x} = \mathbf{0}$ .

$$\text{Nul}(A) = \{\mathbf{x} : \mathbf{x} \in \mathbb{R}^n \text{ and } A\mathbf{x} = \mathbf{0}\}.$$

# Null Spaces

## Theorem

*The null space of an  $m \times n$  matrix  $A$  is a subspace of  $\mathbb{R}^n$ .*

# Null Spaces

## Theorem

*The null space of an  $m \times n$  matrix  $A$  is a subspace of  $\mathbb{R}^n$ .  
Equivalently, the set of all solutions to the system  $A\mathbf{x} = \mathbf{0}$  of  $m$  homogeneous linear equations in  $n$  unknowns is a subspace of  $\mathbb{R}^n$ .*



# Null Spaces

## Theorem

*The null space of an  $m \times n$  matrix  $A$  is a subspace of  $\mathbb{R}^n$ .  
Equivalently, the set of all solutions to the system  $A\mathbf{x} = \mathbf{0}$  of  $m$  homogeneous linear equations in  $n$  unknowns is a subspace of  $\mathbb{R}^n$ .*

**Proof:**  $\text{Nul}(A)$  is a subset of  $\mathbb{R}^n$  since  $A$  has  $n$  columns. We have to verify properties (a), (b), and (c) of the definition of a subspace.

# Null Spaces

## Theorem

*The null space of an  $m \times n$  matrix  $A$  is a subspace of  $\mathbb{R}^n$ .  
Equivalently, the set of all solutions to the system  $A\mathbf{x} = \mathbf{0}$  of  $m$  homogeneous linear equations in  $n$  unknowns is a subspace of  $\mathbb{R}^n$ .*

**Proof:**  $\text{Nul}(A)$  is a subset of  $\mathbb{R}^n$  since  $A$  has  $n$  columns. We have to verify properties (a), (b), and (c) of the definition of a subspace.

**Property (a):** Show that  $\mathbf{0}$  is in  $\text{Nul}(A)$ .

# Null Spaces

## Theorem

*The null space of an  $m \times n$  matrix  $A$  is a subspace of  $\mathbb{R}^n$ .  
Equivalently, the set of all solutions to the system  $A\mathbf{x} = \mathbf{0}$  of  $m$  homogeneous linear equations in  $n$  unknowns is a subspace of  $\mathbb{R}^n$ .*

**Proof:**  $\text{Nul}(A)$  is a subset of  $\mathbb{R}^n$  since  $A$  has  $n$  columns. We have to verify properties (a), (b), and (c) of the definition of a subspace.

**Property (a):** Show that  $\mathbf{0}$  is in  $\text{Nul}(A)$ .

$$A\mathbf{0} = \mathbf{0}.$$

and

$$A \begin{matrix} \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \\ \text{in } \mathbb{R}^n \end{matrix} = \begin{matrix} \begin{bmatrix} 0 \\ \vdots \\ 0 \end{bmatrix} \\ \text{in } \mathbb{R}^m \end{matrix}$$

# Null Spaces

**Property (b):** If  $\mathbf{u}$  and  $\mathbf{v}$  are in  $\text{Nul}(A)$ , show that  $\mathbf{u} + \mathbf{v}$  is also in  $\text{Nul}(A)$ .

# Null Spaces

**Property (b):** If  $\mathbf{u}$  and  $\mathbf{v}$  are in  $\text{Nul}(A)$ , show that  $\mathbf{u} + \mathbf{v}$  is also in  $\text{Nul}(A)$ . Suppose  $A\mathbf{u} = \mathbf{0}$  and  $A\mathbf{v} = \mathbf{0}$ . Then

# Null Spaces

**Property (b):** If  $\mathbf{u}$  and  $\mathbf{v}$  are in  $\text{Nul}(A)$ , show that  $\mathbf{u} + \mathbf{v}$  is also in  $\text{Nul}(A)$ . Suppose  $A\mathbf{u} = \mathbf{0}$  and  $A\mathbf{v} = \mathbf{0}$ . Then

$$A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v} = \mathbf{0} + \mathbf{0} = \mathbf{0}.$$

# Null Spaces

**Property (b):** If  $\mathbf{u}$  and  $\mathbf{v}$  are in  $\text{Nul}(A)$ , show that  $\mathbf{u} + \mathbf{v}$  is also in  $\text{Nul}(A)$ . Suppose  $A\mathbf{u} = \mathbf{0}$  and  $A\mathbf{v} = \mathbf{0}$ . Then

$$A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v} = \mathbf{0} + \mathbf{0} = \mathbf{0}.$$

**Property (c):** If  $\mathbf{u}$  is in  $\text{Nul}(A)$  and  $c$  is a scalar, show that  $c\mathbf{u}$  is also in  $\text{Nul}(A)$ .

# Null Spaces

**Property (b):** If  $\mathbf{u}$  and  $\mathbf{v}$  are in  $\text{Nul}(A)$ , show that  $\mathbf{u} + \mathbf{v}$  is also in  $\text{Nul}(A)$ . Suppose  $A\mathbf{u} = \mathbf{0}$  and  $A\mathbf{v} = \mathbf{0}$ . Then

$$A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v} = \mathbf{0} + \mathbf{0} = \mathbf{0}.$$

**Property (c):** If  $\mathbf{u}$  is in  $\text{Nul}(A)$  and  $c$  is a scalar, show that  $c\mathbf{u}$  is also in  $\text{Nul}(A)$ . Suppose  $A\mathbf{u} = \mathbf{0}$ . Then



# Null Spaces

**Property (b):** If  $\mathbf{u}$  and  $\mathbf{v}$  are in  $\text{Nul}(A)$ , show that  $\mathbf{u} + \mathbf{v}$  is also in  $\text{Nul}(A)$ . Suppose  $A\mathbf{u} = \mathbf{0}$  and  $A\mathbf{v} = \mathbf{0}$ . Then

$$A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v} = \mathbf{0} + \mathbf{0} = \mathbf{0}.$$

**Property (c):** If  $\mathbf{u}$  is in  $\text{Nul}(A)$  and  $c$  is a scalar, show that  $c\mathbf{u}$  is also in  $\text{Nul}(A)$ . Suppose  $A\mathbf{u} = \mathbf{0}$ . Then

$$A(c\mathbf{u}) = c(A\mathbf{u}) = c(\mathbf{0}) = \mathbf{0}.$$

# Null Spaces

Let's restate the theorem.

## Theorem

*The null space of an  $m \times n$  matrix  $A$  is a subspace of  $\mathbb{R}^n$ .  
Equivalently, the set of all solutions to the system  $A\mathbf{x} = \mathbf{0}$  of  $m$  homogeneous linear equations in  $n$  unknowns is a subspace of  $\mathbb{R}^n$ .*

# Null Spaces

Let's restate the theorem.

## Theorem

*The null space of an  $m \times n$  matrix  $A$  is a subspace of  $\mathbb{R}^n$ .  
Equivalently, the set of all solutions to the system  $A\mathbf{x} = \mathbf{0}$  of  $m$  homogeneous linear equations in  $n$  unknowns is a subspace of  $\mathbb{R}^n$ .*

## Remark

- Since properties (a), (b), and (c) hold,  $\text{Nul}(A)$  is a subspace of  $\mathbb{R}^n$ .
- Since  $\text{Nul}(A)$  is a subspace, it is closed under linear combinations.

# Null Spaces

Let's restate the theorem.

## Theorem

*The null space of an  $m \times n$  matrix  $A$  is a subspace of  $\mathbb{R}^n$ .  
Equivalently, the set of all solutions to the system  $A\mathbf{x} = \mathbf{0}$  of  $m$  homogeneous linear equations in  $n$  unknowns is a subspace of  $\mathbb{R}^n$ .*

## Remark

- Since properties (a), (b), and (c) hold,  $\text{Nul}(A)$  is a subspace of  $\mathbb{R}^n$ .
- Since  $\text{Nul}(A)$  is a subspace, it is closed under linear combinations. You can add solutions of  $A\mathbf{x} = \mathbf{0}$  and get a new solution! This is very important.

# Null Spaces

Let's restate the theorem.

## Theorem

*The null space of an  $m \times n$  matrix  $A$  is a subspace of  $\mathbb{R}^n$ . Equivalently, the set of all solutions to the system  $A\mathbf{x} = \mathbf{0}$  of  $m$  homogeneous linear equations in  $n$  unknowns is a subspace of  $\mathbb{R}^n$ .*

## Remark

- Since properties (a), (b), and (c) hold,  $\text{Nul}(A)$  is a subspace of  $\mathbb{R}^n$ .
- Since  $\text{Nul}(A)$  is a subspace, it is closed under linear combinations. You can add solutions of  $A\mathbf{x} = \mathbf{0}$  and get a new solution! This is very important. Not true for  $A\mathbf{x} = \mathbf{b}$  for  $\mathbf{b} \neq \mathbf{0}$ . Here you cannot add solutions!

# Null Spaces

Let's restate the theorem.

## Theorem

*The null space of an  $m \times n$  matrix  $A$  is a subspace of  $\mathbb{R}^n$ .  
Equivalently, the set of all solutions to the system  $A\mathbf{x} = \mathbf{0}$  of  $m$  homogeneous linear equations in  $n$  unknowns is a subspace of  $\mathbb{R}^n$ .*

## Remark

- Since properties (a), (b), and (c) hold,  $\text{Nul}(A)$  is a subspace of  $\mathbb{R}^n$ .
- Since  $\text{Nul}(A)$  is a subspace, it is closed under linear combinations. You can add solutions of  $A\mathbf{x} = \mathbf{0}$  and get a new solution! This is very important. Not true for  $A\mathbf{x} = \mathbf{b}$  for  $b \neq 0$ . Here you cannot add solutions!
- Solving  $A\mathbf{x} = \mathbf{0}$  yields an explicit description of  $\text{Nul}(A)$ .

# Null Spaces

## Example

Find and explicit description of  $\text{Nul}(A)$  where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

# Null Spaces

## Example

Find an explicit description of  $\text{Nul}(A)$  where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

## Solution

We want to find all the solutions to  $A\mathbf{x} = \mathbf{0}$ .



# Null Spaces

## Example

Find an explicit description of  $\text{Nul}(A)$  where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

## Solution

We want to find all the solutions to  $A\mathbf{x} = \mathbf{0}$ . So we need to do Gaussian elimination on the augmented matrix  $[A \mid \mathbf{0}]$ .

# Null Spaces

## Example

Find an explicit description of  $\text{Nul}(A)$  where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

## Solution

We want to find all the solutions to  $A\mathbf{x} = \mathbf{0}$ . So we need to do Gaussian elimination on the augmented matrix  $[A \mid \mathbf{0}]$ .

$$[A \mid \mathbf{0}] =$$

# Null Spaces

## Example

Find an explicit description of  $\text{Nul}(A)$  where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

## Solution

We want to find all the solutions to  $A\mathbf{x} = \mathbf{0}$ . So we need to do Gaussian elimination on the augmented matrix  $[A \mid \mathbf{0}]$ .

$$[A \mid \mathbf{0}] = \left[ \begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 6 & 12 & 13 & 0 & 3 & 0 \end{array} \right]$$

# Null Spaces

## Example

Find an explicit description of  $\text{Nul}(A)$  where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

## Solution

We want to find all the solutions to  $A\mathbf{x} = \mathbf{0}$ . So we need to do Gaussian elimination on the augmented matrix  $[A \mid \mathbf{0}]$ .

$$[A \mid \mathbf{0}] = \left[ \begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 6 & 12 & 13 & 0 & 3 & 0 \end{array} \right] \longrightarrow \left[ \begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right]$$

# Null Spaces

## Example

Find an explicit description of  $\text{Nul}(A)$  where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

## Solution

We want to find all the solutions to  $A\mathbf{x} = \mathbf{0}$ . So we need to do Gaussian elimination on the augmented matrix  $[A \mid \mathbf{0}]$ .

$$[A \mid \mathbf{0}] = \left[ \begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 6 & 12 & 13 & 0 & 3 & 0 \end{array} \right] \longrightarrow \left[ \begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right] \longrightarrow$$

$$\left[ \begin{array}{ccccc|c} 3 & 6 & 0 & 39 & 99 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right]$$

# Null Spaces

## Example

Find an explicit description of  $\text{Nul}(A)$  where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

## Solution

We want to find all the solutions to  $A\mathbf{x} = \mathbf{0}$ . So we need to do Gaussian elimination on the augmented matrix  $[A \mid \mathbf{0}]$ .

$$[A \mid \mathbf{0}] = \left[ \begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 6 & 12 & 13 & 0 & 3 & 0 \end{array} \right] \longrightarrow \left[ \begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right] \longrightarrow$$

$$\left[ \begin{array}{ccccc|c} 3 & 6 & 0 & 39 & 99 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right] \longrightarrow \left[ \begin{array}{ccccc|c} 1 & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right] =$$

# Null Spaces

## Example

Find an explicit description of  $\text{Nul}(A)$  where

$$A = \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix}.$$

## Solution

We want to find all the solutions to  $A\mathbf{x} = \mathbf{0}$ . So we need to do Gaussian elimination on the augmented matrix  $[A \mid \mathbf{0}]$ .

$$[A \mid \mathbf{0}] = \left[ \begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 6 & 12 & 13 & 0 & 3 & 0 \end{array} \right] \longrightarrow \left[ \begin{array}{ccccc|c} 3 & 6 & 6 & 3 & 9 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right] \longrightarrow$$

$$\left[ \begin{array}{ccccc|c} 3 & 6 & 0 & 39 & 99 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right] \longrightarrow \left[ \begin{array}{ccccc|c} 1 & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right] = [U \mid \mathbf{0}].$$

# Null Spaces

$$[U \mid \mathbf{0}] = \left[ \begin{array}{ccccc|c} 1 & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right].$$

This corresponds to the solution:



# Null Spaces

$$[U \mid \mathbf{0}] = \left[ \begin{array}{ccccc|c} 1 & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right].$$

This corresponds to the solution:

$$x_1 = -2x_2 - 13x_4 - 33x_5$$

$$x_3 = 6x_4 + 15x_5.$$

# Null Spaces

$$[U \mid \mathbf{0}] = \left[ \begin{array}{ccccc|c} 1 & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right].$$

This corresponds to the solution:

$$x_1 = -2x_2 - 13x_4 - 33x_5$$

$$x_3 = 6x_4 + 15x_5.$$

Write this as a linear combination:

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} =$$

# Null Spaces

$$[U \mid \mathbf{0}] = \left[ \begin{array}{ccccc|c} 1 & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right].$$

This corresponds to the solution:

$$x_1 = -2x_2 - 13x_4 - 33x_5$$

$$x_3 = 6x_4 + 15x_5.$$

Write this as a linear combination:

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} -2x_2 - 13x_4 - 33x_5 \\ x_2 \\ 6x_4 + 15x_5 \\ x_4 \\ x_5 \end{bmatrix} =$$

# Null Spaces

$$[U \mid \mathbf{0}] = \left[ \begin{array}{ccccc|c} 1 & 2 & 0 & 13 & 33 & 0 \\ 0 & 0 & 1 & -6 & -15 & 0 \end{array} \right].$$

This corresponds to the solution:

$$x_1 = -2x_2 - 13x_4 - 33x_5$$

$$x_3 = 6x_4 + 15x_5.$$

Write this as a linear combination:

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} -2x_2 - 13x_4 - 33x_5 \\ x_2 \\ 6x_4 + 15x_5 \\ x_4 \\ x_5 \end{bmatrix} = x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix}.$$

# Null Spaces

So each vector in  $\text{Nul}(A)$  looks like:

$$x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix} .$$

# Null Spaces

So each vector in  $\text{Nul}(A)$  looks like:

$$x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix}.$$

Thus,

$$\text{Nul}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

In other words,

$$\text{Nul} \left( \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix} \right) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

In other words,

$$\text{Nul} \left( \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix} \right) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

### Remark

If  $\text{Nul}(A) \neq \{\mathbf{0}\}$ , then the number of vectors in the spanning set for  $\text{Nul}(A)$  equals the number of free variables in  $A\mathbf{x} = \mathbf{0}$ .



In other words,

$$\text{Nul} \left( \begin{bmatrix} 3 & 6 & 6 & 3 & 9 \\ 6 & 12 & 13 & 0 & 3 \end{bmatrix} \right) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -13 \\ 0 \\ 6 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -33 \\ 0 \\ 15 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

### Remark

If  $\text{Nul}(A) \neq \{\mathbf{0}\}$ , then the number of vectors in the spanning set for  $\text{Nul}(A)$  equals the number of free variables in  $A\mathbf{x} = \mathbf{0}$ .

In this example, we had **3 free variables** ( $x_2$ ,  $x_4$ , and  $x_5$ ) so there were **3 vectors** in the spanning set for  $\text{Nul}(A)$ . More about this later!