

Math 415 - Lecture 24

Least squares

Wednesday October 21st 2015

Textbook reading: Chapter 3.3

Textbook reading: Chapter 3.3

Suggested practice exercises: Exercises 3, 5, 6, 13, 24, 25

Textbook reading: Chapter 3.3


Suggested practice exercises: Exercises 3, 5, 6, 13, 24, 25

Khan Academy video: Least Squares Approximation, Least Squares Examples, Another Least Squares Example


Review

Let W be a subspace of \mathbb{R}^n and \mathbf{x} in \mathbb{R}^n (but maybe not in W).

Let W be a subspace of \mathbb{R}^n and \mathbf{x} in \mathbb{R}^n (but maybe not in W).




Let W be a subspace of \mathbb{R}^n and \mathbf{x} in \mathbb{R}^n (but maybe not in W).




Let \mathbf{x}_W be the orthonormal projection of \mathbf{x} onto W .

Let W be a subspace of \mathbb{R}^n and \mathbf{x} in \mathbb{R}^n (but maybe not in W).



Let \mathbf{x}_W be the orthonormal projection of \mathbf{x} onto W .
(vector in W as close as possible to \mathbf{x})


Let W be a subspace of \mathbb{R}^n and \mathbf{x} in \mathbb{R}^n (but maybe not in W).



Let \mathbf{x}_W be the orthonormal projection of \mathbf{x} onto W .
(vector in W as close as possible to \mathbf{x})

- If $\mathbf{v}_1, \dots, \mathbf{v}_m$ is an orthogonal basis of W

Let W be a subspace of \mathbb{R}^n and \mathbf{x} in \mathbb{R}^n (but maybe not in W).




Let \mathbf{x}_W be the orthonormal projection of \mathbf{x} onto W .
(vector in W as close as possible to \mathbf{x})

- If $\mathbf{v}_1, \dots, \mathbf{v}_m$ is an orthogonal basis of W then

$$\mathbf{x}_W = \underbrace{\left(\frac{\mathbf{x} \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \right) \mathbf{v}_1}_{\text{proj. of } \mathbf{x} \text{ onto } \mathbf{v}_1} + \dots + \underbrace{\left(\frac{\mathbf{x} \cdot \mathbf{v}_m}{\mathbf{v}_m \cdot \mathbf{v}_m} \right) \mathbf{v}_m}_{\text{proj. of } \mathbf{x} \text{ onto } \mathbf{v}_m}.$$

Let W be a subspace of \mathbb{R}^n and \mathbf{x} in \mathbb{R}^n (but maybe not in W).



Let \mathbf{x}_W be the orthonormal projection of \mathbf{x} onto W .
(vector in W as close as possible to \mathbf{x})

- If $\mathbf{v}_1, \dots, \mathbf{v}_m$ is an orthogonal basis of W then

$$\mathbf{x}_W = \underbrace{\left(\frac{\mathbf{x} \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \right) \mathbf{v}_1}_{\text{proj. of } \mathbf{x} \text{ onto } \mathbf{v}_1} + \dots + \underbrace{\left(\frac{\mathbf{x} \cdot \mathbf{v}_m}{\mathbf{v}_m \cdot \mathbf{v}_m} \right) \mathbf{v}_m}_{\text{proj. of } \mathbf{x} \text{ onto } \mathbf{v}_m}.$$

- The decomposition $\mathbf{x} = \underbrace{\mathbf{x}_W}_{\text{in } W} + \underbrace{\mathbf{x}^\perp}_{\text{in } W^\perp}$ is unique.

Least squares

Least squares

Definition


$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$ if $\hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

Definition

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$ if $\hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.


Definition

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$ if $\hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.




Definition

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$ if $\hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.



Definition


$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$ if $\hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.



- If $A\mathbf{x} = \mathbf{b}$ is consistent, then a least squares solution $\hat{\mathbf{x}}$ is just an ordinary solution.
(in that case, $A\hat{\mathbf{x}} - \mathbf{b} = 0$)

Definition


$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$ if $\hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.



- If $A\mathbf{x} = \mathbf{b}$ is consistent, then a least squares solution $\hat{\mathbf{x}}$ is just an ordinary solution.
(in that case, $A\hat{\mathbf{x}} - \mathbf{b} = 0$)
- Interesting case: $A\mathbf{x} = \mathbf{b}$ is inconsistent.
(in other words: the system is overdetermined)

Definition

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$ if $\hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.




- If $A\mathbf{x} = \mathbf{b}$ is consistent, then a least squares solution $\hat{\mathbf{x}}$ is just an ordinary solution.
(in that case, $A\hat{\mathbf{x}} - \mathbf{b} = 0$)
- Interesting case: $A\mathbf{x} = \mathbf{b}$ is inconsistent.
(in other words: the system is overdetermined)

Idea

$A\mathbf{x} = \mathbf{b}$ is consistent $\iff \mathbf{b}$ is in $\text{Col}(A)$

Definition

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$ if $\hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.




- If $A\mathbf{x} = \mathbf{b}$ is consistent, then a least squares solution $\hat{\mathbf{x}}$ is just an ordinary solution.
(in that case, $A\hat{\mathbf{x}} - \mathbf{b} = 0$)
- Interesting case: $A\mathbf{x} = \mathbf{b}$ is inconsistent.
(in other words: the system is overdetermined)

Idea

$A\mathbf{x} = \mathbf{b}$ is consistent $\iff \mathbf{b}$ is in $\text{Col}(A)$

Definition

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$ if $\hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.



- If $A\mathbf{x} = \mathbf{b}$ is consistent, then a least squares solution $\hat{\mathbf{x}}$ is just an ordinary solution.
(in that case, $A\hat{\mathbf{x}} - \mathbf{b} = 0$)
- Interesting case: $A\mathbf{x} = \mathbf{b}$ is inconsistent.
(in other words: the system is overdetermined)

Idea


$A\mathbf{x} = \mathbf{b}$ is consistent $\iff \mathbf{b}$ is in $\text{Col}(A)$

So if $A\mathbf{x} = \mathbf{b}$ is inconsistent we

- replace \mathbf{b} with its projection $\hat{\mathbf{b}}$ onto $\text{Col}(A)$,

Definition

$\hat{\mathbf{x}}$ is a **least squares solution** of the system $A\mathbf{x} = \mathbf{b}$ if $\hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.



- If $A\mathbf{x} = \mathbf{b}$ is consistent, then a least squares solution $\hat{\mathbf{x}}$ is just an ordinary solution.
(in that case, $A\hat{\mathbf{x}} - \mathbf{b} = 0$)
- Interesting case: $A\mathbf{x} = \mathbf{b}$ is inconsistent.
(in other words: the system is overdetermined)

Idea

$$A\mathbf{x} = \mathbf{b} \text{ is consistent} \iff \mathbf{b} \text{ is in } \text{Col}(A)$$

So if $A\mathbf{x} = \mathbf{b}$ is inconsistent we

- replace \mathbf{b} with its projection $\hat{\mathbf{b}}$ onto $\text{Col}(A)$,
- and solve $A\hat{\mathbf{x}} = \hat{\mathbf{b}}$. (consistent by construction!)

Example

Find the least squares solution to $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}.$$

Solution

Note the columns of A are orthogonal.

Solution

Note the columns of A are orthogonal.

Otherwise, we could not proceed in the same way.

Solution

Note the columns of A are orthogonal.

Otherwise, we could not proceed in the same way.

Hence the projection of $\hat{\mathbf{b}}$ of \mathbf{b} onto $\text{Col}(A)$ is

Solution

Note the columns of A are orthogonal.

Otherwise, we could not proceed in the same way.

Hence the projection of $\hat{\mathbf{b}}$ of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

Solution

Note the columns of A are orthogonal.

Otherwise, we could not proceed in the same way.

Hence the projection of $\hat{\mathbf{b}}$ of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{3}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$$

Solution

Note the columns of A are orthogonal.

Otherwise, we could not proceed in the same way.

Hence the projection of $\hat{\mathbf{b}}$ of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{3}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}.$$

Solution

Note the columns of A are orthogonal.

Otherwise, we could not proceed in the same way.

Hence the projection of $\hat{\mathbf{b}}$ of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{3}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}.$$

We have already solved $A\hat{\mathbf{x}} = \hat{\mathbf{b}}$ in the process:

Solution

Note the columns of A are orthogonal.

Otherwise, we could not proceed in the same way.

Hence the projection of $\hat{\mathbf{b}}$ of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{3}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}.$$

We have already solved $A\hat{\mathbf{x}} = \hat{\mathbf{b}}$ in the process: $\hat{\mathbf{x}} = \begin{bmatrix} 1/2 \\ 3/2 \end{bmatrix}$.

Solution

Note the columns of A are orthogonal.

Otherwise, we could not proceed in the same way.

Hence the projection of $\hat{\mathbf{b}}$ of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{\begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}}{\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} + \frac{3}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}.$$

We have already solved $A\hat{\mathbf{x}} = \hat{\mathbf{b}}$ in the process: $\hat{\mathbf{x}} = \begin{bmatrix} 1/2 \\ 3/2 \end{bmatrix}$.

Question: What to do when the columns of A are not orthogonal?

The normal equations

Theorem

$\hat{\mathbf{x}}$ is a least squares solution of $A\mathbf{x} = \mathbf{b}$

Theorem

$\hat{\mathbf{x}}$ is a least squares solution of $A\mathbf{x} = \mathbf{b} \iff A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$

Theorem

$\hat{\mathbf{x}}$ is a least squares solution of $A\mathbf{x} = \mathbf{b} \iff A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$

Proof.

$\iff A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible

Theorem

$\hat{\mathbf{x}}$ is a least squares solution of $A\mathbf{x} = \mathbf{b} \iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$

Proof.

$\iff A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible

$\iff A\hat{\mathbf{x}} - \mathbf{b}$ is orthogonal to $\text{Col}(A)$

Theorem

$\hat{\mathbf{x}}$ is a least squares solution of $A\mathbf{x} = \mathbf{b} \iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$

Proof.

$\iff A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible

$\iff A\hat{\mathbf{x}} - \mathbf{b}$ is orthogonal to $\text{Col}(A)$

$\overset{\text{FTLA}}{\iff} A\hat{\mathbf{x}} - \mathbf{b}$ is in $\text{Nul}(A^T)$

Theorem

$\hat{\mathbf{x}}$ is a least squares solution of $A\mathbf{x} = \mathbf{b} \iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$

Proof.

$\iff A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible

$\iff A\hat{\mathbf{x}} - \mathbf{b}$ is orthogonal to $\text{Col}(A)$

$\xLeftrightarrow{FTLA} A\hat{\mathbf{x}} - \mathbf{b}$ is in $\text{Nul}(A^T)$

$\iff A^T(A\hat{\mathbf{x}} - \mathbf{b}) = \mathbf{0}$

Theorem

$\hat{\mathbf{x}}$ is a least squares solution of $A\mathbf{x} = \mathbf{b} \iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$

Proof.

$\iff A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible

$\iff A\hat{\mathbf{x}} - \mathbf{b}$ is orthogonal to $\text{Col}(A)$

$\xLeftrightarrow{FTLA} A\hat{\mathbf{x}} - \mathbf{b}$ is in $\text{Nul}(A^T)$

$\iff A^T(A\hat{\mathbf{x}} - \mathbf{b}) = \mathbf{0}$

$\iff A^T A\hat{\mathbf{x}} = A^T \mathbf{b}$



Example (again)

Find the least squares solution to $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}.$$

Solution

Solution

$$A^T A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix}$$

Solution

$$A^T A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

Solution

$$A^T A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}$$

Solution

$$A^T A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

Solution

$$A^T A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

The normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ are

Solution

$$A^T A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

The normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ are $\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$.

Solution

$$A^T A = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ -1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 1 & -1 & 0 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

The normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ are $\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$.

Solving, we find (again) $\hat{\mathbf{x}} = \begin{bmatrix} 1/2 \\ 3/2 \end{bmatrix}$.

Example

Find the least squares solution to $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}.$$

What is the projection of \mathbf{b} onto $\text{Col}(A)$?

Example

Find the least squares solution to $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}.$$

What is the projection of \mathbf{b} onto $\text{Col}(A)$?

Example

Find the least squares solution to $A\mathbf{x} = \mathbf{b}$, where

$$A = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix}, \mathbf{b} = \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}.$$

What is the projection of \mathbf{b} onto $\text{Col}(A)$?

Note that the columns of A are not orthogonal.

Solution

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix}$$

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$$

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix}$$

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$$

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$$

The normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ are

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$$

The normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ are $\begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$.

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$$

The normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ are $\begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$.

Solving, we find $\hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$$

The normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ are $\begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$.

Solving, we find $\hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

The projection of \mathbf{b} onto $\text{Col}(A)$ is

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$$

The normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ are $\begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$.

Solving, we find $\hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

The projection of \mathbf{b} onto $\text{Col}(A)$ is $A\hat{\mathbf{x}}$

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$$

The normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ are $\begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$.

Solving, we find $\hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

The projection of \mathbf{b} onto $\text{Col}(A)$ is $A \hat{\mathbf{x}} = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$

Solution

$$A^T A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix}$$

$$A^T \mathbf{b} = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 11 \end{bmatrix} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$$

The normal equations $A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ are $\begin{bmatrix} 17 & 1 \\ 1 & 5 \end{bmatrix} \hat{\mathbf{x}} = \begin{bmatrix} 19 \\ 11 \end{bmatrix}$.

Solving, we find $\hat{\mathbf{x}} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

The projection of \mathbf{b} onto $\text{Col}(A)$ is $A \hat{\mathbf{x}} = \begin{bmatrix} 4 & 0 \\ 0 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 3 \end{bmatrix}$.

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?
Because, for a least square solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?
Because, for a least square solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

The projection of \mathbf{b} onto $\text{Col}(A)$ is

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?
Because, for a least square solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

The projection of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = A\hat{\mathbf{x}},$$

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?
Because, for a least square solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

The projection of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = A\hat{\mathbf{x}}, \text{ with } \hat{\mathbf{x}} \text{ such that } A^T A\hat{\mathbf{x}} = A^T \mathbf{b}.$$

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?
Because, for a least square solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

The projection of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = A\hat{\mathbf{x}}, \text{ with } \hat{\mathbf{x}} \text{ such that } A^T A\hat{\mathbf{x}} = A^T \mathbf{b}.$$

If A has full column rank,

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?
Because, for a least square solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

The projection of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = A\hat{\mathbf{x}}, \text{ with } \hat{\mathbf{x}} \text{ such that } A^T A\hat{\mathbf{x}} = A^T \mathbf{b}.$$

If A has full column rank, (so the columns of A are independent,)

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?
Because, for a least square solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

The projection of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = A\hat{\mathbf{x}}, \text{ with } \hat{\mathbf{x}} \text{ such that } A^T A\hat{\mathbf{x}} = A^T \mathbf{b}.$$

If A has full column rank, (so the columns of A are independent,) this is

$$\hat{\mathbf{b}} = A(A^T A)^{-1} A^T \mathbf{b}.$$

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?
Because, for a least square solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

The projection of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = A\hat{\mathbf{x}}, \text{ with } \hat{\mathbf{x}} \text{ such that } A^T A\hat{\mathbf{x}} = A^T \mathbf{b}.$$

If A has full column rank, (so the columns of A are independent,) this is

$$\hat{\mathbf{b}} = A(A^T A)^{-1} A^T \mathbf{b}.$$

(In this case $A^T A$ is invertible.)

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?
Because, for a least square solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

The projection of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = A\hat{\mathbf{x}}, \text{ with } \hat{\mathbf{x}} \text{ such that } A^T A\hat{\mathbf{x}} = A^T \mathbf{b}.$$

If A has full column rank, (so the columns of A are independent,) this is

$$\hat{\mathbf{b}} = A(A^T A)^{-1} A^T \mathbf{b}.$$

(In this case $A^T A$ is invertible.)

Hence, the projection matrix for projecting onto $\text{Col}(A)$ is

Just to make sure: why is $A\hat{\mathbf{x}}$ the projection of \mathbf{b} onto $\text{Col}(A)$?
Because, for a least square solution $\hat{\mathbf{x}}$, $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible.

The projection of \mathbf{b} onto $\text{Col}(A)$ is

$$\hat{\mathbf{b}} = A\hat{\mathbf{x}}, \text{ with } \hat{\mathbf{x}} \text{ such that } A^T A\hat{\mathbf{x}} = A^T \mathbf{b}.$$

If A has full column rank, (so the columns of A are independent,) this is

$$\hat{\mathbf{b}} = A(A^T A)^{-1} A^T \mathbf{b}.$$

(In this case $A^T A$ is invertible.)

Hence, the projection matrix for projecting onto $\text{Col}(A)$ is

$$P = A(A^T A)^{-1} A^T.$$

Applications

Least square regression lines

Experimental data: (x_i, y_i) , for $i = 1, 2, 3, \dots$


Experimental data: (x_i, y_i) , for $i = 1, 2, 3, \dots$

Wanted: parameters β_1, β_2 such that $y_i \approx \beta_1 + \beta_2 x_i$ for all i

Least square regression lines

Experimental data: (x_i, y_i) , for $i = 1, 2, 3, \dots$


Wanted: parameters β_1, β_2 such that $y_i \approx \beta_1 + \beta_2 x_i$ for all i



Least square regression lines

Experimental data: (x_i, y_i) , for $i = 1, 2, 3, \dots$

Wanted: parameters β_1, β_2 such that $y_i \approx \beta_1 + \beta_2 x_i$ for all i




The approximation should be so that

Least square regression lines

Experimental data: (x_i, y_i) , for $i = 1, 2, 3, \dots$

Wanted: parameters β_1, β_2 such that $y_i \approx \beta_1 + \beta_2 x_i$ for all i




The approximation should be so that

$$SS_{\text{res}} = \underbrace{\sum_i [y_i - (\beta_1 + \beta_2 x_i)]^2}_{\text{residue sum of squares}} \text{ is as small as possible.}$$

Least square regression lines

Experimental data: (x_i, y_i) , for $i = 1, 2, 3, \dots$

Wanted: parameters β_1, β_2 such that $y_i \approx \beta_1 + \beta_2 x_i$ for all i



The approximation should be so that

$$SS_{\text{res}} = \underbrace{\sum_i [y_i - (\beta_1 + \beta_2 x_i)]^2}_{\text{residue sum of squares}} \text{ is as small as possible.}$$

Example

Find β_1, β_2 such that the line $y = \beta_1 + \beta_2 x$ best fits the data points $(2, 1), (5, 2), (7, 3), (8, 3)$.

Solution

The equations $y = \beta_1 + \beta_2 x$ in matrix form:

Solution

The equations $y = \beta_1 + \beta_2 x$ in matrix form:

$$\underbrace{\begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \\ 1 & x_4 \end{bmatrix}}_{\text{design matrix } X} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}}_{\text{observation vector } y}$$

Solution

The equations $y = \beta_1 + \beta_2 x$ in matrix form:

$$\underbrace{\begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \\ 1 & x_4 \end{bmatrix}}_{\text{design matrix } X} \underbrace{\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix}}_{\text{observation vector } y} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}}_{\text{observation vector } y}$$

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

Calculate $X^T X$ and $X^T \mathbf{y}$ to get the normal equation:

Calculate $X^T X$ and $X^T \mathbf{y}$ to get the normal equation:

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix}$$

Calculate $X^T X$ and $X^T \mathbf{y}$ to get the normal equation:

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

Calculate $X^T X$ and $X^T \mathbf{y}$ to get the normal equation:

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}$$

Calculate $X^T X$ and $X^T \mathbf{y}$ to get the normal equation:

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Calculate $X^T X$ and $X^T \mathbf{y}$ to get the normal equation:

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solving $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 9 \\ 57 \end{bmatrix},$

Calculate $X^T X$ and $X^T \mathbf{y}$ to get the normal equation:

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solving $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$, we find $\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}$.

Calculate $X^T X$ and $X^T \mathbf{y}$ to get the normal equation:

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solving $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$, we find $\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}$.

Hence the least squares line is $y = \frac{2}{7} + \frac{5}{14}x$.

Example

Blood is drawn from volunteers to determine the effects of a new experimental drug designed to lower cholesterol levels.

Example

Blood is drawn from volunteers to determine the effects of a new experimental drug designed to lower cholesterol levels. The following data shows the results of varying the dosage from 0 unit to 1 units in step of 0.2 of a unit.

Example

Blood is drawn from volunteers to determine the effects of a new experimental drug designed to lower cholesterol levels. The following data shows the results of varying the dosage from 0 unit to 1 units in step of 0.2 of a unit. Find a line $C = \beta_1 D + \beta_2$ that best fits the data. What drug usage would you recommend if you want to accomplish a Cholesterol level of 215?

| | | | | | | |
|----------------|-----|-----|-----|-----|-----|-----|
| Drug Dosage: D | 0.0 | 0.2 | 0.4 | 0.6 | 0.8 | 1 |
| Cholesterol: C | 289 | 273 | 254 | 226 | 213 | 189 |

Example

Blood is drawn from volunteers to determine the effects of a new experimental drug designed to lower cholesterol levels. The following data shows the results of varying the dosage from 0 unit to 1 units in step of 0.2 of a unit. Find a line $C = \beta_1 D + \beta_2$ that best fits the data. What drug usage would you recommend if you want to accomplish a Cholesterol level of 215?

| | | | | | | |
|----------------|-----|-----|-----|-----|-----|-----|
| Drug Dosage: D | 0.0 | 0.2 | 0.4 | 0.6 | 0.8 | 1 |
| Cholesterol: C | 289 | 273 | 254 | 226 | 213 | 189 |

Solution

$$\underbrace{\begin{bmatrix} D_1 & 1 \\ D_2 & 1 \\ D_3 & 1 \\ D_4 & 1 \\ D_5 & 1 \\ D_6 & 1 \end{bmatrix}}_{\text{design matrix } D} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \underbrace{\begin{bmatrix} C_1 \\ C_2 \\ C_3 \\ C_4 \\ C_5 \\ C_6 \end{bmatrix}}_{\text{observation vector } \mathbf{c}}$$

Here, we need to find a least squares solution to $D\beta = c$ or

$$\begin{bmatrix} 0 & 1 \\ 0.2 & 1 \\ 0.4 & 1 \\ 0.6 & 1 \\ 0.8 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 289 \\ 273 \\ 254 \\ 226 \\ 213 \\ 189 \end{bmatrix}.$$

Calculate $D^T D$ and $D^T \mathbf{c}$ to get the normal equation:

Here, we need to find a least squares solution to $D\beta = c$ or

$$\begin{bmatrix} 0 & 1 \\ 0.2 & 1 \\ 0.4 & 1 \\ 0.6 & 1 \\ 0.8 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 289 \\ 273 \\ 254 \\ 226 \\ 213 \\ 189 \end{bmatrix}.$$

Calculate $D^T D$ and $D^T \mathbf{c}$ to get the normal equation:

$$D^T D = \begin{bmatrix} 0 & .2 & .4 & .6 & .8 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ .2 & 1 \\ .4 & 1 \\ .6 & 1 \\ .8 & 1 \\ 1 & 1 \end{bmatrix}$$

Here, we need to find a least squares solution to $D\beta = c$ or

$$\begin{bmatrix} 0 & 1 \\ 0.2 & 1 \\ 0.4 & 1 \\ 0.6 & 1 \\ 0.8 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 289 \\ 273 \\ 254 \\ 226 \\ 213 \\ 189 \end{bmatrix}.$$

Calculate $D^T D$ and $D^T \mathbf{c}$ to get the normal equation:

$$D^T D = \begin{bmatrix} 0 & .2 & .4 & .6 & .8 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ .2 & 1 \\ .4 & 1 \\ .6 & 1 \\ .8 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 2.2 & 3 \\ 3 & 6 \end{bmatrix}$$

$$D^T \mathbf{c} = \begin{bmatrix} 0 & .2 & .4 & .6 & .8 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 289 \\ 273 \\ 254 \\ 226 \\ 213 \\ 189 \end{bmatrix}$$

$$D^T \mathbf{c} = \begin{bmatrix} 0 & .2 & .4 & .6 & .8 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 289 \\ 273 \\ 254 \\ 226 \\ 213 \\ 189 \end{bmatrix} = \begin{bmatrix} 651.2 \\ 1444 \end{bmatrix}$$

$$D^T \mathbf{c} = \begin{bmatrix} 0 & .2 & .4 & .6 & .8 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 289 \\ 273 \\ 254 \\ 226 \\ 213 \\ 189 \end{bmatrix} = \begin{bmatrix} 651.2 \\ 1444 \end{bmatrix}$$

Solving $\begin{bmatrix} 2.2 & 3 \\ 3 & 6 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 651.2 \\ 1444 \end{bmatrix},$

$$D^T \mathbf{c} = \begin{bmatrix} 0 & .2 & .4 & .6 & .8 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 289 \\ 273 \\ 254 \\ 226 \\ 213 \\ 189 \end{bmatrix} = \begin{bmatrix} 651.2 \\ 1444 \end{bmatrix}$$

Solving $\begin{bmatrix} 2.2 & 3 \\ 3 & 6 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 651.2 \\ 1444 \end{bmatrix}$, we find $\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} -\frac{65708}{7} \\ \frac{14116}{21} \end{bmatrix}$.

$$D^T \mathbf{c} = \begin{bmatrix} 0 & .2 & .4 & .6 & .8 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 289 \\ 273 \\ 254 \\ 226 \\ 213 \\ 189 \end{bmatrix} = \begin{bmatrix} 651.2 \\ 1444 \end{bmatrix}$$

Solving $\begin{bmatrix} 2.2 & 3 \\ 3 & 6 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 651.2 \\ 1444 \end{bmatrix}$, we find $\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} -\frac{65708}{7} \\ \frac{14116}{21} \end{bmatrix}$.

Hence the least squares line is $c = -\frac{65708}{7}d + \frac{14116}{21}$.