

Math 415 - Lecture 25

Multiple linear regression, Gram Schmidt and Orthogonal matrices

Monday October 26th 2015

Textbook reading: Chapters 3.3,3.4

Textbook reading: Chapters 3.3,3.4

Suggested practice exercises: Chapter 3.3, 3,5,6,13,22,24,25,26
and Chapter 3.4, 10,11,13,14,16,26

Textbook reading: Chapters 3.3,3.4

Suggested practice exercises: Chapter 3.3, 3,5,6,13,22,24,25,26
and Chapter 3.4, 10,11,13,14,16,26

Khan Academy video: Another Least Squares Example,
Gram-Schmidt Example

Textbook reading: Chapters 3.3,3.4

Suggested practice exercises: Chapter 3.3, 3,5,6,13,22,24,25,26
and Chapter 3.4, 10,11,13,14,16,26

Khan Academy video: Another Least Squares Example,
Gram-Schmidt Example

Strang lecture: Orthogonal Matrices and Gram-Schmidt Process.

Review

$\hat{\mathbf{x}}$ is a least squares solution of the system $A\mathbf{x} = \mathbf{b}$

$\hat{\mathbf{x}}$ is a least squares solution of the system $A\mathbf{x} = \mathbf{b}$

$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible

$\hat{\mathbf{x}}$ is a least squares solution of the system $A\mathbf{x} = \mathbf{b}$


$\iff \hat{\mathbf{x}}$ is such that $A\hat{\mathbf{x}} - \mathbf{b}$ is as small as possible

$\xLeftrightarrow{FTLA} A^T A \hat{\mathbf{x}} = A^T \mathbf{b}$ (the **normal equations**)

Application: fitting data


Example

Find β_1, β_2 such that the line $y = \beta_1 + \beta_2 x$ best fits the data points $(2, 1), (5, 2), (7, 3), (8, 3)$.



Comment

As usual in practice, we are minimizing the (the sum of the squares of the) vertical offsets.



Solution

The equations $y = \beta_1 + \beta_2 x$ in matrix form:

The equations $y = \beta_1 + \beta_2 x$ in matrix form:

$$\underbrace{\begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \\ 1 & x_4 \end{bmatrix}}_{\text{design matrix } X} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}}_{\text{observation vector } \mathbf{y}}$$

The equations $y = \beta_1 + \beta_2 x$ in matrix form:

$$\underbrace{\begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \\ 1 & x_4 \end{bmatrix}}_{\text{design matrix } X} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}}_{\text{observation vector } \mathbf{y}}$$

Here, we need to find a least squares solution to

The equations $y = \beta_1 + \beta_2 x$ in matrix form:

$$\underbrace{\begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ 1 & x_3 \\ 1 & x_4 \end{bmatrix}}_{\text{design matrix } X} \underbrace{\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix}}_{\text{observation vector } \mathbf{y}} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix}}_{\text{observation vector } \mathbf{y}}$$

Here, we need to find a least squares solution to

$$\begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} \begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$$

$$X^T X =$$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} =$$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solving $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$, we find $\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} =$

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solving $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$, we find $\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}$.

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solving $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$, we find $\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}$.

Hence the least squares line is

$$X^T X = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 1 & 5 \\ 1 & 7 \\ 1 & 8 \end{bmatrix} = \begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix}$$

$$X^T \mathbf{y} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 5 & 7 & 8 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \\ 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$$

Solving $\begin{bmatrix} 4 & 22 \\ 22 & 142 \end{bmatrix} \hat{\beta} = \begin{bmatrix} 9 \\ 57 \end{bmatrix}$, we find $\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} = \begin{bmatrix} 2/7 \\ 5/14 \end{bmatrix}$.

Hence the least squares line is $y = \frac{2}{7} + \frac{5}{14}x$.

Fitting to other curves

What happens if the data just lie close to any line? We can also fit the experimental data using other curves. Try to find $\beta_1, \beta_2, \beta_3$ such that $y = \beta_1 + \beta_2 x + \beta_3 x^2$ fits the data.

What happens if the data just lie close to any line? We can also fit the experimental data using other curves. Try to find $\beta_1, \beta_2, \beta_3$ such that $y = \beta_1 + \beta_2 x + \beta_3 x^2$ fits the data. **To fit:**
 $y_i \approx \beta_1 + \beta_2 x_i + \beta_3 x_i^2$ with parameters $\beta_1, \beta_2, \beta_3$.

What happens if the data just lie close to any line? We can also fit the experimental data using other curves. Try to find $\beta_1, \beta_2, \beta_3$ such that $y = \beta_1 + \beta_2 x + \beta_3 x^2$ fits the data. **To fit:**
 $y_i \approx \beta_1 + \beta_2 x_i + \beta_3 x_i^2$ with parameters $\beta_1, \beta_2, \beta_3$.
The equations $y_i = \beta_1 + \beta_2 x_i + \beta_3 x_i^2$ in **matrix form**:

What happens if the data just lie close to any line? We can also fit the experimental data using other curves. Try to find $\beta_1, \beta_2, \beta_3$ such that $y = \beta_1 + \beta_2 x + \beta_3 x^2$ fits the data. **To fit:**

$y_i \approx \beta_1 + \beta_2 x_i + \beta_3 x_i^2$ with parameters $\beta_1, \beta_2, \beta_3$.

The equations $y_i = \beta_1 + \beta_2 x_i + \beta_3 x_i^2$ in **matrix form**:

$$\underbrace{\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ \vdots & \vdots & \vdots \end{bmatrix}}_{\text{design matrix } X} \begin{bmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{bmatrix} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ \vdots \end{bmatrix}}_{\text{observation vector } \mathbf{y}}$$

What happens if the data just lie close to any line? We can also fit the experimental data using other curves. Try to find $\beta_1, \beta_2, \beta_3$ such that $y = \beta_1 + \beta_2 x + \beta_3 x^2$ fits the data. **To fit:**

$y_i \approx \beta_1 + \beta_2 x_i + \beta_3 x_i^2$ with parameters $\beta_1, \beta_2, \beta_3$.

The equations $y_i = \beta_1 + \beta_2 x_i + \beta_3 x_i^2$ in **matrix form**:

$$\underbrace{\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \\ \vdots & \vdots & \vdots \end{bmatrix}}_{\text{design matrix } X} \underbrace{\begin{bmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{bmatrix}}_{\text{observation vector } \mathbf{y}} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ \vdots \end{bmatrix}}_{\text{observation vector } \mathbf{y}}$$

Given data (x_i, y_i) , we then find the least squares solution to $X\beta = \mathbf{y}$.

Multiple linear regression

Of course, sometimes the variable y might not just depend on a single variable x , but on two variables, say u and v . So, here you have find the least-squares solution of


$$\underbrace{\begin{bmatrix} 1 & v_1 & w_1 \\ 1 & v_2 & w_2 \\ 1 & v_3 & w_3 \\ \vdots & \vdots & \vdots \end{bmatrix}}_{\text{design matrix}} \begin{bmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{bmatrix} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ \vdots \end{bmatrix}}_{\text{observation vector}}$$

Of course, sometimes the variable y might not just depend on a single variable x , but on two variables, say u and v . So, here you have find the least-squares solution of


$$\underbrace{\begin{bmatrix} 1 & v_1 & w_1 \\ 1 & v_2 & w_2 \\ 1 & v_3 & w_3 \\ \vdots & \vdots & \vdots \end{bmatrix}}_{\text{design matrix}} \begin{bmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{bmatrix} = \underbrace{\begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ \vdots \end{bmatrix}}_{\text{observation vector}}$$

And we again proceed by finding a least squares solution.

Review




Suppose $\mathbf{v}_1, \dots, \mathbf{v}_m$ is an orthonormal basis of W . The **orthogonal projection** of x onto W is :



Suppose $\mathbf{v}_1, \dots, \mathbf{v}_m$ is an orthonormal basis of W . The **orthogonal projection** of \mathbf{x} onto W is :

$$\hat{\mathbf{x}} = \underbrace{\langle \mathbf{x}, \mathbf{v}_1 \rangle \mathbf{v}_1}_{\text{proj. of } \mathbf{x} \text{ onto } \mathbf{v}_1} + \dots + \underbrace{\langle \mathbf{x}, \mathbf{v}_m \rangle \mathbf{v}_m}_{\text{proj. of } \mathbf{x} \text{ onto } \mathbf{v}_m}$$



Suppose $\mathbf{v}_1, \dots, \mathbf{v}_m$ is an orthonormal basis of W . The **orthogonal projection** of \mathbf{x} onto W is :

$$\hat{\mathbf{x}} = \underbrace{\langle \mathbf{x}, \mathbf{v}_1 \rangle \mathbf{v}_1}_{\text{proj. of } \mathbf{x} \text{ onto } \mathbf{v}_1} + \dots + \underbrace{\langle \mathbf{x}, \mathbf{v}_m \rangle \mathbf{v}_m}_{\text{proj. of } \mathbf{x} \text{ onto } \mathbf{v}_m}$$

(To stay agile, we are writing $\langle \mathbf{x}, \mathbf{v}_1 \rangle = \mathbf{x} \cdot \mathbf{v}_1$ for the inner product.)

Gram-Schmidt

Our goal

Our goal

- * In calculating projections we used an *orthogonal basis* and the easy formula for the coefficients.

- * In calculating projections we used an *orthogonal basis* and the easy formula for the coefficients.
- * What if we are given an arbitrary basis, not orthogonal?

- * In calculating projections we used an *orthogonal basis* and the easy formula for the coefficients.
- * What if we are given an arbitrary basis, not orthogonal?
- * Turn the starting basis into an orthogonal (or orthonormal) basis.

- * In calculating projections we used an *orthogonal basis* and the easy formula for the coefficients.
- * What if we are given an arbitrary basis, not orthogonal?
- * Turn the starting basis into an orthogonal (or orthonormal) basis.
- * **Gram-Schmidt Process.**

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\mathbf{b}_1 = \mathbf{a}_1,$$

$$\mathbf{q}_1 = \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|}$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\mathbf{b}_1 = \mathbf{a}_1,$$

$$\mathbf{b}_2 = \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1,$$

$$\mathbf{q}_1 = \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|}$$

$$\mathbf{q}_2 = \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|}$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\mathbf{b}_1 = \mathbf{a}_1,$$

$$\mathbf{b}_2 = \mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1,$$

$$\mathbf{b}_3 = \mathbf{a}_3 - \langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 - \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2,$$

$$\dots$$

$$\mathbf{q}_1 = \frac{\mathbf{b}_1}{\|\mathbf{b}_1\|}$$

$$\mathbf{q}_2 = \frac{\mathbf{b}_2}{\|\mathbf{b}_2\|}$$

$$\mathbf{q}_3 = \frac{\mathbf{b}_3}{\|\mathbf{b}_3\|}$$

$$\dots$$

Example

Find an orthonormal basis for $V = \text{Span}\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right\}$.

Solution

$$\mathbf{b}_1 =$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_2 =$$

$$\mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\mathbf{b}_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} - \left\langle \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix},$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} - \left\langle \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$
$$\mathbf{q}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} - \left\langle \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_3 =$$

$$\mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$
$$\mathbf{q}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} - \left\langle \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} -$$

$$\mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\mathbf{q}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 -$$

$$\mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\mathbf{q}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 - \left\langle \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{q}_2 \right\rangle \mathbf{q}_2 =$$

$$\mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\mathbf{q}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} - \left\langle \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 - \left\langle \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{q}_2 \right\rangle \mathbf{q}_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix},$$

$$\mathbf{q}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\mathbf{q}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

Solution

$$\mathbf{b}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix} - \left\langle \begin{bmatrix} 2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix},$$

$$\mathbf{b}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \left\langle \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{q}_1 \right\rangle \mathbf{q}_1 - \left\langle \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{q}_2 \right\rangle \mathbf{q}_2 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \quad \mathbf{q}_3 = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

We have obtained an orthonormal basis for V : $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3$.

$$\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \frac{1}{\sqrt{2}} \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}.$$

Why does Gram-Schmidt work?

Our goal

Why does Gram-Schmidt work? Recall, if W is a subspace, \mathbf{b} any vector, then

Our goal

Why does Gram-Schmidt work? Recall, if W is a subspace, \mathbf{b} any vector, then

$$\hat{\mathbf{b}} \rightsquigarrow \text{projection to } W, \mathbf{b}^\perp = \mathbf{b} - \hat{\mathbf{b}} \rightsquigarrow \text{orth. to } W$$

Why does Gram-Schmidt work? Recall, if W is a subspace, \mathbf{b} any vector, then

$$\hat{\mathbf{b}} \rightsquigarrow \text{projection to } W, \mathbf{b}^\perp = \mathbf{b} - \hat{\mathbf{b}} \rightsquigarrow \text{orth. to } W$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

Why does Gram-Schmidt work? Recall, if W is a subspace, \mathbf{b} any vector, then

$$\hat{\mathbf{b}} \rightsquigarrow \text{projection to } W, \mathbf{b}^\perp = \mathbf{b} - \hat{\mathbf{b}} \rightsquigarrow \text{orth. to } W$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

Why does Gram-Schmidt work? Recall, if W is a subspace, \mathbf{b} any vector, then

$$\hat{\mathbf{b}} \rightsquigarrow \text{projection to } W, \mathbf{b}^\perp = \mathbf{b} - \hat{\mathbf{b}} \rightsquigarrow \text{orth. to } W$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\mathbf{b}_1 = \mathbf{a}_1,$$

$$\mathbf{q}_1 = \frac{\mathbf{b}_1}{\underbrace{\|\mathbf{b}_1\|}_{\text{normalize}}}$$

Why does Gram-Schmidt work? Recall, if W is a subspace, \mathbf{b} any vector, then

$$\hat{\mathbf{b}} \rightsquigarrow \text{projection to } W, \mathbf{b}^\perp = \mathbf{b} - \hat{\mathbf{b}} \rightsquigarrow \text{orth. to } W$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\mathbf{b}_1 = \mathbf{a}_1,$$

$$\mathbf{q}_1 = \underbrace{\frac{\mathbf{b}_1}{\|\mathbf{b}_1\|}}_{\text{normalize}}$$

$$\mathbf{b}_2 = \underbrace{\mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1}_{\mathbf{a}_2^\perp \rightsquigarrow \text{orth. to } \text{Span}\{\mathbf{q}_1\}},$$

Why does Gram-Schmidt work? Recall, if W is a subspace, \mathbf{b} any vector, then

$$\hat{\mathbf{b}} \rightsquigarrow \text{projection to } W, \mathbf{b}^\perp = \mathbf{b} - \hat{\mathbf{b}} \rightsquigarrow \text{orth. to } W$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\mathbf{b}_1 = \mathbf{a}_1,$$

$$\mathbf{b}_2 = \underbrace{\mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1}_{\mathbf{a}_2^\perp \rightsquigarrow \text{orth. to } \text{Span}\{\mathbf{q}_1\}},$$

$$\mathbf{q}_1 = \underbrace{\frac{\mathbf{b}_1}{\|\mathbf{b}_1\|}}_{\text{normalize}}$$

$$\mathbf{q}_2 = \underbrace{\frac{\mathbf{b}_2}{\|\mathbf{b}_2\|}}_{\text{normalize}}$$

Why does Gram-Schmidt work? Recall, if W is a subspace, \mathbf{b} any vector, then

$$\hat{\mathbf{b}} \rightsquigarrow \text{projection to } W, \mathbf{b}^\perp = \mathbf{b} - \hat{\mathbf{b}} \rightsquigarrow \text{orth. to } W$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\mathbf{b}_1 = \mathbf{a}_1,$$

$$\mathbf{b}_2 = \underbrace{\mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1}_{\mathbf{a}_2^\perp \rightsquigarrow \text{orth. to Span}\{\mathbf{q}_1\}},$$

$$\mathbf{b}_3 = \underbrace{\mathbf{a}_3 - (\langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 + \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2)}_{\mathbf{a}_3^\perp \text{ orth. to Span}\{\mathbf{q}_1, \mathbf{q}_2\}}$$

$$\mathbf{q}_1 = \underbrace{\frac{\mathbf{b}_1}{\|\mathbf{b}_1\|}}_{\text{normalize}}$$

$$\mathbf{q}_2 = \underbrace{\frac{\mathbf{b}_2}{\|\mathbf{b}_2\|}}_{\text{normalize}}$$

Why does Gram-Schmidt work? Recall, if W is a subspace, \mathbf{b} any vector, then

$$\hat{\mathbf{b}} \rightsquigarrow \text{projection to } W, \mathbf{b}^\perp = \mathbf{b} - \hat{\mathbf{b}} \rightsquigarrow \text{orth. to } W$$

Recipe. (Gram-Schmidt orthonormalization)

Given a basis $\mathbf{a}_1, \dots, \mathbf{a}_n$, produce a **orthogonal basis** $\mathbf{b}_1, \dots, \mathbf{b}_n$ and an **orthonormal basis** $\mathbf{q}_1, \dots, \mathbf{q}_n$.

$$\mathbf{b}_1 = \mathbf{a}_1,$$

$$\mathbf{b}_2 = \underbrace{\mathbf{a}_2 - \langle \mathbf{a}_2, \mathbf{q}_1 \rangle \mathbf{q}_1}_{\mathbf{a}_2^\perp \rightsquigarrow \text{orth. to Span}\{\mathbf{q}_1\}},$$

$$\mathbf{b}_3 = \underbrace{\mathbf{a}_3 - (\langle \mathbf{a}_3, \mathbf{q}_1 \rangle \mathbf{q}_1 + \langle \mathbf{a}_3, \mathbf{q}_2 \rangle \mathbf{q}_2)}_{\mathbf{a}_3^\perp \text{ orth. to Span}\{\mathbf{q}_1, \mathbf{q}_2\}}$$

$$\mathbf{q}_1 = \underbrace{\frac{\mathbf{b}_1}{\|\mathbf{b}_1\|}}_{\text{normalize}}$$

$$\mathbf{q}_2 = \underbrace{\frac{\mathbf{b}_2}{\|\mathbf{b}_2\|}}_{\text{normalize}}$$

$$\mathbf{q}_3 = \underbrace{\frac{\mathbf{b}_3}{\|\mathbf{b}_3\|}}_{\text{normalize}}$$

Example

Let $V = \text{Span}\left\{ \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix} \right\}$. Find an orthonormal basis for V .

Check that your basis is actually orthonormal.

Orthogonal matrices

Theorem

Let $A = [\mathbf{a}_1, \dots, \mathbf{a}_n]$ be a matrix. Then $A^T A$ is the matrix of dot products of the columns of A :

Theorem

Let $A = [\mathbf{a}_1, \dots, \mathbf{a}_n]$ be a matrix. Then $A^T A$ is the matrix of dot products of the columns of A :

Theorem

Let $A = [\mathbf{a}_1, \dots, \mathbf{a}_n]$ be a matrix. Then $A^T A$ is the matrix of dot products of the columns of A :

$$A^T A = \begin{bmatrix} \mathbf{a}_1 \cdot \mathbf{a}_1 & \mathbf{a}_1 \cdot \mathbf{a}_2 & \mathbf{a}_1 \cdot \mathbf{a}_3 & \dots \\ \mathbf{a}_2 \cdot \mathbf{a}_1 & \mathbf{a}_2 \cdot \mathbf{a}_2 & \mathbf{a}_2 \cdot \mathbf{a}_3 & \dots \\ \mathbf{a}_3 \cdot \mathbf{a}_1 & \mathbf{a}_3 \cdot \mathbf{a}_2 & \mathbf{a}_3 \cdot \mathbf{a}_3 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Theorem

Let $A = [\mathbf{a}_1, \dots, \mathbf{a}_n]$ be a matrix. Then $A^T A$ is the matrix of dot products of the columns of A :

$$A^T A = \begin{bmatrix} \mathbf{a}_1 \cdot \mathbf{a}_1 & \mathbf{a}_1 \cdot \mathbf{a}_2 & \mathbf{a}_1 \cdot \mathbf{a}_3 & \dots \\ \mathbf{a}_2 \cdot \mathbf{a}_1 & \mathbf{a}_2 \cdot \mathbf{a}_2 & \mathbf{a}_2 \cdot \mathbf{a}_3 & \dots \\ \mathbf{a}_3 \cdot \mathbf{a}_1 & \mathbf{a}_3 \cdot \mathbf{a}_2 & \mathbf{a}_3 \cdot \mathbf{a}_3 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

What happens if the columns of A are orthonormal?

Theorem

The columns of Q are orthonormal $\iff Q^T Q = I$

Theorem

The columns of Q are orthonormal $\iff Q^T Q = I$

Proof.

Let $\mathbf{q}_1, \dots, \mathbf{q}_n$ be the columns of Q .

They orthonormal if and only if $\mathbf{q}_i^T \mathbf{q}_j = \begin{cases} 0, & \text{if } i \neq j, \\ 1, & \text{if } i = j. \end{cases}$

All these products are packaged in $Q^T Q = I$:

$$\begin{bmatrix} \text{---} & \mathbf{q}_1^T & \text{---} \\ \text{---} & \mathbf{q}_2^T & \text{---} \\ & \vdots & \end{bmatrix} \begin{bmatrix} | & | & \\ \mathbf{q}_1 & \mathbf{q}_2 & \cdots \\ | & | & \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \ddots \end{bmatrix}$$



Definition

An **orthogonal matrix** is a square matrix Q with orthonormal columns.

Definition

An **orthogonal matrix** is a square matrix Q with orthonormal columns.

It is historical convention to restrict to square matrices, and to say orthogonal matrix even though “orthonormal matrix” might be better.

Definition

An **orthogonal matrix** is a square matrix Q with orthonormal columns.

It is historical convention to restrict to square matrices, and to say orthogonal matrix even though “orthonormal matrix” might be better.

An $n \times n$ matrix Q is orthogonal $\iff Q^T Q = I$
In other words, $Q^{-1} = Q^T$.

Definition

An **orthogonal matrix** is a square matrix Q with orthonormal columns.

It is historical convention to restrict to square matrices, and to say orthogonal matrix even though “orthonormal matrix” might be better.

An $n \times n$ matrix Q is orthogonal $\iff Q^T Q = I$
In other words, $Q^{-1} = Q^T$.

Example

$P = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$ is orthogonal.

Definition

An **orthogonal matrix** is a square matrix Q with orthonormal columns.

It is historical convention to restrict to square matrices, and to say orthogonal matrix even though “orthonormal matrix” might be better.

An $n \times n$ matrix Q is orthogonal $\iff Q^T Q = I$
In other words, $Q^{-1} = Q^T$.

Example

$P = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$ is orthogonal.

Why?

Why is $P = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$ orthogonal?

Solution

Why is $P = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$ orthogonal?

Solution

Why is $P = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$ orthogonal?

Solution

Because their columns are a permutation of the standard basis.
And so we always have $P^T P = I$.

Why is $P = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$ orthogonal?

Solution

Because their columns are a permutation of the standard basis.
And so we always have $P^T P = I$. So what is P^{-1} ?

Example

$Q = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ is orthogonal, Why?

Example

$Q = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ is orthogonal, Why?

Solution

- $\begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$ is an orthonormal basis of \mathbb{R}^2 .

Example

$Q = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ is orthogonal, Why?

Solution

- $\begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$ is an orthonormal basis of \mathbb{R}^2 .
Just to make sure: why length 1?

Example

$Q = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ is orthogonal, Why?

Solution

- $\begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$ is an orthonormal basis of \mathbb{R}^2 .

Just to make sure: why length 1? Because

$$\left\| \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix} \right\| = \sqrt{\cos^2 \theta + \sin^2 \theta} = 1.$$

Example

$Q = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ is orthogonal, Why?

Solution

- $\begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$ is an orthonormal basis of \mathbb{R}^2 .

Just to make sure: why length 1? Because

$$\left\| \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix} \right\| = \sqrt{\cos^2 \theta + \sin^2 \theta} = 1.$$

- Alternatively:

$$Q^T Q = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Example

$Q = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ is orthogonal, Why?

Solution

- $\begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$ is an orthonormal basis of \mathbb{R}^2 .

Just to make sure: why length 1? Because

$$\left\| \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix} \right\| = \sqrt{\cos^2 \theta + \sin^2 \theta} = 1.$$

- Alternatively:

$$Q^T Q = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

So what is Q^{-1} ?

Example

Is $H = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$ orthogonal?

Example

Is $H = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$ orthogonal?

Solution

No, the columns are orthogonal but not normalized.

Example

Is $H = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$ orthogonal?

Solution

No, the columns are orthogonal but not normalized.

But $\frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$ is an orthogonal matrix.

Example

(Just for fun) an $n \times n$ matrix with entries ± 1 whose columns are orthogonal is called a *Hadamard matrix* of size n .

Example

(Just for fun) an $n \times n$ matrix with entries ± 1 whose columns are orthogonal is called a *Hadamard matrix* of size n .

Example

(Just for fun) an $n \times n$ matrix with entries ± 1 whose columns are orthogonal is called a *Hadamard matrix* of size n .

A size 4 example:
$$\begin{bmatrix} H & H \\ H & -H \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$$

Example

(Just for fun) an $n \times n$ matrix with entries ± 1 whose columns are orthogonal is called a *Hadamard matrix* of size n .

A size 4 example:
$$\begin{bmatrix} H & H \\ H & -H \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$$

Continuing this construction, we get examples of size 8, 16, 32, ...
It is believed that Hadamard matrices exist for all sizes $4n$.

Example

(Just for fun) an $n \times n$ matrix with entries ± 1 whose columns are orthogonal is called a *Hadamard matrix* of size n .

A size 4 example:
$$\begin{bmatrix} H & H \\ H & -H \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$$

Continuing this construction, we get examples of size 8, 16, 32, ...

It is believed that Hadamard matrices exist for all sizes $4n$.

But, no example of size 668 is known yet.

Example

(Just for fun) an $n \times n$ matrix with entries ± 1 whose columns are orthogonal is called a *Hadamard matrix* of size n .

A size 4 example:
$$\begin{bmatrix} H & H \\ H & -H \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$$

Continuing this construction, we get examples of size 8, 16, 32, ...

It is believed that Hadamard matrices exist for all sizes $4n$.

But, no example of size 668 is known yet. If you find one you will be famous!