

Math 415 - Lecture 33

Diagonalization

Monday November 16th 2015

Textbook reading: Chapter 5.2

Textbook reading: Chapter 5.2

Suggested practice exercises: Chapter 5.2: 1, 2, 3, 4, 5, 6, 7, 15,
16, 17, 18, 19, 20, 25, 26, 29, 30, 31, 32, 33

Textbook reading: Chapter 5.2

Suggested practice exercises: Chapter 5.2: 1, 2, 3, 4, 5, 6, 7, 15,
16, 17, 18, 19, 20, 25, 26, 29, 30, 31, 32, 33

Strang lecture: Lecture 22: Diagonalization and powers of A

Review

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$

λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0.$

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$

λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0.$

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$

λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0.$

- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
- At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$

λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0.$

- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
- At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$

λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0.$

- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .

At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$

λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0.$

- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .

At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$,

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$,

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
 At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda = 1, 1$ (i.e. multiplicity 2), eigenspace is

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$

λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0.$

- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda = 1, 1$ (i.e. multiplicity 2), eigenspace is \mathbb{R}^2 .

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
 At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda = 1, 1$ (i.e. multiplicity 2), eigenspace is \mathbb{R}^2 . Any basis is eigen basis.

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
 At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda = 1, 1$ (i.e. multiplicity 2), eigenspace is \mathbb{R}^2 . Any basis is eigen basis.
 - $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$,

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
 At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda = 1, 1$ (i.e. multiplicity 2), eigenspace is \mathbb{R}^2 . Any basis is eigen basis.
 - $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$,

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
 At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda = 1, 1$ (i.e. multiplicity 2), eigenspace is \mathbb{R}^2 . Any basis is eigen basis.
 - $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, $\lambda = 0, 0$,

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda = 1, 1$ (i.e. multiplicity 2), eigenspace is \mathbb{R}^2 . Any basis is eigen basis.
 - $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, $\lambda = 0, 0$, eigenspace is

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$

λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0.$

- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda = 1, 1$ (i.e. multiplicity 2), eigenspace is \mathbb{R}^2 . Any basis is eigen basis.
 - $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, $\lambda = 0, 0$, eigenspace is \mathbb{R}^2 .

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
 At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda = 1, 1$ (i.e. multiplicity 2), eigenspace is \mathbb{R}^2 . Any basis is eigen basis.
 - $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, $\lambda = 0, 0$, eigenspace is \mathbb{R}^2 . Again any basis is an eigenbasis.

These are trivial cases.

- **Eigenvector** equation: $A\mathbf{x} = \lambda\mathbf{x} \iff (A - \lambda I)\mathbf{x} = \mathbf{0}$
 λ is an **eigenvalue** of $A \iff \underbrace{\det(A - \lambda I)}_{\text{characteristic polynomial}} = 0$.
- An $n \times n$ matrix A has up to n eigenvectors for λ .
 - The **eigenspace** of λ is $\text{Nul}(A - \lambda I)$.
 - If λ has **multiplicity** m , then A has up to m (independent) eigenvectors for λ .
 At least one eigenvector is guaranteed (because $\det(A - \lambda I) = 0$).
 - An **Eigenbasis** for an $n \times n$ matrix A is a basis $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ of \mathbb{R}^n so that each \mathbf{v}_i is also an eigenvector: $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$.
- Test yourself! What are the eigenvectors and eigenvalues? Is there an eigen basis?
 - $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\lambda = 1, 1$ (i.e. multiplicity 2), eigenspace is \mathbb{R}^2 . Any basis is eigen basis.
 - $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$, $\lambda = 0, 0$, eigenspace is \mathbb{R}^2 . Again any basis is an eigenbasis.

These are trivial cases. Is there always an eigenbasis?

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations.

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations.

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations. If we want to find eigenvectors, $A\mathbf{x} = \lambda\mathbf{x}$, can we also use row operations?

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations. If we want to find eigenvectors, $A\mathbf{x} = \lambda\mathbf{x}$, can we also use row operations? Try

$$A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}.$$

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations. If we want to find eigenvectors, $A\mathbf{x} = \lambda\mathbf{x}$, can we also use row operations? Try

$$A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}.$$

- What is the echelon form U of A ?

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations. If we want to find eigenvectors, $A\mathbf{x} = \lambda\mathbf{x}$, can we also use row operations? Try

$$A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}.$$

- What is the echelon form U of A ?

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations. If we want to find eigenvectors, $A\mathbf{x} = \lambda\mathbf{x}$, can we also use row operations? Try

$$A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}.$$

- What is the echelon form U of A ?
- What are the characteristic polynomials $\det(A - \lambda I)$ and $\det(U - \lambda I)$? Roots?

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations. If we want to find eigenvectors, $A\mathbf{x} = \lambda\mathbf{x}$, can we also use row operations? Try

$$A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}.$$

- What is the echelon form U of A ?
- What are the characteristic polynomials $\det(A - \lambda I)$ and $\det(U - \lambda I)$? Roots?

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations. If we want to find eigenvectors, $A\mathbf{x} = \lambda\mathbf{x}$, can we also use row operations? Try

$$A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}.$$

- What is the echelon form U of A ?
- What are the characteristic polynomials $\det(A - \lambda I)$ and $\det(U - \lambda I)$? Roots?
- Do A and U have the same eigenvalues? Eigenvectors?

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations. If we want to find eigenvectors, $A\mathbf{x} = \lambda\mathbf{x}$, can we also use row operations? Try

$$A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}.$$

- What is the echelon form U of A ?
- What are the characteristic polynomials $\det(A - \lambda I)$ and $\det(U - \lambda I)$? Roots?
- Do A and U have the same eigenvalues? Eigenvectors?

Example

To solve $A\mathbf{x} = \mathbf{b}$ we use row operations. If we want to find eigenvectors, $A\mathbf{x} = \lambda\mathbf{x}$, can we also use row operations? Try

$$A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}.$$

- What is the echelon form U of A ?
- What are the characteristic polynomials $\det(A - \lambda I)$ and $\det(U - \lambda I)$? Roots?
- Do A and U have the same eigenvalues? Eigenvectors?

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U =$

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.
- Then $\det(A - \lambda I) =$

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.
- Then $\det(A - \lambda I) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) =$
 $= \lambda^2 - 0\lambda + (-4) = \lambda^2 - 4 = (\lambda - 2)(\lambda + 2),$

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.
- Then $\det(A - \lambda I) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) =$
 $= \lambda^2 - 0\lambda + (-4) = \lambda^2 - 4 = (\lambda - 2)(\lambda + 2)$, and
 $\det(U - \lambda I) =$

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.
- Then $\det(A - \lambda I) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) =$
 $= \lambda^2 - 0\lambda + (-4) = \lambda^2 - 4 = (\lambda - 2)(\lambda + 2)$, and
 $\det(U - \lambda I) = \lambda^2 - \text{Tr}(U)\lambda + \det(U) =$
 $= \lambda^2 - 3\lambda + (-4) = (\lambda - 1)(\lambda + 4)$.

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.
- Then $\det(A - \lambda I) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) =$
 $= \lambda^2 - 0\lambda + (-4) = \lambda^2 - 4 = (\lambda - 2)(\lambda + 2)$, and
 $\det(U - \lambda I) = \lambda^2 - \text{Tr}(U)\lambda + \det(U) =$
 $= \lambda^2 - 3\lambda + (-4) = (\lambda - 1)(\lambda + 4)$.
- So the eigenvalues of A and U are

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.
- Then $\det(A - \lambda I) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) =$
 $= \lambda^2 - 0\lambda + (-4) = \lambda^2 - 4 = (\lambda - 2)(\lambda + 2)$, and
 $\det(U - \lambda I) = \lambda^2 - \text{Tr}(U)\lambda + \det(U) =$
 $= \lambda^2 - 3\lambda + (-4) = (\lambda - 1)(\lambda + 4)$.
- So the eigenvalues of A and U are **DIFFERENT!**.

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.
- Then $\det(A - \lambda I) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) =$
 $= \lambda^2 - 0\lambda + (-4) = \lambda^2 - 4 = (\lambda - 2)(\lambda + 2)$, and
 $\det(U - \lambda I) = \lambda^2 - \text{Tr}(U)\lambda + \det(U) =$
 $= \lambda^2 - 3\lambda + (-4) = (\lambda - 1)(\lambda + 4)$.
- So the eigenvalues of A and U are **DIFFERENT!**. Can check that eigenvectors are also different.

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.
- Then $\det(A - \lambda I) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) = \lambda^2 - 0\lambda + (-4) = \lambda^2 - 4 = (\lambda - 2)(\lambda + 2)$, and $\det(U - \lambda I) = \lambda^2 - \text{Tr}(U)\lambda + \det(U) = \lambda^2 - 3\lambda + (-4) = (\lambda - 1)(\lambda + 4)$.
- So the eigenvalues of A and U are **DIFFERENT!**. Can check that eigenvectors are also different.

Upshot:

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.
- Then $\det(A - \lambda I) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) = \lambda^2 - 0\lambda + (-4) = \lambda^2 - 4 = (\lambda - 2)(\lambda + 2)$, and $\det(U - \lambda I) = \lambda^2 - \text{Tr}(U)\lambda + \det(U) = \lambda^2 - 3\lambda + (-4) = (\lambda - 1)(\lambda + 4)$.
- So the eigenvalues of A and U are **DIFFERENT!**. Can check that eigenvectors are also different.

Upshot: **Don't use row operations to deal with eigenvalues and eigenvectors!**

Solution

- If $A = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$ then $U = \begin{bmatrix} 1 & 1 \\ 0 & -4 \end{bmatrix}$.
- Then $\det(A - \lambda I) = \lambda^2 - \text{Tr}(A)\lambda + \det(A) = \lambda^2 - 0\lambda + (-4) = \lambda^2 - 4 = (\lambda - 2)(\lambda + 2)$, and $\det(U - \lambda I) = \lambda^2 - \text{Tr}(U)\lambda + \det(U) = \lambda^2 - 3\lambda + (-4) = (\lambda - 1)(\lambda + 4)$.
- So the eigenvalues of A and U are **DIFFERENT!**. Can check that eigenvectors are also different.

Upshot: **Don't use row operations to deal with eigenvalues and eigenvectors!** (Can use row operations to calculate determinants, though.)

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$.

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

So: $\lambda = 1$ is the only eigenvalue (it has multiplicity 2).

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

So: $\lambda = 1$ is the only eigenvalue (it has multiplicity 2).

- $\lambda = 1$:

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

So: $\lambda = 1$ is the only eigenvalue (it has multiplicity 2).

- $\lambda = 1 : \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies$

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

So: $\lambda = 1$ is the only eigenvalue (it has multiplicity 2).

- $\lambda = 1 : \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

So: $\lambda = 1$ is the only eigenvalue (it has multiplicity 2).

- $\lambda = 1 : \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

So the eigenspace is $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$.

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

So: $\lambda = 1$ is the only eigenvalue (it has multiplicity 2).

- $\lambda = 1 : \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

So the eigenspace is $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$. Only dimension 1!

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

So: $\lambda = 1$ is the only eigenvalue (it has multiplicity 2).

- $\lambda = 1 : \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

So the eigenspace is $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$. Only dimension 1!

- Trouble: We can not find an **Eigenbasis** for this matrix.

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

So: $\lambda = 1$ is the only eigenvalue (it has multiplicity 2).

- $\lambda = 1 : \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

So the eigenspace is $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$. Only dimension 1!

- Trouble: We can not find an **Eigenbasis** for this matrix.
This kind of problem cannot really be fixed.

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

So: $\lambda = 1$ is the only eigenvalue (it has multiplicity 2).

- $\lambda = 1 : \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

So the eigenspace is $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$. Only dimension 1!

- Trouble: We can not find an **Eigenbasis** for this matrix.
This kind of problem cannot really be fixed.
We have to lower our expectations and look for generalized eigenvectors.

Example

Find the eigenvectors and eigenvalues of $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What is the trouble?

Solution

- $\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2$

So: $\lambda = 1$ is the only eigenvalue (it has multiplicity 2).

- $\lambda = 1 : \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \mathbf{x} = \mathbf{0} \implies \mathbf{x}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

So the eigenspace is $\text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$. Only dimension 1!

- Trouble: We can not find an **Eigenbasis** for this matrix.

This kind of problem cannot really be fixed.

We have to lower our expectations and look for generalized eigenvectors.

These are solutions to $(A - \lambda I)^2 \mathbf{x} = \mathbf{0}$, $(A - \lambda I)^3 \mathbf{x} = \mathbf{0}$, ...

Diagonalization

Powers of diagonal matrices

Diagonal matrices are very easy to work with.

Diagonal matrices are very easy to work with.

Example

Let $A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{bmatrix}$. What is A^2 ? What is A^{100} ?

Diagonal matrices are very easy to work with.

Example

Let $A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{bmatrix}$. What is A^2 ? What is A^{100} ?

Solution

$$A^2 =$$

Diagonal matrices are very easy to work with.

Example

Let $A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{bmatrix}$. What is A^2 ? What is A^{100} ?

Solution

$$A^2 = \begin{bmatrix} 2^2 & 0 & 0 \\ 0 & 3^2 & 0 \\ 0 & 0 & 4^2 \end{bmatrix} \text{ and } A^{100} =$$

Diagonal matrices are very easy to work with.

Example

Let $A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{bmatrix}$. What is A^2 ? What is A^{100} ?

Solution

$$A^2 = \begin{bmatrix} 2^2 & 0 & 0 \\ 0 & 3^2 & 0 \\ 0 & 0 & 4^2 \end{bmatrix} \text{ and } A^{100} = \begin{bmatrix} 2^{100} & 0 & 0 \\ 0 & 3^{100} & 0 \\ 0 & 0 & 4^{100} \end{bmatrix}.$$

Powers of generic matrices

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \cdots = (\lambda - 4)(\lambda - 5)$$

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \cdots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$:

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \dots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$: $\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \dots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$: $\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \dots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$: $\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.
- $\lambda_2 = 5$:

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \dots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$: $\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.
- $\lambda_2 = 5$: $\begin{bmatrix} 1 & -1 \\ 2 & -2 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_2 =$

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \dots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$: $\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.
- $\lambda_2 = 5$: $\begin{bmatrix} 1 & -1 \\ 2 & -2 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \dots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$: $\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.
- $\lambda_2 = 5$: $\begin{bmatrix} 1 & -1 \\ 2 & -2 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- Key observation: $A^{100} \mathbf{v}_1 =$

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \dots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$: $\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.
- $\lambda_2 = 5$: $\begin{bmatrix} 1 & -1 \\ 2 & -2 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- Key observation: $A^{100} \mathbf{v}_1 = \lambda_1^{100} \mathbf{v}_1$ and

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \dots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$: $\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.
- $\lambda_2 = 5$: $\begin{bmatrix} 1 & -1 \\ 2 & -2 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- Key observation: $A^{100} \mathbf{v}_1 = \lambda_1^{100} \mathbf{v}_1$ and $A^{100} \mathbf{v}_2 =$

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \dots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$: $\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.
- $\lambda_2 = 5$: $\begin{bmatrix} 1 & -1 \\ 2 & -2 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- Key observation: $A^{100} \mathbf{v}_1 = \lambda_1^{100} \mathbf{v}_1$ and $A^{100} \mathbf{v}_2 = \lambda_2^{100} \mathbf{v}_2$.

Example

If $A = \begin{bmatrix} 6 & -1 \\ 2 & 3 \end{bmatrix}$, then $A^{100} = ?$

Solution

- characteristic polynomial:

$$\begin{vmatrix} 6 - \lambda & -1 \\ 2 & 3 - \lambda \end{vmatrix} = \dots = (\lambda - 4)(\lambda - 5)$$

- $\lambda_1 = 4$: $\begin{bmatrix} 2 & -1 \\ 2 & -1 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$.
- $\lambda_2 = 5$: $\begin{bmatrix} 1 & -1 \\ 2 & -2 \end{bmatrix} \mathbf{v} = \mathbf{0} \implies$ eigenvector $\mathbf{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
- Key observation: $A^{100} \mathbf{v}_1 = \lambda_1^{100} \mathbf{v}_1$ and $A^{100} \mathbf{v}_2 = \lambda_2^{100} \mathbf{v}_2$.
For A^{100} , we need $A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $A^{100} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$.

Solution (continued)

- $\begin{bmatrix} 1 \\ 0 \end{bmatrix} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 = - \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \Rightarrow$

Solution (continued)

$$\bullet \begin{bmatrix} 1 \\ 0 \end{bmatrix} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 = - \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \Rightarrow$$

$$A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix} =$$

Solution (continued)

$$\bullet \begin{bmatrix} 1 \\ 0 \end{bmatrix} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 = - \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \implies$$

$$A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = A^{100} \left(- \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right) =$$

Solution (continued)

$$\bullet \begin{bmatrix} 1 \\ 0 \end{bmatrix} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 = - \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \Rightarrow$$

$$A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = A^{100} \left(- \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right) = -4^{100} \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \cdot 5^{100} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\Rightarrow A^{100} =$$

Solution (continued)

$$\bullet \begin{bmatrix} 1 \\ 0 \end{bmatrix} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 = - \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \Rightarrow$$

$$A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = A^{100} \left(- \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right) = -4^{100} \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \cdot 5^{100} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\Rightarrow A^{100} = \begin{bmatrix} 2 \cdot 5^{100} - 4^{100} & * \\ 2 \cdot 5^{100} - 2 \cdot 4^{100} & * \end{bmatrix}$$

Solution (continued)

$$\bullet \begin{bmatrix} 1 \\ 0 \end{bmatrix} = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 = - \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \implies$$

$$A^{100} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = A^{100} \left(- \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right) = -4^{100} \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 2 \cdot 5^{100} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\implies A^{100} = \begin{bmatrix} 2 \cdot 5^{100} - 4^{100} & * \\ 2 \cdot 5^{100} - 2 \cdot 4^{100} & * \end{bmatrix}$$

- We find the second column of A^{100} likewise. Left as exercise!

The key idea of previous example is to work with respect to an *Eigenbasis*, a basis given by eigenvectors.

The key idea of previous example is to work with respect to an *Eigenbasis*, a basis given by eigenvectors.

- Put the eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_n$ as columns into a matrix P .

The key idea of previous example is to work with respect to an *Eigenbasis*, a basis given by eigenvectors.

- Put the eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_n$ as columns into a matrix P .

The key idea of previous example is to work with respect to an *Eigenbasis*, a basis given by eigenvectors.

- Put the eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_n$ as columns into a matrix P .

$$\begin{aligned}
 A\mathbf{x}_i = \lambda\mathbf{x}_i \implies A \begin{bmatrix} | & & | \\ \mathbf{x}_1 & \cdots & \mathbf{x}_n \\ | & & | \end{bmatrix} &= \begin{bmatrix} | & & | \\ \lambda_1\mathbf{x}_1 & \cdots & \lambda_n\mathbf{x}_n \\ | & & | \end{bmatrix} \\
 &= \begin{bmatrix} | & & | \\ \mathbf{x}_1 & \cdots & \mathbf{x}_n \\ | & & | \end{bmatrix} \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}
 \end{aligned}$$

The key idea of previous example is to work with respect to an *Eigenbasis*, a basis given by eigenvectors.

- Put the eigenvectors $\mathbf{x}_1, \dots, \mathbf{x}_n$ as columns into a matrix P .

$$\begin{aligned} A\mathbf{x}_i = \lambda\mathbf{x}_i \implies A \begin{bmatrix} | & & | \\ \mathbf{x}_1 & \cdots & \mathbf{x}_n \\ | & & | \end{bmatrix} &= \begin{bmatrix} | & & | \\ \lambda_1\mathbf{x}_1 & \cdots & \lambda_n\mathbf{x}_n \\ | & & | \end{bmatrix} \\ &= \begin{bmatrix} | & & | \\ \mathbf{x}_1 & \cdots & \mathbf{x}_n \\ | & & | \end{bmatrix} \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \end{aligned}$$

- In summary $AP = PD$. Such a diagonalization is possible if and only if A has enough eigenvectors.

Motto

Everything in Linear Algebra is a matrix factorization.

So we are going to use eigenvalues and eigenvectors for A to factor A and A^{100} in a useful way. This is called **diagonalization**.

So we are going to use eigenvalues and eigenvectors for A to factor A and A^{100} in a useful way. This is called **diagonalization**.

Definition

A square matrix A is said to be **diagonalizable** if there is a invertible matrix P and a diagonal matrix D such that

$$A = PDP^{-1}.$$

So we are going to use eigenvalues and eigenvectors for A to factor A and A^{100} in a useful way. This is called **diagonalization**.

Definition

A square matrix A is said to be **diagonalizable** if there is a invertible matrix P and a diagonal matrix D such that

$$A = PDP^{-1}.$$

Theorem

An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.

We can express the relation between A and D in terms of change of base matrices.

$$\begin{array}{ccc} \text{coords for } x & \xrightarrow{A} & \text{coords for } Ax \\ \text{in standard basis} & & \text{in standard basis} \end{array}$$

$$\begin{array}{ccc} \text{coords for } x & \xrightarrow{D} & \text{coords for } Ax \\ \text{in eigen-basis} & & \text{in eigen-basis} \end{array}$$

P changes from eigenbasis coordinates to standard coordinates, and P^{-1} goes the other way! Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{B} the basis of eigenvectors of A , then

$$P = I_{\mathcal{E}, \mathcal{B}} \text{ and } P^{-1} = I_{\mathcal{B}, \mathcal{E}}.$$

We can express the relation between A and D in terms of change of base matrices.


$$\begin{array}{ccc} \text{coords for } x & \xrightarrow{A} & \text{coords for } Ax \\ \text{in standard basis} & & \text{in standard basis} \end{array}$$

$$\begin{array}{ccc} \text{coords for } x & \xrightarrow{D} & \text{coords for } Ax \\ \text{in eigen-basis} & & \text{in eigen-basis} \end{array}$$

P changes from eigenbasis coordinates to standard coordinates, and P^{-1} goes the other way! Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{B} the basis of eigenvectors of A , then

$$P = I_{\mathcal{E}, \mathcal{B}} \text{ and } P^{-1} = I_{\mathcal{B}, \mathcal{E}}.$$

We can express the relation between A and D in terms of change of base matrices.




$$D = P^{-1}AP,$$

P changes from eigenbasis coordinates to standard coordinates, and P^{-1} goes the other way! Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{B} the basis of eigenvectors of A , then

$$P = I_{\mathcal{E}, \mathcal{B}} \text{ and } P^{-1} = I_{\mathcal{B}, \mathcal{E}}.$$

We can express the relation between A and D in terms of change of base matrices.




$$D = P^{-1}AP, \quad A = PDP^{-1}$$

P changes from eigenbasis coordinates to standard coordinates, and P^{-1} goes the other way! Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{B} the basis of eigenvectors of A , then

$$P = I_{\mathcal{E}, \mathcal{B}} \text{ and } P^{-1} = I_{\mathcal{B}, \mathcal{E}}.$$

We can express the relation between A and D in terms of change of base matrices.



$$D = P^{-1}AP, \quad A = PDP^{-1}$$

P changes from eigenbasis coordinates to standard coordinates, and P^{-1} goes the other way! Let \mathcal{E} be the standard basis of \mathbb{R}^n and \mathcal{B} the basis of eigenvectors of A , then

$$P = I_{\mathcal{E}, \mathcal{B}} \text{ and } P^{-1} = I_{\mathcal{B}, \mathcal{E}}.$$

Application: Large powers

If A has an eigenbasis, then we can raise it to large powers easily!

Theorem

If $A = PDP^{-1}$, where D is a diagonal matrix, then for any m ,

$$A^m = PD^mP^{-1}$$

If A has an eigenbasis, then we can raise it to large powers easily!

Theorem

If $A = PDP^{-1}$, where D is a diagonal matrix, then for any m ,

$$A^m = PD^mP^{-1}$$

Proof.

$$\begin{aligned} A &= PDP^{-1} \\ A^m &= (PDP^{-1})^m \end{aligned}$$



If A has an eigenbasis, then we can raise it to large powers easily!

Theorem

If $A = PDP^{-1}$, where D is a diagonal matrix, then for any m ,

$$A^m = PD^mP^{-1}$$

Proof.

$$\begin{aligned} A &= PDP^{-1} \\ A^m &= (PDP^{-1})^m \\ &= \underbrace{(PDP^{-1}) \cdot (PDP^{-1}) \cdots (PDP^{-1})}_{m \text{ times}} \end{aligned}$$



If A has an eigenbasis, then we can raise it to large powers easily!

Theorem

If $A = PDP^{-1}$, where D is a diagonal matrix, then for any m ,

$$A^m = PD^mP^{-1}$$

Proof.

$$\begin{aligned} A &= PDP^{-1} \\ A^m &= (PDP^{-1})^m \\ &= \underbrace{(PDP^{-1}) \cdot (PDP^{-1}) \cdots (PDP^{-1})}_{m \text{ times}} \\ &= (PD)(P^{-1} \cdot P)(\overbrace{DP^{-1}}^{m \text{ times}}) \cdots (PDP^{-1}) \end{aligned}$$



If A has an eigenbasis, then we can raise it to large powers easily!

Theorem

If $A = PDP^{-1}$, where D is a diagonal matrix, then for any m ,

$$A^m = PD^mP^{-1}$$

Proof.

$$\begin{aligned} A &= PDP^{-1} \\ A^m &= (PDP^{-1})^m \\ &= \underbrace{(PDP^{-1}) \cdot (PDP^{-1}) \cdots (PDP^{-1})}_{m \text{ times}} \\ &= (PD)(P^{-1} \cdot P)(D^{m \text{ times}}P^{-1}) \cdots (PDP^{-1}) \\ &= PD \cdot DP^{-1} \cdots PDP^{-1} \end{aligned}$$



If A has an eigenbasis, then we can raise it to large powers easily!

Theorem

If $A = PDP^{-1}$, where D is a diagonal matrix, then for any m ,

$$A^m = PD^mP^{-1}$$

Proof.

$$\begin{aligned} A &= PDP^{-1} \\ A^m &= (PDP^{-1})^m \\ &= \underbrace{(PDP^{-1}) \cdot (PDP^{-1}) \cdots (PDP^{-1})}_{m \text{ times}} \\ &= (PD)(P^{-1} \cdot P)(\overbrace{DP^{-1}}^{m \text{ times}}) \cdots (PDP^{-1}) \\ &= PD \cdot DP^{-1} \cdots PDP^{-1} \\ &= PD \cdot D \cdots D \cdot P^{-1} \end{aligned}$$



If A has an eigenbasis, then we can raise it to large powers easily!

Theorem

If $A = PDP^{-1}$, where D is a diagonal matrix, then for any m ,

$$A^m = PD^mP^{-1}$$

Proof.

$$\begin{aligned} A &= PDP^{-1} \\ A^m &= (PDP^{-1})^m \\ &= \underbrace{(PDP^{-1}) \cdot (PDP^{-1}) \cdots (PDP^{-1})}_{m \text{ times}} \\ &= (PD)(P^{-1} \cdot P)(\overset{m \text{ times}}{DP^{-1}}) \cdots (PDP^{-1}) \\ &= PD \cdot DP^{-1} \cdots PDP^{-1} \\ &= PD \cdot D \cdots D \cdot P^{-1} \\ &= PD^mP^{-1} \end{aligned}$$



If A has an eigenbasis, then we can raise it to large powers easily!

Theorem

If $A = PDP^{-1}$, where D is a diagonal matrix, then for any m ,

$$A^m = PD^mP^{-1}$$

Proof.

$$\begin{aligned} A &= PDP^{-1} \\ A^m &= (PDP^{-1})^m \\ &= \underbrace{(PDP^{-1}) \cdot (PDP^{-1}) \cdots (PDP^{-1})}_{m \text{ times}} \\ &= (PD)(P^{-1} \cdot P)(\overset{m \text{ times}}{DP^{-1}}) \cdots (PDP^{-1}) \\ &= PD \cdot DP^{-1} \cdots PDP^{-1} \\ &= PD \cdot D \cdots D \cdot P^{-1} \\ &= PD^mP^{-1} \end{aligned}$$

Only the outside P and P^{-1} remain!



Finding D^m is easy!

$$D^m = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}^m$$

Finding D^m is easy!

$$D^m = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}^m = \begin{bmatrix} (\lambda_1)^m & & \\ & \ddots & \\ & & (\lambda_n)^m \end{bmatrix}$$

Finding D^m is easy!

$$D^m = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}^m = \begin{bmatrix} (\lambda_1)^m & & \\ & \ddots & \\ & & (\lambda_n)^m \end{bmatrix}$$

Finding D^m is easy!

$$D^m = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}^m = \begin{bmatrix} (\lambda_1)^m & & \\ & \ddots & \\ & & (\lambda_n)^m \end{bmatrix}$$

Why?

Example

Let $A = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ -\frac{1}{2} & 1 & 6 \\ 0 & 0 & 2 \end{bmatrix}$.

Example

Let $A = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ -\frac{1}{2} & 1 & 6 \\ 0 & 0 & 2 \end{bmatrix}$. A has eigenvectors and eigenvalues

$$\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \quad \text{with} \quad \lambda_1 = \frac{1}{2}$$

$$\mathbf{x}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad \text{with} \quad \lambda_2 = 1$$

$$\mathbf{x}_3 = \begin{bmatrix} 0 \\ 6 \\ 1 \end{bmatrix} \quad \text{with} \quad \lambda_3 = 2$$

Example

Let $A = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ -\frac{1}{2} & 1 & 6 \\ 0 & 0 & 2 \end{bmatrix}$. A has eigenvectors and eigenvalues

$$\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \quad \text{with} \quad \lambda_1 = \frac{1}{2}$$

$$\mathbf{x}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad \text{with} \quad \lambda_2 = 1$$

$$\mathbf{x}_3 = \begin{bmatrix} 0 \\ 6 \\ 1 \end{bmatrix} \quad \text{with} \quad \lambda_3 = 2$$

Find A^{100} .

Example

Let $A = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ -\frac{1}{2} & 1 & 6 \\ 0 & 0 & 2 \end{bmatrix}$. A has eigenvectors and eigenvalues

$$\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \quad \text{with} \quad \lambda_1 = \frac{1}{2}$$

$$\mathbf{x}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad \text{with} \quad \lambda_2 = 1$$

$$\mathbf{x}_3 = \begin{bmatrix} 0 \\ 6 \\ 1 \end{bmatrix} \quad \text{with} \quad \lambda_3 = 2$$

Find A^{100} . **Hint:** Write $A = PDP^{-1}$.

Solution

Eigenvectors of A form an **Eigenbasis!** So we can write

Solution

Eigenvectors of A form an **Eigenbasis!** So we can write

$$A = PDP^{-1};$$

Matrix of eigenvectors $P = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 6 \\ 0 & 0 & 1 \end{bmatrix}$

Solution

Eigenvectors of A form an **Eigenbasis!** So we can write

$$A = PDP^{-1};$$

Matrix of eigenvectors $P = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 6 \\ 0 & 0 & 1 \end{bmatrix}$

Solution

Eigenvectors of A form an **Eigenbasis!** So we can write

$$A = PDP^{-1};$$

Matrix of eigenvectors $P = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 6 \\ 0 & 0 & 1 \end{bmatrix}$

Find P^{-1}

Solution

Eigenvectors of A form an **Eigenbasis!** So we can write
 $A = PDP^{-1}$:

Matrix of eigenvectors $P = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 6 \\ 0 & 0 & 1 \end{bmatrix}$

Find P^{-1}

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 6 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

Solution

Eigenvectors of A form an **Eigenbasis!** So we can write
 $A = PDP^{-1}$:

Matrix of eigenvectors $P = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 6 \\ 0 & 0 & 1 \end{bmatrix}$

Find P^{-1}

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 6 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R2 \rightsquigarrow R2 - R1} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 6 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

Solution

Eigenvectors of A form an **Eigenbasis**! So we can write
 $A = PDP^{-1}$:

Matrix of eigenvectors $P = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 6 \\ 0 & 0 & 1 \end{bmatrix}$

Find P^{-1}

$$\begin{bmatrix} 1 & 0 & 0 & | & 1 & 0 & 0 \\ 1 & 1 & 6 & | & 0 & 1 & 0 \\ 0 & 0 & 1 & | & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R2 \rightsquigarrow R2 - R1} \begin{bmatrix} 1 & 0 & 0 & | & 1 & 0 & 0 \\ 0 & 1 & 6 & | & -1 & 1 & 0 \\ 0 & 0 & 1 & | & 0 & 0 & 1 \end{bmatrix}$$

$$\xrightarrow{R2 \rightsquigarrow R2 - 6R3} \begin{bmatrix} 1 & 0 & 0 & | & 1 & 0 & 0 \\ 0 & 1 & 0 & | & -1 & 1 & -6 \\ 0 & 0 & 1 & | & 0 & 0 & 1 \end{bmatrix}$$

Solution

Eigenvectors of A form an **Eigenbasis**! So we can write
 $A = PDP^{-1}$:

Matrix of eigenvectors $P = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 6 \\ 0 & 0 & 1 \end{bmatrix}$

Find P^{-1}

$$\begin{array}{c} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 6 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R2 \rightsquigarrow R2 - R1} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 6 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right] \\ \xrightarrow{R2 \rightsquigarrow R2 - 6R3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & -1 & 1 & -6 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right] \end{array}$$

$$P^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & -6 \\ 0 & 0 & 1 \end{bmatrix}$$

Solution (continued)

Eigenvectors of A form a **Eigenbasis!** So we can write

Solution (continued)

Eigenvectors of A form a **Eigenbasis**! So we can write $A = PDP^{-1}$:

Matrix of eigenvalues: $D = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$

Finally, write $A = PDP^{-1}$:

Solution (continued)

Eigenvectors of A form a **Eigenbasis!** So we can write $A = PDP^{-1}$:

Matrix of eigenvalues: $D = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}$

Finally, write $A = PDP^{-1}$:

$$\underbrace{\begin{bmatrix} \frac{1}{2} & 0 & 0 \\ -\frac{1}{2} & 1 & 6 \\ 0 & 0 & 2 \end{bmatrix}}_A = \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 6 \\ 0 & 0 & 1 \end{bmatrix}}_P \underbrace{\begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}}_D \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & -6 \\ 0 & 0 & 1 \end{bmatrix}}_{P^{-1}}$$

Solution (continued)

Take power

$$\begin{aligned}
 \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ -\frac{1}{2} & 1 & 6 \\ 0 & 0 & 2 \end{bmatrix}^{100} &= \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 6 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}^{100} \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & -6 \\ 0 & 0 & 1 \end{bmatrix} \\
 &= \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 6 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{2^{100}} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2^{100} \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -1 & 1 & -6 \\ 0 & 0 & 1 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{1}{2^{100}} & 0 & 0 \\ (\frac{1}{2^{100}} - 1) & 1 & (6 \cdot 2^{100} - 6) \\ 0 & 0 & 2^{100} \end{bmatrix}
 \end{aligned}$$