

# Math 415 - Lecture 19

## Orthonormal basis, orthogonal complement

Friday October 9th 2015

# Math 415 - Lecture 19

## Orthonormal basis, orthogonal complement

Friday October 9th 2015

Textbook reading: Ch 3.1

Textbook reading: Ch 3.1

Suggested practice exercises: Ch 3.1: 7, 8, 9, 10, 11, 12, 14, 15,  
17, 18, 19, 20, 22

Textbook reading: Ch 3.1

Suggested practice exercises: Ch 3.1: 7, 8, 9, 10, 11, 12, 14, 15,  
17, 18, 19, 20, 22

Khan Academy videos:

Textbook reading: Ch 3.1

Suggested practice exercises: Ch 3.1: 7, 8, 9, 10, 11, 12, 14, 15,  
17, 18, 19, 20, 22

Khan Academy videos:

Strang lectures: Lec 10: The Four Fundamental Subspaces / Lec  
14: Orthogonal Vectors and Subspaces

## Review

- $\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \cdots + v_n w_n$  is the **inner product** of  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ .



- $\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \cdots + v_n w_n$  is the **inner product** of  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ .
  - The **length** of  $\mathbf{v}$ ,  $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$ .

- $\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \cdots + v_n w_n$  is the **inner product** of  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ .
  - The **length** of  $\mathbf{v}$ ,  $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$ .

- $\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \cdots + v_n w_n$  is the **inner product** of  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ .
  - The **length** of  $\mathbf{v}$ ,  $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$ .
  - The **distance** between points  $\mathbf{v}, \mathbf{w}$  is  $\|\mathbf{v} - \mathbf{w}\|$ .

- $\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \cdots + v_n w_n$  is the **inner product** of  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ .
  - The **length** of  $\mathbf{v}$ ,  $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$ .
  - The **distance** between points  $\mathbf{v}, \mathbf{w}$  is  $\|\mathbf{v} - \mathbf{w}\|$ .

- $\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \cdots + v_n w_n$  is the **inner product** of  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ .
  - The **length** of  $\mathbf{v}$ ,  $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$ .
  - The **distance** between points  $\mathbf{v}, \mathbf{w}$  is  $\|\mathbf{v} - \mathbf{w}\|$ .
- $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$  are **orthogonal** if  $\mathbf{v} \cdot \mathbf{w} = 0$ .

- $\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \cdots + v_n w_n$  is the **inner product** of  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ .
  - The **length** of  $\mathbf{v}$ ,  $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$ .
  - The **distance** between points  $\mathbf{v}, \mathbf{w}$  is  $\|\mathbf{v} - \mathbf{w}\|$ .
- $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$  are **orthogonal** if  $\mathbf{v} \cdot \mathbf{w} = 0$ .

- $\mathbf{v} \cdot \mathbf{w} = \mathbf{v}^T \mathbf{w} = v_1 w_1 + \cdots + v_n w_n$  is the **inner product** of  $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ .
  - The **length** of  $\mathbf{v}$ ,  $\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$ .
  - The **distance** between points  $\mathbf{v}, \mathbf{w}$  is  $\|\mathbf{v} - \mathbf{w}\|$ .
- $\mathbf{v}, \mathbf{w} \in \mathbb{R}^n$  are **orthogonal** if  $\mathbf{v} \cdot \mathbf{w} = 0$ .
  - This simple criterion is equivalent to Pythagoras' theorem.

## Unit Vectors and Orthonormal basis



## Definition

A vector  $\mathbf{u} \in \mathbb{R}^n$  is called a *unit vector* if

## Definition

A vector  $\mathbf{u} \in \mathbb{R}^n$  is called a *unit vector* if

- $\|\mathbf{u}\| = 1$ , or, equivalently,

## Definition

A vector  $\mathbf{u} \in \mathbb{R}^n$  is called a *unit vector* if

- $\|\mathbf{u}\| = 1$ , or, equivalently,
- $\mathbf{u} \cdot \mathbf{u} = 1$

## Definition

A vector  $\mathbf{u} \in \mathbb{R}^n$  is called a *unit vector* if

- $\|\mathbf{u}\| = 1$ , or, equivalently,
- $\mathbf{u} \cdot \mathbf{u} = 1$

## Example

The standard basis vectors  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$  of  $\mathbb{R}^n$  are all unit vectors.

## Definition

A vector  $\mathbf{u} \in \mathbb{R}^n$  is called a *unit vector* if

- $\|\mathbf{u}\| = 1$ , or, equivalently,
- $\mathbf{u} \cdot \mathbf{u} = 1$

## Example

The standard basis vectors  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$  of  $\mathbb{R}^n$  are all unit vectors.

## Example

If  $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ , then is  $\mathbf{x}$  a unit vector?

## Definition

A vector  $\mathbf{u} \in \mathbb{R}^n$  is called a *unit vector* if

- $\|\mathbf{u}\| = 1$ , or, equivalently,
- $\mathbf{u} \cdot \mathbf{u} = 1$

## Example

The standard basis vectors  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$  of  $\mathbb{R}^n$  are all unit vectors.

## Example

If  $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ , then is  $\mathbf{x}$  a unit vector?

## Solution

Since  $\mathbf{x} \cdot \mathbf{x} = 5$  and  $\|\mathbf{x}\| = \sqrt{5}$ .

## Definition

A vector  $\mathbf{u} \in \mathbb{R}^n$  is called a *unit vector* if

- $\|\mathbf{u}\| = 1$ , or, equivalently,
- $\mathbf{u} \cdot \mathbf{u} = 1$

## Example

The standard basis vectors  $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$  of  $\mathbb{R}^n$  are all unit vectors.

## Example

If  $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ , then is  $\mathbf{x}$  a unit vector?

## Solution

Since  $\mathbf{x} \cdot \mathbf{x} = 5$  and  $\|\mathbf{x}\| = \sqrt{5}$ . However,  $\mathbf{u} = \frac{\mathbf{x}}{\|\mathbf{x}\|} = \frac{\mathbf{x}}{\sqrt{5}}$  is a unit vector. The unit vector  $\mathbf{u}$  is called the *normalization* of  $\mathbf{x}$ .

## Definition

- A bunch of vectors  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p$  is called *orthogonal* if



## Definition

- A bunch of vectors  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p$  is called *orthogonal* if they are all nonzero and  $\mathbf{x}_i \cdot \mathbf{x}_j = 0$  for  $i \neq j$ .

## Definition

- A bunch of vectors  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p$  is called *orthogonal* if they are all nonzero and  $\mathbf{x}_i \cdot \mathbf{x}_j = 0$  for  $i \neq j$ .
- Orthogonal vectors  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_p$  are called *orthonormal*

## Definition

- A bunch of vectors  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p$  is called *orthogonal* if they are all nonzero and  $\mathbf{x}_i \cdot \mathbf{x}_j = 0$  for  $i \neq j$ .
- Orthogonal vectors  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_p$  are called *orthonormal* if they are all unit vectors.

## Example

Let  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{x}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .

## Definition

- A bunch of vectors  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p$  is called *orthogonal* if they are all nonzero and  $\mathbf{x}_i \cdot \mathbf{x}_j = 0$  for  $i \neq j$ .
- Orthogonal vectors  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_p$  are called *orthonormal* if they are all unit vectors.

## Example

Let  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{x}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ . Then they are orthogonal but not orthonormal, since they are not unit vectors.

## Definition

- A bunch of vectors  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p$  is called *orthogonal* if they are all nonzero and  $\mathbf{x}_i \cdot \mathbf{x}_j = 0$  for  $i \neq j$ .
- Orthogonal vectors  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_p$  are called *orthonormal* if they are all unit vectors.

## Example

Let  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{x}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ . Then they are orthogonal but not orthonormal, since they are not unit vectors. We can normalize them to get a orthonormal set  $\mathbf{u}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .

## Definition

- A bunch of vectors  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_p$  is called *orthogonal* if they are all nonzero and  $\mathbf{x}_i \cdot \mathbf{x}_j = 0$  for  $i \neq j$ .
- Orthogonal vectors  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_p$  are called *orthonormal* if they are all unit vectors.

## Example

Let  $\mathbf{x}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{x}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ . Then they are orthogonal but not orthonormal, since they are not unit vectors. We can normalize them to get a orthonormal set  $\mathbf{u}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .

Let  $\mathcal{B} := (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$  be an orthonormal basis for  $\mathbb{R}^n$ , so a basis consisting of unit vectors that are all perpendicular.

Let  $\mathcal{B} := (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$  be an orthonormal basis for  $\mathbb{R}^n$ , so a basis consisting of unit vectors that are all perpendicular. Suppose we want to calculate the coordinates of  $\mathbf{x} \in \mathbb{R}^n$ :

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n.$$



Let  $\mathcal{B} := (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$  be an orthonormal basis for  $\mathbb{R}^n$ , so a basis consisting of unit vectors that are all perpendicular. Suppose we want to calculate the coordinates of  $\mathbf{x} \in \mathbb{R}^n$ :

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n.$$

If this were an arbitrary basis, we would have to solve a system of equations to find the coordinates  $c_1, \dots, c_n$ .

Let  $\mathcal{B} := (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$  be an orthonormal basis for  $\mathbb{R}^n$ , so a basis consisting of unit vectors that are all perpendicular. Suppose we want to calculate the coordinates of  $\mathbf{x} \in \mathbb{R}^n$ :

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n.$$

If this were an arbitrary basis, we would have to solve a system of equations to find the coordinates  $c_1, \dots, c_n$ . Now we know that we have an orthonormal basis things are easier.

Let  $\mathcal{B} := (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$  be an orthonormal basis for  $\mathbb{R}^n$ , so a basis consisting of unit vectors that are all perpendicular. Suppose we want to calculate the coordinates of  $\mathbf{x} \in \mathbb{R}^n$ :

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n.$$

If this were an arbitrary basis, we would have to solve a system of equations to find the coordinates  $c_1, \dots, c_n$ . Now we know that we have an orthonormal basis things are easier. Just calculate

$$\mathbf{u}_1 \cdot \mathbf{x} =$$

Let  $\mathcal{B} := (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$  be an orthonormal basis for  $\mathbb{R}^n$ , so a basis consisting of unit vectors that are all perpendicular. Suppose we want to calculate the coordinates of  $\mathbf{x} \in \mathbb{R}^n$ :

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n.$$

If this were an arbitrary basis, we would have to solve a system of equations to find the coordinates  $c_1, \dots, c_n$ . Now we know that we have an orthonormal basis things are easier. Just calculate

$$\mathbf{u}_1 \cdot \mathbf{x} = \mathbf{u}_1 \cdot (c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n) =$$

Let  $\mathcal{B} := (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$  be an orthonormal basis for  $\mathbb{R}^n$ , so a basis consisting of unit vectors that are all perpendicular. Suppose we want to calculate the coordinates of  $\mathbf{x} \in \mathbb{R}^n$ :

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n.$$

If this were an arbitrary basis, we would have to solve a system of equations to find the coordinates  $c_1, \dots, c_n$ . Now we know that we have an orthonormal basis things are easier. Just calculate

$$\begin{aligned} \mathbf{u}_1 \cdot \mathbf{x} &= \mathbf{u}_1 \cdot (c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n) = \\ &= c_1 \mathbf{u}_1 \cdot \mathbf{u}_1 + c_2 \mathbf{u}_1 \cdot \mathbf{u}_2 + \cdots + c_n \mathbf{u}_1 \cdot \mathbf{u}_n = \end{aligned}$$

Let  $\mathcal{B} := (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$  be an orthonormal basis for  $\mathbb{R}^n$ , so a basis consisting of unit vectors that are all perpendicular. Suppose we want to calculate the coordinates of  $\mathbf{x} \in \mathbb{R}^n$ :

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n.$$

If this were an arbitrary basis, we would have to solve a system of equations to find the coordinates  $c_1, \dots, c_n$ . Now we know that we have an orthonormal basis things are easier. Just calculate

$$\begin{aligned} \mathbf{u}_1 \cdot \mathbf{x} &= \mathbf{u}_1 \cdot (c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n) = \\ &= c_1 \mathbf{u}_1 \cdot \mathbf{u}_1 + c_2 \mathbf{u}_1 \cdot \mathbf{u}_2 + \cdots + c_n \mathbf{u}_1 \cdot \mathbf{u}_n = c_1 \end{aligned}$$

Let  $\mathcal{B} := (\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$  be an orthonormal basis for  $\mathbb{R}^n$ , so a basis consisting of unit vectors that are all perpendicular. Suppose we want to calculate the coordinates of  $\mathbf{x} \in \mathbb{R}^n$ :

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n.$$

If this were an arbitrary basis, we would have to solve a system of equations to find the coordinates  $c_1, \dots, c_n$ . Now we know that we have an orthonormal basis things are easier. Just calculate

$$\begin{aligned} \mathbf{u}_1 \cdot \mathbf{x} &= \mathbf{u}_1 \cdot (c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 + \cdots + c_n \mathbf{u}_n) = \\ &= c_1 \mathbf{u}_1 \cdot \mathbf{u}_1 + c_2 \mathbf{u}_1 \cdot \mathbf{u}_2 + \cdots + c_n \mathbf{u}_1 \cdot \mathbf{u}_n = c_1 \end{aligned}$$

In the same way

$$\mathbf{u}_2 \cdot \mathbf{x} = c_2, \dots, \mathbf{u}_n \cdot \mathbf{x} = c_n$$

### Example

$\mathbf{u}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{u}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  is an orthonormal basis for  $\mathbb{R}^2$ . Let  $\mathbf{x} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ .



### Example

$\mathbf{u}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  is an orthonormal basis for  $\mathbb{R}^2$ . Let  $\mathbf{x} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ .

### Solution

Then

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 =$$

### Example

$\mathbf{u}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  is an orthonormal basis for  $\mathbb{R}^2$ . Let  $\mathbf{x} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ .

### Solution

Then

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 = c_1 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

for

$$c_1 =$$

### Example

$\mathbf{u}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  is an orthonormal basis for  $\mathbb{R}^2$ . Let  $\mathbf{x} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ .

### Solution

Then

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 = c_1 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

for

$$c_1 = \mathbf{u}_1 \cdot \mathbf{x} =$$

### Example

$\mathbf{u}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  is an orthonormal basis for  $\mathbb{R}^2$ . Let  $\mathbf{x} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ .

### Solution

Then

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 = c_1 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

for

$$c_1 = \mathbf{u}_1 \cdot \mathbf{x} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 3 \end{bmatrix} =$$

### Example

$\mathbf{u}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  is an orthonormal basis for  $\mathbb{R}^2$ . Let  $\mathbf{x} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ .

### Solution

Then

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 = c_1 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

for

$$c_1 = \mathbf{u}_1 \cdot \mathbf{x} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \frac{5}{\sqrt{2}}$$

$$c_2 = \mathbf{u}_2 \cdot \mathbf{x} =$$

### Example

$\mathbf{u}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  is an orthonormal basis for  $\mathbb{R}^2$ . Let  $\mathbf{x} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ .

### Solution

Then

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 = c_1 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

for

$$c_1 = \mathbf{u}_1 \cdot \mathbf{x} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \frac{5}{\sqrt{2}}$$

$$c_2 = \mathbf{u}_2 \cdot \mathbf{x} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 3 \end{bmatrix} =$$

### Example

$\mathbf{u}_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ ,  $\mathbf{u}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$  is an orthonormal basis for  $\mathbb{R}^2$ . Let  $\mathbf{x} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ .

### Solution

Then

$$\mathbf{x} = c_1 \mathbf{u}_1 + c_2 \mathbf{u}_2 = c_1 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

for

$$c_1 = \mathbf{u}_1 \cdot \mathbf{x} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \frac{5}{\sqrt{2}}$$

$$c_2 = \mathbf{u}_2 \cdot \mathbf{x} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \frac{-1}{\sqrt{2}}$$

## Theorem

*Let  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  be non-zero and mutually orthogonal. Then  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  are linearly independent.*



## Theorem

*Let  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  be non-zero and mutually orthogonal. Then  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  are linearly independent.*

## Solution

**Proof.** Suppose that

$$c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n = \mathbf{0}.$$

## Theorem

*Let  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  be non-zero and mutually orthogonal. Then  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  are linearly independent.*

## Solution

**Proof.** Suppose that

$$c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n = \mathbf{0}.$$

Take the inner product of  $\mathbf{v}_1$  on both sides.

## Theorem

*Let  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  be non-zero and mutually orthogonal. Then  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  are linearly independent.*

## Solution

**Proof.** Suppose that

$$c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n = \mathbf{0}.$$

Take the inner product of  $\mathbf{v}_1$  on both sides.

$$\mathbf{0} = \mathbf{v}_1 \cdot (c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n)$$

## Theorem

*Let  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  be non-zero and mutually orthogonal. Then  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  are linearly independent.*

## Solution

**Proof.** Suppose that

$$c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n = \mathbf{0}.$$

Take the inner product of  $\mathbf{v}_1$  on both sides.

$$\begin{aligned} 0 &= \mathbf{v}_1 \cdot (c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n) \\ &= c_1\mathbf{v}_1 \cdot \mathbf{v}_1 + c_2\mathbf{v}_1 \cdot \mathbf{v}_2 + \cdots + c_n\mathbf{v}_1 \cdot \mathbf{v}_n \end{aligned}$$

## Theorem

*Let  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  be non-zero and mutually orthogonal. Then  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  are linearly independent.*

## Solution

**Proof.** Suppose that

$$c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n = \mathbf{0}.$$

Take the inner product of  $\mathbf{v}_1$  on both sides.

$$\begin{aligned} \mathbf{0} &= \mathbf{v}_1 \cdot (c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n) \\ &= c_1 \mathbf{v}_1 \cdot \mathbf{v}_1 + c_2 \mathbf{v}_1 \cdot \mathbf{v}_2 + \cdots + c_n \mathbf{v}_1 \cdot \mathbf{v}_n \\ &= c_1 \mathbf{v}_1 \cdot \mathbf{v}_1 = c_1 \|\mathbf{v}_1\|^2 \end{aligned}$$

## Theorem

Let  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  be non-zero and mutually orthogonal. Then  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  are linearly independent.

## Solution

**Proof.** Suppose that

$$c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n = \mathbf{0}.$$

Take the inner product of  $\mathbf{v}_1$  on both sides.

$$\begin{aligned} \mathbf{0} &= \mathbf{v}_1 \cdot (c_1 \mathbf{v}_1 + \cdots + c_n \mathbf{v}_n) \\ &= c_1 \mathbf{v}_1 \cdot \mathbf{v}_1 + c_2 \mathbf{v}_1 \cdot \mathbf{v}_2 + \cdots + c_n \mathbf{v}_1 \cdot \mathbf{v}_n \\ &= c_1 \mathbf{v}_1 \cdot \mathbf{v}_1 = c_1 \|\mathbf{v}_1\|^2 \end{aligned}$$

But  $\|\mathbf{v}_1\| \neq 0$  and so  $c_1 = 0$ .

## Theorem

Let  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  be non-zero and mutually orthogonal. Then  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  are linearly independent.

## Solution

**Proof.** Suppose that

$$c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n = \mathbf{0}.$$

Take the inner product of  $\mathbf{v}_1$  on both sides.

$$\begin{aligned} \mathbf{0} &= \mathbf{v}_1 \cdot (c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n) \\ &= c_1\mathbf{v}_1 \cdot \mathbf{v}_1 + c_2\mathbf{v}_1 \cdot \mathbf{v}_2 + \cdots + c_n\mathbf{v}_1 \cdot \mathbf{v}_n \\ &= c_1\mathbf{v}_1 \cdot \mathbf{v}_1 = c_1\|\mathbf{v}_1\|^2 \end{aligned}$$

But  $\|\mathbf{v}_1\| \neq 0$  and so  $c_1 = 0$ . Similarly, we find that  $c_2 = 0, \dots, c_n = 0$ . Therefore, the vectors are independent.

## Orthogonality and the Fundamental subspaces



### Example

Let  $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$ . Find  $Nul(A)$  and  $Col(A^T)$ .

### Example

Let  $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$ . Find  $Nul(A)$  and  $Col(A^T)$ .

### Solution

### Example

Let  $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$ . Find  $Nul(A)$  and  $Col(A^T)$ .

### Solution

Note that  $Nul(A)$  and  $Col(A^T)$  both are subspace of  $\mathbb{R}^2$ .

$$Nul(A) =$$

### Example

Let  $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$ . Find  $Nul(A)$  and  $Col(A^T)$ .

### Solution

Note that  $Nul(A)$  and  $Col(A^T)$  both are subspace of  $\mathbb{R}^2$ .

$$Nul(A) = span \left\{ \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right\},$$

### Example

Let  $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$ . Find  $Nul(A)$  and  $Col(A^T)$ .

### Solution

Note that  $Nul(A)$  and  $Col(A^T)$  both are subspace of  $\mathbb{R}^2$ .

$$Nul(A) = span \left\{ \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right\}, \quad Col(A^T) =$$

### Example

Let  $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$ . Find  $Nul(A)$  and  $Col(A^T)$ .

### Solution

Note that  $Nul(A)$  and  $Col(A^T)$  both are subspace of  $\mathbb{R}^2$ .

$$Nul(A) = span \left\{ \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right\}, \quad Col(A^T) = span \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$$

### Example

Let  $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 6 \end{bmatrix}$ . Find  $Nul(A)$  and  $Col(A^T)$ .

### Solution

Note that  $Nul(A)$  and  $Col(A^T)$  both are subspace of  $\mathbb{R}^2$ .

$$Nul(A) = span \left\{ \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right\}, \quad Col(A^T) = span \left\{ \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\}$$

The basis vectors for the null and row space are orthogonal.

$$\begin{bmatrix} -2 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \end{bmatrix} = 0$$

### Example

Repeat for  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .



### Example

Repeat for  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

### Solution

$$\begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix} \xrightarrow{RREF}$$

### Example

Repeat for  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

### Solution

$$\begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

### Example

Repeat for  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

### Solution

$$\begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Nul}(A) =$$

### Example

Repeat for  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

### Solution

$$\begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Nul}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\},$$

### Example

Repeat for  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

### Solution

$$\begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Nul}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}, \quad \text{Col}(A^T) =$$

### Example

Repeat for  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

### Solution

$$\begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix} \xrightarrow{RREF} \begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{Nul}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}, \quad \text{Col}(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

Again, the basis for the null space is orthogonal to the basis for the row space.

$$\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} = 0, \quad \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 0.$$

Again, the basis for the null space is orthogonal to the basis for the row space.

$$\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} = 0, \quad \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 0.$$

Since  $\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}$  is orthogonal to both basis vectors for the row space, it's orthogonal to every vector in the row space.



Again, the basis for the null space is orthogonal to the basis for the row space.

$$\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} = 0, \quad \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 0.$$

Since  $\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}$  is orthogonal to both basis vectors for the row space, it's orthogonal to *every* vector in the row space.

It turns out this is true for the null and row space of any matrix  $A$ . That is, vectors in  $Nul(A)$  are orthogonal to vectors in  $Col(A^T)$  for *all* matrices  $A$ .

## Fundamental Theorem of Linear Algebra (Revisited)

## Definition

Let  $W$  be a subspace of  $\mathbb{R}^n$  and  $\mathbf{v} \in \mathbb{R}^n$ .

- $\mathbf{v}$  is **orthogonal** to  $W$  if  $\mathbf{v} \cdot \mathbf{w} = 0$  for all  $\mathbf{w} \in W$ . ( $\iff \mathbf{v}$  is orthogonal to each vector in a basis for  $W$ .)

## Definition

Let  $W$  be a subspace of  $\mathbb{R}^n$  and  $\mathbf{v} \in \mathbb{R}^n$ .

- $\mathbf{v}$  is **orthogonal** to  $W$  if  $\mathbf{v} \cdot \mathbf{w} = 0$  for all  $\mathbf{w} \in W$ . (  $\iff \mathbf{v}$  is orthogonal to each vector in a basis for  $W$ .)
- Another subspace  $V$  is **orthogonal** to  $W$  if every vector in  $V$  is orthogonal to  $W$ .

## Definition

Let  $W$  be a subspace of  $\mathbb{R}^n$  and  $\mathbf{v} \in \mathbb{R}^n$ .

- $\mathbf{v}$  is **orthogonal** to  $W$  if  $\mathbf{v} \cdot \mathbf{w} = 0$  for all  $\mathbf{w} \in W$ . ( $\iff \mathbf{v}$  is orthogonal to each vector in a basis for  $W$ .)
- Another subspace  $V$  is **orthogonal** to  $W$  if every vector in  $V$  is orthogonal to  $W$ .
- The **orthogonal complement** of  $W$  is the space  $W^\perp$

## Definition

Let  $W$  be a subspace of  $\mathbb{R}^n$  and  $\mathbf{v} \in \mathbb{R}^n$ .

- $\mathbf{v}$  is **orthogonal** to  $W$  if  $\mathbf{v} \cdot \mathbf{w} = 0$  for all  $\mathbf{w} \in W$ . ( $\iff \mathbf{v}$  is orthogonal to each vector in a basis for  $W$ .)
- Another subspace  $V$  is **orthogonal** to  $W$  if every vector in  $V$  is orthogonal to  $W$ .
- The **orthogonal complement** of  $W$  is the space  $W^\perp$  of all vectors that are orthogonal to  $W$ .

## Definition

Let  $W$  be a subspace of  $\mathbb{R}^n$  and  $\mathbf{v} \in \mathbb{R}^n$ .

- $\mathbf{v}$  is **orthogonal** to  $W$  if  $\mathbf{v} \cdot \mathbf{w} = 0$  for all  $\mathbf{w} \in W$ . ( $\iff \mathbf{v}$  is orthogonal to each vector in a basis for  $W$ .)
- Another subspace  $V$  is **orthogonal** to  $W$  if every vector in  $V$  is orthogonal to  $W$ .
- The **orthogonal complement** of  $W$  is the space  $W^\perp$  of all vectors that are orthogonal to  $W$ . (Show that the orthogonal complement of any vector space is also a vector space.)

## Example

Let  $V = \text{Span} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$  and  $W = \text{Span} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ .



## Example

Let  $V = \text{Span} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$  and  $W = \text{Span} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ . Then

- $V \perp W$ , because every vector of  $V$  is perp to each vector in  $W$ .

## Example

Let  $V = \text{Span} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$  and  $W = \text{Span} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ . Then

- $V \perp W$ , because every vector of  $V$  is perp to each vector in  $W$ .
- It is not true that  $V^\perp = W$  since  $V^\perp$  consists of *all* vectors in  $\mathbb{R}^3$  perp to  $V$ . Which vectors are missing?

## Example

Let  $V = \text{Span} \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$  and  $W = \text{Span} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ . Then

- $V \perp W$ , because every vector of  $V$  is perp to each vector in  $W$ .
- It is not true that  $V^\perp = W$  since  $V^\perp$  consists of *all* vectors in  $\mathbb{R}^3$  perp to  $V$ . Which vectors are missing?
- $V^\perp = \text{Span} \left( \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right)$

## Example

In the last example,  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

## Example

In the last example,  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

We found that

$$\text{Nul}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\},$$

## Example

In the last example,  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

We found that

$$\text{Nul}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}, \quad \text{Col}(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

are orthogonal subspaces.

## Example

In the last example,  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

We found that

$$\text{Nul}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}, \quad \text{Col}(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

are orthogonal subspaces. Indeed,  $\text{Nul}(A)$  and  $\text{Col}(A^T)$  are orthogonal complements.

## Example

In the last example,  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 0 \\ 3 & 6 & 0 \end{bmatrix}$ .

We found that

$$\text{Nul}(A) = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}, \quad \text{Col}(A^T) = \text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

are orthogonal subspaces. Indeed,  $\text{Nul}(A)$  and  $\text{Col}(A^T)$  are orthogonal complements.

Why?

## Solution

Because  $\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}$ ,  $\begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$  are orthogonal (so independent), and so they're a basis for all of  $\mathbb{R}^3$ .



## Remark

In the last example,  $Nul(A)$  and  $Col(A)$  both happen to be subspaces of  $\mathbb{R}^3$  (because  $A$  was a square  $3 \times 3$  matrix).

$$Nul(A) = span \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\},$$

## Remark

In the last example,  $Nul(A)$  and  $Col(A)$  both happen to be subspaces of  $\mathbb{R}^3$  (because  $A$  was a square  $3 \times 3$  matrix).

$$Nul(A) = span \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}, \quad Col(A) = span \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\}$$

## Remark

In the last example,  $Nul(A)$  and  $Col(A)$  both happen to be subspaces of  $\mathbb{R}^3$  (because  $A$  was a square  $3 \times 3$  matrix).

$$Nul(A) = span \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}, \quad Col(A) = span \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\}$$

However, these spaces are **not** orthogonal. Why?

## Remark

In the last example,  $Nul(A)$  and  $Col(A)$  both happen to be subspaces of  $\mathbb{R}^3$  (because  $A$  was a square  $3 \times 3$  matrix).

$$Nul(A) = span \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right\}, \quad Col(A) = span \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\}$$

However, these spaces are **not** orthogonal. Why?

## Solution

$$\begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \neq 0$$

## Theorem

*Let  $A$  be an  $m \times n$  matrix of rank  $r$ .*

## Theorem

*Let  $A$  be an  $m \times n$  matrix of rank  $r$ .*

## Theorem

*Let  $A$  be an  $m \times n$  matrix of rank  $r$ .*

- $\dim \text{Col}(A) = r$  *(subspace of  $\mathbb{R}^m$ )*

## Theorem

*Let  $A$  be an  $m \times n$  matrix of rank  $r$ .*

- $\dim \text{Col}(A) = r$  *(subspace of  $\mathbb{R}^m$ )*



## Theorem

*Let  $A$  be an  $m \times n$  matrix of rank  $r$ .*

- $\dim \text{Col}(A) = r$  (subspace of  $\mathbb{R}^m$ )
- $\dim \text{Col}(A^T) = r$  (subspace of  $\mathbb{R}^n$ )

## Theorem

*Let  $A$  be an  $m \times n$  matrix of rank  $r$ .*

- $\dim \operatorname{Col}(A) = r$  (subspace of  $\mathbb{R}^m$ )
- $\dim \operatorname{Col}(A^T) = r$  (subspace of  $\mathbb{R}^n$ )

## Theorem

*Let  $A$  be an  $m \times n$  matrix of rank  $r$ .*

- $\dim \operatorname{Col}(A) = r$  (subspace of  $\mathbb{R}^m$ )
- $\dim \operatorname{Col}(A^T) = r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A) = n - r$  (subspace of  $\mathbb{R}^n$ )

## Theorem

*Let  $A$  be an  $m \times n$  matrix of rank  $r$ .*

- $\dim \operatorname{Col}(A) = r$  (subspace of  $\mathbb{R}^m$ )
- $\dim \operatorname{Col}(A^T) = r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A) = n - r$  (subspace of  $\mathbb{R}^n$ )

## Theorem

Let  $A$  be an  $m \times n$  matrix of rank  $r$ .

- $\dim \operatorname{Col}(A) = r$  (subspace of  $\mathbb{R}^m$ )
- $\dim \operatorname{Col}(A^T) = r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A) = n - r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A^T) = m - r$  (subspace of  $\mathbb{R}^m$ )

## Theorem

Let  $A$  be an  $m \times n$  matrix of rank  $r$ .

- $\dim \operatorname{Col}(A) = r$  (subspace of  $\mathbb{R}^m$ )
- $\dim \operatorname{Col}(A^T) = r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A) = n - r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A^T) = m - r$  (subspace of  $\mathbb{R}^m$ )

## Theorem

Let  $A$  be an  $m \times n$  matrix of rank  $r$ .

- $\dim \operatorname{Col}(A) = r$  (subspace of  $\mathbb{R}^m$ )
- $\dim \operatorname{Col}(A^T) = r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A) = n - r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A^T) = m - r$  (subspace of  $\mathbb{R}^m$ )
- $\operatorname{Nul}(A)^\perp = \operatorname{Col}(A^T)$  (both subspaces of  $\mathbb{R}^n$ )

## Theorem

Let  $A$  be an  $m \times n$  matrix of rank  $r$ .

- $\dim \operatorname{Col}(A) = r$  (subspace of  $\mathbb{R}^m$ )
- $\dim \operatorname{Col}(A^T) = r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A) = n - r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A^T) = m - r$  (subspace of  $\mathbb{R}^m$ )
- $\operatorname{Nul}(A)^\perp = \operatorname{Col}(A^T)$  (both subspaces of  $\mathbb{R}^n$ )



## Theorem

Let  $A$  be an  $m \times n$  matrix of rank  $r$ .

- $\dim \operatorname{Col}(A) = r$  (subspace of  $\mathbb{R}^m$ )
- $\dim \operatorname{Col}(A^T) = r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A) = n - r$  (subspace of  $\mathbb{R}^n$ )
- $\dim \operatorname{Nul}(A^T) = m - r$  (subspace of  $\mathbb{R}^m$ )
- $\operatorname{Nul}(A)^\perp = \operatorname{Col}(A^T)$  (both subspaces of  $\mathbb{R}^n$ )

Note that  $\dim \operatorname{Nul}(A) + \dim \operatorname{Col}(A^T) = n$ .

## Theorem

Let  $A$  be an  $m \times n$  matrix of rank  $r$ .

- $\dim \operatorname{Col}(A) = r$  (subspace of  $\mathbb{R}^m$ )
  - $\dim \operatorname{Col}(A^T) = r$  (subspace of  $\mathbb{R}^n$ )
  - $\dim \operatorname{Nul}(A) = n - r$  (subspace of  $\mathbb{R}^n$ )
  - $\dim \operatorname{Nul}(A^T) = m - r$  (subspace of  $\mathbb{R}^m$ )
  - $\operatorname{Nul}(A)^\perp = \operatorname{Col}(A^T)$  (both subspaces of  $\mathbb{R}^n$ )
- Note that  $\dim \operatorname{Nul}(A) + \dim \operatorname{Col}(A^T) = n$ .
- $\operatorname{Nul}(A^T)^\perp = \operatorname{Col}(A)$

## Example

Let  $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ .

- $\text{Col}(A) =$

## Example

Let  $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ .

- $\text{Col}(A) = \text{Span} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .

## Example

Let  $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ .

- $Col(A) = Span \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .
- $Nul(A) =$

## Example

Let  $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ .

- $Col(A) = Span \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .
- $Nul(A) = Span \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .

## Example

Let  $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ .

- $\text{Col}(A) = \text{Span} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .
- $\text{Nul}(A) = \text{Span} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .
- $\text{Col}(A^T) =$

## Example

Let  $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ .

- $\text{Col}(A) = \text{Span} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .
- $\text{Nul}(A) = \text{Span} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .
- $\text{Col}(A^T) = \text{Span} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .



## Example

Let  $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ .

- $\text{Col}(A) = \text{Span} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .
- $\text{Nul}(A) = \text{Span} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .
- $\text{Col}(A^T) = \text{Span} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .
- $\text{Nul}(A^T) =$

## Example

Let  $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ .

- $\text{Col}(A) = \text{Span} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .
- $\text{Nul}(A) = \text{Span} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .
- $\text{Col}(A^T) = \text{Span} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .
- $\text{Nul}(A^T) = \text{Span} \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ .

## Example

Let  $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ .

- $\text{Col}(A) = \text{Span} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .
- $\text{Nul}(A) = \text{Span} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .
- $\text{Col}(A^T) = \text{Span} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ .
- $\text{Nul}(A^T) = \text{Span} \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ .