

Math 415 - Lecture 8

Inverses.

Wednesday September 11th 2015

Textbook: Chapter 1.6

Textbook: Chapter 1.6

Suggested Practice Exercise: Chapter 1.6 Exercise 1, 2, 4, 6, 10,
11, 18, 35, 36, 37, 38, 40, 49, 50

Textbook: Chapter 1.6

Suggested Practice Exercise: Chapter 1.6 Exercise 1, 2, 4, 6, 10,
11, 18, 35, 36, 37, 38, 40, 49, 50

Khan Academy Video: Inverse Matrix (part I), Inverse Matrix (part
II)

Review

- Elementary matrices perform row operations:

$$\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ -2a + d & -2b + e & -2c + f \\ g & h & i \end{bmatrix}$$

- Elementary matrices perform row operations:

$$\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ -2a + d & -2b + e & -2c + f \\ g & h & i \end{bmatrix}$$

- Gaussian elimination on A gives a decomposition $A = LU$:

$$\begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

U is the echelon form, L records the reverse of the row operations we did.

- Elementary matrices perform row operations:

$$\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = \begin{bmatrix} a & b & c \\ -2a + d & -2b + e & -2c + f \\ g & h & i \end{bmatrix}$$

- Gaussian elimination on A gives a decomposition $A = LU$:

$$\begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

U is the echelon form, L records the reverse of the row operations we did.

- LU decomposition lets us solve $A\mathbf{x} = \mathbf{b}$ quickly for many different \mathbf{b} .

Today's goal

- We know how to reverse a single row operation:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- We know how to reverse a single row operation:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Inverting a more complicated matrix is harder:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & b & 1 \end{bmatrix}^{-1} =$$

- We know how to reverse a single row operation:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Inverting a more complicated matrix is harder:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & b & 1 \end{bmatrix}^{-1} =$$

- We know how to reverse a single row operation:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Inverting a more complicated matrix is harder:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & b & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ ab & -b & 1 \end{bmatrix}$$

- We know how to reverse a single row operation:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Inverting a more complicated matrix is harder:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & b & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ ab & -b & 1 \end{bmatrix}$$

Goal today: how to find an “inverse” to any (square!) matrix.

- We know how to reverse a single row operation:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

- Inverting a more complicated matrix is harder:

$$\begin{bmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ 0 & b & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ ab & -b & 1 \end{bmatrix}$$

Goal today: how to find an “inverse” to any (square!) matrix.

Today A will be an $n \times n$ matrix

The inverse of a matrix

The **inverse** of a real number a is denoted by a^{-1} . For example, $7^{-1} = 1/7$ and

$$7 \cdot 7^{-1} = 7^{-1} \cdot 7 = 1.$$

The **inverse** of a real number a is denoted by a^{-1} . For example, $7^{-1} = 1/7$ and

$$7 \cdot 7^{-1} = 7^{-1} \cdot 7 = 1.$$

Remember that the identity matrix I_n is the $n \times n$ -matrix

$$I_n = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

The **inverse** of a real number a is denoted by a^{-1} . For example, $7^{-1} = 1/7$ and

$$7 \cdot 7^{-1} = 7^{-1} \cdot 7 = 1.$$

Remember that the identity matrix I_n is the $n \times n$ -matrix

$$I_n = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

Definition

An $n \times n$ matrix A is said to be **invertible** if there is an $n \times n$ matrix C satisfying

$$CA = AC = I_n$$

The **inverse** of a real number a is denoted by a^{-1} . For example, $7^{-1} = 1/7$ and

$$7 \cdot 7^{-1} = 7^{-1} \cdot 7 = 1.$$

Remember that the identity matrix I_n is the $n \times n$ -matrix

$$I_n = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

Definition

An $n \times n$ matrix A is said to be **invertible** if there is an $n \times n$ matrix C satisfying

$$CA = AC = I_n$$

where I_n is the $n \times n$ identity matrix. We call C the

The **inverse** of a real number a is denoted by a^{-1} . For example, $7^{-1} = 1/7$ and

$$7 \cdot 7^{-1} = 7^{-1} \cdot 7 = 1.$$

Remember that the identity matrix I_n is the $n \times n$ -matrix

$$I_n = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

Definition

An $n \times n$ matrix A is said to be **invertible** if there is an $n \times n$ matrix C satisfying

$$CA = AC = I_n$$

where I_n is the $n \times n$ identity matrix. We call C the **inverse** of A .

Example

We already know that an elementary matrix is invertible:

$$\begin{bmatrix} 1 & 0 & 0 \\ 8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Example

We already know that an elementary matrix is invertible:

$$\begin{bmatrix} 1 & 0 & 0 \\ 8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

In fact:

$$\begin{bmatrix} 1 & 0 & 0 \\ 8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 0 & 0 \\ -8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

(Check this at home!)

Example

We already know that an elementary matrix is invertible:

$$\begin{bmatrix} 1 & 0 & 0 \\ 8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ -8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

In fact:

$$\begin{bmatrix} 1 & 0 & 0 \\ 8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
$$\begin{bmatrix} 1 & 0 & 0 \\ -8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 8 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

(Check this at home!) So the definition works!

Theorem

Let A be an invertible matrix, then its inverse C is unique.

Theorem

Let A be an invertible matrix, then its inverse C is unique.

Proof.

Assume B and C are both inverses of A .

Theorem

Let A be an invertible matrix, then its inverse C is unique.

Proof.

Assume B and C are both inverses of A . Then

$$B = BI_n =$$

Theorem

Let A be an invertible matrix, then its inverse C is unique.

Proof.

Assume B and C are both inverses of A . Then

$$B = BI_n = BAC =$$

Theorem

Let A be an invertible matrix, then its inverse C is unique.

Proof.

Assume B and C are both inverses of A . Then

$$B = BI_n = BAC = I_n C =$$

Theorem

Let A be an invertible matrix, then its inverse C is unique.

Proof.

Assume B and C are both inverses of A . Then

$$B = BI_n = BAC = I_n C = C$$



- We will write A^{-1} for the inverse of A . Multiplying by A^{-1} is like “dividing by A .”

- We will write A^{-1} for the inverse of A . Multiplying by A^{-1} is like “dividing by A .”
- Do not write $\frac{A}{B}$. Why?

- We will write A^{-1} for the inverse of A . Multiplying by A^{-1} is like “dividing by A .”
- Do not write $\frac{A}{B}$. Why?
It is unclear whether this means AB^{-1} or $B^{-1}A$, and these two matrices are *different*.

- We will write A^{-1} for the inverse of A . Multiplying by A^{-1} is like “dividing by A .”
- Do not write $\frac{A}{B}$. Why?
It is unclear whether this means AB^{-1} or $B^{-1}A$, and these two matrices are *different*.
- Fact: if $AB = I$ then $A^{-1} = B$ and so $BA = I$. (Not so easy to show at this stage.)

Remark

Not all $n \times n$ matrices are invertible.

Remark

Not all $n \times n$ matrices are invertible. For example, the 2×2 matrix

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

is not invertible.

Remark

Not all $n \times n$ matrices are invertible. For example, the 2×2 matrix

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

is not invertible. Try to find an inverse!

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Remark

Not all $n \times n$ matrices are invertible. For example, the 2×2 matrix

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

is not invertible. Try to find an inverse!

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} c & d \\ 0 & 0 \end{bmatrix}$$

Remark

Not all $n \times n$ matrices are invertible. For example, the 2×2 matrix

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

is not invertible. Try to find an inverse!

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} c & d \\ 0 & 0 \end{bmatrix} \neq I_2$$

Remark

Not all $n \times n$ matrices are invertible. For example, the 2×2 matrix

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

is not invertible. Try to find an inverse!

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} c & d \\ 0 & 0 \end{bmatrix} \neq I_2$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Remark

Not all $n \times n$ matrices are invertible. For example, the 2×2 matrix

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

is not invertible. Try to find an inverse!

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} c & d \\ 0 & 0 \end{bmatrix} \neq I_2$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & a \\ 0 & c \end{bmatrix}$$

Remark

Not all $n \times n$ matrices are invertible. For example, the 2×2 matrix

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

is not invertible. Try to find an inverse!

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} c & d \\ 0 & 0 \end{bmatrix} \neq I_2$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & a \\ 0 & c \end{bmatrix} \neq I_2$$

Remark

Not all $n \times n$ matrices are invertible. For example, the 2×2 matrix

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

is not invertible. Try to find an inverse!

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} c & d \\ 0 & 0 \end{bmatrix} \neq I_2$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & a \\ 0 & c \end{bmatrix} \neq I_2$$

Definition

A matrix which is *not* invertible is sometimes called a **singular** matrix.

Remark

Not all $n \times n$ matrices are invertible. For example, the 2×2 matrix

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

is not invertible. Try to find an inverse!

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} c & d \\ 0 & 0 \end{bmatrix} \neq I_2$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & a \\ 0 & c \end{bmatrix} \neq I_2$$

Definition

A matrix which is *not* invertible is sometimes called a **singular** matrix. An invertible matrix is also called **nonsingular** matrix.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. If $ad - bc \neq 0$, then A is invertible and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. If $ad - bc \neq 0$, then A is invertible and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

If $ad - bc = 0$, then A is not invertible.

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. If $ad - bc \neq 0$, then A is invertible and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

If $ad - bc = 0$, then A is not invertible.

Proof.

Calculate

$$\frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} =$$

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. If $ad - bc \neq 0$, then A is invertible and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

If $ad - bc = 0$, then A is not invertible.

Proof.

Calculate

$$\frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} =$$

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. If $ad - bc \neq 0$, then A is invertible and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

If $ad - bc = 0$, then A is not invertible.

Proof.

Calculate

$$\begin{aligned} \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} &= \frac{1}{ad - bc} \begin{bmatrix} da - bc & db - bd \\ -ca + ac & -cb + ad \end{bmatrix} \\ &= \end{aligned}$$

Theorem

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. If $ad - bc \neq 0$, then A is invertible and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

If $ad - bc = 0$, then A is not invertible.

Proof.

Calculate

$$\begin{aligned} \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} &= \frac{1}{ad - bc} \begin{bmatrix} da - bc & db - bd \\ -ca + ac & -cb + ad \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

Quick question: when is the 1×1 matrix $[a]$ invertible?

Quick question: when is the 1×1 matrix $[a]$ invertible?
When $a \neq 0$. Its inverse is $[\frac{1}{a}]$.

Quick question: when is the 1×1 matrix $[a]$ invertible?
When $a \neq 0$. Its inverse is $[\frac{1}{a}]$.

Theorem

If A is an invertible $n \times n$ matrix, then for each \mathbf{b} in \mathbb{R}^n , the equation $A\mathbf{x} = \mathbf{b}$ has the unique solution $\mathbf{x} = A^{-1}\mathbf{b}$.

Theorem

If A is an invertible $n \times n$ matrix, then for each \mathbf{b} in \mathbb{R}^n , the equation $A\mathbf{x} = \mathbf{b}$ has the unique solution $\mathbf{x} = A^{-1}\mathbf{b}$.

Proof.

The vector $A^{-1}\mathbf{b}$ is a solution, because

$$A(A^{-1}\mathbf{b}) = (AA^{-1})\mathbf{b} = I_n\mathbf{b} = \mathbf{b}$$

Theorem

If A is an invertible $n \times n$ matrix, then for each \mathbf{b} in \mathbb{R}^n , the equation $A\mathbf{x} = \mathbf{b}$ has the unique solution $\mathbf{x} = A^{-1}\mathbf{b}$.

Proof.

The vector $A^{-1}\mathbf{b}$ is a solution, because

$$A(A^{-1}\mathbf{b}) = (AA^{-1})\mathbf{b} = I_n\mathbf{b} = \mathbf{b}$$

Suppose there is another solution \mathbf{w} , then

Theorem

If A is an invertible $n \times n$ matrix, then for each \mathbf{b} in \mathbb{R}^n , the equation $A\mathbf{x} = \mathbf{b}$ has the unique solution $\mathbf{x} = A^{-1}\mathbf{b}$.

Proof.

The vector $A^{-1}\mathbf{b}$ is a solution, because

$$A(A^{-1}\mathbf{b}) = (AA^{-1})\mathbf{b} = I_n\mathbf{b} = \mathbf{b}$$

Suppose there is another solution \mathbf{w} , then

$$A\mathbf{w} = \mathbf{b}$$

$$A^{-1}A\mathbf{w} = A^{-1}\mathbf{b}$$

$$I_n\mathbf{w} = A^{-1}\mathbf{b}$$

$$\mathbf{w} = A^{-1}\mathbf{b}$$



Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Solution

Matrix form of the linear system:

Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Solution

Matrix form of the linear system:

$$\begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Solution

Matrix form of the linear system:

$$\begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$A^{-1} =$$

Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Solution

Matrix form of the linear system:

$$\begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{14-15} \begin{bmatrix} -2 & -3 \\ -5 & -7 \end{bmatrix}$$

Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Solution

Matrix form of the linear system:

$$\begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{14-15} \begin{bmatrix} -2 & -3 \\ -5 & -7 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix}$$

Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Solution

Matrix form of the linear system:

$$\begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{14-15} \begin{bmatrix} -2 & -3 \\ -5 & -7 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix}$$

$$\mathbf{x} =$$

Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Solution

Matrix form of the linear system:

$$\begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{14-15} \begin{bmatrix} -2 & -3 \\ -5 & -7 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix}$$

$$\mathbf{x} = A^{-1}\mathbf{b}$$

Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Solution

Matrix form of the linear system:

$$\begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{14-15} \begin{bmatrix} -2 & -3 \\ -5 & -7 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix}$$

$$\mathbf{x} = A^{-1}\mathbf{b} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Solution

Matrix form of the linear system:

$$\begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{14-15} \begin{bmatrix} -2 & -3 \\ -5 & -7 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix}$$

$$\mathbf{x} = A^{-1}\mathbf{b} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 7 \\ 17 \end{bmatrix}$$

Example

Use the inverse of $A = \begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix}$ to solve

$$-7x_1 + 3x_2 = 2$$

$$5x_1 - 2x_2 = 1$$

Solution

Matrix form of the linear system:

$$\begin{bmatrix} -7 & 3 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{14-15} \begin{bmatrix} -2 & -3 \\ -5 & -7 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix}$$

$$\mathbf{x} = A^{-1}\mathbf{b} = \begin{bmatrix} 2 & 3 \\ 5 & 7 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 7 \\ 17 \end{bmatrix}$$

Check this works!

Computational rules

Theorem

Suppose A and B are invertible. Then the following results hold:

Theorem

Suppose A and B are invertible. Then the following results hold:

- (a) A^{-1} is invertible and $(A^{-1})^{-1} = A$ (i.e. A is the inverse of A^{-1}).

Theorem

Suppose A and B are invertible. Then the following results hold:

- (a) A^{-1} is invertible and $(A^{-1})^{-1} = A$ (i.e. A is the inverse of A^{-1}).
- (b) AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$

Theorem

Suppose A and B are invertible. Then the following results hold:

- (a) A^{-1} is invertible and $(A^{-1})^{-1} = A$ (i.e. A is the inverse of A^{-1}).
- (b) AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$
- (c) A^T is invertible and $(A^T)^{-1} = (A^{-1})^T$

Theorem

Suppose A and B are invertible. Then the following results hold:

- (a) A^{-1} is invertible and $(A^{-1})^{-1} = A$ (i.e. A is the inverse of A^{-1}).
- (b) AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$
- (c) A^T is invertible and $(A^T)^{-1} = (A^{-1})^T$

Theorem

Suppose A and B are invertible. Then the following results hold:

- (a) A^{-1} is invertible and $(A^{-1})^{-1} = A$ (i.e. A is the inverse of A^{-1}).*
- (b) AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$*
- (c) A^T is invertible and $(A^T)^{-1} = (A^{-1})^T$*

Note: the inverse of the product is the product of inverses **in opposite order**.

Theorem

Suppose A and B are invertible. Then the following results hold:

- (a) A^{-1} is invertible and $(A^{-1})^{-1} = A$ (i.e. A is the inverse of A^{-1}).*
- (b) AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$*
- (c) A^T is invertible and $(A^T)^{-1} = (A^{-1})^T$*

Note: the inverse of the product is the product of inverses **in opposite order**. Think about putting on socks and shoes. How do you undo those two operations?

Proof.

(a) A^{-1} is invertible and $(A^{-1})^{-1} = A$

Proof.

(a) A^{-1} is invertible and $(A^{-1})^{-1} = A$

$$AA^{-1} = I = A^{-1}A \quad \checkmark$$

Proof.

(a) A^{-1} is invertible and $(A^{-1})^{-1} = A$

$$AA^{-1} = I = A^{-1}A \quad \checkmark$$

(b) AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$

Proof.

(a) A^{-1} is invertible and $(A^{-1})^{-1} = A$

$$AA^{-1} = I = A^{-1}A \quad \checkmark$$

(b) AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$

$$(B^{-1}A^{-1})(AB) = B^{-1}IB = B^{-1}B = I \quad \checkmark$$

$$(AB)(B^{-1}A^{-1}) = AIA^{-1} = AA^{-1} = I \quad \checkmark$$

Proof.

(a) A^{-1} is invertible and $(A^{-1})^{-1} = A$

$$AA^{-1} = I = A^{-1}A \quad \checkmark$$

(b) AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$

$$(B^{-1}A^{-1})(AB) = B^{-1}IB = B^{-1}B = I \quad \checkmark$$

$$(AB)(B^{-1}A^{-1}) = AIA^{-1} = AA^{-1} = I \quad \checkmark$$

(c) A^T is invertible and $(A^T)^{-1} = (A^{-1})^T$

Proof.

(a) A^{-1} is invertible and $(A^{-1})^{-1} = A$

$$AA^{-1} = I = A^{-1}A \quad \checkmark$$

(b) AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$

$$(B^{-1}A^{-1})(AB) = B^{-1}IB = B^{-1}B = I \quad \checkmark$$

$$(AB)(B^{-1}A^{-1}) = AIA^{-1} = AA^{-1} = I \quad \checkmark$$

(c) A^T is invertible and $(A^T)^{-1} = (A^{-1})^T$

$$A^T(A^{-1})^T = (A^{-1}A)^T = I^T = I \quad \checkmark$$

$$(A^{-1})^T A^T = (AA^{-1})^T = I^T = I \quad \checkmark$$

An algorithm for computing the inverse matrix

Idea:

- To solve $Ax = b$ we row reduce $[A \mid b]$.

Idea:

- To solve $Ax = b$ we row reduce $[A \mid b]$.
- To solve $AX = I_n$ we row reduce $[A \mid I]$.

Idea:

- To solve $Ax = b$ we row reduce $[A \mid b]$.
- To solve $AX = I_n$ we row reduce $[A \mid I]$.

Theorem

An $n \times n$ matrix A is invertible if and only if A is row equivalent to I_n ,

Idea:

- To solve $Ax = b$ we row reduce $[A \mid b]$.
- To solve $AX = I_n$ we row reduce $[A \mid I]$.

Theorem

An $n \times n$ matrix A is invertible if and only if A is row equivalent to I_n ,

Idea:

- To solve $Ax = b$ we row reduce $[A \mid b]$.
- To solve $AX = I_n$ we row reduce $[A \mid I]$.

Theorem

An $n \times n$ matrix A is invertible if and only if A is row equivalent to I_n , and in this case, any sequence of elementary row operations that reduces A to I_n will also transform I_n to A^{-1} .

So here is the algorithm:

- Place A and I side-by-side to form an augmented matrix $[A \mid I]$.

So here is the algorithm:

- Place A and I side-by-side to form an augmented matrix $[A \mid I]$. This is an $n \times 2n$ matrix (Big Augmented Matrix), instead of $n \times (n + 1)$, for the usual augmented matrix.

So here is the algorithm:

- Place A and I side-by-side to form an augmented matrix $[A \mid I]$.
- Then perform row operations on this matrix (which will produce identical operations on A and I).

So here is the algorithm:

- Place A and I side-by-side to form an augmented matrix $[A \mid I]$.
- Then perform row operations on this matrix (which will produce identical operations on A and I).
- So by Theorem 5:

$$[A \mid I] \text{ will row reduce to } [I \mid A^{-1}]$$

or A is not invertible.

Example

Find the inverse of $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$, if it exists.

Example

Find the inverse of $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$, if it exists.

Example

Find the inverse of $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$, if it exists.

Solution:

$$[A \ I] = \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \sim \dots \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right]$$

Example

Find the inverse of $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$, if it exists.

Solution:

$$[A \ I] = \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \sim \dots \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right]$$

So

$$A^{-1} =$$

Example

Find the inverse of $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$, if it exists.

Solution:

$$[A \ I] = \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right] \sim \dots \sim \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right]$$

So

$$A^{-1} = \begin{bmatrix} \frac{1}{2} & 0 & 0 \\ 0 & 0 & 1 \\ \frac{3}{2} & 1 & 0 \end{bmatrix}$$

Example (Let's do the previous example step by step.)

$$[A \mid I] =$$

Example (Let's do the previous example step by step.)

$$[A \mid I] = \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

Example (Let's do the previous example step by step.)

$$[A \mid I] = \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R1 \rightarrow \frac{1}{2}R1} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

Example (Let's do the previous example step by step.)

$$[A | I] = \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\begin{array}{c} \rightarrow \\ R1 \rightarrow \frac{1}{2}R1 \end{array} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\begin{array}{c} \rightarrow \\ R2 \rightarrow R2 + 3R1 \end{array} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

Example (Let's do the previous example step by step.)

$$[A | I] = \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R1 \rightarrow \frac{1}{2}R1} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R2 \rightarrow R2 + 3R1} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R2 \leftrightarrow R3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right]$$

Example (Let's do the previous example step by step.)

$$[A \mid I] = \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R1 \rightarrow \frac{1}{2}R1} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ -3 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R2 \rightarrow R2 + 3R1} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \end{array} \right]$$

$$\xrightarrow{R2 \leftrightarrow R3} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{1}{2} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & \frac{3}{2} & 1 & 0 \end{array} \right]$$

Check at home that $AA^{-1} = I_3$.

Remark

Why does this algorithm work?

Remark

Why does this algorithm work?

- At each step, we get

$$[A \mid I] \rightsquigarrow [E_1 A \mid E_1] \rightsquigarrow [E_2 E_1 A \mid E_2 E_1] \rightsquigarrow \dots$$

Remark

Why does this algorithm work?

- At each step, we get

$$[A \mid I] \rightsquigarrow [E_1 A \mid E_1] \rightsquigarrow [E_2 E_1 A \mid E_2 E_1] \rightsquigarrow \dots$$

- So each step is of the form

$$[FA \mid F], \quad F = E_r \dots E_3 E_2 E_1$$

Remark

Why does this algorithm work?

- At each step, we get

$$[A \mid I] \rightsquigarrow [E_1 A \mid E_1] \rightsquigarrow [E_2 E_1 A \mid E_2 E_1] \rightsquigarrow \dots$$

- So each step is of the form

$$[FA \mid F], \quad F = E_r \dots E_3 E_2 E_1$$

- If we succeed in row reducing A to I , the final step is

$$[FA \mid F] = [I \mid F]$$

Remark

Why does this algorithm work?

- At each step, we get

$$[A \mid I] \rightsquigarrow [E_1 A \mid E_1] \rightsquigarrow [E_2 E_1 A \mid E_2 E_1] \rightsquigarrow \dots$$

- So each step is of the form

$$[FA \mid F], \quad F = E_r \dots E_3 E_2 E_1$$

- If we succeed in row reducing A to I , the final step is

$$[FA \mid F] = [I \mid F]$$

- So $FA = I$, which means that

Remark

Why does this algorithm work?

- At each step, we get

$$[A \mid I] \rightsquigarrow [E_1 A \mid E_1] \rightsquigarrow [E_2 E_1 A \mid E_2 E_1] \rightsquigarrow \dots$$

- So each step is of the form

$$[FA \mid F], \quad F = E_r \dots E_3 E_2 E_1$$

- If we succeed in row reducing A to I , the final step is

$$[FA \mid F] = [I \mid F]$$

- So $FA = I$, which means that

Remark

Why does this algorithm work?

- At each step, we get

$$[A \mid I] \rightsquigarrow [E_1 A \mid E_1] \rightsquigarrow [E_2 E_1 A \mid E_2 E_1] \rightsquigarrow \dots$$

- So each step is of the form

$$[FA \mid F], \quad F = E_r \dots E_3 E_2 E_1$$

- If we succeed in row reducing A to I , the final step is

$$[FA \mid F] = [I \mid F]$$

- So $FA = I$, which means that $A^{-1} = F$.

Practice Problems. Find the inverse of A :

- $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$

Practice Problems. Find the inverse of A :

- $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{bmatrix}.$ Hint: What is $A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$?

Practice Problems. Find the inverse of A :

- $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{bmatrix}.$ Hint: What is $A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$?

- $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$

Practice Problems. Find the inverse of A :

- $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{bmatrix}.$ Hint: What is $A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$?

- $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 2 & 3 & 0 \\ 4 & 5 & 6 & 0 \\ 6 & 7 & 8 & 0 \\ 9 & 0 & 1 & 0 \end{bmatrix}.$

Practice Problems. Find the inverse of A :

- $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{bmatrix}.$ Hint: What is $A \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$?

- $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 2 & 3 & 0 \\ 4 & 5 & 6 & 0 \\ 6 & 7 & 8 & 0 \\ 9 & 0 & 1 & 0 \end{bmatrix}.$

- $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}.$